

## Temperature-Dependent Growth Kinetics of *Escherichia coli* ML 30 in Glucose-Limited Continuous Culture

K. KOVÁŘOVÁ, A. J. B. ZEHNDER, AND T. EGLI\*

Swiss Federal Institute for Environmental Science and Technology, Swiss Federal Institute of Technology,  
CH-8600 Dübendorf, Switzerland

Received 14 December 1995/Accepted 9 May 1996

Detailed comparison of growth kinetics at temperatures below and above the optimal temperature was carried out with *Escherichia coli* ML 30 (DSM 1329) in continuous culture. The culture was grown with glucose as the sole limiting source of carbon and energy (100 mg liter<sup>-1</sup> in feed medium), and the resulting steady-state concentrations of glucose were measured as a function of the dilution rate at 17.4, 28.4, 37, and 40°C. The experimental data could not be described by the conventional Monod equation over the entire temperature range, but an extended form of the Monod model [ $\mu = \mu_{\max} \cdot (s - s_{\min}) / (K_s + s - s_{\min})$ ], which predicts a finite substrate concentration at 0 growth rate ( $s_{\min}$ ), provided a good fit. The two parameters  $\mu_{\max}$  and  $s_{\min}$  were temperature dependent, whereas, surprisingly, fitting the model to the experimental data yielded virtually identical  $K_s$  values (approximately 33  $\mu\text{g liter}^{-1}$ ) at all temperatures. A model that describes steady-state glucose concentrations as a function of temperature at constant growth rates is presented. In similar experiments with mixtures of glucose and galactose (1:1 mixture), the two sugars were utilized simultaneously at all temperatures examined, and their steady-state concentrations were reduced compared with to growth with either glucose or galactose alone. The results of laboratory-scale kinetic experiments are discussed with respect to the concentrations observed in natural environments.

Knowledge concerning the influence of environmental factors such as temperature, pH, salinity, etc., on microbial growth is of crucial practical importance in the control of bioprocesses, for the safe handling of food (1, 2, 12, 50, 51), in wastewater treatment (7), and in bioremediation (2). In addition, in taxonomy, cardinal temperatures for growth are key characteristics of microbial strains (37).

In recent years, several models for predicting the growth rate of microorganisms as a function of either temperature alone (11, 13, 22, 31, 32, 52, 53) or of temperature in combination with other factors have been proposed (1, 5, 20–22, 36, 50). Surprisingly, few attempts at a better basic understanding have been made to relate the rate of growth and actual substrate concentration. This relationship is traditionally termed growth kinetics (23, 30). (However, note that the same expression has also been used for the description of the time courses of population densities [5].) The current lack of systematic data on the influence of temperature on the kinetics of growth makes the prediction of this effect difficult. Temperature modulation of growth kinetics is to be expected, because both metabolism and cellular composition are affected by cultivation temperature, as was demonstrated by the cellular fatty acid composition (16, 26, 41), the synthesis or degradation of certain proteins (14, 15, 17, 20, 25), changes in protein activity (15, 35), changes in maintenance requirements of cells (19, 27, 39, 42, 47), changes in end products of metabolism (17), and increases in pigment formation (27).

Bacterial metabolism represents a network of reactions. Although these individual biochemical reactions are temperature dependent, the fundamental question of whether the parameters used in growth kinetic models are temperature dependent must be asked. To determine this, a detailed comparison of growth kinetics at temperatures below and above the optimal

temperature was carried out for *Escherichia coli* ML 30 cultivated in continuous culture with glucose and/or galactose. Such investigations are possible only by using an extremely sensitive method for measuring low concentrations of sugars (in micrograms per liter) in culture media (40). The objective of the present study was to compare the experimentally established relationships between growth rate and steady-state substrate concentrations at different constant temperatures and to find out whether the whole set of relationships can be described by a simple mathematical model (Table 1 summarizes the nomenclature used throughout). Additionally, the effect of temperature on steady-state substrate concentrations at constant growth rates (dilution rates in continuous culture) was studied.

**Compendium of the models proposed in the literature. (i) Conventional growth kinetics and models containing an  $s_{\min}$  term.** Various mathematical models have been proposed to quantitatively describe microbial growth kinetics. The Monod model (equation 1) is considered the basic equation (23), which has since been improved by including expressions for, e.g., maintenance, diffusion, or transport limitation (29, 30, 45; for detailed comparison, see reference 40). Microbial growth kinetics in both batch and continuous culture have been investigated. Earlier experiments carried out in batch cultures mostly relied on indirect methods, i.e., growth was measured, whereas the substrate concentrations were not directly determined but were estimated by calculation. In contrast, when growth kinetics in continuous culture were investigated, the actual steady-state concentrations of the growth-limiting substrate were determined as a function of the dilution rate. For such an experimental setup, the  $s = f(D)$  form of the kinetic model (equation 1) correctly expresses the variable dependence, and not the  $\mu = f(s)$  form, in which the models were originally reported (discussed in reference 40).

$$s = K_s \frac{D}{\mu_{\max} - D} \quad (1)$$

Monod's original kinetic equation (equation 1) implies a sub-

\* Corresponding author. Phone: 41-1-823 5158. Fax: 41-1-823 5547. Electronic mail address: egli@eawag.ch.

TABLE 1. Nomenclature

Symbol	Definition	Unit
$a$	Specific maintenance rate (equations 3 and 5)	$\text{h}^{-1}$
$A$	Parameter in Esener model (equation 9)	$\text{h}^{-1}$
$b$	Parameter in Ratkowsky model (equation 10)	$\text{K}^{-1} \text{s}^{-0.5}$
$B$	Model parameter (equation 11)	$\text{K}^2 \mu\text{g liter}^{-1}$
$c$	Parameter in Ratkowsky model (equation 10)	$\text{K}^{-1}$
$C$	Model parameter (equation 11)	$\text{K}^{-1}$
$D$	Dilution rate (specific growth rate in chemostat)	$\text{h}^{-1}$
$k_i, k_j,$ and $k_h$	Rate constants	$\mu\text{g liter}^{-1} \text{h}^{-1}$ or $\mu\text{g}^2 \text{liter}^{-2} \text{h}^{-1}$
$K$	Parameter in Esener model (equation 9)	
$K_m$	Michaelis-Menten substrate saturation constant	$\mu\text{g liter}^{-1}$
$K_s$	Substrate saturation constant	$\mu\text{g liter}^{-1}$
$n$	Number of steady states analyzed (here, number of experiments, not number of datum points collected from a particular steady state)	
$(n - p)$	Number of degrees of freedom	
$R$	Gas constant	$\text{kJ kg}^{-1} \text{mol}^{-1}$
$RR$	Relative residuals (equation 14)	%
$RSS (s)$	Residual sum of squares with respect to $s$ (equation 13)	$\mu\text{g}^2 \text{liter}^{-2}$
$s$	Steady-state substrate concn	$\mu\text{g liter}^{-1}$
$s_{\min}^*, s_{\min}^{**}, s_{\min}^{***}$ , and $s_{\min}^{****}$	Predicted substrate concentration at $D = 0 \text{ h}^{-1}$ for different growth models	$\mu\text{g liter}^{-1}$
$s_{\text{Obs}}, \mu_{\text{Obs}}$	Experimentally established value (for $s$ or $\mu$ )	$\mu\text{g liter}^{-1}$ or $\text{h}^{-1}$
$s_{\text{pred}}$	Value predicted by model equation ( $s$ or $\mu$ )	$\mu\text{g liter}^{-1}$ or $\text{h}^{-1}$
$T$	Cultivation temperature	$\text{K}$ or $^{\circ}\text{C}$
$T_{\min}, T_{\text{opt}}$ , and $T_{\max}$	Minimum, optimum, and maximum temperatures, respectively	$\text{K}$ or $^{\circ}\text{C}$
$x$	Parameter in Westerhoff model (equation 7)	$\text{h}^{-1}$
$y$	Parameter in Westerhoff model (equation 7)	$\text{h}^{-1}$
$\Delta H_1, \Delta H_2$	Enthalpy changes (in Esener model [equation 9])	$\text{kJ mol}^{-1}$
$\mu$	Specific growth rate	$\text{h}^{-1}$
$\mu_{\max}$	Maximum specific growth rate	$\text{h}^{-1}$

strate concentration of 0 at a growth rate of  $0 \text{ h}^{-1}$  (23). This model represents a special case of a more general kinetic expression (equation 2; Fig. 1), in which a term for a finite substrate concentration,  $s = s_{\min}$  at  $D = 0 \text{ h}^{-1}$ , is incorporated. When  $s_{\min} \ll s$ ,  $s_{\min}$  becomes negligible, and equation 2 reduces to equation 1:

$$s = K_s \frac{D}{\mu_{\max} - D} + s_{\min} \quad (2)$$

Expressions having a meaning similar to that of  $s_{\min}$  (and noted




FIG. 1. Example of Monod model with apparent substrate term (equation 2) indicating the model parameters.

here as  $s_{\min}^*, s_{\min}^{**}$ , and  $s_{\min}^{***}$ ) are implicitly also present in other kinetic models (equations 3, 5, and 7). For instance, the original relationship proposed in 1967 by Powell (30) can be easily converted into an  $s = f(D)$  form (equation 3). The first part of this expression is identical to the Monod model (equation 1), and to this a dilution rate-dependent  $s_{\min}^*$  term (equation 4) is added:

$$s = K_s \frac{D}{\mu_{\max} - D} + K_s \frac{a}{\mu_{\max} - D} \quad (3)$$

$$s_{\min}^* = K_s \frac{a}{\mu_{\max} - D},$$

$$\text{hence at } D = 0 \text{ h}^{-1}, s_{\min}^* = K_s \frac{a}{\mu_{\max}} \quad (4)$$

In contrast to equation 2,  $s_{\min}^*$  in equation 4 is dependent on  $D$ . In this model, the contribution of  $s_{\min}^*$  to the steady-state substrate concentrations at high dilution rates is usually negligible, and  $s_{\min}^*$  becomes important at growth rates near 0. Essentially, it can be also assumed that the  $s_{\min}^*$  term is not dependent on the dilution rate (equation 2).

In another analysis of microbial growth, van Uden (45) in 1967 followed up Pirt's original 1965 proposal (28, 29) that only the overall growth rate is reduced by the maintenance rate  $a$  (in its  $s$  form [equation 5]). The resulting substrate concentration at  $D = 0 \text{ h}^{-1}$  is shown in equation 6:

$$s = \frac{K_s (a + D)}{(\mu_{\max} - D - a)} \quad (5)$$

$$s_{\min}^{**} = \frac{a}{\mu_{\max} - a} K_s \quad (6)$$

While all of the previous relationships are extensions of the Monod-type kinetics, a fundamentally different model, based on nonequilibrium thermodynamics, was proposed by Westerhoff and coworkers in 1982 (48). This logarithmic expression [exponential in  $s = f(D)$  form; equation 7] also predicts at  $D = 0 \text{ h}^{-1}$  a positive substrate concentration (equation 8):

$$s = \exp\left(\frac{D - x}{y}\right) \quad (7)$$

$$s_{\min}^{***} = \exp\left(-\frac{x}{y}\right) \quad (8)$$

**(ii) Steady-state substrate concentrations versus temperature at various growth rates.** Because the effect of temperature on steady-state substrate concentrations has not yet been studied systematically, there is no mathematical description of this relationship. Assuming a link between  $\mu$  and  $s$ , we here suggest using a modification of a model which was developed previously to describe the temperature dependence of the maximum specific growth rate. This suggestion follows the approach taken by McMeekin et al. (22), who proposed that the simple Ratkowsky model [i.e.,  $\mu = f(T)$  (equation 10)] might be extended for any set of restricting conditions involving temperature as one of the factors.

Several unsegregated and unstructured phenomenological models (11, 12, 31, 32, 37) have been proposed both to predict the cardinal temperatures and to describe bacterial growth in batch cultures within the growth-permissible temperature range. It has been shown (see, e.g., reference 31) that within this temperature range  $\mu_{\max}$  follows a bell-shaped curve, which has a maximum at the optimum growth temperature ( $T_{\text{opt}}$ ). In the master reaction model proposed by Esener (11) (equation 9),  $\mu_{\max}$  asymptotically approaches the abscissa at both the maximum ( $T_{\max}$ ) and minimum ( $T_{\min}$ ) growth temperatures:

$$\mu_{\max} = \frac{A \exp(-\Delta H_1/RT)}{1 + K \exp(-\Delta H_2/RT)} \quad (9)$$

In contrast, the square root model proposed by Ratkowsky et al. (31) (equation 10) is defined for  $T_{\min} < T < T_{\max}$ ; out of this range  $\mu_{\max}$  is 0:

$$\sqrt{\mu_{\max}} = b(T - T_{\min})\{1 - \exp[c(T - T_{\max})]\} \quad (10)$$

In continuous culture,  $T_{\min}$  and  $T_{\max}$  can be determined for each particular dilution rate, e.g., by washout experiments or from the  $T = f(\mu_{\max})$  relationship. In this contribution, we were primarily interested in how well the minimum and maximum temperatures for growth at particular dilution rates can be predicted, not in the model structure.

For the  $s = f(T)$  model, we assumed that steady-state substrate concentrations are reciprocally proportional to the square of the temperature, i.e., the square root of the concentration is reciprocally proportional to the temperature. This assumption can be justified in the following way. Under optimum conditions, a microbial cell grows with the highest possible  $\mu$  and, therefore, a cell's overall metabolic efficacy should also be at its optimum. This implies that close to  $T_{\text{opt}}$ , the lowest steady-state substrate concentrations should be expected at a particular growth rate, in the same way as the highest  $\mu_{\max}$  is reached at  $T_{\text{opt}}$ . This assumption provides a link between the square root model (equation 10) and the steady-state substrate concentration model (equation 11). For these

reasons, we propose to describe the  $s$  versus  $T$  relationship (equation 11) by a reciprocal form of the maximum specific growth rate versus temperature relationship (equation 10):

$$s = \frac{B}{(T - T_{\min})^2 \{1 - \exp[C(T - T_{\max})]\}^2} \quad (11)$$

where the units of  $C$  are identical to those of  $c$  from the Ratkowsky model (equation 10) and the parameter  $B$  is related to  $b$  from the Ratkowsky model according to equation 12:

$$B \cong \frac{1}{b^2} \quad (12)$$

Since the two regression coefficients ( $B$  and  $C$ ) vary with the dilution rate, it is possible to simply extend equation 11 by introducing terms to describe variations in  $B$  (or  $b$ ) and  $C$  with the growth rate in the chemostat, as determined from experimental data (e.g., see Fig. 7). Additionally, it should be pointed out that in theory,  $T_{\min}$  and  $T_{\max}$  used in equation 11 are equivocally defined because they can be understood as either constant properties of a given strain and medium composition (e.g., its cardinal temperatures) or as variables changing with the dilution rate.

## MATERIALS AND METHODS

**Organism, medium, and culture conditions.** *E. coli* ML 30 (DSM 1329) was grown in mineral medium (40) supplemented either with glucose (100 mg liter<sup>-1</sup> in chemostat; 500 mg liter<sup>-1</sup> in batch culture) or with a mixture of glucose and galactose (each 50 mg liter<sup>-1</sup> in chemostat) as the only sources of carbon and energy. The bioreactor (MBR, Wetzikon, Switzerland), provided with both pH (7.00 ± 0.05) and temperature (±0.1°C) control, was operated in chemostatic mode with a working volume of 1.5 liters. The impeller speed control was set at 1,000 min<sup>-1</sup>, and the oxygen saturation was >90% air saturation. The bioreactors were regularly checked for wall growth to avoid artifacts as reported by Pirt (29).

**Maximum specific growth rates ( $\mu_{\max}$ ).**  $\mu_{\max}$  rates were determined in batch cultures at different temperatures. The cells used as an inoculum were pregrown exponentially for more than 50 generations at the particular temperature. At least duplicate measurements were made at each temperature. The standard deviation obtained for  $\mu_{\max}$  was reproducibly ± 0.05 h<sup>-1</sup>.

**Steady-state substrate concentrations.** Steady-state concentrations of glucose and/or galactose were determined in continuous culture during independent chemostat runs, either as a function of the dilution rate at constant temperatures (17.4, 28.4, 37, and 40°C) or as a function of temperature at constant dilution rates (0.2, 0.3, 0.4, and 0.5 h<sup>-1</sup>). At each temperature, the continuous cultivation was restricted either by the critical dilution rate below which the organism was able to grow without washing out or by wall growth. The entire temperature range within which the organism was able to grow in the chemostat at a particular  $D$  (i.e.,  $\mu_{\max} \geq D$ ) was extrapolated from batch data (for an explanation, see Fig. 2).

Individual steady-state concentrations of glucose or galactose represent the means of approximately 10 measurements determined over a time period of more than 40 generations after the culture had reached steady state with respect to glucose (galactose) concentration. The standard deviations of these values were between ±5% to ±10%, and for clarity they are not given in all figures presented. The sugar analysis has been described in detail elsewhere (40).

**Data processing.** The models were fitted to the experimental data by nonlinear regression (33, 34). The minimum of the least-squares criterion (residual sum of squares [RSS]) (equation 13) was computed with a Simplex algorithm:

$$RSS(s) = \sum (s_{\text{obs}} - s_{\text{pred}})^2 \quad (13)$$

Initial estimates of the model parameters were required, because the structural correlation between parameters made their estimation otherwise difficult. Starting values were chosen from the best extrapolations of the experimental data by exponential and/or polynomial functions.

The quality of the fits was evaluated by standard tools, such as comparing the variance, applying the  $\chi^2$  criterion, or, for the discrimination of competing models, the F test (4), or by analyzing the linear regression between the measured and predicted substrate concentrations. Statistical validation was best visible in plots of relative residuals (equation 14, e.g., Fig. 3):

$$RR = \frac{s_{\text{obs}} - s_{\text{pred}}}{s_{\text{obs}}} \times 100 \quad (14)$$




FIG. 2. Experimentally determined and predicted steady-state glucose concentrations (with the extended Monod model [equation 2]) for growth of *E. coli* ML 30 in glucose-limited chemostat cultures at 17.4, 28.4, 37, and 40°C, as a function of dilution (growth) rate. (A) Bars, steady-state substrate concentrations (height indicates standard deviation [approximately 10%] of the steady-state glucose concentrations estimated as an average of approximately 10 measurements; the horizontal extensions of the bars give the approximate variations in  $D$ ); lines, predictions of steady-state glucose concentrations by different models. (B and C) open circles, data from Senn et al. (40); closed circles and triangles, own steady-state glucose measurements; lines, best fit of equation 2.

## RESULTS

**Temperature dependence of growth kinetics.** The kinetics of growth of *E. coli* ML 30 [i.e., the  $s = f(D)$  relationship] was investigated at four different temperatures (17.4, 28.4, 37, and 40°C, respectively) both above and below the optimum growth temperature (Fig. 2).

For growth at 40°C, the steady-state glucose concentrations were measured as a function of  $D$  during three independent glucose-limited chemostat runs. This cultivation temperature was slightly higher than the calculated optimum growth temperature ( $T_{opt} = 38.7^\circ\text{C}$ ) of *E. coli* ML 30 growing unrestricted in this medium in batch culture (Table 2). The glucose concentrations measured at low and moderate dilution rates (up to approximately  $0.35\text{ h}^{-1}$ ) were distinctly enhanced compared with the previously reported growth kinetic data at 37°C (40) (Fig. 2). Nevertheless, as judged by Student's  $t$  test, the two  $\mu_{max}$  values estimated by fitting equation 2 to the data at 37°C

( $0.76 \pm 0.01$ ) and 40°C ( $0.74 \pm 0.01$ ) were not significantly different (Fig. 2).

In contrast to the predictions made by the Monod model (equation 1), these results clearly demonstrate that the steady-state glucose concentrations of the limiting-substrate glucose did not approach 0. Therefore, the quality of the fit of various mathematical models (equations 2, 5, and 7), which contain implicitly a finite substrate concentration at  $D = 0\text{ h}^{-1}$ , and of the original Monod model (equation 1) was compared with experimentally obtained data (Fig. 2). The quality of the fit of these models (expressed as means of relative residuals [equation 14]) was compared with the theoretical measurement error of  $\pm 10\%$  (Fig. 3). It is obvious that three of the models (equations 2, 5, and 7) exhibited a systematic deviation from the glucose concentrations measured at low dilution rates. The model proposed by Westerhoff (equation 7) could not be successfully fitted to this type of experimental data by the applied minimizing procedure. Only the model, which contains an apparent substrate term (equation 2), exhibited a random distribution of the relative residuals. The mean estimation error for this model (6.7%) was in the same order of magnitude as the measurement errors (10%) and was considerably lower than the mean estimation errors obtained for the other three models (26.8, 20.9, and 23.2% for equations 2, 5, and 7, respectively).

The data at 37, 28.4, and 17.4°C were used to test the validity of equation 2. In Fig. 2, the fits to data from different temperatures are given and the measured steady-state glucose concentrations are compared with predicted curves. It should be pointed out that at 37°C, only data up to  $D = 0.71\text{ h}^{-1}$  were used because data at higher dilution rates can be easily affected by experimental artifacts (40). The excellent correlation between the measured steady-state glucose concentrations and predicted values at all experimental temperatures (Fig. 4) confirms the utility of the model containing  $s_{min}$ . The model parameters computed for equation 2 from experimental data measured at four different temperatures are collected in Table 3. Interestingly, the  $K_s$  values did not vary with temperature, whereas the two parameters  $\mu_{max}$  and  $s_{min}$  were temperature

TABLE 2. Comparison of the Ratkowsky and Esener models describing dependence of maximum specific growth rate of *E. coli* on temperature with data experimentally obtained in batch cultures<sup>a</sup>

Variable	U	Ratkowsky model value	Esener model value
$A$	$\text{h}^{-1}$		$(4.22 \pm 0.35) 10^{11}$
$K$	$\text{h}^{-1}$		$(8.07 \pm 0.11) 10^{49}$
$\Delta H_1$	$\text{kJ mol}^{-1}$		$68.9 \pm 6.2$
$\Delta H_2$	$\text{kJ mol}^{-1}$		$300.6 \pm 25.4$
$b$	$\text{K}^{-1} \text{h}^{-0.5}$	$(2.76 \pm 0.25) 10^{-2}$	
$c$	$\text{K}^{-1}$	$1.12 \pm 0.10$	
$T_{min}$	$\text{K}$	$275.7 \pm 0.3$	
$T_{max}$	$\text{K}$	$315.2 \pm 0.2$	
$T_{opt}$	$\text{K}$	$311.9 \pm 0.2^*$	$311.4 \pm 0.2^*$
$\mu_{max}$	$\text{h}^{-1}$	$0.95 \pm 0.05^*$	$0.92 \pm 0.05^*$
$RSS/(n - p)$	$\text{h}^{-2}$	$2.5 \cdot 10^{-4}^{**}$	$4.7 \cdot 10^{-2}^{**}$

<sup>a</sup> Values of the model parameters (equations 9 and 10), optimum growth temperatures, and statistical parameters were computed for the batch culture data. \*, value calculated from the best-fit parameters.




FIG. 3. Relative residuals for the fits of four different kinetic models to experimentally determined steady-state glucose concentrations during growth of *E. coli* at 40°C. Hatched areas, a systematic underestimation of steady-state glucose concentrations; dashed lines, 10% deviation from experimental steady-state glucose concentrations.

dependent.  $K_s$  remained constant at approximately  $33 \mu\text{g liter}^{-1}$ , which is approximately 1/3 of the value obtained when the original Monod model (equation 1) was fitted to the data (40). The  $\mu_{\max}$  values estimated in chemostat culture were generally some 15% lower than those measured under substrate excess conditions in batch culture (Fig. 5a); this phenomenon has also been observed in another study (40). Interestingly, the increase in  $\mu_{\max}$  values at temperatures lower than the  $T_{\text{opt}}$  is more pronounced than their decrease at the superoptimal range. An inverse pattern was observed for  $s_{\min}$ . The

computed values (Table 3) indicate a slight increase in  $s_{\min}$  with decreasing temperatures and a steep increase at temperatures higher than the  $T_{\text{opt}}$ .

**Modelling substrate concentration as a function of temperature.** Glucose steady-state concentration as a function of cultivation temperature was measured for growth of *E. coli* at a constant dilution rate of  $0.3 \text{ h}^{-1}$  (Fig. 6). The observed parabola-like relationship can be described by equation 11, which is a modification of the Ratkowsky model (equation 10) commonly used to predict the temperature dependency of the maximum specific growth rate. Unfortunately, the original model contains four parameters,  $T_{\min}$ ,  $T_{\max}$ ,  $c$ , and  $b$ . However, the number of parameters in equation 11 could be reduced to two, assuming that  $T_{\min}$  and  $T_{\max}$  are either the cardinal temperatures for growth (predicted by equation 10) or that  $T_{\min}$  and  $T_{\max}$  represent the temperature boundaries within which growth is possible at a particular dilution rate. Therefore, the maximum specific growth rate was determined in batch culture at different temperatures, and the bell-shaped curve in Fig. 5 indicates the temperature boundaries within which growth is possible at a particular dilution rate in continuous culture. By using these temperature boundaries, the experimental data at  $D = 0.3 \text{ h}^{-1}$  could be described by an almost symmetrical




FIG. 4. Correlation between measured and predicted steady-state glucose concentrations by the extended Monod model (equation 2). Line, the calculated regression line for all pairs of  $s_{\text{obs}}$  and  $s_{\text{pred}}$  values for the temperatures of 17.4, 28.4, 37, and 40°C.

TABLE 3. Values of the best-fit parameters of the apparent substrate model<sup>a</sup>

Temp (°C)	$K_s$ ( $\mu\text{g liter}^{-1}$ )	$\mu_{\max}$ ( $\text{h}^{-1}$ )	$s_{\min}$ ( $\mu\text{g liter}^{-1}$ )	$\frac{RSS}{(n-p)}$	$n$
17.4	$33.3 \pm 4.2$	$0.185 \pm 0.02$	$22 \pm 2$	7,503.2	4
28.4	$33.3 \pm 3.3$	$0.54 \pm 0.01$	$18 \pm 2$	302.6	11
37	$32.8 \pm 3.2$	$0.76 \pm 0.01$	$12 \pm 2$	8,122.8	23
40	$33.6 \pm 1.5$	$0.74 \pm 0.01$	$64 \pm 8$	161.8	15

<sup>a</sup> Values were obtained from fitting equation 2 to steady-state glucose concentrations experimentally determined during growth of *E. coli* at the temperatures indicated as a function of the dilution rate.




FIG. 5. Temperature dependency of specific growth rate of *E. coli* ML 30 grown in mineral medium with glucose. (a)  $\mu_{\max}$  as a function of temperature. Closed circles, experimental data from batch cultures; open circles, maximum specific growth rates calculated from chemostat data; curve 1, predictions by the Esener model (equation 9); curve 2, predictions by the Ratkowsky model (equation 10); dashed line, predictions of growth rates in continuous culture by the Ratkowsky model; arrow, the temperature boundary within which growth is possible at  $D = 0.3 \text{ h}^{-1}$ . (b) Relative residuals ( $RR$  [equation 14]) for the two fits of batch culture data. Hatched areas, areas in which the model predictions of the Ratkowsky (right area) and the Esener (right and left areas) equations did not hold; open rectangles, relative residuals of the Ratkowsky model; closed rectangles, relative residuals of the Esener model; dashed line, isoline of  $0.05 \text{ h}^{-1}$  standard deviation of experimental measurements.




FIG. 6. Temperature dependency of steady-state substrate concentrations at a dilution rate of  $0.3 \text{ h}^{-1}$ . Closed circles (glucose) and closed triangles (galactose), experimentally determined steady-state sugar concentrations at  $100 \text{ mg liter}^{-1}$  of glucose (or galactose) was supplied in the inflowing medium as the only substrate; open circles, steady-state glucose concentrations predicted by equation 2 for different temperatures; open (closed) squares, steady-state glucose (galactose) concentrations when 1:1 mixture of glucose and galactose was supplied in the inflowing medium; solid line, model equation 11 fitted with  $T_{\min}$  and  $T_{\max}$  values, which represent experimental boundaries at  $D = 0.3 \text{ h}^{-1}$ ; dashed line, model equation 11 fitted with the cardinal temperature values of *E. coli* ML 30 instead of  $T_{\min}$  and  $T_{\max}$  parameters.

parabola-like curve (Fig. 6). In contrast, using the cardinal temperatures from Table 2 did not lead to a good prediction of the substrate concentrations at lower temperatures (Fig. 6).

As mentioned above, the temperature range at a particular growth rate within which growth in the chemostat is possible (the experimentally accessible range lies below the bell-shaped curve in Fig. 5a) had to be determined in order to obtain  $T_{\min}$  and  $T_{\max}$  for equation 11. The maximum growth rates measured in batch cultures at various temperatures (Fig. 5) were compared with the predictions of the Ratkowsky and the Esener model equations (equations 9 and 10). Neither model described the data well at very low growth rates. Additionally, the Esener model slightly overestimated the maximum growth temperature. Despite these inaccuracies in the extreme low and high temperature ranges, both models allowed a good extrapolation (mean estimation error, less than  $\pm 5\%$ ) of the experimental data (Fig. 5b). In comparison to the  $\mu_{\max}$  values measured under unrestricted growth conditions in batch cultures, those determined from the kinetic investigation made in glucose-limited chemostat cultures were consistently lower (Fig. 5a). However, the  $\mu_{\max}$  values estimated from continuous culture data were also well described by the Ratkowsky model. In this estimation procedure, the two parameters  $b$  and  $c$  were optimized, whereas  $T_{\min}$  and  $T_{\max}$  values were set to those obtained from batch culture data (Table 2). From this curve, the minimum and maximum growth temperatures at particular dilution rates were estimated.

When fitting equation 11 [ $s = f(T)$ ], the two regression coefficients ( $B$  and  $C$ ) were optimized with respect to the

experimental data obtained at dilution rates of 0.2, 0.3, and  $0.4 \text{ h}^{-1}$  and with respect to predicted  $s_{\min}$  values. Both  $T_{\min}$  and  $T_{\max}$  remained fixed during this fitting procedure. From the  $B$  versus  $D$  and  $C$  versus  $D$  plots (Fig. 7), it can be seen that for the range of  $0 \text{ h}^{-1} \leq D \leq 0.4 \text{ h}^{-1}$  the dependency of the two parameters on  $D$  is approximately linear. Hence, steady-state substrate concentrations as a function of temperature and growth rate can be described by replacing these parameters in equation 11 with linear relationships. The resulting model is presented in Fig. 8b. The width of the parabola-like curve is




FIG. 7. Relationships between the two regression coefficients  $B$  and  $C$  and the dilution rate.  $B$  and  $C$  are the best-fit model parameters of the steady-state substrate concentration model (equation 11).

narrower with increasing dilution rates, i.e., the temperature range for growth becomes restricted with increasing dilution rates. Unfortunately, the minimum of this function changed at different growth rates, and this minimum is not always equivalent to the optimum temperature for growth.

**Extrapolation of the simple relationships into more complex systems.** The fact that relationships exist between model parameters and temperature (Table 3; Fig. 7) is of advantage for developing the complex  $s = f(D, T)$  model. In addition, a consistent kinetic description, i.e., using only equation 2 for all temperatures examined, is an advantageous basis for developing additional models, since it is the only equation which accurately described the experimental data at 40°C and, at the same time, the data at lower temperatures. An equation with only two variables ( $T$  and  $D$ ) can be proposed to describe the relationship between steady-state substrate concentration, growth rate/dilution rate, and temperature. However, two different experimental approaches can be used to establish this model. First,  $D$  can be varied and  $s$  can be measured at constant  $T$ , and second,  $T$  can be varied and  $s$  can be determined at constant  $D$ . A three-dimensional projection of the  $s = f(D, T)$  relationship was computed by fitting the best surface to the experimental data measured by the first approach (Fig. 8a). The same relationship, but projected into two dimensions, is shown in Fig. 8b, representing data collected by the second experimental approach.

All of the previous results were obtained with a simple model system with glucose as the only carbon and energy substrate for growth. However, it has been shown recently that *E. coli* cells are able to utilize mixtures of sugars simultaneously when cultivated in carbon-limited continuous culture (10). Therefore, steady-state sugar concentrations were measured as a function of temperature in cultures growing with a 1:1 mixture of glucose and galactose. In Fig. 6, it is shown that at a constant growth rate the steady-state concentrations of glucose were reduced when an additional substrate (e.g., galactose) was utilized simultaneously compared with those during growth with glucose only. Except for the lowest temperature tested, steady-state glucose concentrations were reduced to approximately 50%. For galactose, a similar effect was observed, as judged from the results obtained at  $T = 37^\circ\text{C}$ , where the concentration was reduced from  $41 \mu\text{g liter}^{-1}$  (10) to  $19 \mu\text{g liter}^{-1}$  at  $D = 0.3 \text{ h}^{-1}$ . The data obtained at  $28.4^\circ\text{C}$  indicate that the reduction of individual steady-state sugar concentrations during mixed substrate growth might be even more pronounced at lower temperatures. Although this observation has to be confirmed, it might open a promising road to optimizing biodegradation processes in which pollutants have to be removed to low concentration levels.

## DISCUSSION

**What is the physiological meaning of an  $s_{\min}$  term?** The conventional Monod equation (equation 1) did not hold for the description of the experimental data at all temperatures. This discrepancy was solved by using an extended form of the Monod model (equation 2), which predicts a finite substrate concentration at 0 growth rate. It must be pointed out that  $s_{\min}$  is usually negligible compared with actual steady-state substrate concentrations (i.e.,  $s \gg s_{\min}$  and, therefore,  $s + s_{\min} \cong s$ ), and no statistically significant difference between the Monod model (equation 1) and the model including an apparent substrate term (equation 2) can be observed, for example, at  $37^\circ\text{C}$ . At  $T_{\text{opt}}$  (and perhaps also at lower temperatures), both models predict the experimental data well and, therefore, the original Monod model can be used when a less complex model




FIG. 8. Steady-state glucose concentrations as a function of temperature and dilution rate for the growth of *E. coli* in glucose-limited continuous culture. (a) Three-dimensional relationship describing  $s = f(D, T)$ . Thin-lined three-dimensional surface, the best fit to the experimental data; heavy lines, the particular growth kinetics and the  $s = f(T)$  relationships at 0 and  $0.3 \text{ h}^{-1}$  used in model building. (b) The  $s = f(D, T)$  relationship projected into the  $s$  versus  $T$  plane. Lines, isolines of the same dilution rate. Predictions of residual glucose concentrations as function of temperature at dilution rates of 0, 0.2, 0.3, and  $0.4 \text{ h}^{-1}$  were made by equation 11.

is preferred. On the other hand, the data at  $40^\circ\text{C}$  cannot be predicted well by any of the alternative models, and, therefore, equation 2, in which the Monod relationship was extended with  $s_{\min}$ , must be preferred for describing the  $s = f(D)$  relationship at all temperatures.

The existence of  $s_{\min}$  can be justified on the basis of the maintenance energy concept, and the existence of a cellular maintenance energy requirement can be easily explained by thermodynamic reasoning. A consequence of this concept is the existence of a finite concentration or flux of an energy (carbon) substrate at 0 growth rate. In a system open with respect to the supply of substrate, this results in a finite con-

centration of the energy (carbon) source at  $D = 0 \text{ h}^{-1}$ . For a culture of *E. coli* cells grown with glucose at  $30^\circ\text{C}$ , Schulze and Lipe (39) measured a small rate of substrate consumption, even when no growth was observed. This rate,  $55 \text{ mg}$  of glucose  $(\text{h} \cdot \text{g} [\text{dry weight}])^{-1}$ , which represents a specific maintenance rate of  $0.0286 \text{ h}^{-1}$  when the yield coefficient of  $0.52$  experimentally determined by these authors is used, was just enough to sustain cellular metabolism, but not enough to allow growth and reproduction. Similarly, Shehata and Marr (42) estimated that  $18 \mu\text{g liter}^{-1}$  (which, for the reported  $K_s$  of  $68 \mu\text{g liter}^{-1}$  and  $\mu_{\text{max}} = 0.78 \text{ h}^{-1}$ , represents a specific maintenance rate of  $0.163 \text{ h}^{-1}$ ) was the lowest substrate concentration that allowed maintenance of growth of *E. coli* in batch culture at  $30^\circ\text{C}$ . Wallace and Holms (47) and Mainzer and Hempfling (19) have also found the maintenance requirements of *E. coli* strains affected by temperature.

To compare the  $s_{\text{min}}$  values estimated in this study at different temperatures with previously reported maintenance requirements (19, 39, 42, 47),  $s_{\text{min}}$  values were converted via equation 6 into specific maintenance rates. The resulting specific maintenance rates of  $0.074 \text{ h}^{-1}$  ( $17.4^\circ\text{C}$ ),  $0.191 \text{ h}^{-1}$  ( $28.4^\circ\text{C}$ ),  $0.204 \text{ h}^{-1}$  ( $37^\circ\text{C}$ ), and  $0.485 \text{ h}^{-1}$  ( $40^\circ\text{C}$ ) are in the same order of magnitude as that estimated by Shehata and Marr (42). All of them are at least 1 order of magnitude higher than the maintenance requirement measured by Schulze and Lipe (39).

Additionally, the  $s_{\text{min}}$  estimated from the experimental data ( $12 \mu\text{g}$  of glucose  $\text{liter}^{-1}$  at  $37^\circ\text{C}$ ) can be compared with the threshold substrate concentration of  $10.8 \mu\text{g liter}^{-1}$  computed according to the model proposed by Button (6), using the values of  $K_s = 32.8 \mu\text{g liter}^{-1}$  and  $\mu_{\text{max}} = 0.76 \text{ h}^{-1}$  obtained in the present study and a rate of endogenous metabolism of  $0.25 \text{ h}^{-1}$  (reported in reference 6). The very complex model proposed by Schmidt and coworkers (38) predicts a threshold concentration for growth of  $2.25 \mu\text{g liter}^{-1}$  glucose, which is slightly lower than the value of  $12 \mu\text{g liter}^{-1}$  extrapolated from our experimental data. However, it should be stressed that such comparisons have to be made carefully, because many of the model parameters required for calculation of  $s_{\text{min}}$  in some of the more complex models were not measured in this study but had to be taken from other studies.

The reported threshold ( $s_{\text{min}}$ ) concentrations are difficult to compare, because in each case different experimental systems were used. However, an analysis of the trends exhibited by the data should not be affected by the experimental setup. Interestingly, the trend for the estimates of specific maintenance rates (Fig. 9) from our own data as a function of temperature is comparable with those measured for *E. coli* by Wallace and Holms (47) and also with the effect of temperature on the rate of synthesis of  $\beta$ -galactosidase in *E. coli* (20). A dramatic increase in the specific maintenance rate was also observed in the superoptimal temperature range by Mainzer and Hempfling (19). The dependence of  $s_{\text{min}}$  on temperature did not exactly follow the pattern shown in Fig. 9. This can be due to the fact that temperature affects the maintenance rate at the level both of  $s_{\text{min}}$  and of  $\mu_{\text{max}}$  (see equation 6).

**Is the substrate saturation constant temperature dependent?** Virtually identical  $K_s$  values for glucose (Table 3) were obtained from fitting the extended Monod model (equation 2) to the experimental data at  $17.4$ ,  $28.4$ ,  $37$ , and  $40^\circ\text{C}$ . This is in agreement with data already reported by von Meyenburg (46), who found similar values for  $K_m$  for a mutant of *E. coli* growing with glucose at  $30$  and  $37^\circ\text{C}$ . However, in the few studies which are available on temperature modulation of saturation constants, both positive (7, 43, 49) and negative (7, 16, 18) modulations of  $K_s$  by temperature have been reported. In each of




FIG. 9. Comparison of temperature dependence of maintenance rates and rate of synthesis of  $\beta$ -galactosidase in *E. coli* strains. (Rates are relative values standardized with respect to the minimal value.) Symbols: open circles, specific maintenance rates calculated from  $s_{\text{min}}$  values for *E. coli* growing with glucose (see Table 3, equation 6); diamonds, specific maintenance rates calculated from the experimental data of Wallace and Holms (47) for *E. coli* during growth with glucose; closed circles, differential rates of  $\beta$ -galactosidase synthesis in submaximally induced cryptic strain (20).

these studies, the Arrhenius equation was used to describe the temperature modulation of the  $K_s$  or  $K_m$  constant (7, 49), since it was assumed to be generally valid for defining the temperature dependency of chemical rate constants (8). However, it has never been confirmed that this thermodynamic concept can be applied to such a complex parameter as the Monod saturation constant.

Considering that even for one particular temperature the steady-state glucose concentrations supporting half- $\mu_{\text{max}}$  ( $K_s$  values) have been reported to vary by more than 3 orders of magnitude (40), one has to question the accuracy of such studies of temperature modulation of growth constants. Some investigations indicate (40, 46) that this enormous variability is due to insufficient adaptation of bacteria to low substrate concentrations. To overcome this, standardization of experiments is necessary in comparative studies. In our experimental study, we used cells fully and reproducibly adapted to growth-limiting concentrations in continuous culture. Interestingly, for *Methanosarcina barkeri* it was shown that the temperature dependency of the affinity constant for molecular hydrogen and acetate was reduced when the substrate concentration was lowered to subsaturating levels, i.e., toward concentrations found in balanced ecosystems (49). These data support the results reported here for *E. coli*. For growth at very low substrate concentrations, Button (6) proposed that the ability of bacteria to grow is better described by a specific affinity term which is defined as the ratio of  $\mu_{\text{max}}$  and  $K_s$ . Since saturation constants are temperature independent and  $\mu_{\text{max}}$  varies with temperature, specific affinity will also be temperature dependent.

These observations raise the issue of whether the reported temperature modulation of substrate affinity is an experimental artifact and what is the theoretical background to such behavior. In the appendix, we present a simplified example and suggest possible reasons for a temperature-independent  $K_s$  value. The main difference between  $K_s$  and the other model parameters ( $s_{\text{min}}$  and  $\mu_{\text{max}}$ ) is that because of the complexity of the Monod saturation constant, both the negative and the positive temperature effects may compensate each other. De-

spite the clear evidence obtained in our experiments that the  $K_s$  value for glucose was temperature independent, it would still be premature to generalize this finding. When comparing with other published results (7, 16, 18, 43, 46, 49), one has to keep in mind several factors such as the differences in experimental setup (batch versus continuous culture cultivation, concentration of the feed, etc.), different methods that were used in determining model parameters (linearization or non-linear parameter estimation), and the specific limitations inherent in hyperbolic models, namely, that the parameters strongly influence each other.

**What are the main factors influencing the steady-state substrate concentrations?** In our experience (for comparison, see Fig. 2), steady-state concentrations of the growth-limiting substrate are reproducible under defined conditions in continuous culture. Similarly, in open systems the growth conditions will determine the limit down to which particular compounds can be degraded. Here, specific growth rate, temperature, and composition of the utilized C pool in the feed were shown to be important factors affecting a steady-state concentration of glucose during growth of *E. coli* in the chemostat. Under optimal growth conditions (defined as pH  $\cong$  7, T  $\cong$  37°C), the steady-state glucose concentration of  $32 \pm 4$   $\mu\text{g/liter}$  was measured at a dilution rate of  $0.3 \text{ h}^{-1}$ . This concentration increased when the cultivation temperature was out of the optimum range and at higher growth rates. Vice versa, the steady-state glucose concentration decreased with decreasing growth rates or when the cells were simultaneously utilizing an additional substrate.

We are aware of only two other reports in which parabola-like  $s = f(T)$  dependencies similar to those for glucose described in this study were observed, both of them for nitrifiers in batch reactors (7, 18). The trend in steady-state glucose concentrations affected by temperature was sufficiently well described by the reciprocal Ratkowsky model (equation 11). Unfortunately, the minimum steady-state substrate concentration does not coincide with the optimum temperature for growth. At  $D = 0.3 \text{ h}^{-1}$ , approximately the same glucose concentrations (i.e.,  $31 \pm 4$   $\mu\text{g/liter}$ ) are predicted between the range of 31 and 37°C (Fig. 6); however, according to batch data (Fig. 5), one would expect the minimum to be at approximately 38°C. As such, a simple parabolic function would describe the data similarly well. Such models have been proposed for the description of the effect of pH and temperature on the growth rate (36, 51). Fortunately, the reciprocal Ratkowsky model is only a modification of an already accepted model for temperature dependency of bacterial growth. This modification was also successfully applied for the description of the influence of temperature on lag time (50, 53).

**Are the concentrations observed in real environments comparable to those of laboratory-scale experiments?** In natural waters, threshold concentrations of certain compounds have been observed below which these compounds were not significantly utilized or at which the rate of their degradation slowed down enormously (summarized in reference 2). This phenomenon has been attributed to the fact either that a certain amount of substrate is needed to sustain necessary metabolic functions (i.e., maintenance energy [29]) or that the concentration of a particular compound is too low to provoke induction of the enzymes necessary for its degradation (2, 9). As discussed above, the former phenomenon is based on the kinetic properties of a particular strain. However, the efficiency with which a bacterium takes up substrate is influenced by a variety of environmental factors, an important one being the presence and simultaneous utilization of additional substrates (10). Therefore, one would expect that the concentrations of particular substrates under environmental conditions should

be lower than those observed for laboratory-scale experiments in which these compounds are supplied as the only substrates for growth. The  $s_{\text{min}}$  would, therefore, be the highest expected substrate concentration under conditions of no growth. The fact that the observed maintenance energy requirements of indigenous soil microbial populations were some 3 orders of magnitude lower than those known from pure cultures supports this line of thinking (3, 24). It should be pointed out that such thresholds should not be observed in closed systems such as batch cultures (44), because the maintenance requirement of cells implies continued utilization until all available substrate is exhausted. This might not be the case when, e.g., toxic metabolites are accumulated, or when the culture is limited by an element other than carbon.

#### ACKNOWLEDGMENTS

We acknowledge the advice of A. Hiltbold in processing the experimental data. We are also indebted to U. Lendenmann for introducing K.K. to the handling of the analytical equipment, to M. Snozzi for helpful discussion, and to C. A. Mason for linguistic help.

#### APPENDIX

Some quite contrasting temperature dependencies of  $K_s$  have been reported in the literature (7, 46), although to our knowledge, this problem has not yet been treated theoretically. On the basis of the mathematical analogy between  $K_s$  and the Michaelis-Menten saturation constant ( $K_m$ ), we will try here to explain theoretically our experimental data that, for *E. coli* growing with glucose, have demonstrated that  $K_s$  was independent of temperature. Analogously to  $K_m$  (for its definition, see reference 8), the  $K_s$  can be interpreted as a ratio of rate constants of a single enzymatic step (35) that limits the specific growth rate (equation 15):

$$K_s = \frac{(k_j + k_h)}{k_i} \quad (15)$$

For the special case in which all of the individual rate constants exhibit similar temperature dependencies (equations 16a, b, and c and 17), the apparent  $K_s$  parameter should also be independent of temperature (equation 18).

When

$$k_i(T_1) = q_1 \times k_i(T_2) \quad (16a)$$

$$k_j(T_1) = q_2 \times k_j(T_2) \quad (16b)$$

$$k_h(T_1) = q_3 \times k_h(T_2) \quad (16c)$$

and

$$q = q_1 = q_2 = q_3 \quad (17)$$

then  $K_s$  is independent of temperature (equation 18), i.e.,  $K_s(T_1)$  and  $K_s(T_2)$  are identical:

$$K_s(T_1) = \frac{(k_j + k_h)}{k_i} \text{ and therefore,}$$

$$K_s(T_2) = \frac{q \times k_j + q \times k_h}{q \times k_i} = \frac{q \times (k_j + k_h)}{q \times k_i} = K_s(T_1) \quad (18)$$

Assuming that the rate constants are dependent on temperature according to the Arrhenius equation (equation 19), the parameter  $q$  is expressed as shown in equation 20. For the case in which  $q_1$ ,  $q_2$ , and  $q_3$  are identical, this equation implies that the activation energies are also the same:

$$\ln \frac{k(T_1)}{k(T_2)} = \frac{E_a(T_1 - T_2)}{R T_1 T_2} \quad (19)$$

$$q = \exp\left(\frac{E_a(T_1 - T_2)}{R T_1 T_2}\right) \quad (20)$$

## REFERENCES

- Adams, M. R., C. L. Little, and M. C. Easter. 1991. Modelling the effect of pH, acidulant and temperature on the growth rate of *Yersinia enterocolitica*. *J. Appl. Bacteriol.* **71**:65–71.
- Alexander, M. 1994. Biodegradation and bioremediation. Academic Press, San Diego, Calif.
- Andreson, T.-H., and K. H. Domsch. 1985. Determination of ecophysiological maintenance carbon requirements of soil microorganisms in dormant state. *Biol. Fert. Soils* **1**:81–89.
- Beck, J. V., and K. J. Arnold. 1977. Parameter estimation in engineering and science. Wiley, New York.
- Buchanan, R. L., and L. A. Klawitter. 1992. The effect of incubation temperature, initial pH, and sodium chloride on the growth kinetics of *Escherichia coli* O157:H7. *Food Microbiol.* **9**:185–196.
- Button, D. K. 1985. Kinetics of nutrient-limited transport and microbial growth. *Microbiol. Rev.* **49**:270–297.
- Characklis, W. G., and W. Gujer. 1979. Temperature dependency of microbial reactions. *Prog. Water Technol. Suppl.* **1**:111–130.
- Cornish-Bowden, A. 1979. Fundamentals of enzyme kinetics. Butterworth, London.
- DiMarco, A. A., B. Averhoff, and L. N. Ornston. 1995. Identification of the transcriptional activator *pobR* and characterisation of its role in the expression of *pobA*, the structural gene for *p*-hydroxybenzoate hydrolase in *Acinetobacter calcoaceticus*. *J. Bacteriol.* **175**:4499–4506.
- Egli, T., U. Lendenmann, and M. Snozzi. 1993. Kinetics of microbial growth with mixtures of carbon sources. *Antonie van Leeuwenhoek* **63**:289–298.
- Esener, A. A., J. A. Roels, and N. W. F. Kossen. 1981. The influence of temperature on the maximum specific growth rate of *Klebsiella pneumoniae*. *Biotechnol. Bioeng.* **23**:1401–1405.
- Gibson, A. M., N. Bratchell, and T. A. Roberts. 1987. The effect of sodium chloride and temperature on the rate and extent of growth of *Clostridium botulinum* type A in pasteurized pork slurry. *J. Appl. Bacteriol.* **62**:479–490.
- Heitzer, A., H.-P. E. Kohler, P. Reichert, and G. Hamer. 1991. Utility of phenomenological models for describing temperature dependence of bacterial growth. *Appl. Environ. Microbiol.* **57**:2656–2665.
- Heitzer, A., C. A. Mason, and G. Hamer. 1992. Heat shock gene expression in continuous cultures of *Escherichia coli*. *J. Biotechnol.* **22**:153–170.
- Herendeen, S. L., R. A. VanBogelen, and F. C. Neidhardt. 1979. Levels of major proteins of *Escherichia coli* during growth at different temperatures. *J. Bacteriol.* **139**:185–194.
- Ingraham, J. 1987. Effect of temperature, pH, water activity, and pressure on growth, p. 1543–1554. In F. C. Neidhardt, J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, vol. 2. American Society for Microbiology, Washington, D.C.
- Jones, R. C., and J. S. Hough. 1970. The effect of temperature on the metabolism of baker's yeast growing on continuous culture. *J. Gen. Microbiol.* **60**:107–116.
- Knowles, G., A. L. Downing, and M. J. Barrett. 1965. Determination of kinetic constants for nitrifying bacteria in mixed culture, with the aid of an electronic computer. *J. Gen. Microbiol.* **38**:263–278.
- Mainzer, S. E., and W. P. Hempfling. 1976. Effects of growth temperature on yield and maintenance during glucose-limited continuous culture of *Escherichia coli*. *J. Bacteriol.* **126**:251–256.
- Marr, A. G., J. L. Ingraham, and C. L. Squires. 1964. Effect of the temperature of growth of *Escherichia coli* on the formation of  $\beta$ -galactosidase. *J. Bacteriol.* **87**:356–362.
- McMeekin, T. A., R. E. Chandler, P. E. Doe, C. D. Garland, J. Olley, S. Putro, and D. A. Ratkowsky. 1987. Model for combined effect of temperature and salt concentration/water activity on the growth rate of *Staphylococcus xylosum*. *J. Appl. Bacteriol.* **62**:543–550.
- McMeekin, T. A., J. Olley, and D. A. Ratkowsky. 1988. Temperature effects on bacterial growth rates, p. 75–89. In M. J. Bazin and J. I. Prosser (ed.), *Physiological models in microbiology*, vol. I. CRC Press, Inc., Boca Raton, Fla.
- Monod, J. 1942. Recherches sur la croissance des cultures bactériennes. Hermann et Cie., Paris.
- Morita, R. Y. 1988. Bioavailability of energy and its relationship to growth and starvation survival in nature. *J. Can. Microbiol.* **43**:436–441.
- Ng, H., J. L. Ingraham, and A. G. Marr. 1962. Damage and derepression in *Escherichia coli* resulting from growth at low temperatures. *J. Bacteriol.* **84**:331–339.
- Nishihara, M., M. Ishinaga, M. Kato, and M. Kito. 1976. Temperature-sensitive formation of the phospholipin molecular species in *Escherichia coli* membranes. *Biochim. Biophys. Acta* **431**:54–61.
- Palumbo, S., and L. D. Witter. 1969. Influence of temperature on glucose utilisation by *Pseudomonas fluorescens*. *Appl. Bacteriol.* **18**:137–141.
- Pirt, S. J. 1965. The maintenance energy of bacteria in growing cultures. *Proc. R. Soc. Ser. B* **163**:224–231.
- Pirt, S. J. 1975. Principles of microbe and cell cultivation. Blackwell, London.
- Powell, E. O. 1967. The growth rate of micro-organisms as function of substrate concentration, p. 34–55. In C. G. T. Evans, R. E. Strange, and D. W. Tempest (ed.), *Microbial physiology and continuous culture*. HMSO, London.
- Ratkowsky, D. A., R. K. Lowry, T. A. McMeekin, A. N. Stokes, and R. E. Chandler. 1983. Model for bacterial culture growth rate throughout the entire biokinetic temperature range. *J. Bacteriol.* **154**:1222–1226.
- Ratkowsky, D. A., J. Olley, T. A. McMeekin, and A. Ball. 1982. Relationship between temperature and growth rate of bacterial cultures. *J. Bacteriol.* **149**:1–5.
- Richter, O., and D. Söndgerath. 1990. Parameter estimation in ecology. The link between data and models. Verlag Chemie, Weinheim, Germany.
- Robinson, J. A. 1985. Determining microbial kinetic parameters using non-linear regression analysis. *Adv. Microb. Ecol.* **8**:61–114.
- Ron, E. Z., and M. Shani. 1971. Growth rate of *Escherichia coli* at elevated temperatures: reversible inhibition of homoserin trans-succinylase. *J. Bacteriol.* **107**:397–400.
- Rosso, L., J. R. Lobry, S. Bajard, and J. P. Flandrois. 1995. Convenient model to describe the combined effects of temperature and pH on microbial growth. *Appl. Environ. Microbiol.* **61**:610–616.
- Rosso, L., J. R. Lobry, and J. P. Flandrois. 1993. An unexpected correlation between cardinal temperatures of microbial growth highlighted by a new model. *J. Theor. Biol.* **162**:447–463.
- Schmidt, S. K., M. Alexander, and M. L. Shuler. 1985. Predicting threshold concentrations of organic substrates for bacterial growth. *J. Theor. Biol.* **114**:1–8.
- Schulze, K. L., and R. S. Lipe. 1964. Relationship between substrate concentration, growth rate, and respiration rate of *Escherichia coli* in continuous culture. *Arch. Microbiol.* **48**:1–20.
- Senn, H., U. Lendenmann, M. Snozzi, G. Hamer, and T. Egli. 1994. The growth of *Escherichia coli* in glucose-limited chemostat cultures: a re-examination of the kinetics. *Biochim. Biophys. Acta* **1201**:424–436.
- Shaw, M. K., and J. Ingraham. 1965. Fatty acid composition of *Escherichia coli* as a possible controlling factor of the minimal growth temperature. *J. Bacteriol.* **90**:141–146.
- Shehata, T. E., and A. G. Marr. 1971. Effect of nutrient concentration on the growth of *Escherichia coli*. *J. Bacteriol.* **107**:210–216.
- Topiwala, H., and C. G. Sinclair. 1971. Temperature relationships in continuous culture. *Biotechnol. Bioeng.* **13**:795–813.
- Tros, M. E., G. Schraa, and A. J. B. Zehnder. 1996. Transformation of low concentrations of 3-chlorobenzoate by *Pseudomonas* sp. strain B13: kinetics and residual concentrations. *Appl. Environ. Microbiol.* **62**:437–442.
- van Uden, N. 1967. Transport-limited growth in the chemostat and its competitive inhibition: a theoretical treatment. *Arch. Microbiol.* **58**:145–154.
- von Meyenburg, K. 1971. Transport-limited growth rates in a mutant of *Escherichia coli*. *J. Bacteriol.* **107**:878–888.
- Wallace, R. J., and W. H. Holms. 1986. Maintenance coefficients and rates of turnover of cell material in *Escherichia coli* ML308 at different growth temperatures. *FEMS Microbiol. Lett.* **37**:317–320.
- Westerhoff, H. V., J. S. Lokema, R. Otto, and K. J. Hellingwerf. 1982. Thermodynamics of growth, non-equilibrium thermodynamics of bacterial growth, the phenomenological and the mosaic approach. *Biochim. Biophys. Acta* **683**:181–220.
- Westermann, P., B. K. Ahring, and R. A. Mah. 1989. Temperature compensation in *Methanosarcina barkeri* by modulation of hydrogen and acetate affinity. *Appl. Environ. Microbiol.* **55**:1262–1266.
- Wijtztes, T., P. J. McClure, M. H. Zwietering, and T. A. Roberts. 1993. Modelling bacterial growth of *Listeria monocytogenes* as a function of water activity, pH, and temperature. *Int. J. Food Microbiol.* **18**:139–149.
- Wijtztes, T., J. C. de Wit, J. H. J. Huis in't Veld, K., van't Riet, and M. H. Zwietering. 1995. Modeling of bacterial growth of *Lactobacillus curvatus* as a function of acidity and temperature. *Appl. Environ. Microbiol.* **61**:2533–2539.
- Zwietering, M. H., J. T. de Koos, B. E. Hasenack, C. J. de Wit, and K. van't Riet. 1991. Modelling of bacterial growth as a function of temperature. *Appl. Environ. Microbiol.* **57**:1094–1101.
- Zwietering, M. H., C. J. de Wit, G. A. M. Cuppers, and K. van't Riet. 1994. Modeling of bacterial growth with shifts in temperature. *Appl. Environ. Microbiol.* **60**:204–213.