

## Reversible Phase Variation in the *phnE* Gene, Which Is Required for Phosphonate Metabolism in *Escherichia coli* K-12

Samina Iqbal, George Parker, Helen Davidson, Elham Moslehi-Rahmani, and Robert L. Robson\*

Microbiology Division, School of Animal and Microbial Sciences, University of Reading, Reading, United Kingdom

Received 23 April 2004/Accepted 22 June 2004

It is known that *Escherichia coli* K-12 is cryptic ( $\text{Phn}^-$ ) for utilization of methyl phosphonate (MePn) and that  $\text{Phn}^+$  variants can be selected for growth on MePn as the sole P source. Variants arise from deletion via a possible slip strand mechanism of one of three direct 8-bp repeat sequences in *phnE*, which restores function to a component of a putative ABC type transporter. Here we show that  $\text{Phn}^+$  variants are present at the surprisingly high frequency of  $>10^{-2}$  in K-12 strains. Amplified-fragment length polymorphism analysis was used to monitor instability in *phnE* in various strains growing under different conditions. This revealed that, once selection for growth on MePn is removed,  $\text{Phn}^+$  revertants reappear and accumulate at high levels through reinsertion of the 8-bp repeat element sequence. It appears that, in K-12, *phnE* contains a high-frequency reversible gene switch, producing phase variation which either allows (“on” form) or blocks (“off” form) MePn utilization. The switch can also block usage of other metabolizable alkyl phosphonates, including the naturally occurring 2-aminoethylphosphonate. All K-12 strains, obtained from collections, appear in the “off” form even when bearing mutations in *mutS*, *mutD*, or *dnaQ* which are known to enhance slip strand events between repetitive sequences. The ability to inactivate the *phnE* gene appears to be unique to K-12 strains since the B strain is naturally  $\text{Phn}^+$  and lacks the inactivating 8-bp insertion in *phnE*, as do important pathogenic strains for which genome sequences are known and also strains isolated recently from environmental sources.

*Escherichia coli* can use  $\text{P}^{\text{III}}$  compounds such as phosphite and organophosphonates, e.g., methylphosphonate (MePn) and aminoethylphosphonate (AEPn) as P sources. Their metabolism involves the enzyme C-P lyase, which appears to have a relatively broad substrate specificity (21). Whereas the mechanism of C-P lyase is not well understood, much is known about a cluster of 17 contiguous *phn* genes in *E. coli*, required for utilization of  $\text{P}^{\text{III}}$  compounds (5, 21). The *phnGHIJK* genes within this cluster are thought to encode the core components of C-P lyase, while *phnF* and *phnO* potentially encode regulatory proteins (5). Several *phn* genes appear to encode components of solute transporters, and it has been deduced that, among these, the *phnCDE* genes encode an ABC type transporter. In this transporter, *phnC* encodes the ABC permease component, *phnD* encodes the periplasmic binding protein, and *phnE* encodes the integral membrane component.

An interesting feature of the genetics of phosphonate metabolism in *E. coli* is that the B strain can use phosphonates whereas the K-12 strain is cryptic despite containing the entire *phn* gene cluster (21). The genetic basis for this crypticity was investigated by Makino et al. (14) and traced to an 8-bp insertion in the coding region of the *phnE* gene in the K-12 strain relative to the B strain, causing truncation of the *phnE* product. They also observed that the 8-bp sequence is one element in the direct triply repeated sequence in the K-12 strain comprising two types of octamer variants in the arrangement 5'-ABB-3', where A corresponds to the sequence 5'-CGCTGGCG-3' and B corresponds to the sequence 5'-TGCTGGCG-3' (Fig.

1). Makino et al. isolated variants of *E. coli* K-12 able to use MePn as the sole P source ( $\text{Phn}^+$ ), and these were found to have deletions of octamer B, which, they postulated, occurred via a strand slippage event during DNA replication (14). The nature of the variation in the *phnE* gene in *E. coli* is investigated in more detail in this work.

### MATERIALS AND METHODS

**Bacterial strains and growth conditions.** All the strains of *E. coli* used in this study (Table 1) were routinely cultivated under aerobic conditions at 37°C in Luria-Bertani broth (LB) with agar added to 0.8%, wt/vol, for agar plates except for  $\text{Phn}^+$  variants of K-12 strains (see below) and temperature-sensitive mutants for which the permissive temperature for growth was 30°C. The minimal medium used was modified Neidhardt's medium (MNM) and was based on that of Neidhardt et al. (15), but, in order to study the metabolism of different phosphorous sources, the normal phosphate buffer component was replaced with MOPS (morpholinepropanesulfonic acid). MNM contained MOPS (40 mM), glucose (11.1 mM),  $\text{NH}_4\text{Cl}$  (9.5 mM), Tricine (4 mM), thiamine · HCl (29.6  $\mu\text{M}$ ),  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  (10  $\mu\text{M}$ ),  $\text{CaCl}_2$  (0.5  $\mu\text{M}$ ),  $\text{MnCl}_2$  (0.8 nM),  $\text{CoCl}_2$  (0.3 nM),  $\text{CuSO}_4$  (0.16 nM),  $\text{ZnSO}_4$  (0.1 nM),  $(\text{NH}_4)_6\text{Mn}_7\text{O}_{24}$  (30 pM), and  $\text{H}_3\text{BO}_4$  (4 pM). The medium was adjusted to pH 7.4. MNM was solidified for plates (MNM-agarose) with electrophoresis grade agarose (1%, wt/vol) because it is especially low in phosphate compared to purified agars. MNM and MNM-agarose were supplemented with different P sources at 0.5 mM.  $\text{Phn}^+$  variants of K-12 strains were isolated, purified, and maintained on MNM-agarose containing MePn unless otherwise stated. Culture dilutions were made in MNM lacking added P sources. Phosphorous compounds were obtained from the following sources. MePn was from Fluka Chemie AG; 2-AEPn, aminomethylphosphonate (AMPn), phosphonoacetate, phosphonoformate, phosphonomycin, and *o*-phospho-L-serine were from Sigma Chemical Co.; 1-AEPn, 1-aminopropylphosphonate (1-APPn), 3-APPn, *tert*-butylphosphonate, ethylphosphonate (EPn), phosphonomethylglycine (glyphosate), phenylphosphonate (PhPn), and propylphosphonate (PPn) were from Aldrich Chemical Co.; and *N*-butanephosphonate was from Lancaster Synthesis, Morecambe, United Kingdom.

**Estimation of the frequency of  $\text{Phn}^+$  variants.** The frequency of  $\text{Phn}^+$  variants in populations of *E. coli* strains was estimated by comparing the number of CFU arising when populations were diluted appropriately and plated onto MNM-agarose containing MePn compared to the number on MNM-agarose containing either  $\text{P}_i$  or *o*-phospho-L-serine (positive controls) or no added P source (nega-

\* Corresponding author. Mailing address: Microbiology Division, School of Animal and Microbial Sciences, University of Reading, Reading RG6 6AJ, United Kingdom. Phone: 44-118-9316639. Fax: 44-118-9316562. E-mail: r.l.robson@rdg.ac.uk.




FIG. 1. Repetitive sequence in *phnE* of *E. coli*. The figure shows part of the *phn* gene cluster in *E. coli* and focuses on *phnE* and the location and sequences of a direct triple repeat in the K-12 strain and a direct double repeat in the B strain, where A corresponds to the octamer 5'-CGCTGGCG-3' and B corresponds to the octamer 5'-TGCTGGCG-3'. The relative positions of pairs of primer used in this work to amplify two different segments of *phnE* of *E. coli* K-12 are shown under the gene organization. Amplicon 1 was used in the study in the AFLP analysis, and amplicon 2 was used for gene sequencing.

tive control). The inocula for these experiments were first grown to stationary phase on MNM containing P<sub>i</sub> and then washed in MNM containing no added P source. Dilutions were performed with MNM containing no added P source. Colonies arising on phosphate or phosphonates were counted at 4 and 10 days, respectively, because growth of colonies on phosphonates was generally slower than growth on P<sub>i</sub>.

**Molecular biology techniques.** (i) **Extraction of genomic DNA.** Template DNAs for PCRs were extracted from cultures as follows. One-tenth milliliter of

culture was diluted with 0.9 ml of sterile distilled water in a 1.5-ml microcentrifuge tube. The tube was boiled for 5 min and snap-chilled in ice. The boiled cell suspensions were diluted 100-fold in sterile water, and 5 µl was sufficient to set up PCRs.

(ii) **AFLP analysis.** Deletions or insertions in *phnE* were monitored by amplified-fragment length polymorphism (AFLP) analysis as follows. Fragments of *phnE* of 200 or 192 bp containing the triple or double octameric repeat regions, respectively, were amplified from samples of liquid cultures or colonies by PCR using oligonucleotide primer set 1 (Fig. 1), comprising EcphnEF2 (5'-Cy 5-TTACCAGCCCGTTGCGCGCC-3') and EcphnER2 (5'-CCTTCCACCGGG CCAGGTTCAAT-3'). Amplifications were carried out with Bio-X-Act DNA polymerase (Bioline UK Ltd.) with the following thermal cycle: 30 cycles of 95°C for 30 s, 60°C for 1 min, and 68°C for 1 min. Products were purified with a PCR fragment purification kit (QIAGEN) and checked by gel electrophoresis in a 1% Tris-acetate-EDTA-agarose gel. Fragments were diluted and run together with standard-size fragments (50-bp ladder; Amersham) on a 6%, wt/vol, polyacrylamide sequencing gel in an ALF-Express automated sequencer (Amersham Pharmacia), and the gel image was visualized with ALF-Express software. Fragments were detected, sized, and quantified with AllelLinks, version 1.00 (Amersham Pharmacia). The relative proportions of the two alleles of *phnE* were estimated from a standard curve prepared from AFLP analysis of different mixtures of the K-12 strain MC4100 and the B strain BL21(DE3), prepared by mixing individual cultures grown in LB prior to preparation of templates DNA.

**DNA sequencing.** To sequence the region of *phnE* containing the triply or doubly repeated region, a 600-bp fragment was amplified by PCR with oligonucleotide primer set 2 (Fig. 1), consisting of EcphnES1 (5'-GCGGATCCCGCA GCTG-3') and EcphnER (5'-ACGGTCGCCGAGCGGACGTT-3'). The amplification protocol used was that described for the AFLP analysis above. The DNA was purified with a QIAGEN PCR purification kit and then sequenced with an ALF-Express automated sequencer and a cycle sequencing strategy using Cy 5-labeled EcphnEF and EcphnER primers.

RESULTS

**Occurrence of Phn<sup>+</sup> variants of *E. coli* K-12.** The frequencies of occurrence of Phn<sup>+</sup> variants in populations of common laboratory *E. coli* K-12 strains DH5α and MC4100, grown in LB, were estimated by determining the CFU appearing 10 days after plating washed populations at suitable dilutions on MNM-agarose containing MePn and comparing them to those

TABLE 1. *E. coli* strains used

Strain	Genotype	Source or reference
AB1157	F <sup>-</sup> <i>thi-1 hisG4 Δ(gpt-proA)62 argE3 thr-1 leuB6 kdgK51 rfbD1 ara-14 lacY-1 galK2 xyl-5 mtl-1 tsx-33 supE44 rpsL-31 Rac<sup>-</sup> λ<sup>-</sup></i>	1
BL21 (DE3)	F <sup>-</sup> <i>ompT hsdS<sub>B</sub> (r<sub>B</sub> m<sub>B</sub>) dcm gal λ(DE3)</i>	19
CD4	Hfr <i>metD88 proA3 Δ(lacI-Y)6 tsx-76 λ<sup>-</sup> relA1 malA36(λ<sup>r</sup>) metB1</i>	7
DH5α	φ80 <i>dlacZΔM15 recA1 endA1 gyrA96 thi-1 hsdR17 (r<sub>k</sub><sup>-</sup> m<sub>k</sub><sup>+</sup>) supE44 relA1 deoR Δ(lacZYA-argF)U169</i>	
KA796	<i>ara thi Δpro lac</i>	7
MC4100	<i>araD139 Δ(argF-leu)169 LAM<sup>-</sup> e14<sup>-</sup> FlhD5301 fruA25 relA1 rpsL150 RbsR22 deoC1</i>	4
NR9458	KA796 <i>mutD5 zaf13::Tn10</i>	8
NR9807	CD4 <i>dnaQ49</i>	8
STL1671	AB1157 <i>sbcB15 Δ(slr-recA)304</i>	3
STL2172	AB1157 <i>mutS201::Tn5 Δ(slr-recA)304</i>	13
STL2314	AB1157 <i>dnaQ Δ(slr-recA)304</i>	17
NWA3	Wild-type isolate from sewerage sludge	North West Water UK
NWA4	Wild-type isolate from sewerage sludge	North West Water UK
NWA5	Wild-type isolate from sewerage sludge	North West Water UK
NWD7	Wild-type isolate from sewerage sludge	North West Water UK
NW23341	Wild-type isolate from raw water	North West Water UK
NW211585	Wild-type isolate from raw water	North West Water UK
ST1	Wild-type isolate from water	Severn Trent Water UK
ST8	Wild-type isolate from water	Severn Trent Water UK
ST11	Wild-type isolate from water	Severn Trent Water UK
ST15	Wild-type isolate from water	Severn Trent Water UK




FIG. 2. AFLP analysis of *phnE* in *E. coli*. Primer pair 1 (Fig. 1) was used to amplify a small fragment containing the octameric repetitive element in *phnE* from populations of two strains of *E. coli* grown to log phase in LB. PCR products were analyzed as described in Materials and Methods. Shown are fragments detected as peaks in the AFLP analysis. Tracks 1 and 5, PCR fragments produced from cultures of K-12 strain MC4100 and B strain BL21(DE3), respectively; tracks 2 to 4, artificial mixtures of MC4100 and BL21(DE3), each grown to an optical density at 600 nm of 0.7 and mixed prior to extraction of template and amplification in the ratios 9:1, 6:4, and 2:8; track M, 200-bp marker peak from the 50-bp standard ladder set used to determine fragment lengths.

appearing on MNM-agarose containing  $P_i$  or *o*-phospho-L-serine or no added P source, as a negative control.  $Phn^+$  variants were observed to be present at the surprisingly high levels of 3.4 and 8.6% of the total CFU of DH5 $\alpha$  and MC4100, respectively, of those observed on the plates containing  $P_i$  or *o*-phospho-L-serine. Plates with no added P source contained only pinpoint colonies, presumably growing on traces of utilizable P in the medium.

**AFLP analysis of genotypic events in *E. coli* K-12  $Phn$  variants.** AFLP analysis was used to monitor the status of the *phnE* gene in different populations of strains. DNA fragments spanning that segment of the *phnE* gene in K-12 which contains the triple octameric repeat were amplified with primer set 1 (Fig. 1) from culture lysates of organisms grown in various media. When strains were grown in LB, as expected, a 200-bp fragment was amplified from the K-12 strain (DH5 $\alpha$ ) while a 192-bp fragment was amplified from the B strain [BL21(DE3)] (Fig. 2). When the analysis was performed on various mixtures of cultures of these two strains, both 200- and 192-bp fragments were amplified and the peak areas were approximately proportional to those for the prepared mixtures (Fig. 2). This establishes that AFLP can be used to detect both forms of *phnE* if present in populations. AFLP analysis was then used to




FIG. 3. AFLP analysis of *phnE* in cultures of *E. coli* K-12 MC4100 growing with different phosphonates. Cultures of K-12 MC4100 were grown in MNM with different phosphonates as sole P sources, which had been inoculated from  $Phn^+$  variants isolated and purified on MNM-agarose containing the respective phosphonate. PCR products were analyzed as described in Materials and Methods. Shown are fragments detected as peaks in the AFLP analysis. Tracks correspond to the phosphonates supporting growth as follows: 2, MePn, log phase; 3, MePn, stationary phase; 4, EPn, log phase; 5, AMPn, log phase; 6, 2-AEPn, log phase; 7, 3-APPn, log phase; M, 200-bp marker peak from the 50-bp standard ladder set used to determine fragment lengths.

examine *phnE* in cultures growing in MNM containing alternative P sources. Studies were performed on both MC4100 and DH5 $\alpha$ . Results were essentially identical. Fragments of 200 bp were amplified only from cultures grown to stationary phase in MNM containing  $P_i$  and *o*-phospho-L-serine as the sole added P sources. However, populations of  $Phn^+$  variants which had grown to stationary phase in MNM with MePn as the sole added P source gave rise to two fragment types, the major species being 192 bp and the minor species being 200 bp. Therefore, stationary-phase populations of purified  $Phn^+$  variants unexpectedly contained a mixture of two *phnE* alleles, of which the major form probably contained the expected 8-bp deletion reported by Makino et al. (14). We extended these studies by examining the state of *phnE* in cultures in both logarithmic and stationary phases of growth with MePn as the sole P source. In the log-phase populations we could detect only fragments of 192 bp, but, in stationary-phase populations, both 192- and 200-bp fragments were detected (Fig. 3). This suggested that the deletion in *phnE* may be reversible and of




FIG. 4. AFLP analysis of *phnE* in cultures of *E. coli* K-12 MC4100 growing with different phosphonates. Shown is AFLP analysis of *phnE* at the stationary phase of successive subcultures of a *Phn*<sup>+</sup> variant of *E. coli* K-12 MC4100 selected and purified on MNM-agarose. The inoculum level used for each round of subculture was 0.03%. The tracks show the fragments produced after amplification performed at the end of each subculture in the following succession of media: 1, MNM with MePn as the sole added P source; 2 to 4, LB. Tracks M, 200-bp marker peak from the 50-bp standard ladder set used to determine fragment lengths.

high frequency and that it is potentially linked to later stages of the growth phase.

**Reversible switching in the *phnE* gene.** Further evidence for reversibility of the 8-bp deletion in *phnE* was obtained in the following experiment. *Phn*<sup>+</sup> variants of MC4100 originally selected and purified on MNM containing MePn were first grown in MNM plus MePn and then serially subcultured in LB with a 0.03% inoculum level at each subculture. AFLP analysis was performed on samples removed at stationary phase after each round (Fig. 4). In each set of populations derived from a *Phn*<sup>+</sup>

variant, the initial proportion of the 192-bp fragment was high but subculture in LB resulted in a dramatic decrease in the proportion of the 192-bp forms and a reciprocal increase in the proportion of the 200-bp forms (Fig. 4).

**DNA sequencing of amplified fragments.** To confirm that the 8-bp deletion in *phnE* observed in these experiments was identical to that described by Makino et al. (14), the sequences were determined for 600-bp fragments of *phnE* amplified directly from boiled lysates of logarithmic-phase cultures of BL21(DE3) and MC4100 grown on *P*<sub>i</sub> or on MePn and also a *Phn*<sup>+</sup> variant of MC4100 which had been repeatedly subcultured on LB and that therefore had apparently undergone reversion. These data (Fig. 5) confirmed that one octamer B sequence had been lost in the *Phn*<sup>+</sup> variants selected on MePn and that the B strain contains only the AB repeat. The data also showed for the first time that the apparent reversion event detected by AFLP when *Phn*<sup>+</sup> variants were cultivated in LB and not selected for MePn utilization involves precise restoration of an octamer B lost during the original selection process.

**Behavior of the *phnE* gene in different K-12 strains.** The status and behavior of the *phnE* gene in K-12 derivatives carrying mutations in *recA* (STL1671, STL2172, and STL2314), *mutS* (STL2172), *dnaQ* (STL2314 and NR9807), *mutD* (NR9458), and *sbcB* (STL1671) were examined. When cultures were cultivated in MNM with *P*<sub>i</sub> as the sole added P source, a 200-bp fragment was amplified from all strains with primer set 1. *Phn*<sup>+</sup> variants were then isolated for each strain. All *Phn*<sup>+</sup> variants yielded 192-bp fragments after AFLP analysis using oligonucleotide set 1 (Fig. 5).

***Phn*<sup>+</sup> variants obtained with different phosphonate sources.** A requirement for the 8-bp deletion in *phnE* for utilization of other organophosphonates was also examined. In this experiment, *Phn*<sup>+</sup> variants of MC4100 were isolated and purified on MNM-agarose with alternative phosphonates provided as sole P sources. AFLP analysis was performed on logarithmically growing populations with primer set 1. As with *Phn*<sup>+</sup> variants selected and grown on MePn, fragments of 192 bp were amplified from all populations which grew well using EPn, AMPn, 2-AEPn, and 3-APPn as P sources (Fig. 3). Also, as with *Phn*<sup>+</sup> variants selected on MePn, all these variants gave rise to mixtures of 192- and 200-bp fragments in stationary phase (data not shown). Very slow growth was observed with PhPn and PPn. In the PhPn cultures, a 200-bp fragment was the sole fragment amplified, but in the PPn-grown organisms traces of the 192-bp forms were detected. No significant growth was

	A*****B*****B*****
MC4100 / <i>P</i> <sub>i</sub>	GGCCTCGGCCCGTTTCGCTGGCGTGCTGGCGTGCTGGCGTGTTTATCCACACCACCGGCG
MC4100 /MeP	GGCCTCGGCCCGTTTCGCTGGCGTGCTGGCG-----CTGTTTATCCACACCACCGGCG
MC4100/ MeP⇒LB	GGCCTCGGCCCGTTTCGCTGGCGTGCTGGCGTGCTGGCGTGTTTATCCACACCACCGGCG
BL21	GGTCTCGGACCGTTTCGCTGGCGTGCTGGCG-----CTATTTATCCACACCACCGGCG
K-12	GGCCTCGGCCCGTTTCGCTGGCGTGCTGGCGTGCTGGCGTGTTTATCCACACCACCGGCG
B1	GGTCTCGGACCGTTTCGCTGGCGTGCTGGCG-----CTATTTATCCACACCACCGGCG
O157:H72	GGCCTCGGCCCATTCGCTGGTGTGCTGGCG-----CTGTTTATCCACACCACCGGTG
CFT0733	GGCCTCGGCCCGTTTCGCTGGCGTGCTGGCG-----CTGTTTATCCACACCACCGGTG

FIG. 5. Variation in the sequences of *phnE* genes in various isolates of *E. coli*. (Top) Sequences determined in this work for that part of *phnE* containing the octameric repetitive element for the following strains and conditions: MC4100/*P*<sub>i</sub>, MC4100 grown in MNM with *P*<sub>i</sub> as the sole added P source; MC4100/MePn, a *Phn*<sup>+</sup> variant isolated and purified on MNM-agarose with MePn as the sole P source; MC4100/MePn⇒LB, revertant obtained after successive subculture of a *Phn*<sup>+</sup> variant in LB. Asterisks, positions of the direct repeat sequences. (Bottom) Sequences published for four strains of *E. coli*: K-12 and B (14), the enterohemorrhagic strain O157:H7 (9, 16), and the uropathogenic strain CFT073 (22).



observed with the following phosphonates: *N*-butanephosphonate, *tert*-buty phosphonate, 1-AEPn, 1-EPn, 1-APPn, phosphonoacetate, phosphonoformate, and phosphonomycin.

**Properties and behavior of the *phnE* gene in non K-12 *E. coli* strains.** Eleven strains of *E. coli* originally isolated from raw water or sewerage sludge on nutrient broth or LB were screened for potential crypticity in phosphonate metabolism and for the occurrence of the triple-repeat sequences in *phnE*. Their ability to use MePn as a sole P source on MNM was tested as for the K-12 strains previously. Of 11 independent isolates, four strains, 1, 8, A3, and D7, gave plating efficiencies significantly below 100% (at 5.7, 39, 35, and 1.3%, respectively) on MePn compared to the controls supplied with P<sub>i</sub>. This suggested that these strains may be cryptic and may contain an inactivated *phnE* gene similar to that observed in K-12 strains. Therefore we determined the nucleotide sequences of the 600-bp fragments of the *phnE* genes surrounding the octameric repeats amplified from cultures not exposed to added phosphonates (Fig. 5). However, although a few base substitutions affecting only the third base coding positions were observed in the different *phnE* sequences, all strains contained the double octameric repeat sequence in *phnE*, as found in the B strain (data not shown).

## DISCUSSION

We have established that Phn<sup>+</sup> variants appear to be present at a surprisingly high frequency (>10<sup>-2</sup>) in populations of all the *E. coli* K-12 strains tested here. AFLP analysis proved useful in monitoring the genotypic behavior of *phnE* in populations of various strains grown under different conditions, and significantly it revealed the reappearance of the *phnE* allele in populations of Phn<sup>+</sup> variants, especially in stationary-phase cultures. Possible explanations for persistence of the *phnE* allele include cross-feeding of the Phn<sup>-</sup> strains by the Phn<sup>+</sup> variants and/or genetic heterogeneity in *phnE* in individual cells carrying more than one chromosome copy. However, because our studies were conducted with purified Phn<sup>+</sup> isolates, we conclude that the *phnE* gene in *E. coli* K-12 behaves like a reversible gene switch, causing phase variation in phosphonate metabolism. This appears to be the first example of phase variation in a component of an ABC transporter in a gram-negative bacterium (10) although it is interesting that a high-frequency frameshift phase variation event affects the proposed substrate-binding lipoprotein encoded within an ABC transporter operon in *Mycoplasma fermentans* (20). It is not clear why "off" forms start to accumulate rapidly late in the growth cycle on phosphonates or once the selection for growth on phosphonates is removed. From a large number of such studies with different strains, we estimate that the switch may operate at frequencies as least as high as ~10<sup>-2</sup> per generation in either direction but that the equilibrium of the switch strongly favors the "off" form unless selection for phosphonate utilization is applied.

The octameric sequence involved in this postulated slip strand event is, at 8 bp, relatively long, and interestingly it is the most commonly occurring octamer in the genome of *E. coli* K-12 (2). It also contains the core trimer 5'-CTG-3', thought to be the DnaG primase binding site (11, 18, 23, 24). The apparent high instability in *phnE* in K-12 may be linked to the

potential involvement of this octamer in the initiation of DNA replication.

Mutations in DNA replication, repair, and recombination influence instability in tandem repeat sequences (3, 12, 17). All the K-12 mutants examined prior to selection for MePn utilization exhibited the typical K-12 5'-ABB-3' or "off" form of *phnE* even though some carry mutations in functions known to increase deletions between repetitive sequences, including mutations in *recA* and the *sbcB*-encoded 3' exonuclease I and the *dnaQ49<sup>ts</sup>* mutation, which affects DNA polymerase  $\epsilon$ -subunit exonuclease activity and the physical interaction of the  $\epsilon$ -subunit with the polymerizing  $\alpha$ -subunit.

The physiological significance of the switch in *E. coli* K-12 remains unclear. In the "on" direction, the switch allows *E. coli* K-12 to use not only MePn but also EPn, AMPn, 2-AEPn, and 3-APPn; hence *phnE* is implicated in transport of all these organophosphonates. Elashvili et al. (6) have shown that the *phnE* gene is necessary for uptake of some organophosphates since the 8-bp deletion event in *phnE* enabled the *E. coli* K-12 strain JA221 to utilize diisopropyl fluorophosphate and its hydrolysis product, diisopropyl phosphate.

Surprisingly, the *phnE* switch appears to be confined to K-12 strains. *E. coli* strains for which genomes have been determined appear to contain the "on" form of *phnE*, including the uropathogenic strain CFT073 (22), where the 5'-AB-3' repeat is perfectly conserved, and the enterohemorrhagic strain O157:H7 (9, 16), although here T substitutes for C at the seventh base in octamer A in the 5'-AB-3' sequence (Fig. 5). We also found no evidence for the presence of the "off" form of *phnE* in several *E. coli* strains isolated recently from environmental samples.

Although the *phnE* switch may be an artifact possibly arising from the repeated mutagenesis to which the K-12 strain has been subjected, it is possible that it might protect against the uptake of naturally occurring inhibitory phosphonates present in the natural environment or in some way affect surface receptors required for coliphage or lymphocyte recognition.

## ACKNOWLEDGMENTS

Samina Iqbal was supported by a grant from The Islamic Development Bank of Saudi Arabia, George Parker was supported by a research grant from Zeneca Agrochemicals, and Helen Davidson and Elham Moslehi-Rahmani were supported by summer vacation studentships from The Society for General Microbiology.

We thank Elizabeth Pontin and Mike Taylor for running the sequencer and help with the analysis software packages. We also most grateful for the provision of strains to Malgorzata Bzymek, Susan Lovett, Roel Schaaper, and Robert Wells and the *E. coli* Genetic Stock Center at Yale University.

## REFERENCES

- Bachmann, B. J. 1996. Derivations and genotypes of some mutant derivatives of *Escherichia coli* K-12, p. 2460-2488. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology. AMS Press, Washington, D.C.
- Blattner, F. R., G. Plunkett, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* 277:1453-1461.
- Bzymek, M., C. J. Saveson, V. V. Feschenko, and S. T. Lovett. 1999. Slipped misalignment mechanisms of deletion formation: in vivo susceptibility to nucleases. *J. Bacteriol.* 181:477-482.
- Casadaban, M. J. 1976. Transposition and fusion of the *lac* genes to selected

- promoters in *Escherichia coli* using bacteriophage lambda and mu. *J. Mol. Biol.* **104**:541–555.
5. Chen, C.-M., Q.-Z. Ye, Z. Zhu, B. L. Wanner, and C. T. Walsh. 1990. Molecular biology of carbon-phosphorous bond cleavage. Cloning and sequencing of the *phn* (*psiD*) genes involved in alkylphosphonate uptake and C-P lyase activity in *Escherichia coli* B. *J. Biol. Chem.* **265**:4461–4471.
  6. Elashvili, I., J. D. Joseph, and C. C. Valeria. 1998. *phnE* and *glpT* genes enhance utilization of organophosphates in *Escherichia coli* K-12. *Appl. Environ. Microbiol.* **64**:2601–2608.
  7. Fijalkowska, I. J., R. L. Dunn, and R. M. Schaaaper. 1993. Mutants of *Escherichia coli* with increased fidelity of DNA replication. *Genetics* **134**:1023–1030.
  8. Fijalkowska, I. J., P. Joneczyk, M. M. Tkaczyk, M. Biaolskorska, and R. M. Schaaaper. 1998. Unequal fidelity of leading strand and lagging strand DNA replication on the *Escherichia coli* chromosome. *Proc. Natl. Acad. Sci. USA* **95**:10020–10025.
  9. Hayashi, T., K. Makino, M. Ohnishi, K. Kurokawa, K. Ishii, K. Yokoyama, C.-G. Han, E. Ohtsubo, K. Nakayama, T. Murata, M. Tanaka, T. Tobe, T. Iida, H. Takami, T. Honda, C. Sasakawa, N. Ogasawara, T. Yasunaga, S. Kuhara, T. Shiba, M. Hattori, and H. Shinagawa. 2001. Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* **8**:11–22.
  10. Henderson, I. R., P. Owen, and J. P. Nataro. 1999. Molecular switches—the ON and OFF of bacterial phase variation. **33**:919–932.
  11. Hiasa, H., H. Sakai, K. Tanaka, Y. Honda, T. Komano, and G. N. Godson. 1989. Mutational analysis of the primer RNA template region in the replication origin (*oriC*) of bacteriophage G4: priming signal recognition by *Escherichia coli* primase. *Gene* **84**:9–16.
  12. Iyer, R. R., A. Pluciennik, W. A. Rosche, R. R. Sinden, and R. D. Wells. 2000. DNA polymerase III proofreading mutants enhance the expansion and deletion of triplet repeat sequences in *Escherichia coli*. *J. Biol. Chem.* **275**:2174–2284.
  13. Lovett, S. T., and V. V. Feschenko. 1996. Stabilization of diverged tandem repeats by mismatch repair: evidence for deletion formation via a misaligned replication intermediate. *Proc. Natl. Acad. Sci. USA* **93**:7120–7124.
  14. Makino, K., S.-K. Kim, H. Shinagawa, M. Amemura, and A. Nakata. 1991. Molecular analysis of the cryptic and functional *phn* operons for phosphonate use in *Escherichia coli* K-12. *J. Bacteriol.* **173**:2665–2672.
  15. Neidhardt, F. C., P. L. Bloch, and D. F. Smith. 1974. Culture media for enterobacteria. *J. Bacteriol.* **119**:736–747.
  16. Perna, N. T., G. Plunkett III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grothbeck, N. W. Davis, A. Lim, E. Dimalanta, K. Potamouisis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner. 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
  17. Saveson, C. J., and S. T. Lovett. 1997. Enhanced deletion formation by aberrant DNA replication in *Escherichia coli*. *Genetics* **146**:457–470.
  18. Swart, J. R., and M. A. Griep. 1993. Primase from *Escherichia coli* primes single-stranded templates in the absence of single-stranded DNA-binding protein or other auxiliary proteins. Template sequence requirements based on the bacteriophage G4 complementary strand origin and Okazaki fragment initiation sites. *J. Biol. Chem.* **268**:12970–12976.
  19. Tabor, S., and C. C. Richardson. 1985. A bacteriophage-T7 RNA polymerase promoter system for controlled exclusive expression of specific genes. *Proc. Natl. Acad. Sci. USA* **82**:1074–1098.
  20. Theiss, P., and K. S. Wise. 1997. Localized frameshift mutation generates selective, high-frequency phase variation of a surface lipoprotein encoded by a mycoplasma ABC transporter operon. *J. Bacteriol.* **179**:4013–4022.
  21. Wanner, B. L. 1994. Molecular genetics of carbon-phosphorous bond cleavage in bacteria. *Biodegradation* **5**:175–184.
  22. Welch R. A., V. Burland, G. Plunkett III, P. Redford, P. Roesch, D. Rasko, E. L. Buckles, S. R. Liou, A. Boutin, J. Hackett, D. Stroud, G. F. Mayhew, D. J. Rose, S. Zhou, D. C. Schwartz, N. T. Perna, H. L. Mobley, M. S. Sonnenberg, and F. R. Blattner. 2002. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **99**:17020–17024.
  23. Yoda, K., H. Yasuda, X. W. Xiang, and T. Okazaki. 1988. RNA-primed initiation sites of DNA replication in the origin region of bacteriophage-lambda genome. *Nucleic Acids. Res.* **16**:6531–6546.
  24. Yoda, K., and T. Okazaki. 1991. Specificity of recognition sequence for *Escherichia coli* primase. *Mol. Gen. Genet.* **227**:1–8.