

Molecular Evolution of Typical Enteropathogenic *Escherichia coli*: Clonal Analysis by Multilocus Sequence Typing and Virulence Gene Allelic Profiling^{∇†}

David W. Lacher,¹ Hans Steinsland,^{1,2} T. Eric Blank,³
Michael S. Donnenberg,³ and Thomas S. Whittam^{1*}

Microbial Evolution Laboratory, National Food Safety & Toxicology Center, Michigan State University, East Lansing, Michigan¹; Centre for International Health, University of Bergen, Bergen, Norway²; and Division of Infectious Diseases, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland 21201³

Received 18 September 2006/Accepted 30 October 2006

Enteropathogenic *Escherichia coli* (EPEC) infections are a leading cause of infantile diarrhea in developing nations. Typical EPEC isolates are differentiated from other types of pathogenic *E. coli* by two distinctive phenotypes, attaching effacement and localized adherence. The genes specifying these phenotypes are found on the locus of enterocyte effacement (LEE) and the EPEC adherence factor (EAF) plasmid. To describe how typical EPEC has evolved, we characterized a diverse collection of strains by multilocus sequence typing (MLST) and performed restriction fragment length polymorphism (RFLP) analysis of three virulence genes (*eae*, *bfpA*, and *perA*) to assess allelic variation. Among 129 strains representing 20 O-serogroups, 21 clonal genotypes were identified using MLST. RFLP analysis resolved nine *eae*, nine *bfpA*, and four *perA* alleles. Each *bfpA* allele was associated with only one *perA* allele class, suggesting that recombination has not played a large role in shuffling the *bfpA* and *perA* loci between separate EAF plasmids. The distribution of *eae* alleles among typical EPEC strains is more concordant with the clonal relationships than the distribution of the EAF plasmid types. These results provide further support for the hypothesis that the EPEC pathotype has evolved multiple times within *E. coli* through separate acquisitions of the LEE island and EAF plasmid.

Enteropathogenic *E. coli* (EPEC) infections are a leading cause of infantile diarrhea in developing nations (31, 37). A key characteristic of EPEC strains is the ability to intimately attach to intestinal epithelial cells and create attaching and effacing (AE) lesions (24). The AE phenotype is specified by genes of the locus of enterocyte effacement (LEE), a ~35-kb pathogenicity island located in the bacterial chromosome (23, 41). The LEE island comprises approximately 40 genes and encodes the components of a type III secretion system, various effector molecules, and the intimin adhesin (23, 33, 68). Intimin plays a crucial role in AE lesion formation (15) and is encoded by the highly polymorphic *eae* gene (1, 72), which can be divided into periplasmic, transmembrane, and extracellular domains (39). To date, more than 25 major allelic variants of *eae* have been described (36).

Most typical EPEC strains fall into one of two phylogenetically distinct groups or clonal lineages, designated EPEC 1 and EPEC 2 (69), and differ from atypical EPEC and other types of pathogenic *E. coli* by their ability to form microcolonies on the surface of intestinal epithelial cells (4). This phenotype, termed localized adherence (LA), correlates with the presence of a large virulence plasmid called the EPEC adherence factor (EAF) plasmid (18). The EAF plasmids from different EPEC

strains show considerable variation in size (~85 to 110 kb) (48) and, presumably, gene content. Comparison of the complete EAF plasmid sequences from two prototypical EPEC strains (O127:H6 EPEC 1 strain E2348/69 and O111:NM EPEC 2 strain B171) indicates that the EPEC 2 plasmid of B171 carries fewer genes (80 versus 115 open reading frames) and a greater percentage of intact or partial insertion sequence elements (33% versus 19%) than the pMAR7 plasmid of EPEC 1 strain E2348/69 (11, 64). Nevertheless, certain parts of the plasmid show a high degree of sequence conservation among typical EPEC strains (47), particularly in the region encoding the bundle-forming pilus (BFP), a type IV fimbria whose production is associated with the LA phenotype. An operon of 14 genes is necessary for expression of the BFP (20, 63), with *bfpA* encoding the major structural subunit (bundlin). Sequence comparisons of nine *bfpA* alleles have provided compelling evidence for the action of diversifying selection at the molecular level (7, 8). A second locus on the EAF plasmid implicated in the full virulence of EPEC is the plasmid-encoded regulator (Per), consisting of three genes (*perA*, *perB*, and *perC*). Per has been shown to activate genes within the *bfp* operon (65) and the LEE pathogenicity island (22, 42).

Little is known about the allelic distributions of *eae*, *bfpA*, and *perA* among the EPEC 1, EPEC 2, and other clonal lineages of typical EPEC. In this study we characterized a diverse collection of 129 EPEC, including strains of the classical EPEC serotypes (52), through multilocus sequence typing (MLST) and restriction fragment polymorphism (RFLP) analysis to elucidate the extent to which horizontal transfer of the LEE island and EAF plasmid have contributed to the evolution and diversification of EPEC clones.

* Corresponding author. Mailing address: National Food Safety & Toxicology Center, 165 Food Safety & Toxicology Building, Michigan State University, East Lansing, MI 48824. Phone: (517) 432-3100, ext. 178. Fax: (517) 432-2310. E-mail: whittam@msu.edu.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

[∇] Published ahead of print on 10 November 2006.

TABLE 1. Summary of EPEC strains investigated

Serogroup	No. of isolates	Yr(s) of isolation	Locale(s)	Reference(s)
Classical serogroups				
O55	21	1947–1998	Brazil, Congo, Dutch Guiana, France, Germany, Guinea-Bissau, Mexico, Scotland, United States	8, 44, 47, 52, 56, 57, 67
O86	10	1950–1997	Brazil, Bulgaria, Denmark, Germany, Guinea-Bissau, United Kingdom, United States	8, 9, 34, 52, 56, 67
O111	16	1947–1996	Austria, Brazil, Dutch Guiana, Germany, Mexico, Peru, Scotland, United Kingdom, United States	8, 17, 38, 44, 45, 47, 52, 53, 56
O114	4	1969–1997	Guinea-Bissau, United Kingdom, United States	16, 67
O119	28	1960–1998	Brazil, Chile, Guinea-Bissau, Mexico, Peru, United Kingdom, United States	8, 16, 28, 38, 46, 47, 58, 61, 67
O127	4	1969–1997	Guinea-Bissau, United Kingdom	16, 67
O142	11	<1960–1997	Brazil, Canada, Guinea-Bissau, Indonesia, Peru, Portugal, Scotland, United States	21, 32, 45, 56, 60, 67
Other serogroups				
O2	1	1997	Guinea-Bissau	67
O33	2	1997	Guinea-Bissau	67
O34	1	1997	Guinea-Bissau	67
O49	2	1997	Guinea-Bissau	67
O51	1	1997	Guinea-Bissau	67
O73	1	1997	Guinea-Bissau	67
O76	1	1982	Peru	45
O110	1	1993	Germany	59
O126	3	1962–1964	Egypt, Iran, Pakistan	52
O128	15	1953–1991	Denmark, Germany, Pakistan, United Kingdom, United States	9, 52, 55, 56
O157	3	1983–1998	Brazil, United States	7, 70
OX9	1	1997	Guinea-Bissau	67
O-	3	1996–1997	Guinea-Bissau	67

MATERIALS AND METHODS

Strains. A collection of 95 EPEC strains was assembled based on serotype or their inclusion in one of two studies examining *bfpA* allelic variation (7, 8). These strains represent a variety of serotypes originally isolated between 1947 and 1998 from different regions around the world and were obtained from a number of sources, including the Centers for Disease Control and Prevention, Alejandro Cravioto, Helge Karch, Frits and Ida Ørskov, Phillip I. Tarr, and Luis Trabulsi (Table 1; see also Table S1 in the supplemental material). An additional 34 *eae*⁺ *bfpA*⁺ strains were selected from a cohort study of childhood diarrheal disease in Guinea-Bissau, West Africa (67) (Table 1; see also Table S1). Each strain was grown overnight at 37°C in 10 ml of Luria-Bertani broth with moderate shaking. Genomic DNA was isolated using the Puregene DNA isolation kit (Gentra Systems Inc., Minneapolis, MN). DNA concentrations were determined using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies Inc., Rockland, ME), which were diluted to 25 ng/μl for PCR.

MLST. MLST was performed on seven conserved housekeeping genes (*aspC*, *clpX*, *fadD*, *icdA*, *lysP*, *mdhA*, and *uidA*). A detailed protocol of the MLST procedure, including allelic type and sequence type (ST) assignment methods, can be found at the EcMLST website (<http://www.shigatox.net/mlst>). Sequences were concatenated for phylogenetic analyses.

***eae* fRFLP.** Allelic variation in *eae* was resolved using fluorescent RFLP (fRFLP) as described previously (36).

***bfpA* PCR and DNA sequencing.** PCR primers were designed to target conserved flanking regions of *bfpA* based on nine available allelic sequences (7, 8). Each 25-μl reaction mixture contained 2.5 μl 10× buffer II (Applied Biosystems, Foster City, CA), 2.5 μl 2 mM deoxynucleoside triphosphate, 2.0 μl 25 mM MgCl₂, 0.5 μl 10 μM *bfpA*_{-52F} primer (5'-AGATTATTCCGTGACCTATT-3'), 0.5 μl 10 μM *bfpG*_{-9R} primer (5'-TGCTCTCACATATACCTCCC-3'), 1.5 U AmpliTaq Gold (Applied Biosystems), 1 μl of 25-ng/μl genomic DNA template, and 15.7 μl distilled H₂O (dH₂O). Amplification of the approximately 700-bp fragment utilized an initial denaturing step at 94°C for 10 min, followed by 35 cycles of 92°C for 1 min, 52°C for 1 min, and 72°C for 30 s. A final step of 72°C for 5 min was used to complete any partially extended product. PCR products (5 μl) were visualized on ethidium bromide-stained 1.5% agarose gels by illumination with UV light, purified using the QIAquick PCR purification kit (QIAGEN Inc., Valencia, CA), and quantified. Cycle sequencing reaction mixtures contained 4.0 μl CEQ DTCS Quick Start premix (Beckman Coulter Inc., Fullerton,

CA), 1.0 μl 20 μM *bfpA*_{-52F} or *bfpG*_{-9R} primer, approximately 70 ng of *bfpA* PCR product, and dH₂O to a final volume of 10 μl. Amplification utilized an initial denaturing step at 94°C for 1 min, followed by 35 cycles of 96°C for 30 s, 52°C for 30 s, and 60°C for 2 min. Upon completion of cycle sequencing, samples were purified with Sephadex G-50 fine columns (Amersham Pharmacia Biotech Inc., Piscataway, NJ), dried under vacuum centrifugation (Savant Instruments Inc., Holbrook, NY), suspended in 40 μl of deionized formamide, and run on a CEQ2000XL instrument (Beckman Coulter Inc.). Samples were analyzed using the CEQ2000XL software and then exported for further analysis with the SeqMan module of the Lasergene software (DNASTAR Inc., Madison, WI).

***bfpA* PCR and RFLP.** PCR conditions were identical to those described above for *bfpA* except primers *bfpA*_{-114F} (5'-GTCTGCGTCTGATTCCAATA-3') and *bfpA*_{-521R} (5'-TCAGCAGGAGTAATAGC-3') were used to amplify a 408- to 414-bp internal fragment of the gene. Prior to digestion, each *bfpA* PCR product was purified using the QIAquick PCR purification kit. Three different restriction enzyme digests were used. Digestion with AluI and with BfaI was performed in separate 30-μl reaction mixtures with 10 U of enzyme, 3.0 μl 10× reaction buffer, and 26.0 μl purified PCR product, and samples were incubated overnight at 37°C, while digestion with 10 U of BstNI was performed in 30-μl reaction mixtures with 3.0 μl 10× reaction buffer, 0.3 μl 100× bovine serum albumin, and 25.7 μl purified PCR product followed by an overnight incubation at 60°C. All restriction enzymes were obtained from New England BioLabs Inc. (Ipswich, MA), and the reaction buffer provided with each enzyme was used. After incubation, 15 μl of the digests was visualized on ethidium bromide-stained 1.5% agarose gels by illumination with UV light.

***perA* PCR and DNA sequencing.** Primers were designed to target the conserved flanking and internal regions of *perA* based on 15 available sequences (27, 50, 65). PCR conditions are similar to those described above for *bfpA* except primers *perA*_{-24F} (5'-AACAAACGCGCATGAAGGTG-3') and *perB*_{-222R} (5'-TTCGCCGGTGATGTGGTCT-3') were used with a 58°C annealing temperature and a 1-min extension time. The resulting PCR products (approximately 1.1 kb) were purified and quantified as described above. Cycle sequencing reaction mixtures were similar to those used for *bfpA*, except that 120 ng of *perA* PCR product and primers *perA*_{-24F}, *perA*_{-539F} (5'-AAAACCTGGAACTA GGCGATGTCA-3'), *perA*_{-562R} (5'-TGACATCGCCTAGTTTCCAGTTTT-3'), and *perB*_{-222R} were used at a 58°C annealing temperature.

TABLE 2. Sequence variation among alleles of seven MLST genes, *eae*, *bfpA*, and *perA*

Locus ^a	No. of sites	No. of variable sites	No. of alleles	% GC	$d_S \times 100$ (mean \pm SE)	$d_N \times 100$ (mean \pm SE)	d_N/d_S (95% CI)
<i>aspC</i>	513	21	9	51.4	5.14 \pm 1.23	0.06 \pm 0.06	0.035 (0.002, 0.154)
<i>clpX</i>	567	34	11	53.4	5.51 \pm 1.06	0.05 \pm 0.05	0.014 (0.001, 0.066)
<i>fadD</i>	483	32	11	50.7	8.56 \pm 1.50	0.05 \pm 0.05	0.008 (0.000, 0.036)
<i>icdA</i>	567	30	8	51.3	6.44 \pm 1.25	0.20 \pm 0.15	0.037 (0.006, 0.120)
<i>lysP</i>	477	10	8	53.9	3.44 \pm 1.14	0.00 \pm 0.00	0.000 (0.000, 0.061)
<i>mdh</i>	549	25	12	52.6	3.92 \pm 0.84	0.04 \pm 0.04	0.015 (0.001, 0.065)
<i>uidA</i>	576	26	7	51.9	4.84 \pm 1.10	0.63 \pm 0.25	0.171 (0.080, 0.326)
Avg	533.1	25.4	9.4	52.2	5.41 \pm 1.16	0.15 \pm 0.09	0.040 (0.013, 0.118)
<i>eae</i>	2,853	790	9	42.3	25.22 \pm 1.25	10.89 \pm 0.74	0.310 (0.298, 0.349)
<i>eae</i> _{PPD}	558	67	9	40.7	9.39 \pm 1.47	2.74 \pm 0.55	0.245 (0.171, 0.358)
<i>eae</i> _{TMD}	1,092	84	8 ^b	44.4	8.48 \pm 1.15	0.88 \pm 0.21	0.132 (0.086, 0.204)
<i>eae</i> _{ECD}	1,203	639	9	41.1	59.69 \pm 3.13	27.02 \pm 1.84	0.284 (0.275, 0.343)
<i>bfpA</i>	588	94	11	40.0	9.36 \pm 1.46	6.46 \pm 1.11	0.480 (0.384, 0.608)
<i>perA</i>	825	99	20	28.0	4.91 \pm 0.80	2.03 \pm 0.33	0.400 (0.322, 0.522)

^a Average values for seven MLST loci. PPD, periplasmic domain; TMD, transmembrane domain; ECD, extracellular domain.

^b The κ and μ *eae* alleles analyzed have identical transmembrane domain sequences.

***perA* PCR and RFLP.** PCR conditions were similar to those described for *bfpA* except primers *perA*_{-24F} and *perA*_{562R} were used at an annealing temperature of 58°C. Digestion with DdeI and with Sau96I was performed in separate 30- μ l reaction mixtures with 10 U of enzyme, 3.0 μ l 10 \times reaction buffer, 20.0 μ l unpurified PCR product, and dH₂O to volume; samples were incubated overnight at 37°C and visualized on ethidium bromide-stained 1.5% agarose gels.

***fliC* typing.** Strains that were nonmotile or lacked flagellar serotype data were typed for the *fliC* locus. The entire *fliC* gene was amplified with primers *fliC*_{1F} (5'-ATGGCACAAGTCATTAATACCAA-3') and *fliC*_{1497R} (5'-TTAACCCTGCAGCAGAGACA-3') using the same PCR conditions described for *bfpA* except for an annealing temperature of 55°C and an extension time of 2 min. Amplicons (approximately 2 kb) were either digested with 5 U of DdeI under conditions similar to those described for *perA* or sequenced to determine the allele. H-types that were determined by *fliC* sequencing or RFLP are denoted with a lower case h and are enclosed in square brackets.

Phylogenetic analyses. Sequences were aligned with the ClustalW algorithm using the MegAlign module of the Lasergene software. Neighbor-joining trees were constructed using the Kimura two-parameter model of nucleotide substitution with the MEGA3 software (35), and the inferred phylogenies were each tested with 500 bootstrap replications. Phylogenetic network analysis was conducted with the SplitsTree 4 program (30) using the neighbor-net algorithm (14) and untransformed distances (p distance). The Φ_{st} recombination test (13) as implemented by SplitsTree 4 was used to distinguish recurrent mutation from recombination in generating genotypic diversity. The number of synonymous substitutions per synonymous site (d_S) and the number of nonsynonymous substitutions per nonsynonymous site (d_N) were estimated by the modified Nei-Gojobori method using MEGA3 (35). Allelic sequences were fit to a nucleotide substitution model using the Datamonkey website, and then either the single likelihood ancestor counting (SLAC) or random effects likelihood (REL) method was used to fit a codon model to detect selection on individual codons (54). The SLAC method was also used to calculate the ratio of d_N to d_S and estimate the 95% confidence interval.

Nucleotide sequence accession numbers. The sequences for *bfpA* and *perA* were submitted to GenBank and given the accession numbers EF011027 to EF011059.

RESULTS

MLST analysis. PCR amplification and sequencing of the seven MLST loci in 129 EPEC strains was successful in most (90%) cases. The notable exception was *uidA*, which failed to amplify in 13 strains, including a single O114:H2 strain (380/69) and 12 other strains, 11 of which are serotype O55:[h51]. These 12 *uidA*-negative strains were identical to each other at the six remaining MLST loci. PCR amplification with primers

located in the genes flanking the *uidA* locus produced a truncated amplicon, suggesting that these strains belong to a clonal genotype that has lost most of the *uidA* gene (data not shown). For phylogenetic analysis, the sequenced internal fragments of the seven housekeeping genes were concatenated to yield 3,732 nucleotides. The *uidA* locus was treated as missing data and replaced with alignment gaps in the full concatenated sequence for the 13 *uidA*-negative strains.

MLST analysis resolved an average of 25.4 variable nucleotide sites per locus, which defined a number of alleles, ranging from 7 to 12, at the seven MLST genes (Table 2). The synonymous rate of substitution (d_S) ranged from a low of 3.44% for *lysP* to a high of 8.56% for *fadD*, with an average of 5.41 synonymous substitutions per 100 synonymous sites (Table 2). The nonsynonymous rate (d_N) per 100 nonsynonymous sites was generally 2 orders of magnitude lower than d_S , ranging from 0.00 for *lysP* to 0.63 for *uidA*. Tests for natural selection operating on the allelic variation at each MLST locus based on the SLAC method found no individual sites (codons) that were under significant positive selection and only six codons under negative selection at the 0.1 significance level (one in *clpX* and *mdh* and two in *fadD* and *icdA*). Thus, low values of d_N/d_S at the MLST loci reflect weak negative selection over many codons.

The distinct combinations of alleles across the seven MLST loci were used to define 21 multilocus genotypes or sequence types (STs) among the 129 EPEC strains. Classification of the strains based on the bootstrap analysis indicates that most (77%) of the strains belong to one of four main clonal groups, designated EPEC 1, EPEC 2, EPEC 3, and EPEC 4 (Fig. 1). With the exception of EPEC 3 strains, which were all O86:H34 (or nonmotile relatives), the EPEC groups based on the classification of STs included strains of various O-types (see complete list of serotypes in Table S1 in the supplemental material). There were strains representing three O-types in EPEC 1 (O55, O127, and O142), five O-types in EPEC 2 (O111, O114, O119, O126, and O128), and two O-types in EPEC 4 (O110 and O119). H-types (or the inferred H-type from the *fliC*




FIG. 1. Phylogenetic relationships of 21 EPEC sequence types. (A) An unrooted phylogenetic tree was constructed by the neighbor-joining algorithm based on the Kimura two-parameter model of nucleotide substitution. (B) The phylogenetic (splits) network is based on the neighbor-net algorithm using a *p* distance matrix. The four main clonal groups are indicated by colored boxes or ellipses. The ST and number of isolates are given at the branch tips. Bootstrap values greater than 50% based on 500 replications are given at the internal nodes. The distributions of *eae* alleles and EAF plasmid types are shown on the right (see Table 5, below, for plasmid type definitions).

allele) were conserved among strains of each group: EPEC 1 strains were H6, EPEC 2 were H2, EPEC 3 were H34, and EPEC 4 were H6. These four clonal groups were represented among both the worldwide and Guinea-Bissau strains. The 21 STs differed on average at 1.4% and 0.2% of the nucleotide and amino acid sites, respectively. ST-20 was the most common multilocus genotype (22.5%), followed by ST-13 (19.4%) and ST-1 (14.0%) (Fig. 1).

The splits network (Fig. 1B) revealed several parallel paths indicative of the presence of phylogenetic incompatibilities in the divergence of EPEC clones. Such incompatibilities could arise from recurrent mutation or recombination in the MLST loci. To detect recombination, we used the ϕ_w test, which has been shown to discriminate between recurrent mutation and recombination in a variety of circumstances (13). In application to the concatenated sequences of the 21 STs, there were 129 informative sites, and the ϕ_w test found statistically significant evidence of recombination ($P < 0.001$). The four main clonal groups, however, are separated and intact. Three of the four groups occur at the end of long branches without evidence of multiple paths, suggesting that recombination occurred early in the divergence of EPEC genotypes. With an EPEC phylogenetic framework in place, the allelic distributions of *eae*, *bfpA*, and *perA* were assessed.

Allelic variation in *eae*. The *eae* locus was subtyped by fRFLP (36), and nine alleles (α , β , $\beta 2$, ϵ , η , ι , κ , λ , and μ) were observed among the 129 EPEC strains (Fig. 1). Representative sequences of each fRFLP profile (36) were used for subsequent analyses. In comparison to the MLST loci, the synonymous rate for *eae* was 25.22%, more than 4 times greater than the mean for the seven MLST genes. The nonsynonymous rate

of 10.89% for *eae* was more than 72 times greater the average d_N across all seven MLST genes. When *eae* is divided into its three primary domains, the periplasmic and transmembrane domains have considerably lower synonymous and nonsynonymous rates than the extracellular region (Table 2). However, only a single codon within the extracellular domains of *eae* was found to be under significant positive selection using the SLAC method.

Allelic variation in *bfpA*. The entire *bfpA* gene was amplified and sequenced in 15 strains with STs for which *bfpA* allelic data were not previously available (STs 4, 5, 7, 8, 11, 15, 16, 17, and 18). Comparative sequence analysis revealed the existence of a 10th allele of *bfpA*, which we designated $\beta 7.1$ (Fig. 2). A minor variant of this allele that differed by a single synonymous substitution was designated $\beta 7.2$.

In comparison to the MLST loci, the synonymous rate for *bfpA* was 9.36%, slightly greater than the range and mean for the seven MLST genes. The nonsynonymous rate of 6.46% for *bfpA* was more than 43 times greater the average d_N across all seven MLST genes. There was no evidence for significant positive selection at individual codons within *bfpA* using the SLAC method; however, the less conservative REL method (54) detected 22 codon positions with positive selection, 12 of which were polymorphic for three or more amino acids.

Using the identified *bfpA* sequences, we devised an RFLP-based typing system to subtype *bfpA* alleles based on new PCR primers designed to target the conserved internal regions of the gene. We identified three restriction enzymes (AluI, BfaI, and BstNI) which, when used separately, produced digestion patterns that combined could resolve nine *bfpA* alleles (Table




FIG. 2. Eleven *bfpA* alleles cluster into two major groups. A phylogenetic tree constructed by the neighbor-joining algorithm based on the Kimura two-parameter model of nucleotide substitution is shown on the left. Bootstrap values based on 500 replications are given at the internal nodes. To the right is a graph of the locations of the 39 polymorphic amino acid sites (195 total), which are marked as vertical lines that indicate differences from the consensus of all 11 alleles.

3). In silico analysis with over 500 restriction endonucleases failed to identify an enzyme that could easily distinguish the $\beta 1$ and $\beta 7$ alleles. However, $\beta 1$ and $\beta 7$ *bfpA* strains can be easily differentiated based on their *perA* allele (see below). PCR amplification of *bfpA* was successful in all but 21 isolates. Of the *bfpA*-negative strains, O128:H2 was the most common serotype, with 13 isolates. RFLP analysis of the 108 *bfpA*-positive strains showed that the $\alpha 1$ ($n = 23$) and $\alpha 3$ ($n = 24$) alleles were the most common. The $\alpha 2$ ($n = 15$) and $\alpha 5$ ($n = 17$) alleles were also frequently identified, but the $\beta 2$ ($n = 1$), $\beta 3$ ($n = 4$), $\beta 4$ ($n = 7$), and $\beta 6$ ($n = 2$) alleles were rarely observed. Fourteen strains were classified as $\beta 1/\beta 7$ by RFLP, and DNA sequencing confirmed 11 as $\beta 1$ and 3 as $\beta 7$.

Allelic variation in *perA*. The entire *perA* gene was amplified and sequenced in 33 strains representing a diverse set of EPEC STs and *bfpA* alleles. One strain, 2309-77 (O111:H2), resulted in a PCR product approximately 1 kb larger than expected. DNA sequencing revealed the presence of a 1,055-bp IS element inserted into *perA* at position 414 with significant similarity to IS102 (86%) and IS903 (84%). Sequence analysis also identified eight strains that contained one or more frameshifts within mononucleotide repeats in *perA* that presumably inac-




FIG. 3. Twenty *perA* alleles cluster into four major groups. A phylogenetic tree constructed by the neighbor-joining algorithm based on the Kimura two-parameter model of nucleotide substitution is shown on the left. Bootstrap values for the major groups based on 500 replications are given at the internal nodes. To the right is a graph of the locations of the 46 polymorphic amino acid sites (274 total), which are marked as vertical lines that indicate differences from the consensus of all 20 alleles.

tivate the gene (see Table S2 in the supplemental material). The variability of these frameshifts among closely related alleles indicates their relatively recent occurrence, as there has not been sufficient time for the inactivated alleles to accumulate further mutations. These frameshifts were corrected, and the IS element sequence was excised in silico prior to allele assignment and phylogenetic analyses. The 33 sequences yielded 20 alleles, which clustered into four groups based on phylogenetic sequence analysis and, in keeping with the nomenclature for *eae* and *bfpA*, we designated these four allele classes α , β , γ , and δ (Fig. 3). As with *bfpA*, each distinct translated *perA* sequence was given an allele designation, resulting in 11 major α types, 3 β , 2 γ , and a single δ . Two of the α alleles had variants resulting from synonymous substitutions, and each variant was given its own subtype designation ($\alpha 1.1$, $\alpha 1.2$, $\alpha 1.3$, $\alpha 5.1$, and $\alpha 5.2$).

In comparison to the MLST loci, the synonymous rate for *perA* was 4.91%, within the range and slightly less than the mean for the seven MLST genes. The nonsynonymous rate of 2.03% for *perA* was more than 13 times greater the average d_N across all seven MLST genes. However, only six codons within *perA* were found to be under significant negative selection, and there was no evidence for significant positive selection at individual codons using the SLAC method.

Based on the sequence data, an RFLP method using DdeI and Sau96I was designed to detect the four *perA* allele classes (Table 4). PCR amplification of *perA* was successful in all but 17 isolates. As with the *bfpA*-negative strains, O128:H2 was the most common serotype among the *perA*-negative isolates ($n = 12$). RFLP analysis of the 112 *perA*-positive strains showed that

TABLE 3. Expected restriction fragment length polymorphisms of *bfpA* PCR amplicons

<i>bfpA</i> allele	Digestion pattern (bp) ^a		
	AluI	BfaI	BstNI
$\alpha 1$	408	408	155, 253
$\alpha 2$	408	75, 333	155, 253
$\alpha 3$	408	75, 333	408
$\beta 1$	16, 54, 113, 231	39, 375	414
$\beta 2$	16, 170, 228	414	414
$\beta 3$	16, 54, 344	39, 375	414
$\beta 4$	16, 177, 215	39, 369	408
$\beta 5$	16, 54, 123, 215	39, 369	408
$\beta 6$	16, 392	39, 369	408
$\beta 7$	16, 54, 113, 225	39, 369	408

^a Underlined fragments are not detectable under standard electrophoretic conditions.

TABLE 4. Expected restriction fragment length polymorphisms of *perA* PCR amplicons

<i>perA</i> allele class	Digestion pattern (bp) ^a	
	DdeI	Sau96I
α	13, 74, 82, 417	586
β	13, 74, 499	586
γ	82, 87, 417	174, 186, 226
δ	87, 499	226, 360

^a Underlined fragments are not detectable under standard electrophoretic conditions.

the α allele was the most common ($n = 74$), followed by β ($n = 29$), γ ($n = 8$), and δ ($n = 1$).

Association between EAF types, virulence factor alleles, and STs. By combining the *bfpA* and *perA* allelic data, we found that each *bfpA* allele was associated with only one *perA* allele class, resulting in 11 distinct EAF plasmid types, which we have designated as EAF types 1 to 11 (Table 5). EAF plasmid types 4 and 8 appear to be the most promiscuous, being found in five and four clonal groups, respectively. Interestingly, the EAF type represented by the fully sequenced plasmid from O111:NM EPEC 2 strain B171 (64) is among the least promiscuous, being found in only one serotype (O111:H2) of a single sequence type (ST-20).

The α1 and α2 alleles of *bfpA*, as well as β4 and β5, differ by only one nonsynonymous nucleotide substitution. These two sets of closely related *bfpA* alleles were found in divergent EPEC lineages: EPEC 1 contains α1 and β5, whereas EPEC 2 has α2 and β4 (Table 6). In addition, multiple *bfpA* alleles were found within the same sequence type: α3, β2, and β5 within ST-13, and α2, β1, and β4 within ST-20 (Table 6).

As shown previously (1), the EPEC 1 and EPEC 2 clonal groups possess the α and β alleles of *eae*, respectively. In contrast to the distribution of the EAF types, strains of the same ST had identical *eae* alleles as resolved by rFLP. The α-*eae* allele had the widest distribution among the EPEC clones, being found in strains with O51:[h49], O73:[h34], O142:[h34], and O157:[h45] serotypes in addition to the EPEC 1 group (O55:H6, O127:H6, and O142:H6). The rarest *eae* alleles were ε, η, ι, and λ and, combined, these alleles account for less than 5% of the strains examined.

TABLE 5. EAF plasmid types and distribution among EPEC clones

EAF type	<i>bfpA</i> allele	<i>perA</i> allele	No. of isolates	No. of STs	No. of clonal groups
1	α1	α	23	4	2
2	α2	α	15	1	1
3	α3	β	24	4	3
4	β1	α	11	6	5
5	β2	δ	1	1	1
6	β3	γ	4	1	1
7	β4	α	7	3	3
8	β5	α	18	4	4
9	β6	β	2	1	1
10	β7	β	3	2	2
11	Neg ^a	γ	4	1	1
Neg	Neg	Neg	17	4	3

^a Neg, negative by PCR.

TABLE 6. Characteristics of four common EPEC clonal groups

Clonal group and ST	<i>eae</i> allele	EAF type ^a	Serotype ^b	No. of isolates
EPEC 1 STs				
1	α	1	O55:H6	8
1	α	1	O127:[h6]	4
1	α	1	O142:[h6]	1
1	α	1	O-:[h6]	1
1	α	8	O142:H6	4
2	α	1	O55:H6	2
3	α	6	O142:H6	4
EPEC 2 STs				
19	β		O128:H2	11
20	β	2	O111:[h2]	15
20	β	4	O114:[h2]	4
20	β	4	O126:H2	1
20	β	7	O128:H2	2
20	β	Neg	O111:H2	1
20	β	Neg	O126:H2	1
20	β	Neg	O128:H2	1
20	β	11	O119:H2	3
20	β	11	O128:H2	1
21	β	4	O126:H2	1
EPEC 3 STs				
9	κ	1	O86:[h34]	6
9	κ	Neg	O86:[h34]	1
10	κ	1	O86:H34	1
EPEC 4 STs				
13	β2	3	O119:H6	20
13	β2	5	O110:H6	1
13	β2	8	O119:H6	2
13	β2	Neg	O119:H6	2
14	β2	3	O119:H6	1

^a Neg, negative for *bfpA* and *perA* by PCR.

^b Lowercase H-types in square brackets were inferred from the *flhC* allele.

DISCUSSION

Common EPEC clones. This is the first study to take a comprehensive look at the evolution of typical EPEC by combining clonal relatedness based on MLST with the allelic distributions of three important virulence factors. Previous clonal studies of EPEC have focused primarily on two main groups, EPEC 1 and EPEC 2, which were first described and defined based on the genetic relatedness of strains as determined by multilocus enzyme electrophoresis (69). EPEC 1 was described to comprise strains with O55:H6, O86:H34, O127:H6, and O142:H6 serotypes, while EPEC 2 included O111:H2, O114:H2, O126:H2, and O128:H2. O119:H6 strains have also been regarded as members of EPEC 1, because they share a number of genetic traits with the group (H6 flagellar antigen, *eae*⁺, and EAF⁺) (66), even though multilocus enzyme electrophoresis places them just outside of EPEC 1 (69). However, we feel that there are sufficient genetic differences to warrant the removal of O86:H34 and O119:H6 from EPEC 1, and we have reclassified them as EPEC 3 and EPEC 4, respectively (Fig. 1; Table 6). Our data indicate that EPEC 1 strains (O55:H6, O127:H6, and O142:H6) all possess α-*eae*, whereas the κ and β2 *eae* alleles are associated with EPEC 3 and EPEC 4, respectively. O86:H34 strains have also been shown to possess cytolethal distending toxin, while EAF⁺ O55:H6, O119:H6, O127:H6,

TABLE 7. EAF plasmid changes within four common EPEC clonal groups

Clonal group	Recombination (change in <i>bfpA</i>)	Plasmid replacement (change in <i>bfpA</i> and <i>perA</i>)	Plasmid loss
EPEC 1	EAF 1 to EAF 8	EAF 1 to EAF 6	Not observed
EPEC 2	EAF 4 to EAF 2 or EAF 7	EAF 4 to EAF 11	Observed
EPEC 3	Not observed	Not observed	Observed
EPEC 4	Not observed	EAF 3 to EAF 5 or EAF 8	Observed

and O142:H6 strains are negative (29). O119:H6 strains, on the other hand, are negative for *espC*, which encodes an enterotoxin, whereas EPEC 1 strains are positive (43).

Other EPEC clones. Thirty of the 129 strains (23%) that were analyzed did not belong to any of the above-mentioned major lineages, and most of these had unusual serotypes for EPEC. Of these serotypes, only O2:[h2], O49:[h10], and O51:[h49] have previously been reported to possess *eae* and/or express the AE phenotype (2, 5, 12), and only O33:[h34], O142:[h34], and O157:[h45] have been previously described as *eae*⁺ *bfpA*⁺ and therefore classified as typical EPEC (5, 25, 26, 40, 51, 62). Literature searches on the remaining serotypes, including O34:[h45], O73:[h34], O76:H51, O86:[h8], O142:[h21], OX9:[h7], O-:[h7], and O-:[h34], failed to find any association with typical EPEC.

An interesting finding of this study was the prevalence of the O55:[h51] clone among the strains isolated in Guinea-Bissau. Strains with this serotype have previously been described as relatively minor members of the O55 serogroup and have been reportedly isolated only in South America (6, 57). It is possible that O55:[h51] strains are simply common among children in Guinea-Bissau, or their higher prevalence may be due to sampling bias. Alternatively, the abundance of these strains among the West African isolates could indicate that O55:[h51] is an emerging clone which is increasing in frequency and spreading geographically, possibly because of a distinct combination of μ intimin and EAF type 8.

Virulence gene distribution. Recombination appears to have played a role in the initial generation of the EAF plasmid types. The highly divergent α 1/ α 2 and β 4/ β 5 *bfpA* alleles are all associated with α -*perA*, whereas the closely related β 1 and β 7 *bfpA* alleles are each found with different *perA* alleles. However, since *bfpA* and *perA* are in complete linkage disequilibrium (each *bfpA* allele is associated with only one *perA* allele class), it does not appear that recombination has played a large role in assorting allele combinations. Our results also indicate varying degrees of promiscuity among the different EAF types (Table 5). Recently, the complete sequence of a derivative of the wild-type EAF plasmid (pMAR7) from prototypical EPEC 1 strain E2348/69 was determined (11). In comparison to the EAF of O111:NM strain B171, the primary difference is the presence of the *tra* locus in pMAR7 (EPEC 1, EAF type 1) and its absence from pB171 (EPEC 2, EAF type 2). The *tra* genes, which are responsible for conjugal transfer in plasmids F and R100, were found to have varying degrees of conservation among the EAF plasmids from different EPEC strains (11). This finding has intriguing implications for our results. In addition to EAF type 2, plasmid types 5, 6, 9, and 11 were all found in only a single ST. Like pB171, these plasmids may lack the entire *tra* locus or possess defective conjugation machinery,

thereby preventing their transfer to other EPEC clones. In contrast to the distribution of the EAF plasmid types, the distribution of *eae* alleles among EPEC strains is more consistent with their clonal relationships. This suggests that the EAF plasmid is more mobile than the LEE island.

Inferred evolution of EAF plasmids and EPEC clones. Based on the MLST and EAF plasmid type data, ancestral or primitive clonal types within the four main EPEC groups can be inferred under the parsimony principle, that is, positing a simple evolution model based on minimizing the number of evolutionary genetic events. The principal events in the evolutionary change of an ancestral EPEC clone are EAF plasmid recombination, plasmid replacement, and plasmid loss. Here, we define recombination of the EAF plasmid as a change in *bfpA* allele, since each *bfpA* allele is associated with a single *perA* allele class. Plasmid replacement, presumably resulting from the horizontal transfer of an EAF plasmid, is believed to have occurred when both *bfpA* and *perA* differ from the primitive condition. Plasmid loss is inferred when an isolate is PCR negative for both *bfpA* and *perA*.

Under the parsimony principle, we have deduced the types of genetic events underlying the evolution of each of the EPEC groups (Table 7). The EPEC 3 group is the most homogenous, with only plasmid loss being inferred. Two of the three possible plasmid changes were found in the EPEC 1 and EPEC 4 clonal groups. For EPEC 1, the inferred ancestral type possessed EAF type 1 (α 1-*bfpA*, α -*perA*). Plasmid loss was not observed, but plasmid recombination (EAF type 8; β 5-*bfpA*, α -*perA*) and replacement (EAF type 6; β 3-*bfpA*, γ -*perA*) have occurred. For EPEC 4, plasmid recombination was not observed, but plasmid loss and replacement were detected. EAF type 3 (α 3-*bfpA*, β -*perA*) was replaced with EAF type 5 (β 2-*bfpA*, δ -*perA*) and with EAF type 8 (β 5-*bfpA*, α -*perA*). EPEC 2 is the most variable clonal group. With an inferred ancestral state of EAF type 4 (β 1-*bfpA*, α -*perA*), all three types of plasmid changes were observed. Two different recombination events involving EAF type 2 (α 2-*bfpA*, α -*perA*) and EAF type 7 (β 4-*bfpA*, α -*perA*), plasmid replacement with EAF type 11 (*bfpA* negative, γ -*perA*), and plasmid loss have shaped the diversity of this clonal group.

EAF type 11 is unusual in that it was the only plasmid type that did not contain *bfpA* according to our PCR screening. Bortolini and colleagues (10) reported a similar finding when they described EAF⁺ O119:H2 and O128:H2 strains in which most (~13 kb) of the *bfp* operon had been deleted and replaced with an IS66-like element. We suspected that since the deletion encompasses the 3' end of *bfpA*, this could explain why plasmid type 11 is *bfpA* negative. To confirm this, we designed new primers to target the 5' end of *bfpA* and the IS66-like element. All EAF type 11 strains yielded the ex-

pected amplicon, indicating that this plasmid type possesses a similar *bfp* operon structure as that described by Bortolini and colleagues (10). Aside from the EAF type 11 strains, 17 additional isolates were *bfpA* negative. These strains, however, were also *perA* negative, suggesting that they did not possess the EAF plasmid. Of the EAF-negative isolates, most (82%) were part of the EPEC 2 clonal group, with the O128:H2 serotype being the most common. This finding was not unexpected, as O128:H2 strains are often reported to be EAF negative and, therefore, classified as atypical EPEC (66).

Relationship between typical and atypical EPEC. Our work shows that at least some atypical EPEC strains, such as those that lost both *bfpA* and *perA* (and presumably the whole EAF plasmid), evolved from typical EPEC, rather than typical EPEC evolving from atypical EPEC by acquisition of the plasmid. It has been shown that typical EPEC can lose the EAF plasmid at a surprisingly high rate during passage through adult volunteers (19, 38), and so there appears to be selective pressure to lose the plasmid and convert from typical to atypical EPEC. This is interesting, given the recent reports of atypical EPEC in human clinical isolates, some of which belong to typical EPEC O-serogroups (3, 5, 49).

Concluding remarks. The evolution of EPEC appears to be a dynamic process involving repeated acquisition of the LEE island and transfer of the EAF plasmid. The work presented here is the first to classify EAF plasmids into types based on *bfpA* and *perA* allelic data, and we identified 11 distinct plasmid types among the EPEC strains investigated. Nevertheless, it remains unclear what level of conservation exists among plasmids of the same EAF type. Given the amount of IS elements present within the two fully sequenced EAF plasmids (11, 64), there could be considerable variation within each plasmid type, and further characterization of the EAF types is warranted. Another area of future research could focus on the insertion site of the LEE island. Although it is known that the EPEC 1 LEE is within the *selC* site, whereas EPEC 2 has LEE inserted into *pheU* (61, 71), the insertion sites of the LEE island in the other EPEC clonal types remain to be elucidated. The further characterization of pathogenic strains will improve our understanding of the processes that underlie microbial evolution. The identification of unique genetic determinants in these strains may then be used to facilitate the detection of specific epidemic clones during outbreaks of disease.

ACKNOWLEDGMENTS

We thank Lindsey Ouellette and Weihong Qi for technical assistance and Shannon Manning for comments on a previous version of the manuscript.

This project has been funded in part with federal funds from the NIAID, NIH, DHHS, under NIH research contract N01-AI-30058 (T.S.W.) and NIH research grant AI-37606 (M.S.D.).

REFERENCES

- Adu-Bobie, J., G. Frankel, C. Bain, A. G. Goncalves, L. R. Trabulsi, G. Douce, S. Knutton, and G. Dougan. 1998. Detection of intimins α , β , γ , and δ four intimin derivatives expressed by attaching and effacing microbial pathogens. *J. Clin. Microbiol.* **36**:662–668.
- Albert, M. J., K. Alam, M. Ansaruzzaman, J. Montanaro, M. Islam, S. M. Faruque, K. Haider, K. Bettelheim, and S. Tzipori. 1991. Localized adherence and attaching-effacing properties of nonenteropathogenic serotypes of *Escherichia coli*. *Infect. Immun.* **59**:1864–1868.
- Alikhani, M. Y., A. Mirsalehian, and M. M. Aslani. 2006. Detection of typical and atypical enteropathogenic *Escherichia coli* (EPEC) in Iranian children with and without diarrhoea. *J. Med. Microbiol.* **55**:1159–1163.
- Bieber, D., S. W. Ramer, C. Y. Wu, W. J. Murray, T. Tobe, R. Fernandez, and G. K. Schoolnik. 1998. Type IV pili, transient bacterial aggregates, and virulence of enteropathogenic *Escherichia coli*. *Science* **280**:2114–2118.
- Blanco, M., J. E. Blanco, G. Dahbi, M. P. Alonso, A. Mora, M. A. Coira, C. Madrid, A. Juarez, M. I. Bernardez, E. A. Gonzalez, and J. Blanco. 2006. Identification of two new intimin types in atypical enteropathogenic *Escherichia coli*. *Int. Microbiol.* **9**:103–110.
- Blanco, M., J. E. Blanco, G. Dahbi, A. Mora, M. P. Alonso, G. Varela, M. P. Gadea, F. Schelotto, E. A. Gonzalez, and J. Blanco. 2006. Typing of intimin (*eae*) genes from enteropathogenic *Escherichia coli* (EPEC) isolated from children with diarrhoea in Montevideo, Uruguay: identification of two novel intimin variants (μ B and ξ R/ β 2B). *J. Med. Microbiol.* **55**:1165–1174.
- Blank, T. E., D. W. Lacher, I. C. Scaletsky, H. Zhong, T. S. Whittam, and M. S. Sonnenberg. 2003. Enteropathogenic *Escherichia coli* O157 strains from Brazil. *Emerg. Infect. Dis.* **9**:113–115.
- Blank, T. E., H. Zhong, A. L. Bell, T. S. Whittam, and M. S. Sonnenberg. 2000. Molecular variation among type IV pilin (*bfpA*) genes from diverse enteropathogenic *Escherichia coli* strains. *Infect. Immun.* **68**:7028–7038.
- Bokete, T. N., T. S. Whittam, R. A. Wilson, C. R. Clausen, C. M. O'Callahan, S. L. Moseley, T. R. Fritsche, and P. I. Tarr. 1997. Genetic and phenotypic analysis of *Escherichia coli* with enteropathogenic characteristics isolated from Seattle children. *J. Infect. Dis.* **175**:1382–1389.
- Bortolini, M. R., L. R. Trabulsi, R. Keller, G. Frankel, and V. Sperandio. 1999. Lack of expression of bundle-forming pili in some clinical isolates of enteropathogenic *Escherichia coli* (EPEC) is due to a conserved large deletion in the *bfp* operon. *FEMS Microbiol. Lett.* **179**:169–174.
- Brinkley, C., V. Burland, R. Keller, D. J. Rose, A. T. Boutin, S. A. Klink, F. R. Blattner, and J. B. Kaper. 2006. Nucleotide sequence analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid pMAR7. *Infect. Immun.* **74**:5408–5413.
- Broes, A., R. Drolet, M. Jacques, J. M. Fairbrother, and W. M. Johnson. 1988. Natural infection with an attaching and effacing *Escherichia coli* in a diarrheic puppy. *Can. J. Vet. Res.* **52**:280–282.
- Bruen, T. C., H. Philippe, and D. Bryant. 2006. A simple and robust statistical test for detecting the presence of recombination. *Genetics* **172**:2665–2681.
- Bryant, D., and V. Moulton. 2004. Neighbor-net: an agglomerative method for the construction of phylogenetic networks. *Mol. Biol. Evol.* **21**:255–265.
- Cleary, J., L. C. Lai, R. K. Shaw, A. Straatman-Iwanowska, M. S. Sonnenberg, G. Frankel, and S. Knutton. 2004. Enteropathogenic *Escherichia coli* (EPEC) adhesion to intestinal epithelial cells: role of bundle-forming pili (BFP), EspA filaments and intimin. *Microbiol.* **150**:527–538.
- Cravioto, A., R. J. Gross, S. M. Scotland, and B. Rowe. 1979. An adhesive factor found in strains of *Escherichia coli* belonging to the traditional infantile enteropathogenic serotypes. *Curr. Microbiol.* **3**:95–99.
- Cravioto, A., A. Tello, A. Navarro, J. Ruiz, H. Villafan, F. Uribe, and C. Eslava. 1991. Association of *Escherichia coli* HEP-2 adherence patterns with type and duration of diarrhoea. *Lancet* **337**:262–264.
- Donnenberg, M. S., J. A. Giron, J. P. Nataro, and J. B. Kaper. 1992. A plasmid-encoded type IV fimbrial gene of enteropathogenic *Escherichia coli* associated with localized adherence. *Mol. Microbiol.* **6**:3427–3437.
- Donnenberg, M. S., C. O. Tacket, S. P. James, G. Losonsky, J. P. Nataro, S. S. Wasserman, J. B. Kaper, and M. M. Levine. 1993. Role of the *eaeA* gene in experimental enteropathogenic *Escherichia coli* infection. *J. Clin. Invest.* **92**:1412–1417.
- Donnenberg, M. S., H. Z. Zhang, and K. D. Stone. 1997. Biogenesis of the bundle-forming pilus of enteropathogenic *Escherichia coli*: reconstitution of fimbriae in recombinant *E. coli* and role of DsbA in pilin stability. *Gene* **192**:33–38.
- Drolet, R., J. M. Fairbrother, J. Harel, and P. Helie. 1994. Attaching and effacing and enterotoxigenic *Escherichia coli* associated with enteric colibacillosis in the dog. *Can. J. Vet. Res.* **58**:87–92.
- Elliott, S. J., V. Sperandio, J. A. Giron, S. Shin, J. L. Mellies, L. Wainwright, S. W. Hutcheson, T. K. McDaniel, and J. B. Kaper. 2000. The locus of enterocyte effacement (LEE)-encoded regulator controls expression of both LEE- and non-LEE-encoded virulence factors in enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **68**:6115–6126.
- Elliott, S. J., L. A. Wainwright, T. K. McDaniel, K. G. Jarvis, Y. K. Deng, L. C. Lai, B. P. McNamara, M. S. Sonnenberg, and J. B. Kaper. 1998. The complete sequence of the locus of enterocyte effacement (LEE) from enteropathogenic *Escherichia coli* E2348/69. *Mol. Microbiol.* **28**:1–4.
- Finlay, B. B., I. Rosenshine, M. S. Sonnenberg, and J. B. Kaper. 1992. Cytoskeletal composition of attaching and effacing lesions associated with enteropathogenic *Escherichia coli* adherence to HeLa cells. *Infect. Immun.* **60**:2541–2543.
- Franzolin, M. R., R. C. Alves, R. Keller, T. A. Gomes, L. Beutin, M. L. Barreto, C. Milroy, A. Strina, H. Ribeiro, and L. R. Trabulsi. 2005. Prevalence of diarrheagenic *Escherichia coli* in children with diarrhea in Salvador, Bahia, Brazil. *Mem. Inst. Oswaldo Cruz* **100**:359–363.
- Ghilardi, A. C., T. A. Gomes, W. P. Elias, and L. R. Trabulsi. 2003. Virulence factors of *Escherichia coli* strains belonging to serogroups O127 and O142. *Epidemiol. Infect.* **131**:815–821.

27. Gomez-Duarte, O. G., and J. B. Kaper. 1995. A plasmid-encoded regulatory region activates chromosomal *eaeA* expression in enteropathogenic *Escherichia coli*. *Infect. Immun.* **63**:1767–1776.
28. Goncalves, A. G., L. C. Campos, T. A. Gomes, J. Rodrigues, V. Sperandio, T. S. Whittam, and L. R. Trabulsi. 1997. Virulence properties and clonal structures of strains of *Escherichia coli* O119 serotypes. *Infect. Immun.* **65**:2034–2040.
29. Guth, B. E., R. Giraldo, T. A. Gomes, and L. R. Marques. 1994. Survey of cytotoxin production among *Escherichia coli* strains characterized as enteropathogenic (EPEC) by serotyping and presence of EPEC adherence factor (EAF) sequences. *Can. J. Microbiol.* **40**:341–344.
30. Huson, D. H., and D. Bryant. 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* **23**:254–267.
31. Kaper, J. B. 1994. Molecular pathogenesis of enteropathogenic *Escherichia coli*, p. 173–195. *In* V. L. Miller, J. B. Kaper, D. A. Portnoy, and R. R. Isberg (ed.), *Molecular genetics of bacterial pathogenesis*. American Society for Microbiology, Washington, D.C.
32. Kennedy, D. H., G. H. Walker, R. J. Fallon, J. F. Boyd, R. J. Gross, and B. Rowe. 1973. An outbreak of infantile gastroenteritis due to *E. coli* O142. *J. Clin. Pathol.* **26**:731–737.
33. Kenny, B. 2002. Mechanism of action of EPEC type III effector molecules. *Int. J. Med. Microbiol.* **291**:469–477.
34. Knutton, S., T. Baldwin, P. H. Williams, and A. S. McNeish. 1989. Actin accumulation at sites of bacterial adhesion to tissue culture cells: basis of a new diagnostic test for enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **57**:1290–1298.
35. Kumar, S., K. Tamura, and M. Nei. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* **5**:150–163.
36. Lacher, D. W., H. Steinsland, and T. S. Whittam. 2006. Allelic subtyping of the intimin locus (*eae*) of pathogenic *Escherichia coli* by fluorescent RFLP. *FEMS Microbiol. Lett.* **261**:80–87.
37. Levine, M. M., and R. Edelman. 1984. Enteropathogenic *Escherichia coli* of classic serotypes associated with infant diarrhea: epidemiology and pathogenesis. *Epidemiol. Rev.* **6**:31–51.
38. Levine, M. M., J. P. Nataro, H. Karch, M. M. Baldini, J. B. Kaper, R. E. Black, M. L. Clements, and A. D. O'Brien. 1985. The diarrheal response of humans to some classic serotypes of enteropathogenic *Escherichia coli* is dependent on a plasmid encoding an enteroadhesiveness factor. *J. Infect. Dis.* **152**:550–559.
39. Luo, Y., E. A. Frey, R. A. Pfuetzner, A. L. Creagh, D. G. Knoechel, C. A. Haynes, B. B. Finlay, and N. C. Strynadka. 2000. Crystal structure of enteropathogenic *Escherichia coli* intimin-receptor complex. *Nature* **405**:1073–1077.
40. Makino, S., H. Asakura, T. Shirahata, T. Ikeda, K. Takeshi, K. Arai, M. Nagasawa, T. Abe, and T. Sadamoto. 1999. Molecular epidemiological study of a mass outbreak caused by enteropathogenic *Escherichia coli* O157:H45. *Microbiol. Immunol.* **43**:381–384.
41. McDaniel, T. K., K. G. Jarvis, M. S. Sonnenberg, and J. B. Kaper. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. USA* **92**:1664–1668.
42. Mellies, J. L., S. J. Elliott, V. Sperandio, M. S. Sonnenberg, and J. B. Kaper. 1999. The Per regulon of enteropathogenic *Escherichia coli*: identification of a regulatory cascade and a novel transcriptional activator, the locus of enterocyte effacement (LEE)-encoded regulator (Ler). *Mol. Microbiol.* **33**:296–306.
43. Mellies, J. L., F. Navarro-Garcia, I. Okeke, J. Frederickson, J. P. Nataro, and J. B. Kaper. 2001. *espC* pathogenicity island of enteropathogenic *Escherichia coli* encodes an enterotoxin. *Infect. Immun.* **69**:315–324.
44. Moyenuddin, M., I. K. Wachsmuth, S. L. Moseley, C. A. Bopp, and P. A. Blake. 1989. Serotype, antimicrobial resistance, and adherence properties of *Escherichia coli* strains associated with outbreaks of diarrheal illness in children in the United States. *J. Clin. Microbiol.* **27**:2234–2239.
45. Nataro, J. P., M. M. Baldini, J. B. Kaper, R. E. Black, N. Bravo, and M. M. Levine. 1985. Detection of an adherence factor of enteropathogenic *Escherichia coli* with a DNA probe. *J. Infect. Dis.* **152**:560–565.
46. Nataro, J. P., J. B. Kaper, R. Robins-Browne, V. Prado, P. Vial, and M. M. Levine. 1987. Patterns of adherence of diarrheagenic *Escherichia coli* to HEp-2 cells. *Pediatr. Infect. Dis. J.* **6**:829–831.
47. Nataro, J. P., K. O. Maher, P. Mackie, and J. B. Kaper. 1987. Characterization of plasmids encoding the adherence factor of enteropathogenic *Escherichia coli*. *Infect. Immun.* **55**:2370–2377.
48. Nataro, J. P., I. C. Scaletsky, J. B. Kaper, M. M. Levine, and L. R. Trabulsi. 1985. Plasmid-mediated factors conferring diffuse and localized adherence of enteropathogenic *Escherichia coli*. *Infect. Immun.* **48**:378–383.
49. Nguyen, R. N., L. S. Taylor, M. Tauschek, and R. M. Robins-Browne. 2006. Atypical enteropathogenic *Escherichia coli* infection and prolonged diarrhea in children. *Emerg. Infect. Dis.* **12**:597–603.
50. Okeke, I. N., J. A. Borneman, S. Shin, J. L. Mellies, L. E. Quinn, and J. B. Kaper. 2001. Comparative sequence analysis of the plasmid-encoded regulator of enteropathogenic *Escherichia coli* strains. *Infect. Immun.* **69**:5553–5564.
51. Okeke, I. N., A. Lamikanra, H. Steinruck, and J. B. Kaper. 2000. Characterization of *Escherichia coli* strains from cases of childhood diarrhea in provincial southwestern Nigeria. *J. Clin. Microbiol.* **38**:7–12.
52. Ørskov, F., T. S. Whittam, A. Cravioto, and I. Ørskov. 1990. Clonal relationships among classic enteropathogenic *Escherichia coli* (EPEC) belong to different O groups. *J. Infect. Dis.* **162**:76–81.
53. Paulozzi, L. J., K. E. Johnson, L. M. Kamahle, C. R. Clausen, L. W. Riley, and S. D. Helgeson. 1986. Diarrhea associated with adherent enteropathogenic *Escherichia coli* in an infant and toddler center, Seattle, Washington. *Pediatrics* **77**:296–300.
54. Pond, S. L., and S. D. Frost. 2005. Datamonkey: rapid detection of selective pressure on individual sites of codon alignments. *Bioinformatics* **21**:2531–2533.
55. Reid, S. D., D. J. Betting, and T. S. Whittam. 1999. Molecular detection and identification of intimin alleles in pathogenic *Escherichia coli* by multiplex PCR. *J. Clin. Microbiol.* **37**:2719–2722.
56. Robins-Browne, R. M., W. C. Yam, L. E. O'Gorman, and K. A. Bettelheim. 1993. Examination of archetypal strains of enteropathogenic *Escherichia coli* for properties associated with bacterial virulence. *J. Med. Microbiol.* **38**:222–226.
57. Rodrigues, J., I. C. Scaletsky, L. C. Campos, T. A. Gomes, T. S. Whittam, and L. R. Trabulsi. 1996. Clonal structure and virulence factors in strains of *Escherichia coli* of the classic serogroup O55. *Infect. Immun.* **64**:2680–2686.
58. Rothbaum, R., A. J. McAdams, R. Giannella, and J. C. Partin. 1982. A clinicopathologic study of enterocyte-adherent *Escherichia coli*: a cause of protracted diarrhea in infants. *Gastroenterology* **83**:441–454.
59. Schremmer, C., J. E. Lohr, U. Wasthuber, J. Kisters, K. Ravelshofer, H. Steinruck, and L. H. Wieler. 1999. Enteropathogenic *Escherichia coli* in *Psittaciformes*. *Avian Pathol.* **28**:349–354.
60. Sohel, I., J. L. Puente, W. J. Murray, J. Vuopio-Varkila, and G. K. Schoolnik. 1993. Cloning and characterization of the bundle-forming pilin gene of enteropathogenic *Escherichia coli* and its distribution in *Salmonella* serotypes. *Mol. Microbiol.* **7**:563–575.
61. Sperandio, V., J. B. Kaper, M. R. Bortolini, B. C. Neves, R. Keller, and L. R. Trabulsi. 1998. Characterization of the locus of enterocyte effacement (LEE) in different enteropathogenic *Escherichia coli* (EPEC) and Shiga-toxin producing *Escherichia coli* (STEC) serotypes. *FEMS Microbiol. Lett.* **164**:133–139.
62. Stephan, R., N. Borel, C. Zweifel, M. Blanco, and J. E. Blanco. 2004. First isolation and further characterization of enteropathogenic *Escherichia coli* (EPEC) O157:H45 strains from cattle. *BMC Microbiol.* **4**:10.
63. Stone, K. D., H. Z. Zhang, L. K. Carlson, and M. S. Sonnenberg. 1996. A cluster of fourteen genes from enteropathogenic *Escherichia coli* is sufficient for the biogenesis of a type IV pilus. *Mol. Microbiol.* **20**:325–337.
64. Tobe, T., T. Hayashi, C. G. Han, G. K. Schoolnik, E. Ohtsubo, and C. Sasakawa. 1999. Complete DNA sequence and structural analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid. *Infect. Immun.* **67**:5455–5462.
65. Tobe, T., G. K. Schoolnik, I. Sohel, V. H. Bustamante, and J. L. Puente. 1996. Cloning and characterization of *bfpTVW*, genes required for the transcriptional activation of *bfpA* in enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **21**:963–975.
66. Trabulsi, L. R., R. Keller, and T. A. Tardelli Gomes. 2002. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg. Infect. Dis.* **8**:508–513.
67. Valentin-Branth, P., H. Steinsland, T. K. Fischer, M. Perch, F. Scheutz, F. Dias, P. Aaby, K. Molbak, and H. Sommerfelt. 2003. Cohort study of Guinean children: incidence, pathogenicity, conferred protection, and attributable risk for enteropathogens during the first 2 years of life. *J. Clin. Microbiol.* **41**:4238–4245.
68. Vallance, B. A., and B. B. Finlay. 2000. Exploitation of host cells by enteropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **97**:8799–8806.
69. Whittam, T. S., and E. A. McGraw. 1996. Clonal analysis of EPEC serogroups. *Rev. Microbiol.* **27**(Suppl. 1):7–16.
70. Whittam, T. S., and R. A. Wilson. 1988. Genetic relationships among pathogenic *Escherichia coli* of serogroup O157. *Infect. Immun.* **56**:2467–2473.
71. Wieler, L. H., T. K. McDaniel, T. S. Whittam, and J. B. Kaper. 1997. Insertion site of the locus of enterocyte effacement in enteropathogenic and enterohemorrhagic *Escherichia coli* differs in relation to the clonal phylogeny of the strains. *FEMS Microbiol. Lett.* **156**:49–53.
72. Zhang, W. L., B. Kohler, E. Oswald, L. Beutin, H. Karch, S. Morabito, A. Caprioli, S. Suerbaum, and H. Schmidt. 2002. Genetic diversity of intimin genes of attaching and effacing *Escherichia coli* strains. *J. Clin. Microbiol.* **40**:4486–4492.