

Characterization of an Inducible Oxidative Stress System in *Bacillus subtilis*

DAVID K. BOL AND RONALD E. YASBIN*

Department of Biological Sciences, Program in Molecular and Cellular Biology,
University of Maryland Baltimore County, Baltimore, Maryland 21228

Received 25 September 1989/Accepted 20 March 1990

Exponentially growing cells of *Bacillus subtilis* demonstrated inducible protection against killing by hydrogen peroxide when prechallenged with a nonlethal dose of this oxidative agent. Cells deficient in a functional *recE*⁺ gene product were as much as 100 times more sensitive to the H₂O₂ but still exhibited an inducible protective response. Exposure to hydrogen peroxide also induced the *recE*⁺-dependent DNA damage-inducible (*din*) genes, the resident prophage, and the product of the *recE*⁺ gene itself. Thus hydrogen peroxide is capable of inducing the SOS-like or SOB system of *B. subtilis*. However, the induction of this DNA repair system by other DNA-damaging agents is not sufficient to activate the protective response to hydrogen peroxide. Therefore, at least one more regulatory network (besides the SOB system) that responds to oxidative stress must exist. Furthermore, the data presented indicate that a functional catalase gene is necessary for this protective response.

Following exposure of many bacteria to nonlethal doses of a variety of environmental stresses, the cells demonstrate an enhanced resistance to the noxious compounds. These responses are generally defined by the induction of a unique set of genes. An example of such a system is the regulon in *Bacillus subtilis* that is induced following exposure of the bacteria to UV light and other DNA-damaging agents (5, 16, 23). This system, termed the SOS-like or SOB response (23), is analogous to the SOS regulon of *Escherichia coli* (12). The SOB system is a set of coordinately controlled genes which regulate responses such as prophage induction, Weigle reactivation, and error-prone repair (15). In *Salmonella typhimurium* and *E. coli* there exists a separate regulon, OxyR, that responds to oxidative stress (1, 4). Similarly, the heat shock regulon in *E. coli* is a set of operons that are transcribed following a temperature upshift (21). Interactions between stimulated regulons (stimulons) can often be demonstrated by the induction of one response rendering a cell population more tolerant to another stimulon-inducing agent (20). For instance, starved *E. coli* are more resistant to oxidative stress and heat (11). Additionally, some of the proteins produced during the heat shock response are considered oxidative stress proteins because they are under OxyR control (19). Furthermore, the competence regulon of *B. subtilis* (3), which consists of those genes required for the binding, uptake, and incorporation of exogenous DNA, has been shown to overlap with the SOB regulon (14). This interaction was demonstrated by the observation that genes controlled by the SOB system are induced in the absence of DNA-damaging agents when the organism reaches its natural competent state (14). Finally, the competence-controlled induction of the SOB genes is under the control of the regulatory genes *spo0A*⁺ and *spo0H*⁺, the products of which are also essential for sporulation to proceed correctly (22, 24).

When *B. subtilis* is grown to late stationary phase or exposed to hydrogen peroxide, a specific set of genes is induced (2). The transcription of some of these genes is also dependent on the products of *spo0A*⁺ and *spo0H*⁺ genes (2).

Because competence, a growth-phase dependent event, also induces the SOB system and requires the products of *spo0A*⁺ and *spo0H*⁺, it is considered possible that there exists an interaction between the oxidative stress-inducible system and the other stress-related regulons. Accordingly, we began to elucidate the molecular and genetic mechanisms that control the inducible oxidative stress system and to define the interactions between this system and the SOB response. Results presented in this report demonstrate that inducible resistance to oxidative stress appears to be independent of *recE*⁺ but dependent on a functional catalase gene.

(This research was submitted by D. K. Bol in partial fulfillment of the requirements for the Ph.D. degree.)

Cell survival and induced protection. Resistance to H₂O₂ in three *B. subtilis* strains was measured by exposing mid-exponentially growing cultures of each strain to various concentrations of peroxide for 15 min before the survivors were plated. Of the strains tested, repair-proficient strain YB886 showed the greatest resistance to H₂O₂ (Fig. 1). Recombination- and repair-deficient strain YB1015 (*recE4*) was 100-fold more sensitive to low concentrations of H₂O₂ than was wild-type strain YB886. However, both strains were killed with equal efficiency at higher concentrations of the oxidative agent.

Two catalases, one of which has been shown to be inducible by hydrogen peroxide (*katA*) (13), have recently been identified in *B. subtilis*. A mutation in the *katA* gene was moved into strain YB886 by congression (8), yielding strain YB2001, and sensitivity to H₂O₂ was tested as described above for strain YB886. Without a functional inducible catalase, *B. subtilis* was significantly more sensitive to H₂O₂ (Fig. 1); thus, it appears as though the vegetative catalase gene *katA*⁺ is necessary to provide resistance of growing *B. subtilis* to H₂O₂.

Repair-proficient *B. subtilis* YB886 grown in the presence of 50 μM H₂O₂ for two generations prior to a subsequent challenge with more substantial doses of peroxide and compared with naive cultures (Fig. 1) showed increased resistance to H₂O₂. Killing of the prechallenged cells decreased 2 to 3 orders of magnitude (at the higher doses) compared with

* Corresponding author.




FIG. 1. Cell survival and reduced sensitivity. Strains were grown to mid-exponential phase (50 CFU/ml) in Penassay antibiotic medium no. 3 (Difco Laboratories) with aeration at 37°C before challenge for 15 min with various concentrations of hydrogen peroxide (Fisher Scientific Co.) (open symbols). For induction of the oxidative stress system (closed symbols), cultures were grown to the same phase, diluted 1:10, and exposed to 50 μ M H₂O₂. Subsequent challenge with H₂O₂ occurred when the cultures reached 50 Klett units. Cell survival was determined by promptly diluting challenged cells into Spizizen salts (GIBCO Laboratories) and plating samples on tryptose blood agar base (Difco) immediately. The strains examined here are YB886 (○, ●), YB1015 (△, ▲), and YB2001 (□, ■).

untreated cells. Although the data shown here are from one representative experiment, cell killing at any one dose in an individual experiment deviated less than 1/2 order of magnitude from the average killing at that dose. In order to determine whether the protection induced by H₂O₂ in strain

YB886 was RecE-dependent, strain YB1015 was similarly prechallenged with H₂O₂ for two generations. This pretreatment was sufficient to reduce cell killing by 2 to 3 orders of magnitude at higher concentrations of peroxide. Thus, the protective response induced by small doses of H₂O₂ is not dependent on a functional *recE*⁺ gene product. Alternatively, the *kata* mutant strain YB2001 failed to show the response of reduced killing of pretreated cells shown in naive cultures (Fig. 1). Dependence on catalase for a protective response to H₂O₂ has been reported for *E. coli* (9, 10). The catalase-peroxidase enzyme HPI of *E. coli* is responsible for the protective response that is inducible by H₂O₂-ascorbate and is part of the *oxyR* regulon (7). A proposed mechanism for enhanced resistance during exposure to high concentrations of H₂O₂ is the protection of cellular transport components from H₂O₂ by catalase activity (4). The data presented in this paper demonstrate that *B. subtilis* has a similar system. Although a regulon like the OxyR system has not yet been discovered in *B. subtilis*, increased resistance to H₂O₂ and the correlation of this resistance with elevated catalase levels (13) indicate that a mechanism of activation or derepression of the catalase gene must exist. Since the *oxyR* gene product regulates genes identified in the heat shock response as well as resistance to oxidants, recent data suggest that the *kata*⁺ gene of *B. subtilis* may be under the control of multiple stress regulons (2).

Induction of SOB characteristics. *B. subtilis* cells treated with H₂O₂ showed elevated levels of the RecE protein within 30 min after exposure (data not shown). The product of the *recE*⁺ gene is a protein necessary for SOB induction, postreplication repair, and homologous recombination (17). RecE levels obtained after H₂O₂ treatment were comparable to those observed when cells were treated with other DNA-damaging agents that are known to induce the SOB response (14). Since induction of the *recE*⁺ gene in the presence of DNA damage (17) or during competence (14) is correlated with the activation of the SOB response, the ability of H₂O₂ to induce the SOB system was tested by assaying for the production of β -galactosidase from transcriptional fusions at three DNA-damage-inducible (*din*) loci (6). Essentially, pro-

TABLE 1. SOB induction profile for hydrogen peroxide

Strain	Din induction ^a after treatment with:		Prophage induction ^b after treatment with:			Filamentation ^c after treatment with:	
	H ₂ O ₂	MC ^d	NT ^e	H ₂ O ₂	MC	H ₂ O ₂	MC
YB886 DinA	+++	+++					
YB886 DinB	+++	+++					
YB886 DinC	+++	+++					
YB1015 DinA	—	—					
YB886 ϕ 105			6.8 × 10 ⁶	3 × 10 ⁹	1.2 × 10 ⁹		
YB1015 ϕ 105			5 × 10 ³	4 × 10 ³	5.1 × 10 ³		
YB886						+++	+++
YB1015						+++	+++

^a Strains carrying Tn917::lacZ fusions to damage-inducible genes (*din*) were maintained on nutrient broth no. 2 (Oxoid Ltd.) with 1.5% purified Oxoid agar. Erythromycin and lincomycin (Sigma Chemical Co.) were added to maintain selection for the transposon Tn917 in concentrations of 1 and 25 μ g/ml, respectively. Cultures to be assayed were taken from overnight growth on Oxoid nutrient and in GM1 (0.5% glucose, 0.1% yeast extract, 0.05% casein hydrolysate, and Spizizen salts supplemented with tryptophan and methionine [50 μ g/ml] [24]). Exponentially growing cells were induced with 3 mM H₂O₂ or 50 ng of mitomycin C per ml, and samples were assayed every 15 min for 2 h. β -Galactosidase levels were obtained as described previously (14). Basal levels were consistently less than 3 U per optical density of cells at 600 nm, and induction was as high as 40 U per optical density of cells at 600 nm.

^b Values are in PFU per milliliter. Strains lysogenic for ϕ 105 were grown to 35 Klett units in Oxoid nutrient broth no. 2 supplemented with 0.5% yeast extract and were induced with mitomycin C (0.5 μ g/ml) or selected concentrations of hydrogen peroxide. After 20 min, cultures were diluted 1/100, incubated for 90 min, and centrifuged for 5 min (13,500 rpm [10,000 × g]), and supernatant titers were then determined (25). +++, Maximum level; —, no induction.

^c Filamentation was determined by exposing mid-exponential-phase cells to 1 mM H₂O₂ or 50 ng of mitomycin C per ml and observing the cells 30 min later under a light-phase microscope (magnification, ×100).

^d MC, Mitomycin C.

^e NT, No treatment.




FIG. 2. Sensitivities of mitomycin C-treated cells to H₂O₂. Repair-proficient strain YB886 was grown exponentially for two generations in the presence of 50 ng of mitomycin C per ml prior to exposure to the indicated concentrations of H₂O₂. Survival of pretreated cells (■) is compared with that of naive cells (□).

motors of *dinA76*, *dinB7*, and *dinC17* were induced with hydrogen peroxide as well as with mitomycin C (Table 1). The results demonstrated that lack of a functional *recE*⁺ gene renders all of the promoters noninducible by both H₂O₂ and mitomycin C (16). *din* gene induction was also examined in a dose-dependent fashion by exposing strain YB886, carrying an insertion at the *dinB17* locus, to various amounts of H₂O₂ (data not shown). In this case, β-galactosidase production was induced maximally from this fusion by 3 mM H₂O₂. These levels of enzyme activity were comparable to those induced by mitomycin C and were not increased by higher concentrations of H₂O₂. Another characteristic of the activation of the SOB response is the induction of resident prophage (15). *B. subtilis* prophage φ105 was easily induced by exposure of lysogenic strain YB886 to 1 mM H₂O₂ (Table 1). Again, the induction of the prophage did not occur in the absence of a functional *recE*⁺ gene product (Table 1). However, H₂O₂ induced filamentation in both wild-type strain YB886 and *recE4* mutant strain YB1015 (Table 1). It has been demonstrated that filamentation is a *recE*⁺-independent component of the SOB response (15). Taken collectively, these data demonstrate that hydrogen peroxide is an inducer of the SOB response. Since H₂O₂ is capable of inducing the SOB response, the possibility exists that H₂O₂ protection was merely the result of the induction of an unidentified repair mechanism that is *recE*⁺ independent. However, when cells were treated with mitomycin C to fully induce the SOB system, no corresponding induction of oxidative stress resistance was observed (Fig. 2). There was a small increase in resistance; however, this is probably due to filamentation, in which killing kinetics are obscured by failure to septate.

The results clearly demonstrate that the inducible oxidative stress resistance of *B. subtilis* is independent of a functional *recE*⁺ gene product. However, this system does require an inducible *kata*⁺ gene. The regulatory components involved in the induction of *kata* and other oxidative stress resistance genes remain to be defined.

We thank Peter Loewen for his generous gift of the catalase mutation in *B. subtilis* 168 and Hal Schreier and Ken Bayles for their editorial comments.

This research was supported by Public Health Service grant R01DE08506 from the National Institutes of Health.

LITERATURE CITED

- Christman, M. F., R. W. Morgan, F. S. Jacobson, and B. N. Ames. 1985. Positive control of a regulon for defense against oxidative stress and some heat shock proteins in *Salmonella typhimurium*. *Cell* **41**:753–762.
- Dowds, B. C. A., P. Murphy, D. J. McConnell, and K. M. Devine. 1987. Relationship among oxidative stress, growth cycle, and sporulation in *Bacillus subtilis*. *J. Bacteriol.* **169**:5771–5775.
- Dubnau, D. 1989. The competence regulon of *Bacillus subtilis*, p. 147–166. In I. Smith, R. A. Slepecky, and P. Setlow (ed.), Regulation of procaryotic development: structural and functional analysis of bacterial sporulation and germination. American Society for Microbiology, Washington, D.C.
- Farr, S. B., D. Touati, and T. Kogoma. 1988. Effects of oxygen stress on membrane functions in *Escherichia coli*: role of HPI catalase. *J. Bacteriol.* **170**:1837–1842.
- Friedman, B. F., and R. E. Yasbin. 1983. The genetics and specificity of the constitutive excision repair system of *Bacillus subtilis*. *Mol. Gen. Genet.* **190**:481–486.
- Gillespie, K., and R. E. Yasbin. 1987. Chromosomal locations of three *Bacillus subtilis* *din* genes. *J. Bacteriol.* **169**:3372–3374.
- Greenberg, J. T., and B. Dimple. 1988. Overproduction of peroxide-scavenging enzymes in *Escherichia coli* suppress spontaneous mutagenesis and sensitivity to redox-cycling agents in *OxyR*[−] mutants. *EMBO J.* **7**:2611–2617.
- Hadden, C., and E. W. Nester. 1968. Purification of competent cells in the *Bacillus subtilis* transformation system. *J. Bacteriol.* **95**:876–885.
- Imlay, J. A., and S. Linn. 1986. Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. *J. Bacteriol.* **166**:519–527.
- Imlay, J. A., and S. Linn. 1987. Mutagenesis and stress responses induced in *Escherichia coli* by hydrogen peroxide. *J. Bacteriol.* **169**:2967–2976.
- Jenkins, D. E., J. E. Schultz, and A. Matin. 1988. Starvation-induced cross protection against heat or H₂O₂ challenge in *Escherichia coli*. *J. Bacteriol.* **170**:3910–3914.
- Kenyon, C., and G. C. Walker. 1980. DNA-damaging agents stimulate gene expression at specific loci in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **77**:2819–2823.
- Loewen, P. C., and J. Switala. 1987. Multiple catalases in *Bacillus subtilis*. *J. Bacteriol.* **169**:3601–3607.
- Love, P. E., M. J. Lyle, and R. E. Yasbin. 1985. DNA damage-inducible loci are transcriptionally activated in competent *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **82**:6201–6205.
- Love, P. E., and R. E. Yasbin. 1984. Genetic characterization of the inducible SOS-like system of *Bacillus subtilis*. *J. Bacteriol.* **160**:910–920.
- Love, P. E., and R. E. Yasbin. 1986. Induction of the *Bacillus subtilis* SOS-like response by *E. coli* RecA protein. *Proc. Natl. Acad. Sci. USA* **83**:5204–5208.
- Marrero, R., and R. E. Yasbin. 1988. Cloning of the *Bacillus subtilis* *recE*⁺ gene and functional expression of *recE*⁺ in *B. subtilis*. *J. Bacteriol.* **170**:335–344.
- Miller, J. H. 1972. Experiments in molecular genetics, p. 352–355. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Morgan, R. W., M. F. Christman, F. S. Jacobson, G. Storz, and B. N. Ames. 1986. Hydrogen peroxide inducible proteins in *Salmonella typhimurium* overlap with heat shock and other stress proteins. *Proc. Natl. Acad. Sci. USA* **83**:8059–8063.
- Murphy, P., B. C. A. Dowds, D. J. McConnell, and K. M. Devine. 1987. Oxidative stress and growth temperature in *Bacillus subtilis*. *J. Bacteriol.* **169**:5766–5770.
- Neidhardt, F. C., and R. A. VanBogelen. 1987. Heat shock

- response, p. 1334–1345. In J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, vol. 1. American Society for Microbiology, Washington, D.C.
22. Piggot, P. J., A. Moir, and D. A. Smith. 1981. Advances in the genetics of *Bacillus subtilis* differentiation, p. 29–39. In H. S. Levinson, A. L. Sonenshein, and D. J. Tipper (ed.), Sporulation and germination. American Society for Microbiology, Washington, D.C.
23. Yasbin, R. E. 1977. DNA repair in *Bacillus subtilis*. The presence of an inducible system. *Mol. Gen. Genet.* **153**:211–218.
24. Yasbin, R. E., J. Jackson, P. Love, and R. Marrero. 1988. Dual regulation of the RecE gene, p. 109–113. In A. T. Ganesan and J. A. Hoch (ed.), Genetics and biotechnology of bacilli, vol. 2. Academic Press, Inc. (London), Ltd., London.
25. Yasbin, R. E., G. A. Wilson, and F. E. Young. 1975. Transformation and transfection in lysogenic strains of *Bacillus subtilis*: evidence for selective induction of prophage in competent cells. *J. Bacteriol.* **121**:296–304.