

## A Highly Conserved Repeated Chromosomal Sequence in the Radioresistant Bacterium *Deinococcus radiodurans* SARK

EILEEN LENNON,† PABLO D. GUTMAN, HANLONG YAO, AND KENNETH W. MINTON\*

Department of Pathology, F. E. Hébert School of Medicine, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Road, Bethesda, Maryland 20814-4799

Received 10 September 1990/Accepted 5 January 1991

**A DNA fragment containing a portion of a DNA damage-inducible gene from *Deinococcus radiodurans* SARK hybridized to numerous fragments of SARK genomic DNA because of a highly conserved repetitive chromosomal element. The element is of variable length, ranging from 150 to 192 bp, depending on the absence or presence of one or two 21-bp sequences located internally. A putative translational start site of the damage-inducible gene is within the reiterated element. The element contains dyad symmetries that suggest modes of transcriptional and/or translational control.**

*Deinococcus radiodurans* is the type species of the family *Deinococcaceae*, a small eubacterial family whose members share in common extreme resistance to the lethal and mutagenic effects of ionizing and UV radiation and many genotoxic chemicals. Although repair of DNA damage by these organisms is known to be unusually efficient, the molecular mechanisms responsible remain largely unidentified (6). We recently reported a method that uses transformation of *D. radiodurans* SARK to generate random chromosomal gene fusions with a 5'-truncated *lacZ* (*lac'Z*) gene (5). Several of these gene fusions were inducible following exposure of *D. radiodurans* to the DNA-damaging agent mitomycin C, and these gene fusions were cloned (5).

**SARK sequence in pEL21.** A SARK DNA fragment from one cloned mitomycin C-inducible gene fusion (pEL21) hybridized to numerous fragments in Southern blots of SARK genomic DNA, suggesting the presence of a reiterated sequence (Fig. 1). The SARK DNA fragment in pEL21 is 419 bp (Fig. 2). There are two translational terminators, both TGA, in the SARK fragment in frame with *lac'Z* (Fig. 2). Downstream from these terminators, and in frame with *lac'Z*, are seven translational start codons: two ATGs, two GTGs, and three TTGs. One ATG (bases 255 to 257 [Fig. 2]) appears most likely to be the translation start site by virtue of its proximity to Shine-Dalgarno (S-D) sequences. The deduced protein-coding sequence for this possible start site encodes 55 amino acids prior to fusion with *lac'Z*, and both this start site and part of the coding sequence are within the reiterated element (Fig. 2 and 3). There is a 31-nucleotide hairpin with almost perfect dyad symmetry, 62 nucleotides upstream from this ATG, which is also within the reiterated element. The ATG itself also falls within a region compatible with formation of an mRNA hairpin with the ATG located on the upstream portion of the stem (Fig. 2).

**Sequence responsible for the multiple hybridization pattern is highly conserved.** When SARK genomic DNA fully digested with *EcoRI* was used to construct a library in the *EcoRI* site of pUC18, about 3% of all *Escherichia coli* colonies hybridized with the *D. radiodurans* insert in pSRE1 (pSRE1 is a subclone of pEL21 [Table 1]; SRE stands for

SARK repeated element). Four of the isolates detected by colony hybridization that contained SARK DNA fragments of different sizes, pSRE4, pSRE5, pSRE11, and pSRE29 (Table 1), were selected for sequence analysis of the reiterated element. Sequencing was by the dideoxy method using <sup>35</sup>S-dATP. When needed, dITP was used to resolve band compressions. pSRE1 and pSRE11 were sequenced by using forward and reverse primers for pUC18. Once these sequences were determined, two custom oligonucleotides




FIG. 1. Southern blot of *D. radiodurans* DNA probed with insert of pSRE1. *D. radiodurans* SARK genomic DNA was cleaved with *EcoRI* (lane 1), *HindIII* (lane 2), *SmaI* (lane 3), and *SmaI* (lane 4). Each lane contains 1 µg of DNA. The probe was the 360-bp *BglII*-*PvuII* SARK DNA insert from pSRE1 (a subclone of pEL21 [Table 1]). The fragment was radiolabeled with [<sup>32</sup>P]dCTP by nick translation. Hybridization was at 10<sup>5</sup> cpm/ml for 18 h in 50% formamide at 37°C. The final washing was at high stringency (65°C in 0.1× SSC [1× SSC is 0.15 M NaCl plus 0.015 M sodium citrate] for 1 h). *HindIII*-cleavable sites in *D. radiodurans* DNA are rare, and the majority of *HindIII*-cleaved DNA migrates uniformly at the upper limit of resolution (4, 10). The pattern of hybridization to the *HindIII* digest reflects this bias.

\* Corresponding author.

† Present address: National Center for Research Resources, National Institutes of Health, Bethesda, MD 20892.




FIG. 2. Map of pEL21, DNA sequence of its *D. radiodurans* portion, and likely RNA secondary structures. The pRF1 fusion vector contains the kanamycin resistance determinant *aphA* (■) and the 5'-truncated *lacZ* gene deleted of its promoter and first seven codons (*lacZ*, ■■■). A unique *Bgl*III site is present adjacent to the eighth codon (5, 7). Inserted into the *Bgl*III site is the *D. radiodurans* fragment (■■■■) that contains the mitomycin C-inducible gene translationally fused to *lacZ* (5). Arrows indicate direction of transcription. Restriction sites are as follows: B, *Bgl*III; C, *Cla*I; E, *Eco*RI; H, *Hind*III; P, *Pvu*II; Ps, *Pst*I; RV, *Eco*RV; S, *Sal*I; Sa/B, *Sau*3AI-*Bgl*III fusion; Sm, *Sma*I; Ss, *Sst*II; and St, *Stu*I. The DNA sequence extends from the *Bgl*III site at the 5' end of the *D. radiodurans* fragment to just beyond the *Bgl*III-*Sau*3AI site at the eighth codon of *lacZ*. The *Bgl*III, *Sma*I, and *Pvu*II sites in the sequence are underlined. At the 3' end of the sequence, *lacZ* bases and corresponding amino acids are indicated in italics. Upstream translational terminators that are in frame with *lacZ* are underlined and labeled End. The two ATG codons that are in frame with *lacZ* are underlined. S-D sequences are underlined, using the criterion (3) of at least four consecutive nucleotides, including three G's, from the GGAGG consensus (8). The deduced amino acid sequence begins with the second ATG. The two brackets in the DNA sequence indicate the ends of the repeated sequence detailed in Fig. 3. The two dyad symmetries in the DNA sequence, one near the 5' end and the other near the 3' end of the reiterated element, are indicated by opposing arrows above the relevant bases. Hypothetical RNA structures corresponding to the reiterated DNA element are shown with dyad symmetries in stem-loop conformation. Free energies of formation were calculated essentially as previously described (12).

TABLE 1. Bacteria and plasmids

Strain or plasmid	Description (reference or source <sup>a</sup> )
<b>Strains</b>	
<i>Deinococcus radiodurans</i> SARK <sup>b</sup>	Wild type (2); UWO 298 <sup>c</sup>
<i>Escherichia coli</i> DH5 $\alpha$ <sup>d</sup>	<i>recA1 endA1 hsdR17</i> ( $r_K^- m_K^+$ ) F <sup>-</sup> $\phi$ 80 <i>lacZ</i> $\Delta$ M15 $\Delta$ ( <i>lacZYA-argF</i> )U169 (Bethesda Research Laboratories)
<b>Plasmids</b>	
pEL21	419-bp fragment of SARK chromosomal DNA ligated to a 5'-truncated <i>lacZ</i> gene in the <i>E. coli lacZ</i> translational fusion vector pRF1, Km <sup>r</sup> (5)
pUC18	2,686-bp pBR322 derivative, Ap <sup>r</sup> , multiple cloning site in proximal <i>lacZ'</i> protein-coding sequence (14)
pSRE1	360-bp subclone of SARK DNA that contains reiterated sequence from pEL21; pUC18 <i>Bam</i> HI- <i>Sma</i> I-A::pEL21 <i>Bgl</i> III- <i>Pvu</i> II-D
pSRE4	pUC18 <i>Eco</i> RI::4-kb <i>Eco</i> RI SARK fragment
pSRE5	pUC18 <i>Eco</i> RI::1.2-kb <i>Eco</i> RI SARK fragment
pSRE11	pUC18 <i>Eco</i> RI::2.0-kb <i>Eco</i> RI SARK fragment
pSRE29	pUC18 <i>Eco</i> RI::1.6-kb <i>Eco</i> RI SARK fragment
pS19	Full-length clone of natural <i>D. radiodurans</i> SARK plasmid pUE11 (45 kb) in the <i>Eco</i> RV site of Km <sup>r</sup> ColE1 derivative pS27 (9)
pS28	Full-length clone of natural <i>D. radiodurans</i> SARK plasmid pUE10 (37 kb) in the <i>Eco</i> RV site of pS27 (9)

<sup>a</sup> Plasmids for which a reference is not noted were constructed during this study.

<sup>b</sup> *D. radiodurans* was grown at 32°C in TGY broth or on TGY plates solidified with 1.5% agar (10).

<sup>c</sup> University of Western Ontario culture collection, London, Ontario, Canada.

<sup>d</sup> *E. coli* was grown in LB broth or LB agar at 37°C. Selection of *E. coli* strains transformed with pUC18 was on LB agar supplemented with 50  $\mu$ g of ampicillin per ml. Selection of *E. coli* containing pEL21 was on agar containing 20  $\mu$ g of kanamycin per ml.

were employed to obtain the sequence and flanking sequences of the repetitive elements in the other clones. One custom oligonucleotide, 5'-AGAGCAGTTCTCCGAAT-3', is complementary to the sequence of the repeated element from nucleotides 175 to 191 (as numbered in Fig. 3) and was used to sequence the upper strand of the repeated element; the other custom primer, 5'-GAGCATTGACAAAAG-3', is the sequence of the repeated element from nucleotides 4 to 20 (as numbered in Fig. 3) and was used to sequence the lower strand. In all cases, both upper and lower strands were sequenced. The sizes of the five repeated elements are 151 nucleotides for SRE1, 171 nucleotides for SRE4 and SRE29, and 192 nucleotides for SRE11 and SRE5 (Fig. 3). The difference in the sizes is accounted for by the presence or absence of one or two 21-nucleotide sequences inserted after nucleotide 90 and the presence of an additional base at nucleotide 157 in SRE1. The two 21-bp inserts are direct repeats. Within the entire reiterated sequence, including the 21-bp inserts, there are only eight nucleotides that show point variations (Fig. 3). The repeated elements have a mean GC content of 56%, which is lower than the 67 to 70% GC content characteristic of *D. radiodurans* SARK and all other members of the family *Deinococcaceae* (2). All versions of the repeat contain dyad symmetry A (DS-A) at the 5' end, as shown in the pEL21 (SRE1) gene fusion in Fig. 2; however,




FIG. 3. Reiterated chromosomal DNA sequence. Sequences flanking the reiterated element are shown in lowercase letters, while the reiterated element is in uppercase letters. Numbering begins at the first base of the reiterated sequence and continues through base 190; the reiterated sequence ends at base 192. This numbering scheme differs from that employed in Fig. 2. S-D sequences and the ATG start codon as proposed for SRE1 are underlined. The eight asterisks indicate the sites of point nucleotide differences among the five sequences. The dashes in the sequences of SRE4, SRE29, and SRE1 indicate the absence of one or both 21-bp inserts that are present in SRE11 and SRE5. Continuous lines below the sequences indicate dyad symmetries (DS-A, DS-B, DS-C, and DS-C), while the double dashed lines indicate direct repeats (DR-I, DR-II, and DR-III). Homologies between DR-I and DR-II, DR-I and DR-III, and DR-II and DR-III are 81, 88, and 77%, respectively.

there is a single base change from C to A within DS-A in the other four sequences (Fig. 3).

Hypothetical RNA secondary structure corresponding to the 3' end of the element depends on the number of 21-bp inserts, with the potential formation of either one stem-loop structure or two closely adjacent stem-loop structures (Fig. 3). We speculate that the location of S-D sequences in and adjacent to this region of potential secondary structure may have regulatory significance for translation. Modifications in this portion of the repeated element in other copies compared with SRE1 include elimination of several canonical S-D sequences and an additional potential hairpin structure (SRE4 and SRE29). In SRE5 and SRE11 there is the possible formation of mutually exclusive hairpin structures (DS-B versus DS-C) (Fig. 3). All five of the sequenced elements contain DS-A and DS-D (Fig. 2 and 3). We suggest that DS-A may be a transcriptional terminator because of its nearly perfect dyad symmetry of 26 nucleotides (13 bp; one G-T base pair) in pEL21 (SRE1), a feature common in transcriptional terminators (1).

**Possible functions of reiterated element.** Since the reiterated element was detected as part of a mitomycin C-inducible gene fusion, it may be a highly conserved consensus sequence for a DNA damage-inducible regulon in *D. radio-*

*durans*. If so, our estimate of  $\geq 30$  copies would suggest a large number of genes within the regulon, comparable with the number of genes in the SOS regulon of *E. coli* (13). Using appropriate oligonucleotide primers for this element and polymerase chain reaction conditions sufficient to amplify sequences up to 1.5 kb, we did not find any polymerase chain reaction-amplifiable sequences larger than the sizes of the elements described here (not shown), suggesting that this element is not part of a transposon. Strain SARK contains two cryptic plasmids, pUE11 and pUE10 (9). On the basis of strong hybridization of pUE11 against SARK chromosomal DNA, we have suggested that pUE11 is an episome (11). Neither pUE11 nor pUE10 hybridizes with the pSRE1 insert (Fig. 4); consequently, homology between pUE11 and the chromosome cannot be ascribed to the reiterated element. *Deinococcus* spp. are unique in their ability to mend several hundred double-strand breaks per chromosome without any loss of viability (6). In this regard, a novel hypothesis for the occurrence of these conserved repeated sequences is that they function as specific substrate sites for recombinational repair of DNA. Frequent exchange of sequence information by recombination at these sites may serve to retain the repeated elements' high degree of sequence conservation.

**Nucleotide sequence accession number.** The GenBank ac-




FIG. 4. Southern blot of *D. radiodurans* SARK plasmids probed with insert of pSRE1. pS28 (pUE10 [Table 1]), pS19 (pUE11 [Table 1]), and SARK total DNA were cleaved with *EcoRI*, electrophoresed in a 1% agarose gel, stained with 0.1  $\mu$ g of ethidium bromide per ml, and blotted. Hybridization and washing were as described in the legend to Fig. 1. (A) Ethidium bromide-stained agarose gel. (B) Autoradiogram of Southern blot.

cession number for the SARK fragment in pEL21 is M59306. The GenBank accession numbers for SRE1, SRE4, SRE5, SRE11, and SRE29 are M59306, M59307, M59308, M59309, and M59310, respectively.

This work was supported by Public Health Service grant GM39933.

#### REFERENCES

- Brendel, V., and E. N. Trifonov. 1984. A computer algorithm for testing potential prokaryotic terminators. *Nucleic Acids Res.* **12**:4411–4427.
- Brooks, B. W., and R. G. E. Murray. 1981. Nomenclature for “*Micrococcus radiodurans*” and other radiation-resistant cocci: *Deinococcaceae* fam. nov., including five species. *Int. J. Syst. Bacteriol.* **31**:353–360.
- Gold, L., and G. Stormo. 1987. Translational initiation, p. 1302–1307. In F. C. Niedhardt, J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, vol. 2. American Society for Microbiology, Washington, D.C.
- Gutman, P. D., H. Yao, and K. W. Minton. *Mutat. Res.*, in press.
- Lennon, E., and K. W. Minton. 1990. Gene fusions with *lacZ* by duplication insertion in the radioresistant bacterium *Deinococcus radiodurans*. *J. Bacteriol.* **172**:2955–2961.
- Moseley, B. E. B. 1983. Photobiology and radiobiology of *Micrococcus (Deinococcus) radiodurans*. *Photochem. Photobiol. Rev.* **7**:223–275.
- Robinson, G. W., C. M. Nicolet, D. Kalaninov, and E. C. Friedberg. 1986. A yeast excision-repair gene is inducible by DNA damaging agents. *Proc. Natl. Acad. Sci. USA* **83**:1842–1846.
- Shine, J., and L. Dalgarno. 1974. The 3'-terminal sequence of *Escherichia coli* 16S ribosomal RNA: complementarity to non-sense triplets and ribosome binding sites. *Proc. Natl. Acad. Sci. USA* **71**:1342–1346.
- Smith, M. D., R. Abrahamson, and K. W. Minton. 1989. Shuttle plasmids constructed by the transformation of an *Escherichia coli* cloning vector into two *Deinococcus radiodurans* plasmids. *Plasmid* **22**:132–142.
- Smith, M. D., E. Lennon, L. B. McNeil, and K. W. Minton. 1988. Duplication insertion of drug resistance determinants in the radioresistant bacterium *Deinococcus radiodurans*. *J. Bacteriol.* **170**:2126–2135.
- Smith, M. D., L. B. McNeil, and K. W. Minton. 1990. Plasmid genetics and gene expression in *Deinococcus radiodurans*, p. 125–131. In L. O. Butler, C. Harwood, and B. E. B. Moseley (ed.), *Recent advances in genetic transformation of animal and plant cells*. Intercept, Ltd., Andover, Hants, U.K.
- Tinoco, I., P. N. Borer, B. Dengler, M. D. Levine, O. C. Uhlenbeck, D. M. Crothers, and J. Gralla. 1973. Improved estimation of secondary structure in ribonucleic acids. *Nature (London) New Biol.* **246**:40–41.
- Walker, G. C. 1987. The SOS response of *Escherichia coli*, p. 1346–1357. In J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology, vol. 2. American Society for Microbiology, Washington, D.C.
- Yanisch-Perron, C., J. Vieira, and J. Messing. 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:109–119.