

FIG. 1. Elution profile of trichloroacetic acid-extracted cells of *B. alvei* separated on a Sephadex G-25 fine column. I to VI indicate peaks obtained; void volume is 90 ml.

modes were accumulated for 5 million and 10 million fission events, respectively.

Mass calibration was based on H^+ and on the $[M+H]^+$ ion of the matrix 3-(3-pyridyl) acrylic acid (PAA) (m/z 150,0555, monoisotopic) in the positive ionization mode and on H^- and the $[M-H]^-$ ion of the matrix (m/z 148,0398, monoisotopic) in the negative ionization mode.

PAA surfaces for sample adsorption in plasma desorption mass spectrometry were prepared by electrospraying as described elsewhere (2). The sample was dissolved in 8 μ l of *n*-hexane and was spin deposited (17) on a PAA-covered target foil.

Hydrolytic conditions. Hydrolytic conditions were as follows: condition a (amino sugars), 4 M HCl, 16 h, 100°C; condition b (neutral sugars), 2 M HCl, 2 h, 100°C; and condition c (mild acid hydrolysis [lipid pyrophosphates] [8, 15]), 1 drop of 1-propanol in 0.01 M HCl, 5 min, 100°C. For sugar-1-phosphates, conditions were 0.01 M HCl, 30 min, 100°C (condition d) and 0.01 M HCl, 10 min, 100°C (condition e). Condition f (4 M HCl, 30 min, 100°C) was used for partial hydrolysis. Condition g (chloroform-methanol-10 M NaOH [10:10:3 by volume]) was used for dolichyl monophosphate (21). The lipids were extracted from the hydrolysates (conditions c and g) with chloroform.

Standards. Dolichol- C_{80-105} , dolichol- C_{55} , dolichyl- C_{80-105} monophosphate, dolichyl- C_{55} monophosphate, and undecaprenol were obtained from Sigma, Deisenhofen, Germany. GlcNAc, Gal, and Glc were obtained from Serva, Heidelberg, Germany. ManN was obtained from Fluka, Neu Ulm, Germany. PAA was obtained from Aldrich, Steinheim, Germany.

RESULTS

Isolation of the nucleotide-activated precursors. Cells were extracted with aqueous trichloroacetic acid (12), and the extracts were separated on a Sephadex G-25 column (8). The elution profile showed six distinct peaks (Fig. 1). Peaks I and II containing ManNAc or GlcNAc were further separated on cellulose F254 aluminum sheets (solvent c; Table 1). The UV-absorbing bands were eluted and finally purified on polyethyleneimine-cellulose plates (solvents a and b; Table 1).

Characterization of nucleotide-activated monomers. Four nucleotide-activated monomers could be purified from peak II (compounds 3 to 6) by TLC (Table 1). Two compounds consisted of uracil, phosphate, and Glc (compound 3) or Gal (compound 4). Compound 5 was composed of ManNAc, guanine, and phosphate; compound 6 was composed of GlcNAc, uracil, and phosphate. N acetylation of the amino sugars was indicated by a positive direct Morgan-Elson test after removal of UDP (hydrolytic condition e) and a negative ninhydrin reaction. The data (Table 1) indicate that the four compounds are nucleotide-activated sugar monomers.

Characterization of nucleotide-activated oligosaccharides. Two nucleotide-activated oligosaccharides (compounds 1 and 2; Table 1) could be isolated by TLC from peak I (Fig. 1). Compound 1 was composed of guanine, phosphate, ManNAc, GlcNAc, and Gal. Compound 2 additionally contained Glc. Both compounds did not react with ninhydrin. This finding indicates that ManN and GlcN should be N acetylated. The molar ratio of the components (Table 1) indicates that compound 1 is a GDP-activated trisaccharide and compound 2 is a GDP-activated tetrasaccharide (Table 1). Both compounds have ManNAc at the reducing end (see Materials and Methods). After partial hydrolysis, two disaccharides (compounds 1a and 1b) and three disaccharides (compounds 2a, 2b, and 2c) could be isolated by TLC from compounds 1 and 2, respectively (solvent d; Table 2). The molar ratios of the components of compounds 1 and 2 (Table 1), the isolated oligosaccharides (Table 2), structural analysis, and chromatographic behavior indicate that compound 1 is a GDP-activated trisaccharide and that compound 2 is a GDP-activated tetrasaccharide (Table 1). Both oligosaccharides contain additional GlcNAc residues compared with the intact glycan of the mature S-layer glycoprotein (3).

Isolation and characterization of a lipid-activated intermediate. The butanol extracts (9) of cells disrupted by shaking with glass beads were applied to silica plates (solvent e). Bands were visualized with iodine vapor, eluted, and subsequently run on silica plates for a second time, using solvent g. One lipid- and ManNAc-containing band could be purified (Table 1, compound 7). Quantitative and structural analyses of compound 7 (Table 1 and 2) were performed as described for compounds 1 and 2. The results indicate that compound 7 is a dolichyl-activated hexasaccharide consisting of the repeating trisaccharide unit of the intact glycan (3) and three additional GlcNAc residues. As only one GlcNAc residue was destroyed with periodate, the three GlcNAc residues should be linearly arranged in a 1 \rightarrow 3 or 1 \rightarrow 4 linkage. The lipid carrier of compound 7 was identified as dolichol- C_{55} by cochromatography with standards as described recently (8) and by mass spectrometry after splitting off of the carbohydrate and phosphate residues (hydrolytic condition g) (8).

As shown in Fig. 2, the mass spectrometric analysis revealed a protonated molecular ion ($[M+H]^+$) at m/z 770.26, a sodiated molecular ion ($[M+Na]^+$) at m/z 792.58, and a cluster molecular ion ($[M+PAA+H]^+$) with high abundance at m/z 919.75, consistent with the dolichol- C_{55} structure. The calculated average relative molecular mass of dolichol- C_{55} is 769.35. The relative molecular mass was determined to be 769.40 (+0.0065%), calculated as the average of those for $[M+H]^+$, $[M+Na]^+$, and $[M+PAA+H]^+$. Additionally, the negative-ion plasma desorption mass spectrum (not shown) exhibited a deprotonated molecular ion ($[M-H]^-$) at m/z 767.79 and a deprotonated cluster molecular ion ($[M+PAA-H]^-$) at m/z 917.94. Weak signals at m/z 850.14, 872.48, and 999.75 were also present. These ions correspond to the protonated and sodiated molecular

TABLE 1. Analysis of isolated nucleotide-activated precursors (compounds 1 to 6) and an isolated lipid-activated oligosaccharide (compound 7)

Com- pound no.	Sephadex column no. ^a	TLC (R _f)						Absorption		Composition (molar ratio)						Proposed structure ^b		
		Solvent a	Solvent b	Solvent c	Solvent e	Solvent f	Solvent	260/ 280 nm	250/ 260 nm	Dolichol- C ₅₅	Guanine	Uracil	Phosphate	MannAc	GlcNAc		Glc	Gal
1	I	0.35 ^d	0.82	0.24 ^e	ND	ND	ND	0.69	0.99	—	1.0	— ^c	2.3	1.1	1.0	—	0.9	GDP-MannAc→Gal←GlcNAc
																		↓ MannAc ↓ Gal
																		↓ GlcNAc ↓ Glc
2	I	0.10 ^d	0.75	0.09 ^e	ND	ND	ND	0.71	1.03	—	1.0	—	2.0	0.9	0.9	1.0	0.9	GDP-MannAc→Gal←GlcNAc
																		↓ Glc ↓ MannAc→Glc ↓ Gal
																		↓ GlcNAc ↓ UDP-Glc
3	II	0.50	ND ^d	0.25	ND	ND	ND	0.42	0.78	—	—	1.0	2.1	—	—	1.1	—	GlcNAc ↓ UDP-Glc
4	II	0.49	ND	0.22	ND	ND	ND	0.40	0.79	—	—	1.0	2.0	—	—	0.9	—	Glc ↓ UDPGal
5	II	0.50	ND	0.32	ND	ND	ND	0.61	0.96	—	1.0	—	1.9	1.2	—	—	—	GDP-MannAc ↓ Gal
6	II	0.51	ND	0.33	ND	ND	ND	0.40	0.80	—	—	1.0	2.1	—	0.8	—	—	MannAc ↓ UDP-GlcNAc ↓ GlcNAc
7	ND	ND	ND	ND	0.42	0.49	ND	ND	Present	—	—	2.04	1.18	3.20	1.00	0.91	—	Dolichol-C ₅₅ -PP _i -MannAc→Gal←GlcNAc→GlcNAc→GlcNAc

^a See Fig. 1.^b See Table 2.^c —, not present.^d ND, not determined.

TABLE 2. Analysis of purified oligosaccharides obtained after partial hydrolysis of the isolated precursors

Compound no. ^a	Derived oligosaccharide no.	R_f^b	Composition (molar ratio)				Proposed structure
			ManN	GlcN	Gal	Glc	
1	1a	0.44	1.0 ^c	— ^d	0.9	—	ManNAc←Gal
	1b	0.34	—	1.0	1.0 ^c	—	Gal←GlcNAc
2	2a	0.45	1.0 ^c	—	1.1	—	ManNAc←Gal
	2b	0.40	1.0 ^c	—	—	1.2	ManNAc←Glc
7	2c	0.35	—	0.9	1.0 ^c	—	Gal←GlcNAc
	7a	0.45	1.0 ^c	—	1.1	—	ManNAc←Gal
	7b	0.41	1.0 ^c	—	—	1.2	ManNAc←Glc
	7c	0.35	—	0.9	1.0 ^c	—	Gal←GlcNAc
	7d	0.45	—	1:1 ^e	—	—	GlcNAc←GlcNAc
	7e	0.27	—	1:2 ^e	—	—	GlcNAc←GlcNAc←GlcNAc

^a See Table 1.^b Value in solvent d after partial hydrolysis and N acetylation of the amino sugars.^c Reducing sugar (determined as described in Materials and Methods).^d —, not present.^e Molar ratio of reducing sugar (=1) to nonreducing sugar.

ion as well as to the cluster molecular ion ($[M+PAA+H]^+$) of the dolichyl- C_{55} monophosphate. This observation clearly indicates incomplete hydrolysis of the sample and also helps to verify the precursor structure. However, no fragment ions at this level of deposited sample amount (approximately 5 μ g) could be detected.

DISCUSSION

From analysis of the isolated precursors, we believe that the biosynthesis (Fig. 3) of the S-layer glycoprotein of *B. alvei* starts with the formation of nucleotide-activated monosaccharides of Gal, Glc, ManNAc, and GlcNAc. The nucleotide is GDP in the case of ManNAc and UDP in the case of the other sugars. Three of the nucleotide-activated monosaccharides (GDP-ManNAc, UDP-Gal, and UDP-GlcNAc) are linked together, forming a GDP-activated trisaccharide with ManNAc at the reducing end. In the next step, UDP-Glc is transferred to the trisaccharide, forming a GDP-activated branched tetrasaccharide. This tetrasaccharide shows the same structure as does the repeating unit of the intact glycan (3), but it contains one additional GlcNAc residue. In a next step, the GDP residue of the tetrasaccharide should be replaced by dolichyl- C_{55} -PP_i. This lipid-activated precursor contains two additional GlcNAc residues compared with the GDP-activated tetrasaccharide.

Whether the GlcNAc residues are added before or after the transfer of the oligosaccharide to the lipid remains to be determined. No GlcNAc is found in the intact glycoprotein (3), which indicates that the additional GlcNAc residues are removed during further processing.

Transient occurrence of sugar residues is also the case with eukaryotic glycoproteins (4), the S-layer glycoprotein of *Methanothermobacter feravidus* (8), and the S-layer glycoprotein of *Halobacterium halobium* (15). Like undecaprenol, the dolichol derivative consists of 55 carbons. It is smaller than most eukaryotic dolichol derivatives but of the same size as the one found in *M. feravidus* (8). The proposed pathway has to be verified by further studies with labeled compounds and enzyme preparations.

Recently, nucleotide-activated oligosaccharides were proposed to be involved in the biosynthesis of different cell envelope polymers (glycoprotein [8], pseudomurein [9], and polysaccharide [10]) of methanogens. Therefore, the formation of nucleotide-activated oligosaccharides seems to be a typical and common feature of the biosynthesis of methanobacterial cell wall glycans, while in eubacteria, they occur only as intermediates in glycoprotein biosynthesis.

Since dolichol functions as a lipid carrier in the biosynthesis of two archaeobacterial glycoproteins (8, 14), a eubacterial glycoprotein, and eukaryotic glycoproteins (4), it seems most likely that dolichol is the universal lipid carrier for glycoproteins, while undecaprenol is the common lipid carrier of other cell wall glycans (9, 10, 20).




FIG. 2. Positive-ion plasma desorption mass spectrum of the dolichyl monophosphate fraction after partial hydrolysis.




FIG. 3. Proposed scheme of the biosynthesis of the S-layer glycoprotein of *B. alvei*. Dol, dolichyl; Dol_{C55}, dolichyl- C_{55} .

Except for a saturated α -C-C bond, dolichol-C₅₅ has the same structure as does undecaprenol. Nevertheless, the consequences for the further biosynthetic route of nucleotide-activated precursors are significant.

ACKNOWLEDGMENTS

We gratefully acknowledge support from the Deutsche Forschungsgemeinschaft (DFG Ko 785/4-3), Fonds der Chemischen Industrie, and University of Ulm (grant 8974/92/1423 to E.H.). The mass spectrometric part of this work was supported by the Hochschuljubiläumstiftung der Stadt Wien (grant H-00155/92 to G.A.).

REFERENCES

1. Albersheim, P., D. J. Nevins, P. D. English, and A. Karr. 1967. A method for the analysis of sugars in plant cell-wall polysaccharides by gas liquid chromatography. *Carbohydr. Res.* **5**:340-345.
2. Allmaier, G., E. Pittenauer, and E. R. Schmid. 1991. A chloroform-resistant, low-molecular-mass matrix for californium-252 plasma desorption mass spectrometry. *Org. Mass Spectrom.* **26**:804-806.
3. Altman, E., J.-R. Brisson, P. Messner, and U. B. Sleytr. 1991. Structure of the glycan chain from the surface layer glycoprotein of *Bacillus alvei* CCM 2051. *Biochem. Cell Biol.* **69**:72-78.
4. Berger, E. G., E. Buddecke, J. P. Kamerling, A. Kobata, I. C. Paulson, and J. F. G. Vliegenthart. 1982. Structure, biosynthesis and function of glycoprotein glycans. *Experientia* **38**:1129-1163.
5. Bergmeyer, H. U. 1974. *Methoden der enzymatischen Analyse*. Verlag Chemie, Weinheim, Germany.
6. Chen, P. S., T. Y. Toribara, and H. Warner. 1957. Microdetermination of phosphorus. *Anal. Chem.* **28**:1756-1758.
7. Dunn, D. B., and R. H. Hall. 1975. Purines, pyrimidines, nucleosides and nucleotides: physical constants and spectral properties, p. 65-215. *In* G. D. Fasman and H. A. Sober (ed.), *Handbook of biochemistry*. Selected data for molecular biology, 3rd ed. CRC Press, Cleveland.
8. Hartmann, E., and H. König. 1989. Uridine and dolichyl diphosphate activated oligosaccharide intermediates are involved in the biosynthesis of the surface layer glycoprotein of *Methanothermus fervidus*. *Arch. Microbiol.* **151**:274-281.
9. Hartmann, E., and H. König. 1990. Comparison of the biosynthesis of the methanobacterial pseudomurein and the eubacterial murein. *Naturwissenschaften* **77**:472-475.
10. Hartmann, E., and H. König. 1991. Nucleotide-activated oligosaccharides are intermediates of the cell wall polysaccharide of *Methanosarcina barkeri*. *Biol. Chem. Hoppe-Seyler* **372**:971-974.
11. Kandler, O., and H. König. 1985. Cell envelopes of archaebacteria, p. 413-457. *In* I. C. Gunsalus, J. R. Sokatch, and L. N. Ornston (ed.), *The bacteria. A treatise of structure and function*, vol. 8. Academic Press, New York.
12. König, H., O. Kandler, and W. Hammes. 1989. Biosynthesis of pseudomurein: isolation of putative precursors from *Methanobacterium thermoautotrophicum*. *Can. J. Microbiol.* **35**:176-181.
13. Laskowski, M. 1971. Venom exonuclease, p. 313-328. *In* P. D. Boyer (ed.), *The enzymes*, vol. IV, 3rd ed. Academic Press, New York.
14. Lechner, J., and F. Wieland. 1989. Structure and biosynthesis of prokaryotic glycoproteins. *Annu. Rev. Biochem.* **58**:173-194.
15. Lechner, J., F. Wieland, and M. Sumper. 1985. Transient methylation of dolichyl oligosaccharides is an obligatory step in halobacterial sulfated glycoproteins. *J. Biol. Chem.* **260**:8984-8989.
16. Messner, P., and U. B. Sleytr. 1991. Bacterial surface layer glycoproteins. *Glycobiology* **1**:545-551.
17. Nielsen, P. F., K. Klarskov, P. Hojrup, and P. Roepstorff. 1988. Optimization of sample preparation for plasma desorption mass spectrometry of peptides and proteins using a nitrocellulose matrix. *Biomed. Environ. Mass Spectrom.* **17**:355-362.
18. Reissig, J. L., J. L. Strominger, and L. F. Leloir. 1955. A modified colorimetric method for the estimation of N-acetyl amino sugars. *J. Biol. Chem.* **217**:959-966.
19. Sumper, M. 1987. Halobacterial glycoprotein biosynthesis. *Biochim. Biophys. Acta* **906**:69-79.
20. Tipper, D. J., and A. Wright. 1979. Structure and biosynthesis of bacterial cell walls, p. 201-426. *In* I. C. Gunsalus, J. R. Sokatch, and L. N. Ornston (ed.), *The bacteria. A treatise on structure and function*, vol. 7. Academic Press, New York.
21. Wedgwood, J. E., J. L. Strominger, and C. D. Warren. 1974. Transfer of sugars from nucleoside diphosphosugar compounds to endogenous and synthetic dolichyl phosphate in human lymphocytes. *J. Biol. Chem.* **249**:6316-6324.
22. Wheat, R. W. 1966. Analysis of hexosamines in bacterial polysaccharides by chromatographic procedures. *Methods Enzymol.* **8**:60-78.
23. Wieland, F., J. Lechner, and M. Sumper. 1982. The cell wall glycoprotein of halobacteria: structural, functional and biosynthetic aspects. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Abt. 1 Orig. Reihe C* **3**:161-170.