

Cloning, Characterization, and Functional Expression of *acs*, the Gene Which Encodes Acetyl Coenzyme A Synthetase in *Escherichia coli*

SUMAN KUMARI,¹ RINNAT TISHEL,² MICHAEL EISENBACH,² AND ALAN J. WOLFE^{1*}

Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois 60153,¹ and Department of Membrane Research and Biophysics, The Weizmann Institute of Science, 76100 Rehovot, Israel²

Received 23 November 1994/Accepted 9 March 1995

Acetyl coenzyme A synthetase (Acs) activates acetate to acetyl coenzyme A through an acetyladenylate intermediate; two other enzymes, acetate kinase (Ack) and phosphotransacetylase (Pta), activate acetate through an acetyl phosphate intermediate. We subcloned *acs*, the *Escherichia coli* open reading frame purported to encode Acs (F. R. Blattner, V. Burland, G. Plunkett III, H. J. Sofia, and D. L. Daniels, *Nucleic Acids Res.* 21:5408–5417, 1993). We constructed a mutant allele, $\Delta acs::Km$, with the central 0.72-kb *BclI-BclI* portion of *acs* deleted, and recombined it into the chromosome. Whereas wild-type cells grew well on acetate across a wide range of concentrations (2.5 to 50 mM), those deleted for *acs* grew poorly on low concentrations (≤ 10 mM), those deleted for *ackA* and *pta* (which encode Ack and Pta, respectively) grew poorly on high concentrations (≥ 25 mM), and those deleted for *acs*, *ackA*, and *pta* did not grow on acetate at any concentration tested. Expression of *acs* from a multicopy plasmid restored growth to cells deleted for all three genes. Relative to wild-type cells, those deleted for *acs* did not activate acetate as well, those deleted for *ackA* and *pta* displayed even less activity, and those deleted for all three genes did not activate acetate at any concentration tested. Induction of *acs* resulted in expression of a 72-kDa protein, as predicted by the reported sequence. This protein immunoreacted with antiserum raised against purified Acs isolated from an unrelated species, *Methanoxobacterium soehngenii*. The purified *E. coli* Acs then was used to raise anti-*E. coli* Acs antiserum, which immunoreacted with a 72-kDa protein expressed by wild-type cells but not by those deleted for *acs*. When purified in the presence, but not in the absence, of coenzyme A, the *E. coli* enzyme activated acetate across a wide range of concentrations in a coenzyme A-dependent manner. On the basis of these and other observations, we conclude that this open reading frame encodes the acetate-activating enzyme, Acs.

Escherichia coli cells activate acetate to acetyl coenzyme A (acetyl-CoA) by two distinct pathways (Fig. 1). One pathway, catalyzed by the enzyme acetyl-CoA synthetase (Acs) (acetate:CoA ligase [AMP forming]; EC 6.2.1.1), proceeds through an acetyladenylate (AcAMP) intermediate (3, 8). The second pathway, catalyzed by the enzymes acetate kinase (Ack) [acetyl-CoA(CoA):P; acetyltransferase; EC 2.7.2.1] and phosphotransacetylase (Pta) (ATP:acetate phosphotransferase; EC 2.3.1.8), proceeds through an acetyl phosphate intermediate (41). Acs first converts acetate and ATP to AcAMP, producing PP_i; it then reacts AcAMP with CoA to form acetyl-CoA, releasing AMP. Similarly, Ack first converts acetate and ATP to acetyl phosphate, producing ADP; Pta then reacts acetyl phosphate with CoA to form acetyl-CoA, releasing P_i.

Brown et al. (6) hypothesized that the Acs pathway functions as a catabolite-repressible, acetate-inducible, high-affinity acetate uptake system that scavenges acetate present extracellularly at relatively low concentrations. They also proposed that the Ack-Pta pathway functions primarily in a catabolic role, excreting acetate and generating ATP during mixed-acid fermentation and aerobic growth on excess glucose or other glycolytic intermediates. Finally, they argued that the low-affinity

Ack-Pta pathway activates acetate only when that molecule is present extracellularly in large quantity.

In addition to their role in acetate metabolism, the acetate activation pathways have been implicated in the regulation of signal transduction by two-component regulatory systems in several bacterial species (reviewed in references 32 and 54; see also references 2, 9, 39, and 55), the regulation of the glucose starvation stimulon of *E. coli* (35), and the synthesis of adenosine 5'-tetrphosphate and adenosine 5'-pentaphosphate in *Saccharomyces cerevisiae* (17).

Although *E. coli* mutants deficient for Ack and Pta activities have existed for some time, those deficient for Acs activity have not. Recently, Blattner et al. (5) reported an open reading frame, located within min 92 of the *E. coli* chromosome, that bears significant homology to those which encode Acs in other organisms. We now report the subcloning of the putative *E. coli acs* gene, the construction and characterization of mutant cells deleted for a portion of that gene, and the purification of its protein product. On the basis of a variety of biological, immunological, and biochemical assays, we conclude that this open reading frame does indeed encode Acs and that *E. coli* cells require both acetate activation pathways for optimal growth across a wide range of acetate concentrations.

* Corresponding author. Mailing address: Department of Microbiology and Immunology, Loyola University Chicago, Stritch School of Medicine, 2160 S. First Ave., Maywood, IL 60153. Phone: (708) 216-5814. Fax: (708) 216-9574. Electronic mail address: awolfe@wpo.it.luc.edu.

MATERIALS AND METHODS

Chemicals. The enzymes and substrates used in the enzyme assays were obtained from Boehringer Mannheim (Indianapolis, Ind.) or Sigma Chemical Company (St. Louis, Mo.). Radiolabelled materials were purchased from Amersham

TABLE 1. Bacterial strains used in this study

Strain	Relevant genotype ^a	Source or reference
AJW377	<i>cheA503I</i> (Am) <i>recA</i> ::Cm <i>thi-1 thr-1</i> (Am) <i>leuB6 metF159</i> (Am) <i>his-4 rpsL136 lacY1 xyl-5 ara-14 tonA31 tsx-78</i>	56
AJW763	<i>polA</i> (Ts) <i>rha zig</i> ::Tn10 <i>thi-1 thr-1</i> (Am) <i>leuB6 metF159</i> (Am) <i>his-4 rpsL136 lacY1 xyl-5 ara-14 tonA31 tsx-78</i>	This study
AJW803	CP875 Φ (Δ <i>acs</i> ::Km-1)	This study
AJW804	CP875 Φ (Δ <i>acs</i> ::Km-1) Δ (<i>ackA pta hisJ hisP dhu</i>)	This study
AJW807	AJW804 <i>recA</i> :: <i>cml</i>	This study
AJW810	AJW807 pSR31	This study
AJW811	AJW807 pSR10	This study
BL21(λ DE3)	<i>hsdS gal ompT r_B⁻ m_B⁻ [λC1857(Ts) <i>ind1 Sam7 nin5 lacUV5-T7 gene 1]</i></i>	50
BMH71-18 <i>mutS</i>	<i>thi supE</i> Δ (<i>lac-proAB</i>) <i>mutS</i> ::Tn10/ <i>F'</i> <i>proA⁺B⁺ lacI^qZ</i> Δ M15	26
CP875	Δ <i>lacX74 thi-1 thr-1</i> (Am) <i>leuB6 metF159</i> (Am) <i>rpsL136 lacY</i>	39
CP911	CP875 Δ (<i>ackA pta hisJ hisP dhu</i>)	39
DH5 α	<i>supE44</i> Δ (<i>argF-lac</i>)U169 (ϕ 80 <i>lacZ</i> Δ M15) <i>hsdR17 recA1 endA1 gyrA96 thi-1 relA1</i>	18
JM109	<i>supE44</i> Δ (<i>lac-proAB</i>) <i>hsdR17 recA1 endA1 gyrA96 thi-1 relA1</i>	58
SE3001	Δ (<i>malK-lamB</i>)1 <i>araD139</i> Δ (<i>argF-lac</i>)U169 <i>rpsL150 relA1 flhD5301 deoC1 ptsF25 rbsR</i>	44

^a Δ (*ackA pta hisJ hisP dhu*) is linked to *zej*::Tn10, which confers Tc^r.

ing conditions, the cells were harvested by centrifugation at $3,000 \times g$ for 20 min at 4°C. Purification of the fusion protein and cleavage of His-Tag were performed according to the manufacturer's instructions (Novagen [20]) except that CoA (533 μ M) was added to the elution and storage buffers to ensure that the purified protein retained activity.

SDS-polyacrylamide gel electrophoresis and immunoblotting. A 50- μ g sample of total protein was loaded onto a 10% polyacrylamide gel in the presence of 0.1% sodium dodecyl sulfate (SDS) and subjected to a current of 30 mA at 4 to 10°C in a Bio-Rad minigel apparatus (27) until the loading dye was about 1 cm from the bottom. The proteins were transferred electrophoretically overnight to a 0.45- μ m-pore-size nitrocellulose filter by using a Trans-Blot Cell (Bio-Rad, Hercules, Calif.). The nitrocellulose was blocked with 5% (wt/vol) nonfat dried milk in PBST (0.01 M sodium phosphate [pH 7.4], 150 mM NaCl, and 0.1% Tween 20) for 2 h at room temperature. The filter was washed for 5 min in PBST before being subjected to sequential incubation with rabbit anti-*Methanohalobium soehngenii* Acs (a gift from R. Eggen, Wageningen Agricultural University, Wageningen, The Netherlands) or rabbit anti-*E. coli* Acs (obtained by standard procedures [19]) for 1 h at room temperature and goat anti-rabbit immunoglobulin G (heavy- and light-chain specific) at appropriate dilutions for 1 h at room temperature. Finally, it was washed three times for 5 min each in PBST between each incubation step. Color development was achieved with 0.5 mg of 4-chloro-1-naphthol ml⁻¹ in PBS with 17% (vol/vol) methanol at a final concentration of 0.015% H₂O₂.

Southern blot analysis. DNA-DNA hybridization was performed by the method of Southern (47) as described by Davis et al. (10). Two probes, the 0.72-kb *BclI-BclI* fragment of pSR9 (Fig. 2) and the 1.3-kb *BamHI-BamHI* kanamycin cassette derived from pUC4K, were labelled with [³²P]dCTP by using the Ready-To-Go DNA labelling kit (Pharmacia). The 0.72-kb *BclI-BclI* probe was used to hybridize the *E. coli* Gene Mapping Membrane (TaKaRa Shuzo Co., Ltd.), which represents the entire *E. coli* chromosome. The 0.72-kb *BclI-BclI* and 1.3-kb *BamHI-BamHI* probes also were used to hybridize chromosomal DNA digested with either *EcoRV* or *EcoRI*. After hybridization for 16 h at 42°C, the blots were exposed to X-ray film at -70°C for 24 h.

Sequence analysis. Double-stranded DNA was purified with the Magic Mini-prep purification system (Promega) and denatured as described elsewhere (7). Single-stranded DNA was prepared by the method of Messing (33) with the Magic M13 DNA purification system (Promega). Both single- and double-stranded DNAs were sequenced by the chain termination method (43) using Sequenase, version 2.0 (U.S. Biochemicals, Cleveland, Ohio). Sequence analysis was performed with the software package (version 7.2) of the Genetics Computer Group (13a).

RESULTS

Genetic and molecular characterization. To determine whether the putative *acs* open reading frame actually encodes Acs (5), we subcloned the 2.2-kb *MunI-MunI* restriction fragment of the recombinant phage λ 638 of Kohara et al. (24) into pUC19 (53) to yield the plasmid pSR9. To verify the identity of the 2.2-kb *MunI-MunI* region of pSR9, we performed restriction enzyme analysis using seven different enzymes and sequenced about 300 bases from each of the 5' and 3' ends of the

fragment. The results of both analyses corresponded identically to those predicted by the sequence of Blattner et al. (5).

To provide genetic evidence that the putative *acs* open reading frame encodes Acs, we constructed the deletion allele Δ *acs*::Km and recombined it into the chromosome of the *PolA*(Ts) strain AJW763. To determine whether Δ *acs*::Km had recombined into the 92-min region of the chromosome, we mapped it relative to the neighboring genes, *malK* and *lamB*, which contribute to the ability of cells to utilize maltose as their sole carbon source and which map to the 91.5-min region (42). To do so, we used cells of the putative *acs*-deficient strain, AJW803, as the source of donor DNA to P1 transduce cells of the *LamB*- and *MalK*-deficient strain, SE3001, selecting for transductants resistant to kanamycin. We then screened kanamycin-resistant transductants for the ability to grow on minimal media supplemented with maltose (0.2%). The phenotypes Km^r and Mal⁺ cotransduced with a frequency of about 25% (46 of 179 Km^r transductants were also Mal⁺), indicating that Δ *acs*::Km mapped to the 90- to 93-min region.

Wu et al. (57) reported the sequence of a partial open reading frame called *orfX*, located 3' to *ubiG*, that shares about 1 kb of 100% identity with that reported by Blattner et al. (5). Whereas *ubiG* maps within the 50-min region of the chromosome (42), Blattner and coworkers located the putative *acs* open reading frame within the 92-min region. To distinguish between these two possible locations, we used the 0.72-kb *BclI-BclI* restriction fragment of pSR9 to probe the library of recombinant bacteriophages of Kohara et al. (24). This 0.72-kb restriction fragment, which was deleted to construct the Δ *acs*::Km allele, hybridized only with phages which carried the 92.3- to 92.4-min region, i.e., λ 638 and λ 639 (data not shown).

To determine whether the resultant recombinants carried a single putative *acs* allele, we performed a Southern blot analysis. The 0.72-kb *BclI-BclI* probe (Fig. 3A) hybridized to 4.5- and 2.5-kb *EcoRV* (lane 1) and 17.3-kb *EcoRI* (lane 3) chromosomal fragments of the wild-type strain, CP875; it did not hybridize with chromosomal fragments of the Δ *acs*::Km strain, AJW803 (lanes 2 and 4). In contrast, the 1.3-kb *BamHI-BamHI* probe (Fig. 3B), derived from plasmid pUC4K and used to mark the *acs* deletion, did not hybridize to chromosomal fragments of strain CP875 (lanes 1 and 3). However, it did hybridize to 7.6-kb *EcoRV* (lane 2) and 17.9-kb *EcoRI* (lane 4) chromosomal fragments from strain AJW803.




FIG. 3. Southern hybridization analysis demonstrating the location of the kanamycin-resistant deletion allele $\Delta acs::Km$. Chromosomal DNA, isolated from cells wild type for acetate activation (strain CP875) or their $\Delta acs::Km$ derivatives (strain AJW803), was digested with *EcoRV* or *EcoRI*, hybridized with the 0.72-kb *BclI-BclI* fragment of pSR9 (A), stripped, and rehybridized with the 1.1-kb *BamHI-BamHI* fragment of pUC4K (B). Lanes 1 and 2, *EcoRV* digests of DNA derived from strains CP875 and AJW803, respectively; lanes 3 and 4, *EcoRI* digests of DNA from strains CP875 and AJW803, respectively. Fragment sizes (in kilobases) are indicated on the left.

Phenotypic characterization. To characterize the phenotype of cells deleted for *acs*, we grew wild-type (strain CP875) cells or cells deficient for either *Acs* (strain AJW803) or *Ack* and *Pta* (strain CP911) at 37°C in TB to mid-exponential phase ($OD_{610} \sim 0.3$), washed and diluted them 10^{-5} in PBS (pH 7.4), plated them onto M63 minimal salts medium supplemented with a range of acetate concentrations (0 to 50 mM) as the sole carbon source, incubated them at 37°C for 60 h, and compared the resultant colony sizes (Fig. 4). Wild-type cells grew relatively well at all concentrations of acetate tested. Although cells of the putative *Acs*-deficient strain grew about as well as wild-type cells at higher concentrations (≥ 25 mM), they grew poorly on lower concentrations (≤ 10 mM). In contrast, cells of the *Ack*- and *Pta*-deficient strain grew best at lower concentrations (≤ 10 mM). Finally, cells deficient for *Acs* and *Ack*, *Acs* and *Pta*, or *Acs*, *Ack*, and *Pta* did not grow on any of the concentrations tested (data not shown).

We inoculated TB with wild-type (strain CP875) cells or cells deficient for *Acs* (strain AJW803), *Ack* and *Pta* (strain CP911), or *Acs*, *Ack*, and *Pta* (strain AJW804), aerated the cultures at 37°C, monitored their growth rates by optical density, and measured their extracellular acetate concentrations (Fig. 5). Wild-type and *Acs*-deficient cells grew at similar rates (~ 1.7 generations per h) and excreted similar amounts of acetate (~ 1.2 mM) prior to mid-exponential phase ($OD_{610} \leq 0.4$). Although the two cell types continued to grow at similar rates, their behavior with respect to extracellular acetate differed. Whereas the concentration of extracellular acetate of wild-type cells declined to a minimum of < 0.1 mM, that of the *Acs*-deficient cells did not decrease. In contrast, the growth of cells




FIG. 4. Growth on acetate as the sole carbon source of cells wild type for acetate activation (strain CP875) or deficient for either *Acs* (strain AJW803) or *Ack* and *Pta* (strain CP911). Cells were grown at 37°C in TB to an OD_{610} of ~ 0.3 , washed and resuspended in PBS, spread onto M63 minimal plates supplemented with the indicated concentration of acetate, incubated at 37°C for 72 h, and photographed.




FIG. 5. Optical density (A) and extracellular acetate concentration (B) plotted as a function of time for cells wild type for acetate activation (● and ○; strain CP875) or deficient for Acs (▼ and ▽; strain AJW803), Ack and Pta (■ and □; strain CP911), or Acs, Ack, and Pta (◆ and ◇; strain AJW804) grown in TB at 37°C.

deficient for Ack and Pta or for Acs, Ack, and Pta lagged substantially (between 2 and 3 h) before somewhat lower rates of growth (~ 1.4 generations per h) were attained. Neither cell type excreted appreciable amounts of acetate.

Next, we attempted to rescue growth on acetate of cells deficient for Acs, Ack, and Pta. We transformed cells deficient for Acs, Ack, Pta, and RecA (strain AJW807) with plasmids which expressed *acs* under the control of P_{acs} (pSR10) or P_{lac} (pSR31) to yield strains AJW811 and AJW810, respectively. We grew cells of the parental and transformant strains at 37°C in TB to mid-exponential phase (OD_{610} , ~ 0.3), washed and diluted them with PBS, and plated them onto M63 minimal medium supplemented with acetate (10 mM) and a range of IPTG concentrations (0 to 200 μ M). After 60 h of incubation at 37°C, the plates were removed and the resultant colony sizes

were compared (Fig. 6). In contrast to cells which expressed Acs from its native promoter, P_{acs} (strain AJW811), those of the parental strain (AJW807) did not grow. Cells which expressed Acs from the IPTG-inducible promoter, P_{lac} (strain AJW810), formed colonies only when grown in the presence of 25 to 100 μ M IPTG, forming the largest colonies in the presence of 50 μ M IPTG. They did not form colonies in the absence of IPTG or in the presence of 200 μ M IPTG (data not shown).

Enzymatic and immunological characterization. To test for Acs activity, we grew wild-type (strain CP875) cells or cells deficient for Acs (strain AJW803), Ack and Pta (strain CP911), or Acs, Ack, and Pta (strain AJW804) at 37°C in either TB or M63 minimal medium supplemented with acetate and glycerol. At various times, we harvested and washed cells and prepared crude lysates which were centrifuged and dialyzed to remove cellular debris and small molecules and assayed for the ability to convert acetate to one of its activated forms.

When grown in TB, cells which were wild type (strain CP875) or deficient for Acs (strain AJW803) activated acetate (present in the reaction mixture at 2.5, 5.0, or 10.0 mM) across the range of optical densities tested, i.e., OD_{610} s of 0.2, 0.4, and 0.6 (Table 2). Cells deficient for Ack and Pta (strain CP911), however, activated acetate only upon passing through mid-exponential phase ($OD_{610} = 0.4$). This activity was significantly reduced relative to those exhibited by wild-type cells and cells deficient for Acs. In contrast, cells deficient for all three enzymes (strain AJW804) exhibited no detectable acetate-activating activity.

Under all conditions tested, the activity of Ack and Pta seemed to be partially dependent on the presence of CoA. For example, when CoA was excluded from the reaction mixture, the activity of cells which expressed either Acs, Ack, and Pta (strain AJW875) or Ack and Pta (strain AJW803) decreased, on average, by about 70% relative to that observed when CoA was included. In contrast, the activity of Acs was completely dependent on the presence of CoA. Cells that expressed only Acs exhibited no detectable activity in the absence of CoA (data not shown).

When grown overnight in M63 minimal medium supplemented with acetate (50 mM) and glycerol (2.5 mM), cells which were wild type (strain CP875) or deficient for either Acs (strain AJW803) or Ack and Pta (strain CP911) activated acetate (present in the reaction mixture at 2.5, 5.0, or 10.0 mM; Table 3). Wild-type cells exhibited the highest activity, while cells deficient for Ack and Pta displayed the lowest. Cells deficient for Acs, i.e., those that expressed Ack and Pta, increased their activity relative to acetate concentration. In contrast, cells deficient for Ack and Pta, i.e., those that expressed Acs, did not change their activity relative to acetate concentration. Under all conditions tested, the activity exhibited by wild-type cells was approximately equal to the sum of the Ack-Pta and Acs activities exhibited by cells deficient for Acs or for Ack and Pta, respectively. In contrast, cells deficient for all three enzymes (strain AJW804) exhibited no detectable acetate-activating activity.

To verify the presence or absence of Acs expression in wild-type strains or strains deficient for Acs, we required a polyclonal antibody specific for *E. coli* Acs. To prepare such an antibody, we purified, to $>95\%$ purity, a 72-kDa His-Tag-tagged fusion protein (Fig. 7A). We verified the identity of this protein by Western immunoblot analysis using rabbit anti-*M. soehngenii* Acs as the primary antibody (Fig. 7B). Using this purified 72-kDa Acs protein, we generated rabbit anti-*E. coli* Acs antibody, which we used for Western immunoblot analysis (Fig. 8) of lysates derived from cells which were wild type




FIG. 6. Growth on acetate as the sole carbon source of cells deficient for Acs, Ack, and Pta (strain AJW807) transformed with either pSR10 (strain AJW811) or pSR31 (strain AJW810), plasmids that carry *acs* expressed by either its native promoter or the *lac* promoter, respectively. Cells were grown at 37°C in TB to an OD₆₁₀ of ~0.3, washed and resuspended in PBS, spread onto M63 minimal plates supplemented with 10 mM acetate and the indicated concentration of IPTG, incubated at 37°C for 72 h, and photographed.

(strain CP875) or deficient for Acs (strain AJW803), Ack and Pta (strain CP911), or Acs, Ack, and Pta (strain AJW804) and grown at 37°C in M63 minimal medium supplemented with acetate (10 mM) and glycerol (2.5 mM) to an OD₆₁₀ of ~0.2. Cells of strains that carried the wild-type *acs* allele, i.e., CP875 (Fig. 8, lane 1) and CP911 (lane 3), expressed a 72-kDa protein. In contrast, cells of strains that carried the deletion allele Δ *acs*::Km, i.e., AJW803 (Fig. 8, lane 2) and AJW804 (lane 4), did not.

Finally, we determined that the purified 72-kDa protein can activate acetate across a wide range of acetate concentrations (Fig. 9). This activity required the presence of CoA and was stable only if CoA (533 μ M) was present during purification (data not shown).

DISCUSSION

We conclude that *E. coli* cells possess a single gene that encodes Acs and that it maps within the 92.3- to 92.4-min region of the chromosome, as proposed by Blattner et al. (5). The following observations support this hypothesis. (i) The open reading frame, as predicted by the reported sequence, encodes a 72-kDa protein which cross-reacts with polyclonal antibody specific for Acs isolated from an unrelated species, *M. soehngenii*. (ii) Cells that express Ack and Pta, but not the

72-kDa protein, grow poorly on low concentrations of acetate; those that express the 72-kDa protein, but not Ack and Pta, grow poorly on high concentrations of acetate; and those that do not express any of the three proteins do not grow on any amount of acetate. (iii) Relative to wild-type cells, those that express Ack and Pta, but not the 72-kDa protein, do not activate acetate as well; those which express the 72-kDa protein, but not Ack and Pta, display even less activity; and those that do not express any of the three proteins exhibit no observable activity at all. (iv) Plasmids that express the 72-kDa protein restore growth on acetate to cells deficient for all three proteins. (v) The 0.72-kb *BclI-BclI* probe derived from pSR9 hybridized to only those recombinant phages, λ 638 and λ 639, that carry the 92.3- to 92.4-min region of the chromosome.

The unrelated species *E. coli* and *Bacillus subtilis* each possess and utilize both the Acs and the Ack-Pta pathways (3, 6, 8, 15, 16, 41). In contrast, some prokaryotes, e.g., *Alcaligenes eutrophus* (49, 52) and the *Methanothrix* spp. (25, 37), and apparently all eukaryotes (29) utilize only the Acs pathway. Other prokaryotes, e.g., the *Methanosarcina* spp. (1, 12, 28, 31, 51), use exclusively the Ack-Pta pathway.

If other organisms survive perfectly well using only one acetate activation pathway, then why do cells of *E. coli* and *B. subtilis* possess both pathways? Cells of both organisms excrete acetate during exponential aerobic growth in the presence of

TABLE 2. Acetate-activating activity in tryptone-grown cells which were either wild type or impaired for acetate activation^a

Strain	Sp act of cells at the indicated OD ₆₁₀ in acetate concn of:								
	2.5 mM			5.0 mM			10.0 mM		
	0.2	0.4	0.6	0.2	0.4	0.6	0.2	0.4	0.6
CP875 (Ack ⁺ Pta ⁺ Acs ⁺)	1.1 ± 0.1	2.6 ± 0.1	3.1 ± 0.1	2.9 ± 0.2	4.6 ± 0.1	6.8 ± 0.1	5.8 ± 0.2	7.4 ± 0.1	7.6 ± 0.1
AJW803 (Ack ⁺ Pta ⁺ Acs ⁻)	1.2 ± 0.2	2.2 ± 0.0	2.2 ± 0.1	2.9 ± 0.0	3.8 ± 0.1	4.8 ± 0.1	5.3 ± 0.1	7.6 ± 0.3	6.9 ± 0.2
CP911 (Ack ⁻ Pta ⁻ Acs ⁺)	BLD ^b	0.1 ± 0.0	0.2 ± 0.0	BLD	<0.1	0.4 ± 0.1	BLD	1.0 ± 0.2	1.3 ± 0.1
AJW804 (Ack ⁻ Pta ⁻ Acs ⁻)	BLD	BLD	BLD	BLD	BLD	BLD	BLD	BLD	BLD

^a Cells were harvested at OD₆₁₀s of 0.2, 0.4, and 0.6 after growth at 37°C in TB. Crude lysates were prepared and Acs activities were measured as described in Materials and Methods. Mean specific activities and standard errors of the means of independent duplicate experiments are expressed as milliunits per milligram, where 1 mU of activity catalyzes the formation of 1 nmol of acetohydroxamate per min.

^b BLD, below the level of detection, i.e., 0.01 mU mg⁻¹.

TABLE 3. Acetate-activating activity in acetate-grown cells which were wild type or impaired for acetate activation^a

Strain	Sp act at acetate concn (mM) of:		
	2.5	5.0	10.0
CP875 (Ack ⁺ Pta ⁺ Acs ⁺)	2.3 ± 0.4	4.3 ± 0.4	7.1 ± 0.5
AJW803 (Ack ⁺ Pta ⁺ Acs ⁻)	1.8 ± 0.2	3.5 ± 0.4	6.2 ± 0.0
CP911 (Ack ⁻ Pta ⁻ Acs ⁺)	0.9 ± 0.0	1.1 ± 0.0	0.9 ± 0.1
AJW804 (Ack ⁻ Pta ⁻ Acs ⁻)	BLD ^b	BLD	BLD

^a Cells were harvested at an OD₆₁₀ of 0.2 after growth at 37°C in M63 salts medium supplemented with 50 mM acetate and 2.5 mM glycerol. Crude lysates were prepared and Acs activities were measured as described in Materials and Methods. Mean specific activities and standard errors of the means of four independent experiments are expressed as milliunits per milligram, where 1 mU of activity catalyzes the formation of 1 nmol of acetoxyhydroxamate per min.

^b BLD, below the level of detection, i.e., 0.01 mU mg⁻¹.

excess carbon (11, 21, 23, 30, 38, 48). Acetate excretion (Fig. 5) and intracellular AcP accumulation by cells of *E. coli* throughout early-exponential-phase growth in tryptone broth depends on the presence of both Pta and Ack (39). Similarly, early-exponential-phase growth of *B. subtilis* cells in media containing high glucose concentrations results in both elevated *ackA* expression (15) and elevated Pta levels (40). Midway through exponential growth, cells of *E. coli* begin utilizing the acetate they had excreted previously, a process that requires Acs (6) (Fig. 5). Similarly, *B. subtilis* cells induce *acsA* expression as they enter stationary phase (14).

Whereas *B. subtilis* cells require Acs for growth or sporulation on acetate as the sole carbon source, *E. coli* cells can utilize either pathway to grow on acetate as the sole carbon source, albeit with significantly different affinities for acetate. Although *E. coli* cells which are wild type for both pathways grow about equally well across a 20-fold range of acetate con-




FIG. 7. Purified *E. coli* Acs isolated from strain BL21(DE3) transformed with a plasmid (pSR30) which expresses Acs from the T7 promoter. Lane 1, whole-cell extract prior to loading onto the nickel column; lane 2, column flowthrough; lane 3, column eluent; lane 4, column eluent with His-Tag cleaved with thrombin; lane 5, purified Acs from *Saccharomyces cerevisiae* mitochondria (Sigma). Molecular mass standards are labeled in kilodaltons to the right of the gels. (A) SDS-10% polyacrylamide gels stained with Coomassie blue. (B) Immunoblot analysis using rabbit anti-*M. soehngenii* Acs following SDS-PAGE (10% polyacrylamide). Note (lane 4) that cleavage by thrombin of the His-Tag-Acs fusion protein results in the spurious production of two additional, smaller products (molecular masses between 50 and 60 kDa).




FIG. 8. Immunoblot analysis using rabbit anti-*E. coli* Acs polyclonal antibody after SDS-PAGE (10% polyacrylamide) of lysates from cells wild type for acetate activation (strain CP875 [lane 1]) or deficient for Acs (strain AJW803 [lane 2]), Ack and Pta (strain CP911 [lane 3]), or Acs, Ack, and Pta (strain AJW804 [lane 4]). Purified *E. coli* Acs (lane 5) was also used. Molecular mass standards are labeled in kilodaltons to the right of the gel.

centrations, i.e., from 2.5 to 50 mM, those deficient for either pathway grow poorly at one end or the other of that range. Ack binds acetate poorly ($K_m = 7$ mM) (13), while Acs binds with much higher affinity ($K_m = 0.2$ mM) (6). Thus, one might predict that, on low acetate concentrations, cells which express only Ack and Pta would grow poorly, while those that express only Acs would grow well. We have shown this prediction to be correct. One also might expect that cells which express only Acs would grow about as well on higher concentrations of acetate as they do on lower ones. However, high concentrations seemingly inhibit growth of these cells. Such acetate concentrations are much larger than that required to achieve V_{max} . Perhaps Acs cannot process acetate sufficiently fast to reduce the toxicity associated with high acetate concentrations. Perhaps possession of both acetate activation pathways permits *E.*




FIG. 9. Activity of purified *E. coli* Acs plotted as a function of acetate concentration. The purification and assay conditions are described in Materials and Methods. In the absence of CoA, no activity was detected at all concentrations of acetate tested. Acs act., Acs activity.

coli cells to survive in environments that contain widely varying concentrations of acetate.

ACKNOWLEDGMENTS

We thank C. Park for alerting us to the work of Blattner and coworkers, F. R. Blattner for providing recombinant bacteriophage, R. Eggen for providing antiserum, Z. Livneh, Y. Blat, and R. Stewart for technical advice, and S. Maloy, T. Henkin, R. Barak, H. Falk-Krzesinski, and B. P. McNamara for critical reading of the manuscript.

This work was supported by the United States-Israel Binational Science Foundation (grant 93-00211). A.J.W. also was supported by Public Health Service grant GM46221 from the National Institute of General Medical Sciences, and S.K.R. was supported by a Dean's Fellowship in Molecular Biology from Loyola University Chicago.

REFERENCES

- Aceti, D., and J. G. Ferry. 1988. Purification and characterization of acetate kinase from acetate grown *Methanosarcina thermophila*. Evidence for regulation of synthesis. *J. Biol. Chem.* **263**:15444–15448.
- Barak, R., M. Welch, A. Yanovsky, K. Oosawa, and M. Eisenbach. 1992. Acetyladenylate or its derivative acetylates the chemotaxis protein CheY in vitro and increases its activity at the flagellar switch. *Biochemistry* **31**:10099–10107.
- Berg, P. 1956. Acyl adenylates: an enzymatic mechanism of acetate activation. *J. Biol. Chem.* **222**:991–1013.
- Bergmeyer, H. U., and H. Möllering. 1966. Enzymatische Bestimmung von Acetat. *Biochem. Z.* **344**:167–189.
- Blattner, F. R., V. Burland, G. Plunkett III, H. J. Sofia, and D. L. Daniels. 1993. Analysis of the *Escherichia coli* genome. IV. DNA sequence of the region from 89.2 to 92.8 minutes. *Nucleic Acids Res.* **21**:5408–5417.
- Brown, T. D. K., M. C. Jones-Mortimer, and H. L. Kornberg. 1977. The enzymic interconversion of acetate and acetyl-coenzyme A in *Escherichia coli*. *J. Gen. Microbiol.* **102**:327–336.
- Chen, E. Y., and P. H. Seeberg. 1985. Laboratory methods. Supercoil sequencing: a fast and simple method for sequencing plasmid DNA. *DNA* **4**:165–170.
- Chou, T. C., and F. Lipmann. 1952. Separation of acetyl transfer enzymes in pigeon liver extract. *J. Biol. Chem.* **196**:89–103.
- Dailey, F. E., and H. C. Berg. 1993. Change in direction of flagellar rotation in *Escherichia coli* mediated by acetate kinase. *J. Bacteriol.* **175**:3236–3239.
- Davis, L. G., M. D. Dibner, and J. F. Battey. 1986. Basic methods in molecular biology. Elsevier Science Publishing Co., Inc., New York.
- El-Mansi, E. M. T., and W. H. Holms. 1989. Control of carbon flux to acetate excretion during growth of *Escherichia coli* in batch and continuous culture. *J. Gen. Microbiol.* **135**:2875–2883.
- Fischer, R., and R. K. Thauer. 1988. Methane formation from acetyl phosphate in cell extracts of *Methanosarcina barkeri*. *FEBS Lett.* **228**:249–253.
- Fox, D. K., and S. Roseman. 1986. Isolation and characterization of homogeneous acetate kinase from *Salmonella typhimurium* and *Escherichia coli*. *J. Biol. Chem.* **261**:13487–13497.
- Genetics Computer Group. 1991. Software package, version 7.2. University of Wisconsin, Madison.
- Grundy, F. J., A. J. Turinsky, and T. M. Henkin. 1994. Catabolite regulation of *Bacillus subtilis* acetate and acetoin utilization genes by CcpA. *J. Bacteriol.* **176**:4527–4533.
- Grundy, F. J., D. A. Waters, H. G. Allen, and T. M. Henkin. 1993. Regulation of the *Bacillus subtilis* acetate kinase gene by CcpA. *J. Bacteriol.* **175**:7348–7355.
- Grundy, F. J., D. A. Waters, T. Y. Takova, and T. M. Henkin. 1993. Identification of genes involved in utilization of acetate and acetoin in *Bacillus subtilis*. *Mol. Microbiol.* **10**:259–271.
- Guranoski, A., M. A. G. Sillero, and A. Sillero. 1994. Adenosine 5'-tetraphosphate and adenosine 5'-pentaphosphate are synthesized by yeast acetyl coenzyme A synthetase. *J. Bacteriol.* **176**:2986–2990.
- Hanahan, D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**:557–580.
- Harlow, E., and D. Lane. 1988. Antibodies: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Hoffmann, A., and R. G. Roeder. 1991. Purification of his-tagged proteins in non-denaturing conditions suggests a convenient method for protein interaction studies. *Nucleic Acids Res.* **19**:6337–6338.
- Holms, W. H. 1986. The central metabolic pathways of *Escherichia coli*: relationship between flux and control at a branch point, efficiency of conversion to biomass, and excretion of acetate. *Curr. Top. Cell. Regul.* **28**:69–105.
- Jones, M. E., and F. Lipmann. 1955. Aceto-CoA-kinase. *Methods Enzymol.* **1**:585–595.
- Kleman, G. L., and W. R. Strohl. 1994. Acetate metabolism by *Escherichia coli* in high-cell-density fermentation. *Appl. Environ. Microbiol.* **60**:3952–3958.
- Kohara, Y., K. Akiyama, and K. Isono. 1987. The physical map of the whole *Escherichia coli* chromosome: application of a new strategy for rapid analysis and sorting of a large genomic library. *Cell* **50**:495–508.
- Kohler, H. P. E., and A. J. B. Zehnder. 1984. Carbon monoxide dehydrogenase and acetate thiokinase in *Methanotrix soehngenii*. *FEMS Microbiol. Lett.* **21**:287–292.
- Kramer, B., W. Kramer, and H.-J. Fritz. 1984. Different base/base mismatches are corrected with different efficiencies by the methyl-directed DNA mismatch-repair system of *E. coli*. *Cell* **38**:879–887.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680–685.
- Laufer, K., B. Eikmanns, U. Frimmer, and R. K. Thauer. 1987. Methanogenesis from acetate by *Methanosarcina barkeri*: catalysis of acetate formation from methyl iodide, CO₂, and H₂ by the enzyme system involved. *Z. Naturforsch. Teil C* **42**:360–372.
- Lipmann, F. 1945. Acetylation of sulfanilamide by liver homogenate and extracts. *J. Biol. Chem.* **160**:173–190.
- Luli, G. W., and W. R. Strohl. 1990. Comparison of growth, acetate production, and acetate inhibition of *Escherichia coli* strains in batch and fed-batch fermentations. *Appl. Environ. Microbiol.* **56**:1004–1011.
- Lundie, L. L., Jr., and J. G. Ferry. 1989. Activation of acetate by *Methanosarcina thermophila*. *J. Biol. Chem.* **264**:18392–18396.
- McClary, W. R., J. B. Stock, and A. J. Ninfa. 1993. Is acetyl phosphate a global signal in *Escherichia coli*? *J. Bacteriol.* **175**:2793–2798.
- Messing, J. 1983. New M13 vectors for cloning. *Methods Enzymol.* **101**:20–78.
- Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Nyström, T. 1994. The glucose-starvation stimulon of *Escherichia coli*: induced and repressed synthesis of enzymes of central metabolic pathways and role of acetyl phosphate in gene expression and starvation survival. *Mol. Microbiol.* **12**:833–843.
- Park, C., and G. L. Hazelbauer. 1986. Mutation specifically affecting ligand interaction of the Trg chemosensory transducer. *J. Bacteriol.* **167**:101–109.
- Pellerin, P., B. Gruson, G. Prensier, G. Albagnac, and P. Debeire. 1987. Glycogen in *Methanotrix*. *Arch. Microbiol.* **146**:377–381.
- Prüß, B. M., J. M. Nelms, C. Park, and A. J. Wolfe. 1994. Mutations in NADH:ubiquinone oxidoreductase of *Escherichia coli* affect growth on mixed amino acids. *J. Bacteriol.* **176**:2143–2150.
- Prüß, B. M., and A. J. Wolfe. 1994. Regulation of acetyl phosphate synthesis and degradation, and the control of flagellar expression in *Escherichia coli*. *Mol. Microbiol.* **12**:973–984.
- Rado, T. A., and J. A. Hoch. 1973. Phosphotransacetylase from *Bacillus subtilis*: purification and physiological studies. *Biochim. Biophys. Acta* **321**:114–125.
- Rose, I. A., M. Grunberg-Manago, S. R. Korey, and S. Ochoa. 1954. Enzymatic phosphorylation of acetate. *J. Biol. Chem.* **211**:737–756.
- Rudd, K. E. 1992. Alignment of *E. coli* DNA sequences to a revised, integrated genomic restriction map, p. 2.3–2.43. In J. H. Miller (ed.), A short course in bacterial genetics: a laboratory manual and handbook for *Escherichia coli* and related bacteria. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
- Silhavy, T. J., M. L. Berman, and L. W. Enquist. 1984. Experiments with gene fusions. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Simons, F. H., and N. Kleckner. 1987. Improved single and multicopy lac-based cloning vectors for protein and operon fusions. *Gene* **53**:85–96.
- Smith, P. K., R. J. Krohen, G. T. Hermansen, A. K. Mallia, F. H. Gartner, M. D. Provenzano, E. K. Fujimoto, N. M. Goekle, B. J. Olsen, and D. C. Klenk. 1985. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **150**:76–85.
- Southern, E. M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* **98**:503–517.
- Speck, E. L., and E. Freese. 1973. Control of metabolite secretion in *Bacillus subtilis*. *J. Gen. Microbiol.* **78**:261–275.
- Steinbüchel, A., C. Fründ, D. Jendrossek, and H. G. Schlegel. 1987. Isolation of mutants of *Alcaligenes eutrophus* unable to derepress the fermentative alcohol dehydrogenase. *Arch. Microbiol.* **148**:178–186.
- Studier, F. W., A. H. Rosenberg, J. J. Dunn, and J. W. Dubendorff. 1990. Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.* **185**:60–89.
- Terlesky, K. C., M. J. Barber, D. J. Aceti, and J. G. Ferry. 1987. EPR properties of the Ni-Fe-C center in an enzyme complex with carbon monoxide dehydrogenase activity from acetate-grown *Methanosarcina thermophila*. *J. Biol. Chem.* **262**:15392–15395.
- Trüper, H. G. 1965. Tricarboxylic acid cycle and related enzymes in *Hydrogenomonas* strain H16G⁺ grown on various carbon sources. *Biochim. Biophys. Acta* **111**:565–568.
- Vieira, J., and J. Messing. 1982. The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* **19**:259–268.
- Wanner, B. L. 1992. Is cross regulation by phosphorylation of two-compo-

- nent response regulator proteins important in bacteria? *J. Bacteriol.* **174**: 2053–2058.
55. **Wolfe, A. J., M. P. Conley, and H. C. Berg.** 1988. Acetyladenylate plays a role in controlling the direction of flagellar rotation. *Proc. Natl. Acad. Sci. USA* **85**:6711–6715.
56. **Wolfe, A. J., B. P. McNamara, and R. C. Stewart.** 1994. The short form of CheA couples chemoreception to CheA phosphorylation. *J. Bacteriol.* **176**: 4483–4491.
57. **Wu, G., H. D. Williams, M. Zamanian, F. Gibson, and R. K. Poole.** 1992. Isolation and characterization of *Escherichia coli* mutants affected in aerobic respiration: the cloning and nucleotide sequence of *ubiG*. Identification of an *S*-adenosylmethionine-binding motif in protein, RNA, and small-molecule methyltransferases. *J. Gen. Microbiol.* **138**:2101–2112.
58. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:103–119.