

FIG. 3. *tra* box disruptions abolish TraR-dependent transcription. Primer extension reactions were as described for Fig. 2. (A) *traA* reverse transcripts detected by using RNA isolated from A136(pCF251)(pCF336) (lane 2) and A136(pCF251)(pCF278) (lane 4) cultured in ATGN supplemented with 0.4 μ M AAI. (B) *traC* reverse transcripts detected by using the same RNA as used for panel A. (C) *traI* reverse transcripts detected by using RNA prepared from A136(pCF251)(pCF278) (lane 2) and A136(pCF251)(pCF291) (lane 4). Lanes 1 and 3 in all panels are *tetA*-specific reverse transcription reactions, although in some cases, full-length *tetA* transcripts are not shown.

cultures after normalization against the control mRNA (Fig. 2B, lane 4).

Sequences showing some similarity to *Escherichia coli* σ^{70} promoters were identified upstream of the *traA* and *traC* transcriptional start sites (Fig. 1). Adjacent to the possible -35 site of each promoter was a perfect 18-bp inverted repeat, designated *tra* box I. This sequence is similar to the *lux* operator identified in *V. fischeri*, which provides a binding site for LuxR (8, 37). The position of this motif with respect to the *traA* and *traC* promoters (-43 and $-43/-45$, respectively) is highly similar to the position of the *V. fischeri lux* box with respect to the *luxI* promoter (-44 [11]).

A third, unlinked *tra* promoter is required for expression of an operon containing *traI* and 11 genes believed to direct synthesis of the conjugal mating pore (1, 14, 22) (Fig. 1). We localized this promoter by primer extension analysis using RNA isolated from A136(pVK211)(pCF251) and A136(pVK211)(pDH99). pVK211 is a cosmid containing *traI*, while pCF251 is a derivative of pDH99 that overexpresses TraR (14, 27). A 39-nucleotide reverse transcript was synthesized by using RNA from the former strain (Fig. 2C, lane 2) but was not detectable from the latter strain (Fig. 2C, lane 1). The apparent *traI* transcription start site is directly downstream from sequences resembling a σ^{70} promoter (Fig. 1). Like the *traC* and *traA* promoters, the putative -35 sequence for *traI* overlaps by one base an 18-bp inverted repeat centered at position -43 (Fig. 1). This sequence is identical to *tra* box I at all positions except one, and we designate this element *tra* box II.

A third *tra* box-like sequence was found upstream from *tra* box II (designated *tra* box III; Fig. 1). The possibility that this sequence has a role in *traI* expression is examined below. It also seemed possible that this site could have some role in expression of the nearby *rep* operon, which is expressed divergently from *traI* and is required for vegetative plasmid replication (14, 39). However, primer extension analysis did not reveal a transcript in this region divergent from *traI* (data not shown).

Disruptions of *tra* boxes I and II abolish TraR-AAI-dependent transcriptional activation. To determine whether *tra* boxes are required for induction of *traA*, *traC*, and *traI*, we mutagenized *tra* boxes I and II of the broad-host-range plasmids pCF336 and pCF278, respectively (see Materials and Methods). pCF336 contains the *Bam*HI-20a fragment of pTiA6, which contains the *traC* and *traA* promoters, while pCF278

contains the *Eco*RI-6 fragment, which contains the *traI* promoter. Both plasmids have unique *Bg*II sites at the middle of *tra* box I and *tra* box II, respectively. They were treated with *Bg*II, DNA polymerase, and ligase to add four nucleotides at these sites. The resulting plasmids (pCF338 and pCF291) and their wild-type parents were introduced into the Ti plasmidless strain A136(pCF251) and assayed for production of *tra* transcripts in the presence of AAI. These mutations abolished detectable production of all three transcripts (Fig. 3, compare lanes 2 and 4 of each panel). The effect of the *tra* box II mutation was also tested by comparing A136(pCF251)(pCF278) and A136(pCF251)(pCF291) for the production of AAI. A culture supernatant of the former strain contained approximately 30 μ M AAI activity, while a supernatant from the latter strain contained less than 0.3 μ M AAI activity. The residual expression of *traI* from the mutant plasmid did not require TraR. This finding suggests strongly that *tra* box II is essential for TraR-dependent production of AAI.

Deletion analysis of the *traI* upstream region. The experiments described above did not rule out the possibility that a site upstream of these *tra* boxes was required for regulation and that the four-nucleotide insertions altered the helical phase of such a site with respect to its target promoter. To address this issue and to assess the relative roles of the two *tra* boxes in *traI* expression, we used PCR amplification to construct five resections of the *traI* promoter (Fig. 1, bottom) and used these resected promoters to express a *traI-lacZ* translational fusion. Three of the resulting plasmids (pCF372, pCF369, and pCF370) contained *tra* box II, and all three expressed similar levels of β -galactosidase in the presence of AAI (Table 1), indicating that the presence or absence of *tra* box III did not significantly affect *traI* expression. Plasmid pCF373, which lacks a portion of *tra* box II, expressed approximately 1% as much β -galactosidase as the three larger plasmids, although this low-level expression required TraR and AAI. Plasmid pCF371, which is missing virtually all of *tra* box II but retains an intact -35 element, did not express detectable β -galactosidase (Table 1). Note that plasmids pCF370, pCF373, and pCF371 span a region of only 12 nucleotides, thereby providing precise localization of this site. We conclude that *tra* box II is the sole *cis*-acting site essential for *traI* expression.

Ti plasmid conjugation efficiency is strongly stimulated by high numbers of conjugal donors. We sought to test directly the model that the promoters described above were regulated in a cell density-dependent manner. Strain R10(pCF240.113) contains wild-type Ti plasmid pTiR10 and a multicopy plasmid bearing a *traA-lacZ* fusion (14). We performed a series of filter matings in which different number of donors (from 10^4 to 10^8 donors per filter) were mixed with a large excess of conjugal recipients (see Materials and Methods). In the absence of

TABLE 1. Effects of upstream deletions on *traI-lacZ* expression

Plasmid ^a	Deletion end point	β -Galactosidase sp act (Miller units) in the presence of AAI ^b
pCF372	-136	831
pCF369	-103	580
pCF370	-50	833
pCF373	-40	5
pCF371	-33	<1

^a Upstream deletion construct tested in *A. tumefaciens* A136 harboring pCF218.

^b Crude AAI added to approximately 0.4 μ M (1%). In all cases, specific activity in the absence of AAI was <1 Miller unit.




FIG. 4. Donor density dependence of *traI* expression and conjugal transfer. (A) Strain R10(pCF240.113) was cultured in ATGN, serially diluted, mixed with 10^9 cells of strain C58CIRS, deposited on nitrocellulose filters, and incubated for 15 h on ATO agar (■) or ATO supplemented with 0.4 μ M AAI (●). Titers of donor bacteria and transconjugants were determined for filters. (B) Donors from the mating experiment in panel A were assayed for *traA-lacZ* expression by measuring total β -galactosidase activity per recovered donor and calculating total activity per 10^9 donor cells.

exogenous AAI, conjugation was undetectable when conjugal donors were inoculated at fewer than 3×10^7 cells per filter and was at least 10,000-fold more efficient at higher donor cell densities (Fig. 4A). In the presence of AAI, donor cells transferred the Ti plasmid with approximately equal efficiencies at all tested cell densities (Fig. 4A).

β -Galactosidase specific activity in donor cells was measured in the same experiment and plotted as a function of inoculated donor cells per filter. In the absence of exogenous AAI, β -galactosidase was weakly expressed when fewer than 3×10^7 donors were applied to the filter (Fig. 4B) and increased more than 25-fold by higher donor densities. Similar patterns of *tra* expression were found in the absence of conjugal recipients, while expression of a *lac* promoter control was not cell density stimulated (data not shown). This induction closely paralleled the conjugation efficiencies shown in Fig. 4A. Addition of AAI strongly stimulated β -galactosidase activity at low donor densities (Fig. 4B). Unexpectedly, however, *traA* expression was moderately stimulated by high donor densities even in the presence of exogenous AAI.

DISCUSSION

We have identified the transcriptional start sites of the *traA*, *traC*, and *traI* genes and found that all three promoters show an architecture strikingly similar to that of the *luxI* promoter of *V. fischeri*. All three *tra* promoters contain sequences similar to *E. coli* σ^{70} promoters and *tra* boxes that overlap rather weak

possible -35 elements (8, 15, 19). The *tra* boxes are similar in sequence and position to the LuxR binding site upstream of the *luxI* promoter and to a possible binding site for the *Pseudomonas aeruginosa* LasR protein that is just upstream of the *lasB* promoter (15, 19). TraR and other LuxR homologs have homology in their DNA binding regions with a larger class of transcription factors, the FixJ/NarL superfamily (7, 15). Several of these proteins are known to bind to sites upstream of their target promoters to activate transcription (16, 28, 33, 37). In contrast to the conserved positions of the *cis*-acting sites for TraR, LuxR, and LasR, the locations of binding sites for other members of this superfamily are highly variable.

The existence of a single *tra* box between *traA* and *traC* suggested that these genes could have the same *cis*-acting element. Although it remains possible that the *tra* box insertion mutations that we made acted by changing the phase of some site further upstream, we favor the interpretation that *tra* box I is the only essential *cis*-acting site required for *traA* and *traC* expression. If so, this is the first example of a LuxR-type protein strongly activating divergent promoters from a single operator. Although this may seem similar to the situation for the divergent *luxR* and *luxI* promoters, both of which can be activated by LuxR (10, 34), the organization of the *traA* and *traC* promoters is quite different. The *luxICDABEG* promoter is adjacent to the *lux* box, such that this site overlaps the -35 element by one nucleotide (8, 11). In contrast, the *luxR* promoter is located 144 bp away from the *lux* box. Furthermore, LuxR protein activates the *luxR* promoter weakly and only under certain conditions (34, 36). In contrast, the *traA* and *traC* promoters have perfect symmetry centered around *tra* box I and are both strongly activated by TraR (Fig. 1). This compact spacing of two promoters and a shared *cis*-acting site is highly unusual (2).

A truncated *V. fischeri* LuxR protein and RNA polymerase (RNAP) bind synergistically to the 20-bp *lux* box (8, 37). Assuming that *tra* boxes provide binding sites for TraR, the dyad symmetry of *tra* boxes suggests that TraR could bind as a dimer, as has been found for many other regulatory proteins (6, 14). The fact that the *traA* and *traC* promoters are positioned symmetrically with respect to *tra* box I leads us to speculate that RNAP bound to *traA* could make contacts to one bound TraR protomer, while RNAP bound to *traC* could make identical contacts to the other bound TraR protomer. RNAP might transcribe these promoters simultaneously, although the close spacing of these promoters might prevent their simultaneous occupancy. The position of *tra* boxes suggests that TraR may be a class II-type transcriptional activator as defined by Ishihama (23). However, detailed understanding of the activation mechanism and putative interactions with RNAP await further studies.

The conserved spacing between *tra* boxes and the *traA*, *traC*, and *traI* promoters suggests that interactions between putatively bound TraR and RNAP may be similar. A plasmid lacking the distal seven nucleotides of *tra* box II (pCF373) expressed *traI* at extremely low but still detectable levels, indicating that TraR can weakly recognize the promoter proximal half of this *tra* box. Similar data have been obtained for *V. fischeri*: 5' deletions removing part of the *lux* box reduced but did not abolish transcriptional activation via LuxR (8). Another possible TraR binding site, *tra* box III, is centered 132 bp upstream of the *traI* start site (Fig. 2). *tra* box III is the least similar to the consensus sequence, with one divergent nucleotide in each arm of the repeat. There is precedent for the existence of distal, regulatory sequences in *V. fischeri* *lux* gene regulation (8, 35). However, deletion of *tra* box III did not affect TraR-dependent activation of *traI*. Therefore, the func-

tion of *tra* box III, if any, remains unresolved. The proximity of this element to the *rep* operon suggested that it might have some role in vegetative replication, especially since coregulation of *tra* genes and genes required for vegetative replication has been reported in other plasmids (24). However, we could not detect a *rep* transcript originating near *tra* box III.

LuxR-LuxI-type regulatory systems are often hypothesized to activate target genes at high cell densities (25, 30, 38). Culturing *A. tumefaciens* on a solid medium dramatically increases *tra* gene expression and AAI synthesis (14, 42). We report here that TraR stimulated conjugation only at high densities of donor cells and that this density dependence was overcome by providing AAI exogenously. As expected, *traA* expression in the absence of exogenous AAI was strongly expressed at densities sufficient for efficient conjugation and poorly expressed at lower densities. Similar patterns of *tra* gene expression were observed in the absence of recipient cells, indicating that in contrast to pheromone-regulated conjugation systems (9), Ti plasmid conjugation is not responsive to signals released from recipient cells. Exogenous AAI caused *traA* to be expressed at lower cell densities, although expression was still somewhat stimulated by high donor densities (Fig. 4B). This was somewhat surprising, especially since AAI rendered conjugation by these same cells density independent. Residual cell density dependence of gene expression in the presence of AAI was observed in the absence of recipient cells for the *traA*, *traI*, and *traC* promoters (data not shown). However, this stimulation by high cell densities in the presence of AAI is evidently too mild to result in elevated conjugation efficiencies (Fig. 4A). It should be pointed out that these conjugation assays were quantitated over 4 orders of magnitude, while *traA* expression in the presence of AAI varied by less than 1 order of magnitude.

ACKNOWLEDGMENTS

We acknowledge Steve Farrand, John Helmann, and Valley Stewart for helpful discussions during the course of this study. We also thank Lu Wang for technical advice with the primer extensions.

This study was supported by National Institutes of Health grant GM42893 (S.C.W.) and postdoctoral fellowship GM15128 (C.F.).

REFERENCES

- Alt-Mörbe, J., J. S. Stryker, C. Fuqua, Stephen K. Farrand, and S. C. Winans. Unpublished data.
- Beck, C. F., and R. A. J. Warren. 1988. Divergent promoters, a common form of gene organization. *Microbiol. Rev.* **52**:318–326.
- Casadaban, M. J., J. Chou, and S. N. Cohen. 1980. In vitro gene fusions that join an enzymatically active β -galactosidase segment to amino-terminal fragments of exogenous proteins: *Escherichia coli* plasmid vectors for the detection and cloning of translational initiation signals. *J. Bacteriol.* **143**:971–980.
- Chen, C.-Y., and S. C. Winans. 1991. Controlled expression of the transcriptional activator gene *virG* in *Agrobacterium tumefaciens* by using the *Escherichia coli lac* promoter. *J. Bacteriol.* **173**:1139–1144.
- Close, T. J., D. Zaitlin, and C. I. Kado. 1984. Design and development of amplifiable broad-host-range cloning vectors: analysis of *Agrobacterium tumefaciens* plasmid pTiC58. *Plasmid* **12**:111–118.
- Collado-Vides, J., B. Magasanik, and J. D. Gralla. 1991. Control site location and transcriptional regulation in *Escherichia coli*. *Microbiol. Rev.* **55**:371–394.
- Da Re, S., S. Bertagnoli, J. Fourment, J.-M. Reyrat, and D. Kahn. 1994. Intramolecular signal transduction within the FixJ transcriptional activator: in vitro evidence for the inhibitory effect of the phosphorylatable regulatory domain. *Nucleic Acids Res.* **9**:1555–1561.
- Devine, J. H., G. S. Shadel, and T. O. Baldwin. 1989. Identification of the operator of the *lux* regulon from the *Vibrio fischeri* strain ATCC7744. *Proc. Natl. Acad. Sci. USA* **86**:5688–5692.
- Dunny, G. M., B. A. B. Leonard, and P. J. Hedberg. 1995. Pheromone-inducible conjugation in *Enterococcus faecalis*: interbacterial and host-parasite chemical communication. *J. Bacteriol.* **177**:871–876.
- Engebrecht, J., K. Nealon, and M. Silverman. 1983. Bacterial bioluminescence: isolation and genetic analysis of functions from *Vibrio fischeri*. *Cell* **32**:773–781.
- Engebrecht, J., and M. Silverman. 1987. Nucleotide sequence of the regulatory locus controlling expression of bacterial genes for bioluminescence. *Nucleic Acids Res.* **15**:10455–10467.
- Farrand, S. K. 1993. Conjugal transfer of *Agrobacterium* plasmids, p. 255–291. In D. B. Clewell (ed.), *Bacterial conjugation*. Plenum Publishing Corp., New York.
- Fuqua, C., M. Burbea, and S. C. Winans. 1995. Activity of the *Agrobacterium* Ti plasmid conjugal transfer regulator TraR is inhibited by the product of the *traM* gene. *J. Bacteriol.* **177**:1367–1373.
- Fuqua, W. C., and S. C. Winans. 1994. A LuxR-LuxI type regulatory system activates *Agrobacterium* Ti plasmid conjugal transfer in the presence of a plant tumor metabolite. *J. Bacteriol.* **176**:2796–2806.
- Fuqua, W. C., S. C. Winans, and E. P. Greenberg. 1994. Quorum sensing in bacteria: the LuxR/LuxI family of cell density-responsive transcriptional regulators. *J. Bacteriol.* **176**:269–275.
- Galinier, A., A.-M. Garnerone, J.-M. Reyrat, D. Kahn, J. Batut, and P. Boistard. 1994. Phosphorylation of the *Rhizobium meliloti* FixJ protein induces its binding to a compound regulatory region at the *fixK* promoter. *J. Biol. Chem.* **269**:23784–23789.
- Gambello, M. J., and B. H. Iglewski. 1991. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *J. Bacteriol.* **173**:3000–3009.
- Genetello, C., N. Van Larebeke, M. Holsters, A. De Picker, M. Van Montagu, and J. Schell. 1977. Ti plasmids of *Agrobacterium* as conjugative plasmids. *Nature (London)* **265**:561–563.
- Gray, K. M., L. Passador, B. H. Iglewski, and E. P. Greenberg. 1994. Interchangeability and specificity of components from the quorum-sensing regulatory systems of *Vibrio fischeri* and *Pseudomonas aeruginosa*. *J. Bacteriol.* **176**:3076–3080.
- Habeeb, L. F., L. Wang, and S. C. Winans. 1991. Transcription of the octopine catabolism operon of the *Agrobacterium* tumor-inducing plasmid pTiA6 is activated by a LysR homolog. *Mol. Plant-Microbe Interact.* **4**:379–385.
- Hwang, I., D. M. Cook, and S. K. Farrand. 1995. A new regulatory element modulates homoserine lactone-mediated autoinduction of Ti plasmid conjugal transfer. *J. Bacteriol.* **177**:449–458.
- Hwang, I., P.-L. Li, L. Zhang, K. R. Piper, D. M. Cook, M. E. Tate, and S. K. Farrand. 1994. TraI, a LuxI homologue, is responsible for production of conjugation factor, the Ti plasmid *N*-acylhomoserine lactone autoinducer. *Proc. Natl. Acad. Sci. USA* **91**:4639–4643.
- Ishihama, A. 1993. Protein-protein communication within the transcription apparatus. *J. Bacteriol.* **175**:2483–2489.
- Jagura-Burdzy, G., and C. M. Thomas. 1994. KorA protein of the promiscuous plasmid RK2 controls a transcriptional switch between divergent operons for plasmid replication and conjugative transfer. *Proc. Natl. Acad. Sci. USA* **91**:10571–10575.
- Jones, S., B. Yu, N. J. Bainton, M. Birdsall, B. W. Bycroft, S. R. Chhabra, A. J. R. Cox, P. Golby, P. J. Reeves, S. Stephens, M. K. Winson, G. P. C. Salmund, G. S. A. B. Stewart, and P. Williams. 1993. The *lux* autoinducer regulates the production of exoenzyme virulence determinants in *Erwinia carotovora* and *Pseudomonas aeruginosa*. *EMBO J.* **12**:2477–2482.
- Kingston, R. E. 1987. Primer extension, p. 4.8.1–4.8.3. In F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Stuhl (ed.), *Current protocols in molecular biology*. John Wiley & Sons, New York.
- Knauf, V. C., and E. W. Nester. 1982. Wide host range cloning vectors: a cosmid clone bank of an *Agrobacterium* Ti plasmid. *Plasmid* **8**:45–54.
- Li, J., S. Kustu, and V. Stewart. 1994. In vitro interaction of nitrate-responsive regulatory protein NarL with DNA target sequences in the *fdnG*, *narG*, *narK*, and *frdA* operon control regions of *Escherichia coli* K-12. *J. Mol. Biol.* **241**:150–165.
- McGowan, S., M. Sebahia, S. Jones, B. Yu, N. Bainton, P. F. Chan, B. Bycroft, G. S. A. B. Stewart, P. Williams, and G. P. C. Salmund. 1995. Carbapenem antibiotic production in *Erwinia carotovora* is regulated by CarR, a homologue of the LuxR transcriptional activator. *Microbiology* **141**:541–550.
- Passador, L., J. M. Cook, M. J. Gambello, L. Rust, and B. H. Iglewski. 1993. Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science* **260**:1127–1130.
- Pierson, L. S., V. D. Keppenne, and D. W. Wood. 1994. Phenazine antibiotic biosynthesis in *Pseudomonas aeruginosa* 30-84 is regulated by PhzR in response to cell density. *J. Bacteriol.* **176**:3966–3974.
- Piper, K. R., S. Beck von Bodman, and S. K. Farrand. 1993. Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction. *Nature (London)* **362**:448–450.
- Roggiani, M., and D. Dubnau. 1993. ComA, a phosphorylated response regulator protein of *Bacillus subtilis*, binds to the promoter region of *sfA*. *J. Bacteriol.* **175**:3182–3187.
- Shadel, G. S., and T. O. Baldwin. 1991. The *Vibrio fischeri* LuxR protein is capable of bidirectional stimulation of transcription and both positive and negative regulation of the *luxR* gene. *J. Bacteriol.* **173**:568–574.
- Shadel, G. S., and T. O. Baldwin. 1992. Identification of a distantly located

- regulatory element in the *luxD* gene required for negative autoregulation of the *Vibrio fischeri* *luxR* gene. *J. Biol. Chem.* **267**:7690–7695.
36. **Shadel, G. S., and T. O. Baldwin.** 1992. Positive autoregulation of the *Vibrio fischeri* *luxR* gene. *J. Biol. Chem.* **267**:7696–7702.
37. **Stevens, A. M., K. M. Dolan, and E. P. Greenberg.** 1994. Synergistic binding of the *Vibrio fischeri* LuxR transcriptional activator domain and RNA polymerase to the *lux* promoter region. *Proc. Natl. Acad. Sci. USA* **91**:12619–12623.
38. **Swift, S., M. K. Winson, P. F. Chan, N. J. Bainton, M. Birdsall, P. J. Reeves, C. E. D. Rees, S. R. Chhabra, P. J. Hill, J. P. Throup, B. W. Bycroft, G. P. C. Salmond, P. Williams, and G. S. A. B. Stewart.** 1993. A novel strategy for the isolation of *luxI* homologues: evidence for the widespread distribution of a LuxR:LuxI superfamily in enteric bacteria. *Mol. Microbiol.* **10**:511–520.
39. **Tabata, S., P. J. J. Hooykaas, and A. Oka.** 1989. Sequence determination and characterization of the replicator region in the tumor-inducing plasmid pTiB6S3. *J. Bacteriol.* **171**:1665–1672.
40. **Tempé, J., A. Petit, M. Holsters, M. Van Montagu, and J. Schell.** 1977. Thermosensitive step associated with transfer of the Ti plasmid during conjugation: possible relation to transformation in crown gall. *Proc. Natl. Acad. Sci. USA* **74**:2848–2849.
41. **Wang, L., J. D. Helmann, and S. C. Winans.** 1992. The *Agrobacterium tumefaciens* transcriptional activator OccR causes a bend at a target promoter, which is partially relaxed by a plant tumor metabolite. *Cell* **69**:659–667.
42. **Zhang, L., and A. Kerr.** 1991. A diffusible compound can enhance conjugal transfer of the Ti plasmid in *Agrobacterium tumefaciens*. *J. Bacteriol.* **173**:1867–1872.
43. **Zhang, L., P. J. Murphy, A. Kerr, and M. E. Tate.** 1993. *Agrobacterium* conjugation and gene regulation by *N*-acyl-homoserine lactones. *Nature (London)* **362**:446–448.