

Sequencing, Expression, and Genetic Characterization of the *Helicobacter pylori* *ftsH* Gene Encoding a Protein Homologous to Members of a Novel Putative ATPase Family

ZHONGMING GE AND DIANE E. TAYLOR*

Department of Medical Microbiology and Immunology, University of Alberta, Edmonton, Alberta, Canada T6G 2H7

Received 6 June 1996/Accepted 26 August 1996

In this study, we isolated and sequenced a *Helicobacter pylori* gene, designated *ftsH*, coding for a 632-amino-acid protein which displayed striking similarity throughout its full length to FtsH proteins identified in *Escherichia coli*, *Lactococcus lactis*, and *Bacillus subtilis*. *H. pylori* FtsH also possessed a ~200-amino-acid region containing a putative ATPase module which is conserved among members of the AAA protein family (AAA, ATPase associated with diverse cellular activities). The *H. pylori* *ftsH* product was overexpressed in *E. coli* and reacted immunologically with an anti-*E. coli* FtsH serum (T. Tomoyasu, K. Yamanaka, K. Murata, T. Suzaki, P. Bouloc, A. Kato, H. Niki, S. Hiraga, and T. Ogura, *J. Bacteriol.* 175:1352-1357, 1993). FtsH was also shown to be present in the membrane fraction of *H. pylori*, suggesting that it is membrane bound. Disruption of the *ftsH* gene led to the loss of viability of *H. pylori*, demonstrating that this gene is essential for cell growth. Overproduction of both *H. pylori* FtsH and *E. coli* FtsH together tremendously reduced the growth rate of the *E. coli* host cells, whereas the growth of the *E. coli* cells carrying the wild-type *E. coli* *ftsH* operon on the chromosome was not significantly affected by overproduction of *H. pylori* FtsH itself. This result suggests that the abnormal growth of cells results from interaction between *H. pylori* FtsH and *E. coli* FtsH.

Helicobacter pylori is an important cause of gastritis and has been associated with duodenal ulcer, gastric ulcers, and gastric cancer in some individuals (6, 7). Several factors have been implicated in the pathogenesis of *H. pylori*, including urease (12), flagella (20, 26, 36), a cytotoxin-associated gene A protein (CagA) (10, 42), an adhesin (14), and a vacuolating toxin (VacA) (11, 29, 33, 37). These proteins are either surface exposed (for example, flagella, CagA, and adhesin) or extracellular (for example, ureases and VacA). Hence, the process of how these proteins are either translocated into the membrane or secreted from the cells is essential for *H. pylori* pathogenesis. Recently, it has been reported that two *H. pylori* genes, *picA* and *picB*, play a role in the induction of an inflammatory response (43). On the basis of the sequence similarity between the *picB* gene product and the *Bordetella pertussis* toxin secretion protein, it was suggested that the *picB* gene product is involved in the induction of interleukin 8 expression in gastric epithelial cells by secreting an interleukin 8 inducer present in the *H. pylori* cells (43). However, the general processes of membrane translocation or protein secretion by *H. pylori* are poorly understood at present.

The FtsH (filamentation-temperature-sensitive) protein of *Escherichia coli* is membrane bound and essential for the viability of *E. coli* (28, 40, 41). This protein is thought to be involved in secretory protein export (1, 2) and has been demonstrated to play an important role in degradation of transcriptional regulatory factors, σ^{32} and λ CII (21, 22, 39), and SecY (an essential protein translocase subunit) (23). In addition, the *ftsH* gene, which is involved in tolerance to both high temperature and salt, has also been identified in *Bacillus subtilis* (13, 18) and *Lactococcus lactis* (27). Data indicating that the FtsH

proteins have multiple functions which are essential for the cell survival have accumulated increasingly. Moreover, these bacterial FtsH proteins belong to a new family of putative ATPases, known as the AAA (ATPases associated with diverse cellular activities) protein family (8, 9, 24). Members of this AAA protein family, found in both prokaryotes and eukaryotes, are believed to have diverse cellular activities (8, 24).

The goals of this study were (i) to clone and sequence an *H. pylori* *ftsH* gene which encodes a protein homologous to members of the AAA family, (ii) to study gene expression in *E. coli*, (iii) to localize FtsH in *H. pylori* by using an anti-*E. coli* serum, and (iv) to determine if this gene product is essential for the viability of *H. pylori*.

MATERIALS AND METHODS

Bacterial strains and media. A clinical *H. pylori* isolate, UA802, was used in this study and cultured as described previously (16). *E. coli* strains are listed in Table 1.

LB medium (31), M9 medium (31), and a supplemented M9 medium (32) were used for growth of bacteria. Ampicillin (100 μ g/ml), chloramphenicol (20 μ g/ml), and kanamycin (20 μ g/ml) were added to these media for growth of plasmid-containing cells and for screening transformants.

Techniques for DNA manipulations. Chromosomal DNA from *H. pylori* was extracted as described by Ezaki et al. (15). Plasmid DNA was prepared by the method of Birnboim and Doly (5). Sequential deletions of recombinant plasmid DNA were then made with the Erase-a-Base system as described by the supplier (Promega, Madison, Wis.). Nucleotide sequence was determined by a thermocycling sequencing method with *Taq* DNA polymerase as described in the supplier's manual (Gibco BRL, Gaithersburg, Md.). Primers were also employed for such sequencing. PCR amplification was used to detect the *H. pylori* *ftsH* gene within other *H. pylori* strains as described previously (16).

Plasmid constructs. A genomic DNA library of *H. pylori* UA802 chromosomal DNA prepared with the λ DASH vector (Stratagene, La Jolla, Calif.) was detailed previously (16). A clone, designated P9, containing a fragment of ~20 kb from *H. pylori* UA802 is shown in Fig. 1. A ~4.6-kb fragment of interest was then subcloned into pBluescript SK⁺ (Stratagene) by digestion with *Xba*I and *Eco*RI, followed by ligation with the appropriately digested vector. A plasmid clone, pBShpC10, carrying the *H. pylori* *ftsH* gene and the partial *H. pylori* *copA* gene was selected for further study. Various plasmid constructs derived from pBShpC10 are listed in Table 1.

Expression of the *H. pylori* *ftsH* gene in *E. coli*. Expression of the *H. pylori* *ftsH* gene in CSR603 was carried out as described previously (32) but in CSRDE3

* Corresponding author. Mailing address: Department of Medical Microbiology and Immunology, 1-28 Medical Sciences Building, University of Alberta, Edmonton, Alberta, Canada. Phone: (403) 492-4777. Fax: (403) 492-7521. Electronic mail address: diane-taylor@ualberta.ca.

TABLE 1. Bacterial strains and plasmids

Strain or plasmid	Characteristics	Source or reference
<i>E. coli</i>		
DH10B	F ⁻ <i>mcrA</i> Δ[<i>mrr-hsdRMS-mcrBC</i>] φ80 <i>dlacZ</i> ΔM15 Δ <i>lacX74 deoR ecA1 ara</i> Δ139 Δ(<i>ara leu</i>)7697 <i>galU galK</i> λ ⁻ <i>rpsL endA1 nupG</i>	Gibco BRL
CSR603	<i>rescA1 uvrA6 phr1</i>	32
CSRDE3	CSR603 with DE3, λ prophage carrying the T7 RNA polymerase under the <i>lac</i> promoter control	48
YJ430	F ⁻ <i>araD139</i> Δ(<i>argF-lac</i>)U169 <i>rpsL150</i> (Str ^r) <i>relA1 flbB5301 deoC1 ptsF25 rbsR</i> Δ(<i>srl-recA</i>) 306::Tn10 <i>ftsH3::kan</i> [F' <i>lac</i> ⁺ <i>lacI</i> ^q /pSTD401 <i>ftsH</i> Cm ^r]	35
Plasmid		
pSTD401	pHSG575 with <i>E. coli ftsH</i> under the <i>lac</i> promoter control	35
pBluescript SK ⁺	Cloning vector (Ap ^r)	Stratagene
pBluescript KS ⁻	Cloning vector (Ap ^r)	Stratagene
pBShpC10	pBluescript SK ⁺ with a ~4.6-kb <i>XbaI-EcoRI</i> fragment (<i>H. pylori ftsH copA</i>)	This study
pBShpC10Nm	pBShpC10 with the insertion of a CAT cassette at a <i>NaeI</i> site	This study
pBShpC10Bm	pBShpC10 with the insertion of a CAT cassette at a <i>BamHI</i> site	This study
pBShpC10Nhm	pBShpC10 with the insertion of a CAT cassette at a <i>NheI</i> site	This study
pBShpC10H	pBluescript SK ⁺ with a ~2.6-kb <i>XbaI-HindIII</i> fragment containing <i>H. pylori ftsH</i> under the <i>lac</i> promoter control	This study
pBKHpC10	pBluescript KS ⁻ with a ~4.6-kb <i>XbaI-EcoRI</i> fragment containing <i>H. pylori ftsH</i> and <i>copA</i> under the T7 promoter control	This study
pBKHpC10H	pBluescript KS ⁻ with a ~2.6-kb <i>XbaI-HindIII</i> fragment containing <i>H. pylori ftsH</i> under the T7 promoter control	This study
pBKHpC10HN	pBKHpC10 with a deletion of a ~0.6-kb <i>NheI-HindIII</i> fragment	This study

with some modifications. CSRDE3 cells harboring a plasmid carrying the *H. pylori ftsH* gene (pBKHpC10H, pBKHpC10NH, or pBKHpC10) were grown in LB broth supplemented with 100 μg of ampicillin per ml to an optical density at 620 nm and then collected. For [³⁵S]methionine-labeled *H. pylori* FtsH synthesis, cells were washed twice with 1× M9 medium and resuspended in 1 ml of supplemented M9 medium. The cells were incubated for 1 h at 37°C with agitation and induced by IPTG (isopropyl-β-D-thiogalactopyranoside; 1 mM) for 30 min. Subsequently, 10 μl of a 20-mg/ml concentration of rifampin was added to the respective samples, and after 30 min, 5 μl of [³⁵S]methionine was added to the respective samples and cell growth was continued for 1 h. For the synthesis of nonradioactively labeled FtsH protein, cells were induced with 1 mM IPTG for 1 h in the presence of 50 μg of cold methionine per ml instead of [³⁵S]methionine and then harvested. The cells were boiled for 5 min in 1× cracking buffer containing 50 mM Tris-HCl (pH 6.8), 1% sodium dodecyl sulfate (SDS), 20 mM disodium ethylenediaminetetraacetic acid, 1% mercaptoethanol, and 10% glycerol and stored at -20°C.

Insertional mutagenesis and natural transformation. Plasmid mutants carrying the disrupted *H. pylori ftsH* gene were created by inserting the chloramphenicol acetyltransferase (CAT) cassette (47) at *NaeI* and *BamHI* sites (Table 1 and Fig. 1). An additional mutant was constructed by the insertion of the CAT cassette at a *NheI* site located immediately downstream of the *ftsH* gene. Subsequently, plasmid mutants were introduced into *H. pylori* UA802 by a natural transformation procedure (46).

Preparation of membrane and soluble fractions of *H. pylori* cells. Cells were grown to an optical density of 0.3 and harvested by centrifugation. The cells were then washed three times with 1× MOPS buffer (50 mM morpholinepropanesulfonic acid [pH 7.0], 5 mM EDTA, 1 mM dithiothreitol) and resuspended in the same buffer. The cells were cracked with a French press at 8,000 lb/in². The cell

extracts were centrifuged at 12,900 × g at 4°C for 10 min. The supernatants were collected and recentrifuged in a Beckman TL 100(rotor 100.2) ultracentrifuge at 60,000 rpm at 4°C for 1 h. The supernatant was collected as the cytoplasmic fraction. The membrane fraction was treated with 1 mM NaCl as described previously (27).

Immunoblotting. The protein samples were separated by a 10% SDS polyacrylamide gel (31) and electrophoretically blotted onto a nitrocellulose membrane (MSI). The membrane was probed with antiserum against *E. coli* FtsH with an enhanced chemiluminescence system (Amersham Canada Ltd., Ontario, Canada). The antiserum was kindly provided by Y. Akiyama and K. Ito (Institute for Virus Research, Kyoto University, Kyoto, Japan). Protein bands were visualized on X-ray film.

Analysis and accession number of the nucleotide sequence. The nucleotide sequence reported in this study was analyzed with the software package (version 8) of Genetics Computer Group (Madison, Wis.) and has been deposited in GenBank under accession number U59452.

RESULTS

Cloning and nucleotide sequence of the *H. pylori ftsH* gene.

A λDASH clone P9 was demonstrated to contain the *H. pylori copA* gene (16, 17) and its upstream sequence. Subsequently, the *EcoRI-XbaI* fragments derived from P9 were cloned into pBluescript SK⁺. A subclone, namely pBShpC10, was obtained and sequenced (Fig. 1). Figure 2 depicts the sequence containing an open reading frame (ORF) and its downstream 162 nucleotides. The ORF coding for 632 amino acid residues starts at nucleotide 59 and terminates at nucleotide 1938. A putative ribosome binding site or Shine-Dalgarno sequence (SD) was identified (34). A putative -10 region of a promoter was located in the upstream sequence of the start codon of the predicted ORF. A potential stem-loop structure followed by a string of T's (nucleotides 2056 to 2091) was located downstream of this ORF, which may represent a rho-independent transcription terminator (30).

Since the amino acid sequence deduced from the predicted ORF was highly homologous to various bacterial FtsH proteins as described below, this ORF was designated the *H. pylori ftsH* gene. A hydropathy profile of the deduced amino acid sequence of *H. pylori ftsH* calculated by the method of Kyte and Doolittle (25) showed that *H. pylori* FtsH contains two highly hydrophobic regions (Fig. 3, underlined residues 9 to 31 and 121 to 140). These regions could form two transmembrane




FIG. 1. Cloning of the *H. pylori ftsH* gene. A plasmid recombinant, pBShpC10, was generated by subcloning the *XbaI-EcoRI* fragment derived from a λDASH clone P9 into the vector pBluescript SK⁺. The corresponding locations of the gene *ftsH* (this study) and the copper-exporting gene *copA* (16, 17) are indicated as are the restriction enzymes used for constructing various plasmids.

```

XbaI
TCTAGAATTGCTTAAAAACAACCAATAAATTAAGGATTATAATGAAACCAACGAACCTAAAAAACCTTTTTTCAAAGTCCCATTTATCCTCGGG
- 10 SD FtsH M K P T N E P K K P F F Q S P I I L A V
100
TTCTTGGAGGGATTTTACTCAITTTTTTCTCGCTCTTTCAATTCGATGCGAGTTTTTCGGACAATTTCTTAGCTTCTAGCACTAAAAATGTGAGCTA
L G G I L L I F F L R S F N S D G S F S D N F L A S S T K N V S Y
200
CCATGAGATCAAAACAGCTCATCAGCAATAATGAAGTGGAAAAATGTGAGTATCGGTCAAACCTTTGATCAAAAGCCAGCCATAAAGAGGGCAACAATCGTGTG
H E I K Q L I S N N E V E N V S I G Q T L I K A S H K E G N N R V
300
ATTTATATCGCTAAACGAGTGCCTGATTTGACCTTAGTGCCTTTTGTAGACGAGAAAAAATCAATTTATCTGGTTTTAGCGAGTCTAACTTTTTTACGG
I Y I A K R V P D L T L V P L L D E K K I N Y S G F S E S N F F T D
400
ACATGCTAGGGTGGCTCATGCCATTTTATGTGATTTTAGGGCTATGGATGTTTTATGGCAAACCGCATGCAAAAAAATATGGTGGGGGTATTTTTGGCAT
M L G W L M P I L V I L G L W M F M A N R M Q K N M G G G I F G M
500
GGGAGTGGCAAAAACTCATTAACGCTGAAAAACCAATGTGCGTTTTAAATGACATGGCAGCAATGAAGAAGCCAAAGAAGTGGTAGAAATCGTA
G S A K K L I N A E K P N V R F N D M A G N E E A K E E V V E I V
600
GATTTCTTAAATACCCGTAACGATACGCCAATTTAGGGGCTAAAAATCCCTAAGGCGTGTATTTAGTAGGGCTCCAGGAACGGGTAAACCCCTTTTAG
D F L K Y P E R Y A N L G A K I P K G V L L V G P P G T G K T L L A
800
CCAAAGCGGTAGCCGGCAAGCGCATGTGCGTTTTTCTCTATGGGAGGAGCAGTTTTTCATTTGAAATGTTTTGTGGCTTAGGGGCAAGCAGGGTTAGGGA
K A V A G E A H V P F F S M G G S S F I E M F V G L G A S R V R D
900
TTTGTTTGAAACCGCTAAAAACAAGCCCTTAGCATCATTTTTTATGATGAAATGATGCAATACCGAAGAGCAGCGGCTGGAGGCATGATAAGCGGG
L F E T A K K Q A P S I I F I D E I D A I P K S R A A G G M I S G
1000
AACGATGAAAGAGAGCAAACTTAAACCGCTCTTAGCCGAAATGGATGGTTTTGGGAGCGAAAAACCGCCTGTGATTTCTTAGCTGCAACGAACCGCC
N D E R E Q T L N Q L L A E M D G F G S E N A P V I V L A A T N R P
1100
CTGAAATCTTGGATCCGGCTTTAATGCGTCCAGGGCGCTTTGATAGCAGGTTTTAGTGGATAAGCCTGATTTTAAATGGCAGGGTGGAAATCTTAAAGT
E I L D P A L M R P G R F D R Q V L V D K P D F N G R V E I L K V
1200
CGATATTAAGGCGTGAACCTCGCTAATGATGTAATTTGGAAGAAGTCCGCAAACTCACCAGCGGCTTGGCGAGCGGATTTAGCGAATATCATCAAT
D I K G V K L A N D V N F E E V A K L T A G L A G A D L A N I I N
1300
GAAGCCGCACTTTTAGCAGGACGAAACAACCAAAAAGAAGTCAAAACAGCATTAAAAAGAAGCGGTTGAAAGAGGGATTCAGGGTTAGAAAAGAAAA
E A A L L A G R N N Q K E V K Q Q H L K E A V E R G I A G L E K K S
1400
GCAGGCGCATCAGTCTTAAAGAAAAGAAAATCGTCGCCCTACCATGAAAAGCGGCATGCCGTGATTTCTGAAAATGACTAAAAGGAGCGCTAGGGTGAATAA
R R I S P K E K K I V A Y H E S G H A V I S E M T K G S A R V N K
1500
AGTTTCTATCATTTCCAAGGGCATGGCGCTTTAGGCTACACCCCTTAAACAGCCTGAAGAAAACAAATACTTGTGCAAAAACGCAACTCATCGCTGAA
V S I I P R G M A A L G Y T L N T P E E N K Y L M Q K H E L I A E
1600
ATTGATGTGCTTTTAGCGGGAGAGCGGCTGAAGATGCTTTTTGGAAGAAAATTTCTACCGGTGCGAGCAACGATTTAGAAAGGGGACTGATATTTATTA
I D V L L G R A A E D V F L E E I S T G A S N D L I I K
1700
AAGGCATGGTGTGTTACTACGGCATGAGCAGTGTGACGGGCTTATGTTATGAAAAGCAACGGAACCGCTTTTTAGGAGGTGGTACCGAAGCAGTAG
G M V S Y Y G M S S V S G L M V L E K Q R N A F L G G G Y G S S R
1800
GGAATTTAGCGAAAAAACCGCAGAAGAAATGGATCTTTTCAATTAATAATTTGCTAGAAGAGCGCTATGAGCATGTCAAACAAACCTTTAGCGCATATAGA
E F S E K T A E E M D L F I K N L L E E R Y E H V K Q T L S D Y R
1900
GAAGCGATTGAAATCATGGTCAAAGAAATGTTTGACAAAAGATGATTAACAGCGAAAGGGTGGCTGAAATCATCAGCGAATACGAAGCCGCCAACAAAT
E A I E I M V K E L F D P N V R E I T G E R V R E I I S E Y E A A N N L
2000
TAGAAAAGCGCTTTGATCCCTTTAGAAGAGCAAGCGATTAAGTCTAGCTTACAAACAGATTTTTGATAAGGGTTTAAAGCCTTATGATAAACATTTCT
E S R L I P L E E Q A S * NheI
2100
GTTTGTTTAAAGTTTTTTTTTAGGTTTTGTTTTCTAAAAACATGCTTATCAACAGCGTTATAAAGCGTTTGGCTTTTAAAGCCTTTTTTTTTGTGAGTTTT
Stem-loop

```

FIG. 2. Nucleotide sequence and the predicted amino acid residues of the cloned *H. pylori ftsH* gene. Several putative transcription and translation elements, including the -10 region of a promoter, ribosome binding sites (SD), and a transcription terminator, are underlined. The nucleotides are numbered on the right. The restriction recognition sites of *Xba*I and *Nhe*I are indicated in bold letters.

segments, which would correspond to the transmembrane segments found in the FtsH proteins of *E. coli*, *L. lactis*, and probably *B. subtilis* (Fig. 3) (27, 40, 41). In addition, a model of the membrane topology of *H. pylori* FtsH was generated by a method described previously (44, 45). In this model, two transmembrane helices were connected by a periplasmic loop and the C-terminal domain containing about 480 residues was exposed to the cytoplasm (data not shown). The cytoplasmic domain carries the putative ATP-binding motifs (19) and the conserved region of the AAA protein family (9, 24). This predicted membrane topology of *H. pylori* FtsH is consistent with those of the *E. coli* and *L. lactis* FtsH proteins (27, 40).

Sequence similarity of *H. pylori* FtsH protein to other proteins. The amino acid sequence of *H. pylori* FtsH was applied to search homologous proteins available in the databases. *H. pylori* FtsH displays a striking sequence identity (50 to 60%) to the AAA protein family (ranging from bacterial to human)

within a ~200-amino-acid ATPase module (8, 9, 24). Several members of this proposed protein family have been identified in bacteria including *E. coli*, *L. lactis*, *B. subtilis*, *Haemophilus influenzae*, *Synechocystis* sp. PCC6803, *Mycoplasma genitalium*, and *Mycobacterium leprae* (nucleotide sequence accession numbers P28691, P46469, P37476, U32758, D64000, U39732 and U00017, respectively). Among these known bacterial AAA proteins, *E. coli*, *L. lactis*, and *B. subtilis* FtsH proteins have been functionally characterized and the others were deduced from their nucleotide sequences. Therefore, in Fig. 3, only a sequence comparison of *H. pylori* FtsH with the *E. coli*, *L. lactis*, and *B. subtilis* FtsH proteins was presented. *H. pylori* FtsH exhibits a similar degree of sequence identity to each of these three FtsH proteins, namely, 41% identity and 60% similarity to *B. subtilis* FtsH, 41% identity and 58% similarity to *L. lactis* FtsH, and 40% identity and 61% similarity to *E. coli* FtsH. It is apparent that the amino acid residues in the putative

HpFtsH	-MKPTNEPKK	PFQSPILLA	VLGGILLIFF	LRSPNSDGSF	SDNFLASSTK	NVSYHEIKQL	59		
LlFtsH	MNNKQPKQG	NFVKNLIMWV	ILAIWVVG	NFFPSSN	-----	KISYSQMTK	52		
EcFtsH	-----	-MAKNLILN	LVIATVLMV	FQSGPGS	-----	ESNGR	KVDYSYTFQE	40	
BsFtsH	-----	M	NRVRFNTIFY	LLILLVIVG	VSYFQTS	-----	NPKTE	NMSYSTPIKN	43
HpFtsH	ISNNEVENVS	IGQTLIKASH	KEGNNRVYI	AKR---VPD	-----	-----	LTL	98	
LlFtsH	LDGNKINVT	MQPSDSLITV	TGEYKEPVK	KGTNNPFLG	NSSSEVKNPQ	AYIIPDTSV	112		
EcFtsH	VNNDQVREAR	INGREINVT	KDSNRYTTY	PVQ---DPK	-----	-----	LL	78	
BsFtsH	LDDGKVDVSV	VQPVRGVYEV	KQGLKNYDK	Q---YFLTH	VPEG-----	-----	KGA	86	
HpFtsH	VPLLDEK---	KINYSQFS	ESNFFDMLG	WLMPIILVIG	LWMPANRMQ	KN---MGGGI	150		
LlFtsH	KDIQNAAKSN	DVKLSVQVQS	SSGMVQLIS	YIIPMLLVG	IFWLMGGMQ	ARGGGGGNP	172		
EcFtsH	DNLLTKM---	VKVVGEFPE	EPFLLASIFI	SWFPMLLIG	WIFPMRQMQ	G---GGKGA	131		
BsFtsH	DQIFNALKKT	DVKVEPAQET	SG--WVTLT	TIIFVVIPI	LFFFLNQAQ	G---GGSRV	140		
HpFtsH	FCMG-SAKKL	INAEPNVRP	NMDAGNEAK	EEVVEIVDFL	KYPERYANLG	AKIPKGVLLV	209		
LlFtsH	MSFGKSRKQ	QDGKTSKVR	ADVAGSEEK	QELVEVDFL	RNPKYHDLG	ARIPAGVILL	232		
EcFtsH	MSFGKSKARM	LTEDQIKTTF	ADVAGDEAK	EEVAVLVEYL	REPSRFQKLG	GKIPKGVLLV	191		
BsFtsH	MNFGKSKAKL	YTEKKRVRK	KDVAGADEEK	QELVEVDFL	KDPRKFAELG	ARIPKGVLLV	200		
motif A				motif B					
HpFtsH	GFPGTGRKLL	AKAVAGEAHV	PFPSMGSSGF	IEMFVGLGAS	RVRDLFETAK	KQAPSIIFID	269		
LlFtsH	GFPGTGRKLL	AKAVAGEAGV	PFYISGSDP	VEMFVGVGAS	RVRDLFENAK	KTAPSIIFID	292		
EcFtsH	GFPGTGRKLL	AKA IAGEAKV	PFPTTSGSDP	VEMFVGVGAS	RVRDMFENAK	KAAPCIIFID	251		
BsFtsH	GFPGTGRKLL	AKACAGEAGV	PFPSISGSDP	VEMFVGVGAS	RVRDLFENAK	KNAPCLIFID	260		
HpFtsH	BIDAIKPKRA	AGGMISGNDE	REQTLNQLLA	EMDGFSGENA	PVIVLAATNR	PEILDPALMR	329		
LlFtsH	BIDAVGRQRG	AGLGG-GNDE	REQTLNQLLV	EMDGFQDDGN	SVIVIAATNR	SDVLDPALLR	351		
EcFtsH	BIDAVGRQRG	AGLGG-GHDE	REQTLNQLLV	EMDGFQGN-E	GIIIVIAATNR	PDVLDPALLR	309		
BsFtsH	BIDAVGRQRG	AGLGG-GHDE	REQTLNQLLV	EMDGFQAN-E	GIIIVIAATNR	ADILDPALLR	318		
HpFtsH	PGRFDRQVLV	DKPDPNGRVE	ILKVDIKGVK	LANDVNFEEV	AKLTAGLAGA	DLANIINEAA	389		
LlFtsH	PGRFDRKVLV	GAPDVKGRKA	VLKVHAKRNP	LASDVLHNV	ATQTPGYVGA	DLNVLNEAA	410		
EcFtsH	PGRFDRQVVV	GLPDRVGRGQ	ILKVHMRVVP	LAPIDAAAI	ARGTGPGSGA	DLANLVNEAA	369		
BsFtsH	PGRFDRQITV	DRPDIQVREA	VLKVHAKRNP	LDETWNKLSI	AMRTGPGSGA	DLNVLNEAA	378		
HpFtsH	LILAGRNQKE	VKQQLHKEAV	ERGIAGLEKK	SRISPKPEKK	IVAYHEAGHA	VISEMPTKGA	449		
LlFtsH	LVAARQNKKE	INAADIDEGM	DRAMAGPAK	DRIQSMRRE	IVAYHEAGHA	IVGLVLENGS	470		
EcFtsH	LFAARGNKRV	VSMVEFEKAK	DKTMMGAERR	SMVMTAOKK	STAYHEAGHA	IIGRLVPEHD	429		
BsFtsH	LVAARQNKKK	IDARDIDEAT	DRVIACPAK	SRVISKKERN	IVAYHEGGHT	VIGLVLEDEAD	438		
HpFtsH	RVNKVSITPR	GMAALGYTLN	TPEENKYLMO	KHELIAEIDV	LLGGRAAEDV	FLE--EISTG	507		
LlFtsH	TVRNVTVVPR	-GRIGGYMLA	LPDEIMQPT	NFLHQDLAS	LMGGRLGEEI	VFG--VATPG	527		
EcFtsH	PVHKVITVPR	-GRALGVVFP	LPEGDALISAS	RQKLESQIST	LYGGRLAEEI	IYGPHEVSTG	488		
BsFtsH	MVHKVITVPR	-GQAGGYAVM	LFREDRVYQT	KPELLDKIVG	LLGGRVAEEI	IFG--EVSTG	495		
HpFtsH	ASNDLERATD	IIGMVSYYG	MSSVGLMVL	EK--QRNFL	GQYGGSSREF	SEKTAEMDML	565		
LlFtsH	ASNDIEKATH	IARSMVTEYG	MSKKLMVSY	EG--DHQVFI	GRDYGQTKTY	SEATAVMIDD	585		
EcFtsH	ASNDIKVATN	IARNMVTQMG	DRAMAGPLLY	AE-EDEGEVFL	GRSVAKAKHM	SDETARIIDD	547		
BsFtsH	AHNDFORATN	IARRMVTQFG	MSEKGLPLQF	GQSGGGQVFL	GRDPNNEQNY	SQIATVEIDD	555		
HpFtsH	PIKNLLEERY	EHVKQTLSDV	REAIEMVKE	LPDKVEITGE	RVREIISEYE	AAMN-----	619		
LlFtsH	EVRLIGIEAY	DRAKEALPETH	REORATAEA	LLKVEITLDAK	QIMSLFVTKG	MPDEAAAEV	645		
EcFtsH	EVKALIERNY	NRARQLLTDN	MDTLHAMKDA	LMKVETIDAP	QIDLMARRD	VRPAGWREP	607		
BsFtsH	EIQRIKIECV	ERAKQLTLEN	RDKLELIAGT	LLKVEITLDAE	QIKHLTDHGT	LPERNFSDDE	615		
HpFtsH	-----LES-	-----RLIPLLE	QAS-----	-----	-----	-----	632		
LlFtsH	PEPKTFEESL	KDANANVDDP	SNINLYNGDE	KTDSKPEENK	EKSEDETAE	-----	695		
EcFtsH	KASNSGSD--	-----NG	SPKARPVDE	PRTPNNGTVM	SEQLDGK--	-----	644		
BsFtsH	GN-----	DD-----	VKNVLLTKTE	EKKDDTKE-	-----	-----	637		

FIG. 3. Sequence comparison of *H. pylori* FtsH with three other bacterial FtsH proteins. Amino acid sequences were aligned with the program of CLUSTAL W (38). Hp, *H. pylori*; Ll, *L. lactis*; Ec, *E. coli*; Bs, *B. subtilis*. The identical residues at the corresponding positions of these proteins are indicated by asterisks, whereas the conservative replacements are marked by dots. Gaps are introduced to optimize alignment. Two putative ATP-binding motifs (A and B) are indicated by letters in bold type (19). The putative transmembrane segments of *H. pylori* FtsH corresponding to those of the other FtsH proteins are underlined. The ~200-amino-acid ATPase module conserved by the AAA protein family is indicated by arrows.

ATPase module (Fig. 3) (~200 amino acid residues), which are well conserved by the AAA protein family (9, 24), are highly homologous among these four FtsH proteins, with up to 57% identity and 73% similarity. In addition, the putative Zn-binding motif HEXXH located in the C-terminal region of *E. coli* FtsH (39) was identified in *H. pylori* FtsH (residues 434 to 438) (Fig. 3).

Expression of *H. pylori* ftsH in *E. coli* and localization of *H. pylori* FtsH in *H. pylori*. A maxicell system (32) was used to detect a protein encoded by the *H. pylori* ftsH gene. By using a conventional CSR603 strain as a host, a protein with the molecular mass of approximately 68 kDa was produced in cells harboring plasmid pBShpC10H containing the *H. pylori* ftsH gene, but the protein was poorly expressed (data not shown). This result could be due to the fact that the upstream sequence of the *H. pylori* ftsH gene is too short to contain complete




FIG. 4. Expression of *H. pylori* FtsH in *E. coli* CSRDE3. The plasmid-encoded proteins were synthesized as detailed in Materials and Methods. Equal amounts of the protein extracts adjusted by the turbidity of the cultures were separated on a 10% SDS-polyacrylamide gel. Lanes: 1, pBKHpC10H; 2, no plasmid DNA; 3, pBKHpC10; 4, pBKHpC10NH; 5, pBluescript KS⁻. *H. pylori* FtsH (~68 kDa) is present in lanes 1, 3, and 4. Molecular mass markers (BRL) are labeled on the left.

transcription elements (Fig. 2). Subsequently, a modified CSR603 strain (designated CSRDE3), which carries a *plac*-controlled T7 DNA polymerase gene (48) on the chromosome, was used for *H. pylori* FtsH expression. Various plasmids in which the *H. pylori* ftsH gene was controlled by a T7 promoter were constructed (Table 1). The expression results are shown in Fig. 4. The plasmid constructs pBKHpC10H (Fig. 4, lane 1), pBKHpC10NH (lane 3), and pBKHpC10 (lane 4), which carried the entire *H. pylori* ftsH gene (detailed in Table 1), all gave rise to a specific protein band of ~68 kDa, whereas there was no such protein expressed in the cells with pBluescript KS⁻ (Fig. 4, lane 5) or without any plasmids (lane 2). The molecular mass of this protein is in agreement with that (67.5 kDa) of the deduced *H. pylori* FtsH. In addition, a protein of about 50 kDa was also weakly produced from the cells containing the *H. pylori* ftsH gene. This protein may result from nonspecific translation initiation in the region of nucleotides 510 to 550, where two possible translation signals exist (Fig. 2, SD? and AUG). Two lines of evidence support this possibility: (i) translation initiation in this region is capable of encoding ~500 amino acid residues, which is consistent with the size of the ~50-kDa protein; and (ii) this protein reacted immunologically with the anti-FtsH serum as described below. Furthermore, a protein of ~35 kDa was present in the cells harboring both a pBluescript plasmid and its derivatives but not in the cells without the plasmid (Fig. 4, lane 2), indicating that this likely is β -lactamase.

Since putative *H. pylori* FtsH and *E. coli* FtsH exhibit a striking degree of sequence similarity, they should contain some identical antigenic epitopes. An antiserum, which was raised against a synthetic peptide corresponding to amino acid residues 297 to 312 of *E. coli* FtsH (40), was applied to detect the *H. pylori* FtsH protein in the maxicells. The *H. pylori* FtsH was expressed with cold methionine instead of [³⁵S]methionine, separated on a SDS-polyacrylamide gel, and blotted onto a nitrocellulose membrane. Although there are three substitutions in the 15 residues used to produce the anti-FtsH




FIG. 5. Immunological detection of *H. pylori* FtsH in the cell fractions of *H. pylori* and in the *H. pylori ftsH*-containing *E. coli* cells with the anti-*E. coli* FtsH serum. Ten micrograms of proteins prepared from *H. pylori* cells (lanes 1 to 3) were subjected to immunoblotting, whereas fixed amounts (determined by the turbidity of the cells) of protein extracts prepared from *E. coli* cells (lanes 4 to 6) were used. The protein samples were separated on a 10% SDS-polyacrylamide gel. Lanes: 1, cytoplasmic proteins; 2, salt-treated membrane supernatant; 3, membrane proteins; 4, pBKHpC10H; 5, pBluescriptSK⁻; 6, pSTD401. The positions of *H. pylori* FtsH (~68 kDa) and *E. coli* FtsH (~72 kDa) are indicated by an arrowhead and an arrow, respectively. Molecular mass markers (Gibco-BRL) are labeled on the left.

serum (Fig. 3, D by E, V by I, and L by M) between *H. pylori* and *E. coli* FtsH proteins, the antiserum still strongly reacted with the ~68-kDa *H. pylori* FtsH (Fig. 5, lane 4), whereas cells harboring pBluescript KS⁻ without the insert did not give rise to a band of the corresponding protein (Fig. 5, lane 5). Another protein, whose molecular mass appeared larger (~72 kDa in this gel) than that of *H. pylori* FtsH, was detected in all the *E. coli* cells by using this antiserum (Fig. 5, lanes 4 to 6). This 72-kDa protein must be *E. coli* FtsH, since this protein is identical to that generated from *E. coli* YJ430 harboring the plasmid pSTD401 (Fig. 5, lane 6). The minor protein of ~50 kDa expressed in cells containing the *H. pylori ftsH* gene was also detected by the same antiserum, indicating that this protein could represent a portion of the *H. pylori* FtsH protein as proposed above.

To investigate the localization of FtsH in the *H. pylori* cells, membrane and cytoplasmic fractions were prepared and probed with anti-*E. coli* FtsH serum. A protein of ~68 kDa was identified in the membrane fraction (Fig. 5, lane 3) but not in either the cytoplasmic fraction (Fig. 5, lane 1) or the salt-treated membrane supernatant (Fig. 5, lane 2). This result demonstrated that *H. pylori* FtsH is membrane bound, which is in agreement with that of *L. lactis* (27) and *E. coli* FtsH (40) proteins. The molecular mass of the *H. pylori* FtsH is consistent with that of the protein encoded by the predicted ORF (Fig. 5, lanes 3 and 4), indicating that the cloned gene represents the *H. pylori ftsH* locus.

***ftsH* is essential for the viability of *H. pylori*.** Three plasmid mutants, pBShpC10Nm, pBShpC10Bm, and pBShpC10Nhm, were constructed as detailed in Table 1. pBShpC10Nm and pBShpC10Bm were constructed by *H. pylori ftsH* inactivation with the CAT cassette, and pBShpC10Nhm was constructed by the insertion of the CAT cassette at a *NheI* site located immediately downstream of *ftsH*. By using a procedure for natural transformation as described previously (46), *H. pylori* mutants with the disrupted *ftsH* gene were not obtained, whereas the plasmid pBShpC10pNhm was integrated into the UA802 chromosome and an *H. pylori* mutant was obtained




FIG. 6. Effects of overexpressing *H. pylori* FtsH and *E. coli* FtsH on growth of cells. Cells of YJ430 ($\Delta ftsH::kan/pSTD401$) with or without an additional plasmid were grown at 37°C with agitation overnight in M9 glucose medium supplemented with Casamino Acids (0.2%), thiamine (2 μ g/ml), chloramphenicol (20 μ g/ml), ampicillin (50 μ g/ml), if appropriate, and IPTG (1 mM). Cells were then treated as described previously (35). Thereafter, the cells were grown in the same medium in the presence of 1 mM IPTG, and the turbidity of the cells was measured as optical densities (at 620 nm) with a microtiter plate reader. Symbols: squares, YJ430; triangles, YJ430 with pBluescript KS⁻; circles, YJ430 with pBShpC10H.

with a Cm^r phenotype. In addition, various *copA*-disrupted mutants of *H. pylori* have been created by the same technique (16, 17), and the *copA* gene is located at about 1 kb downstream of *ftsH* (Fig. 1). Therefore, our failure to generate the *H. pylori ftsH*-disrupted mutants is likely due to the fact that the normal function of *ftsH* is essential for the viability of *H. pylori*. In addition, conservation of the *FtsH* gene was investigated by PCR. Two primers were generated from the regions corresponding to nucleotides 1479 to 1499 and 1944 to 1962 (Fig. 2), respectively. Eighteen strains of *H. pylori* were subjected to PCR, and all gave rise to a specific product of ~640 nucleotides (data not shown). This result indicated that this gene is common to all *H. pylori* strains.

Cooverexpression of *H. pylori* and *E. coli ftsH* genes interferes with the normal growth of the *E. coli* host cells. Since *H. pylori* FtsH displays very strong sequence similarity to *E. coli* FtsH, we were interested in whether the *H. pylori ftsH* gene is functional in *E. coli* cells. The plasmids pBShpC10H and pBKHpC10H were introduced into *E. coli* DH10B and CSRDE3, respectively, which carry a normal *ftsH* gene on their chromosome. Overproduction of *H. pylori* FtsH in these strains did not significantly affect the growth of the host cells (data not shown). In contrast, pBShpC10H was introduced into *E. coli* YJ430 harboring a compatible pSC101-based plasmid, pST401, containing the *E. coli ftsH* gene under the *lac* promoter control. It was evident that such cells grew much more slowly than the cells harboring either pST401 alone or pST401 and pBluescript KS⁻ on LB agar plates supplemented with 1 mM IPTG. Growth rates of cells containing the respective plasmids in the presence of 1 mM IPTG, therefore, were monitored (Fig. 6). In comparison with cells harboring pST401, the pST401- and pBShpC10H-containing cells grew well in the first hour and




FIG. 7. Abundance of *H. pylori* FtsH and *E. coli* FtsH in cells induced with 1 mM IPTG as described in the legend to Fig. 6. Cells were harvested 8 h after the addition of IPTG. Fixed amounts of cell extracts adjusted by the turbidity of cultures were subjected to electrophoresis and then detected with the anti-*E. coli* FtsH antiserum. Lanes: 1, YJ430; 2, YJ430 with pBluescript KS⁻; 3, YJ430 with pBShpC10H.

then began to grow very slowly. In contrast, the growth curve of cells harboring pST401 and pBluescript KS⁻ was similar to that of cells containing pSTD401.

The levels of *H. pylori* FtsH and *E. coli* FtsH in these plasmid-containing cells were detected immunologically with the anti-*E. coli* FtsH serum as shown in Fig. 7. Approximately equal amounts of *E. coli* FtsH existed in the cells harboring pST401, pST401-pBShpC10pH, and pST401-pBluescript KS⁻. However, *H. pylori* FtsH of ~68 kDa was also accumulated in cells containing a *H. pylori* *ftsH* gene (Fig. 7, lane 3). The decrease of the cell growth rate is therefore associated with the overproduction of both of the FtsH proteins.

DISCUSSION

The *H. pylori* *ftsH* gene appears to encode a membrane-bound protein (FtsH) of 632 amino acid residues with the predicted two transmembrane segments. *H. pylori* FtsH exhibits a striking full-length sequence similarity to known bacterial FtsH proteins and carries a ~200-amino-acid ATPase module which has been highly conserved by the AAA protein family (8). As a result, the *H. pylori* FtsH protein belongs to this protein family.

The *H. pylori* *ftsH* gene plays an essential role in the growth of *H. pylori*, since disruption of this gene leads to a loss in viability of *H. pylori* cells. Similarly, the *E. coli* *ftsH* gene has been shown to be essential for cell viability (41). However, the *ftsH* genes isolated from *L. lactis* and *B. subtilis* seem dispensable for growth since their *ftsH*-knockout mutants are still viable (13, 18, 27). It should be noted that the sites used for the inactivation of the *L. lactis* and *B. subtilis* *ftsH* genes were close to their C termini so that the truncated FtsH proteins could still remain functional for cell growth. Thus, the results obtained from the knockout mutagenesis of the *L. lactis* and *B. subtilis* *ftsH* genes did not completely rule out the possibility that the *ftsH* genes are essential for the growth of *L. lactis* and *B. subtilis*.

Precise functions of *H. pylori* FtsH are unclear at present. Lethality associated with *ftsH* inactivation indicates that FtsH plays a crucial role in the life cycle of *H. pylori*. A high degree of sequence identity and similar patterns of the putative functional domains such as transmembrane segments and putative ATP-binding motifs between *H. pylori* FtsH and other bacterial FtsH proteins suggest that they have similar functions in these organisms. In *E. coli*, FtsH has been implicated in multiple cellular activities. Accumulation of penicillin-binding protein 3 and β -lactamase precursors in the *E. coli* *ftsH1* mutant indi-

icates that the maturation of these proteins is dependent on FtsH (4, 41). It was suggested that FtsH is involved in folding and assembly of proteins into and through the membrane, since mutations in FtsH lead to a dominant Std (stop-transfer-defective) phenotype, reduced retention of a cytoplasmic reporter SecY-PhoA fusion protein, and translocation retardation of some export proteins (1, 2). Recently, it has been shown that *E. coli* FtsH plays a crucial role in rapid degradation of three proteins, including the heat shock transcription factor σ^{32} (22, 39), the λ CII transcriptional activator (21), and the uncomplexed forms of SecY (3). In addition, the proteolytic activity of *E. coli* FtsH as an ATP- and Zn-dependent metalloprotease was demonstrated in vitro (39). On the other hand, *L. lactis* FtsH is able to functionally replace the *E. coli* FtsH to some extent and is involved in stress responses such as increased temperature and salt level (27). Similarly, an increased amount of the *B. subtilis* FtsH is essential for the cells in response to osmotic and temperature upshift (13).

Our results showed that overproduction of both *H. pylori* FtsH and *E. coli* FtsH significantly reduced the growth rate of host cells (Fig. 7). *E. coli* CSRDE3 cells containing the *E. coli* *ftsH* gene under its own promoter control are able to grow normally when *H. pylori* FtsH is overexpressed. How can this result be explained? It has been found that the high-level overexpression of *E. coli* FtsH is toxic to host cells (1, 3). A similar effect on cell viability may, therefore, be produced by the overaccumulation of the FtsH proteins encoded by both *H. pylori* *ftsH* and *E. coli* *ftsH*. This might be expected, since *H. pylori* FtsH is structurally and probably functionally related to *E. coli* FtsH. Alternatively, the decrease in growth rate could result from interference of the *H. pylori* FtsH protein with the formation of an FtsH complex, possibly located in the cytoplasmic membrane of *E. coli* (3). The previous study has shown that expression of the mutated *E. coli* FtsH even in the presence of the chromosomal wild-type *ftsH* gene reduces cytoplasmic retention of the alkaline phosphatase moiety attached to cytoplasmic regions of the membrane protein SecY, which is defined as a dominant Std phenotype (2). Recently, formation of the *E. coli* FtsH protein complex in the cytoplasmic membrane of *E. coli* has been demonstrated, indicating that *E. coli* FtsH may function as such a complex (3). Hence, it is possible that the *H. pylori* FtsH proteins in the cells are heterologously associated with *E. coli* FtsH to form inactive or malfunctioning complexes which interfere with the growth of the host cells. On the other hand, excess *H. pylori* FtsH in the *E. coli* cells may tend to form homomultimeric complexes. The host cell could maintain normal growth, which is regulated by the *E. coli* FtsH complexes and probably by the *H. pylori* FtsH multimeric complexes. Additional analysis of components of the FtsH complexes in the cells cooverproducing *H. pylori* FtsH and *E. coli* FtsH would be required to shed light on this hypothesis.

In summary, we cloned the *H. pylori* *ftsH* gene which has been highly conserved in *H. pylori* strains and is essential for the viability of *H. pylori*. More importantly, *H. pylori* FtsH could, like *E. coli* FtsH, have essential and multiple roles such as protein assembly, secretory protein export, and proteolytic activity. As a result, *H. pylori* FtsH would be involved in the export of some extracellular virulence factors so that this may greatly contribute to the pathogenesis of *H. pylori*. This point will be able to be addressed by using *H. pylori* Std and temperature-sensitive mutants created by mutations within the *H. pylori* *ftsH* gene.

ACKNOWLEDGMENTS

We thank K. Ito and Y. Akiyama for kindly providing the anti-*E. coli* FtsH serum and Q. Jiang for technical assistance.

This work was supported by funding from the Canada Bacterial Disease Network (Centers for Excellence Program). D.E.T. is a recipient of a Medical Scientific Award from the Alberta Heritage Foundation for Medical Research.

REFERENCES

- Akiyama, Y., T. Ogura, and K. Ito. 1994. Involvement of FtsH in protein assembly into and through the membrane. I. Mutations that reduce retention efficiency of a cytoplasmic reporter. *J. Biol. Chem.* **269**:5218–5224.
- Akiyama, Y., Y. Shirai, and K. Ito. 1994. Involvement of FtsH in protein assembly into and through the membrane. II. Dominant mutations affecting FtsH function. *J. Biol. Chem.* **269**:5225–5229.
- Akiyama, Y., T. Yoshihisa, and K. Ito. 1995. FtsH, a membrane-bound ATPase, forms a complex in the cytoplasmic membrane of *Escherichia coli*. *J. Biol. Chem.* **270**:23485–23490.
- Begg, K. J., T. Tomoyasu, W. D. Donachie, M. Khattar, H. Niki, K. Yamanaka, S. Hiraga, and T. Ogura. 1992. *Escherichia coli* mutant Y16 is a double mutant carrying thermosensitive *ftsH* and *ftsJ* mutations. *J. Bacteriol.* **174**:2416–2417.
- Birnboim, J. C., and J. Doly. 1979. Rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* **7**:1513–1523.
- Blaser, M. J. 1992. Hypothesis on the pathogenesis and natural history of *Helicobacter pylori*-induced inflammation. *Gastroenterology* **102**:720–727.
- Blaser, M. J., and J. Parsonnet. 1994. Parasitism by a "slow" bacterium *Helicobacter pylori* leads to altered homeostasis and neoplasia. *J. Clin. Invest.* **94**:4–8.
- Confalonieri, F., and M. Duguet. 1995. A 200-amino acid ATPase module in search of a basic function. *BioEssays* **17**:639–650.
- Confalonieri, F., J. Marsault, and M. Duguet. 1994. Sav, an archaeobacterial gene with extensive homology to a family of highly conserved eukaryotic ATPases. *J. Mol. Biol.* **235**:396–401.
- Covacci, A., S. Censini, M. Bugnoli, R. Petracca, D. Burroni, G. Macchia, A. Massone, E. Papini, Z. Xiang, N. Figura, and R. Rappuoli. 1993. Molecular characterization of the 128-kDa immunodominant antigen of *Helicobacter pylori* associated with cytotoxicity and duodenal ulcer. *Proc. Natl. Acad. Sci. USA* **90**:5791–5795.
- Cover, T. L., M. K. R. Tummuru, P. Cao, S. A. Thompson, and M. J. Blaser. 1994. Divergence of genetic sequences for the vacuolating cytotoxin among *Helicobacter pylori* strains. *J. Biol. Chem.* **269**:10566–10573.
- Cussac, V., R. L. Ferrero, and A. Labigne. 1992. Expression of *Helicobacter pylori* urease genes in *Escherichia coli* grown under nitrogen-limiting conditions. *J. Bacteriol.* **174**:2466–2473.
- Deuerling, E., B. Paeslack, and W. Schumann. 1995. The *ftsH* gene of *Bacillus subtilis* is transiently induced after osmotic and temperature upshift. *J. Bacteriol.* **177**:4105–4112.
- Evans, D. G., T. K. Karjalainen, D. J. Evans, D. Y. Graham, and C. Lee. 1993. Cloning, nucleotide sequence, and expression of a gene encoding an adhesin subunit protein of *Helicobacter pylori*. *J. Bacteriol.* **175**:674–683.
- Ezaki, T., N. Takeuchi, S. Liu, A. Kai, H. Yamamoto, and E. Yabuuchi. 1988. Small-scale DNA preparation for rapid genetic identification of *Campylobacter* species without radioisotope. *Microbiol. Immunol.* **32**:141–150.
- Ge, Z., K. Hiratsuka, and D. E. Taylor. 1995. Nucleotide sequence and mutational analysis indicate that two *Helicobacter pylori* genes encode a P-type ATPase and a cation-binding protein associated with copper transport. *Mol. Microbiol.* **15**:97–106.
- Ge, Z., and D. E. Taylor. Submitted for publication.
- Geisler, U., and W. Schumann. 1993. Isolation of stress mutants of *Bacillus subtilis* by a novel genetic method. *FEMS Microbiol. Lett.* **108**:251–254.
- Gorbalenya, A. E., E. V. Koonin, A. P. Domchenko, and V. M. Blinov. 1989. Two related superfamilies of putative helicases involved in replication, recombination, repair, and expression of DNA and RNA genomes. *Nucleic Acids Res.* **17**:4713–4729.
- Haas, R., T. F. Meyer, and J. P. M. van Puttern. 1993. A flagellated mutant of *Helicobacter pylori* generated by genetic transformation of naturally competent strains using transposon shuttle mutagenesis. *Mol. Microbiol.* **8**:753–760.
- Herman, C., T. Ogura, T. Tomoyasu, S. Hiraga, Y. Akiyama, K. Ito, R. Thomas, R. D'Ari, and P. Bouloc. 1993. Cell growth and λ phage development controlled by the same essential *Escherichia coli* gene, *ftsH/hflB*. *Proc. Natl. Acad. Sci. USA* **90**:10861–10865.
- Herman C., D. Thévenet, R. D'Ari, and P. Bouloc. 1995. Degradation of σ^{32} , the heat shock regulator in *Escherichia coli*, is governed by HflB. *Proc. Natl. Acad. Sci. USA* **92**:3516–3520.
- Kihara, A., Y. Akiyama, and K. Ito. 1995. FtsH is required for proteolytic elimination of uncomplexed forms of SecY, an essential protein translocase subunit. *Proc. Natl. Acad. Sci. USA* **92**:4532–4536.
- Kunau, W. H., A. Beyer, T. Franken, K. Götte, M. Marzochio, J. Saidowsky, A. Skalez-Rorowski, and F. F. Wiebel. 1993. Two complementary approaches to study peroxisome biogenesis in *Saccharomyces cerevisiae*: forward and reversed genetics. *Biochimie* **75**:209–224.
- Kyte, J., and R. F. J. Doolittle. 1982. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**:105–132.
- Leying, H., S. Suerbaum, G. Geis, and R. Haas. 1992. Cloning and genetic characterization of a *Helicobacter pylori* flagellin gene. *Mol. Microbiol.* **6**:2863–2874.
- Nilsson, D., A. A. Lauridsen, T. Tomoyasu, and T. Ogura. 1994. A *Lactococcus lactis* gene encodes a membrane protein with putative ATPase activity that is homologous to the essential *Escherichia coli* *ftsH* gene product. *Microbiology* **140**:2601–2610.
- Ogura, T., T. Tomoyasu, T. Yuki, S. Morimura, K. J. Gegg, W. D. Donachie, H. Mori, H. Noki, and S. Hiraga. 1991. Structure and function of the *ftsH* gene in *Escherichia coli*. *Res. Microbiol.* **142**:279–282.
- Phadnis, S. H., D. Ilver, L. Janzon, S. Normark, and T. U. Westblom. 1994. Pathological significance and molecular characterization of the vacuolating toxin gene of *Helicobacter pylori*. *Infect. Immun.* **62**:1557–1565.
- Platt, T. 1986. Transcription termination and the regulation of gene expression. *Annu. Rev. Biochem.* **55**:339–372.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Sancar, A., A. M. Hack, and W. D. Rupp. 1979. Simple method for identification of plasmid-coded proteins. *J. Bacteriol.* **137**:692–693.
- Schimitt, W., and R. Haas. 1994. Genomic analysis of the *Helicobacter pylori* vacuolating cytotoxin: structural similarities with the IgA protease type of exported protein. *Mol. Microbiol.* **12**:307–319.
- Shine, J., and L. Dalgarno. 1974. The 3'-terminal sequence of *Escherichia coli* 16S ribosomal RNA: complementarity to nonsense triplets and ribosome binding sites. *Proc. Natl. Acad. Sci. USA* **71**:1342–1346.
- Shirai, Y., Y. Akiyama, and K. Ito. 1996. Suppression of *ftsH* mutant phenotypes by overproduction of molecular chaperones. *J. Bacteriol.* **178**:1141–1145.
- Suerbaum, S., C. Josenhans, and A. Labigne. 1993. Cloning and genetic characterization of the *Helicobacter pylori* and *Helicobacter mustelae* flagellin genes and construction of *H. pylori* *flaA*- and *flaB*-negative mutants by electroporation-mediated allelic exchange. *J. Bacteriol.* **175**:3278–3288.
- Telford, J. L., P. Ghiara, M. Dell'Orco, M. Comanducci, D. Burroni, M. Bugnoli, M. F. Tecce, S. Censini, A. Covacci, Z. Xiang, E. Papini, C. Montecucco, L. Parente, and R. Rappuoli. 1994. Genome structure of the *Helicobacter pylori* cytotoxin and evidence of its key role in gastric disease. *J. Exp. Med.* **179**:1653–1658.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
- Tomoyasu, T., J. Gamer, B. Bukau, M. Kanemori, H. Mori, A. J. Rutman, A. B. Oppenheim, T. Yura, K. Yamanaka, H. Niki, S. Hiraga, and T. Ogura. 1995. *Escherichia coli* FtsH protein is a membrane-bound, ATP-dependent protease which degrades the heat-shock transcription factor σ^{32} . *EMBO J.* **14**:2551–2560.
- Tomoyasu, T., K. Yamanaka, K. Murata, T. Suzuki, P. Bouloc, A. Kato, H. Niki, S. Hiraga, and T. Ogura. 1993. Topology and subcellular localization of FtsH protein in *Escherichia coli*. *J. Bacteriol.* **175**:1352–1357.
- Tomoyasu, T., T. Yuki, S. Morimura, H. Mori, K. Yamanaka, H. Niki, S. Hiraga, and T. Ogura. 1993. The *Escherichia coli* FtsH protein is a prokaryotic member of a protein family of putative ATPases involved in membrane functions, cell cycle control, and gene expression. *J. Bacteriol.* **175**:1344–1351.
- Tummuru, M. K., T. L. Cover, and M. J. Blaser. 1993. Cloning and expression of a high-molecular-mass major antigen of *Helicobacter pylori*: evidence of linkage to cytotoxin production. *Infect. Immun.* **61**:1799–1809.
- Tummuru, M. K., S. A. Sharma, and M. J. Blaser. 1995. *Helicobacter pylori* *picB*, a homologue of the *Bordetella pertussis* toxin secretion protein, is required for induction of IL-8 in gastric epithelial cells. *Mol. Microbiol.* **18**:867–876.
- Turner, R., and J. H. Weiner. 1993. Evaluation of transmembrane helix prediction methods using the recently defined NMR structures of the coat proteins from bacteriophages M13 and Pf1. *Biochim. Biophys. Acta* **1202**:161–168.
- von Heijne, G. 1992. Membrane protein structure prediction: hydrophobicity analysis and the positive-inside rule. *J. Mol. Biol.* **225**:487–492.
- Wang, Y., K. P. Roos, and D. E. Taylor. 1993. Transformation of *Helicobacter pylori* by chromosomal metronidazole resistance and by a plasmid with a selectable chloramphenicol resistance marker. *J. Gen. Microbiol.* **139**:2485–2493.
- Wang, Y., and D. E. Taylor. 1990. Chloramphenicol resistance in *Campylobacter coli*: nucleotide sequence, expression, and cloning vector construction. *Gene* **94**:23–28.
- Whelan, K. F., R. K. Sherburne, and D. E. Taylor. Submitted for publication.