

An Unexpected *flaA* Homolog Is Present and Expressed in *Borrelia burgdorferi*

YIGONG GE AND NYLES W. CHARON*

Department of Microbiology and Immunology, Robert C. Byrd Health Sciences Center,
West Virginia University, Morgantown, West Virginia 26506-9177

Received 31 July 1996/Accepted 7 November 1996

Most investigators have assumed that the periplasmic flagella (PFs) of *Borrelia burgdorferi* are composed of only one flagellin protein. The PFs of most other spirochete species are complex: these PFs contain an outer sheath of FlaA proteins and a core filament of FlaB proteins. During an analysis of a chemotaxis gene cluster of *B. burgdorferi* 212, we were surprised to find a *flaA* gene homolog with a deduced polypeptide having 54 to 58% similarity to FlaA from other spirochetes. Like other FlaA proteins, *B. burgdorferi* FlaA has a conserved signal sequence at its N terminus. Based on reverse transcription-PCR and primer extension analysis, this *flaA* homolog and five chemotaxis genes constitute a motility-chemotaxis operon. Immunoblots using anti-FlaA serum from *Treponema pallidum* and a lysate of *B. burgdorferi* showed strong reactivity to a protein of 38.0 kDa, which is consistent with the expression of *flaA* in growing cells.

Lyme disease, which is caused by *Borrelia burgdorferi*, is the most common arthropod-borne human infection in the United States (1). *B. burgdorferi* has a morphology similar to that of other spirochetes; it has a protoplasmic cell cylinder and periplasmic flagella (PFs) surrounded by an outer membrane sheath (15, 17). Approximately seven PFs are attached at each end of the protoplasmic cell cylinder, and these PFs have been shown to be involved in motility (16, 17, 34). *B. burgdorferi* motility is likely to play a role in the development of Lyme disease. These organisms are highly motile in viscous gel-like environments such as connective tissue and thus can penetrate tissues when other bacteria fail to invade (16, 19, 20, 37). In addition, an analysis of a spontaneously occurring PF mutant has indicated that motility augments cell penetration (34). Finally, over 35 motility genes spanning approximately 35 kb have been identified in *B. burgdorferi* (GenBank accession no. L76303, L75945, U28962, U43739, U61498, U62900, U62901, and U66699) (12, 14); we have estimated that at least 3% of its genome, which is approximately 945 kb (7, 35), is involved in motility and chemotaxis.

The structure of PFs is distinct from that of the flagella of other motile bacteria. Specifically, the PFs of most spirochetes are composed of multiple protein species referred to as FlaA and FlaB proteins (4, 8, 21, 29, 30, 38). FlaA proteins form the PF sheath, and FlaB proteins comprise the core (4, 8, 21, 30). Within a given species, there are one or two different FlaA proteins and three or four different FlaB proteins (8, 21, 29, 30, 33, 38). FlaB proteins show amino acid sequence similarities to the flagellin proteins of rod-shaped bacteria (29, 30, 41). Because FlaB proteins contain no typical signal sequences, these proteins are evidently excreted by the flagellum-specific pathway (8, 27). On the other hand, FlaA proteins possess an N-terminal signal sequence recognized by the typical SecA-dependent excretion pathway (5, 8, 18, 22).

In contrast to other spirochetes, only one flagellin protein of 41 kDa has been reported in *B. burgdorferi* (3, 8, 9, 40). This

protein, encoded by the *fla* gene (hereafter referred to as the *flaB* gene), serves as a major antigen in the diagnosis of Lyme disease; high titers of anti-FlaB antibodies are detected in all disease stages (2, 9, 40). Based on electron microscopic observations (17) and sodium dodecyl sulfate-polyacrylamide gel electrophoresis profiles of purified PFs, no apparent PF sheath or proteins other than the 41-kDa protein have been identified. During an analysis of a chemotaxis gene cluster in *B. burgdorferi* 212, we were surprised to find a *flaA* homolog directly upstream of the putative *cheA* gene. Here we present our characterization of this *flaA* gene. We show that this gene is part of a putative motility-chemotaxis operon and that it is expressed in growing cells.

Trueba et al. recently sequenced and mapped a *cheA* homolog in *B. burgdorferi* B31 which mapped at 722 to 737 kb from the 0 telomere (39). We cloned and analyzed the region downstream of *cheA* in *B. burgdorferi* 212 and found four other putative chemotaxis genes, including *cheW*, *cheR*, *cheX*, and *cheY* (GenBank accession no. U61498) (11). Since motility and chemotaxis genes are generally clustered to form a functional operon, we hypothesized that there would be additional chemotaxis genes upstream of *cheA*. To clone and sequence the DNA segment upstream of the *cheA* gene, we used a method referred to as "semi-random PCR chromosome walking," which has been successfully used in the analysis of other motility operons of *B. burgdorferi* (10, 13). To sequence the upstream region, we first amplified cellular DNA under low-stringency conditions by using a primer (pcheA3) derived from the B31 *cheA* sequence (Fig. 1). We obtained a 0.7-kb fragment, and sequence analysis indicated that it was upstream of *cheA*. We then did a second PCR amplification under low-stringency conditions by using a primer, pflaA1, based on the sequence data obtained from the first 0.7-kb clone; we obtained a 0.9-kb fragment which was also upstream of *cheA*. Thus, the region extending approximately 1.6 kb upstream of *cheA* was cloned and sequenced. Immediately upstream of *cheA*, we identified a 1,023-bp open reading frame encoding a 341-amino-acid protein which has an estimated size of 38.0 kDa (Fig. 1). The GC content of this gene was 34%, which is similar to that found for the *B. burgdorferi* genomic DNA. The results of multiple alignments indicated that the protein encoded by this open reading frame was a well-conserved *flaA*

* Corresponding author. Mailing address: Department of Microbiology and Immunology, West Virginia University, Box 9177, Robert C. Byrd Health Sciences Center, Morgantown, WV 26506-9177. Phone: (304) 293-4170. Fax: (304) 293-7823. E-mail: charon@wvnet.wvnet.edu.


```

Sh MK_K L FVVLT SIFIAASAYGLTN ST LIDFA LTGNADNLQAGEGDTNEVVPVAENLYND
Sa MK_K AVVLSAVALLSGVCAVADES V LIDFAKLNADIMADKSGGMTHNRRRTVLDYASLADTSYTDEQ
Tp MKRF FFAILGAALFVGN SGAFAEQAT LIDFSKLVGE GNTGLHAPTTIDYSRQAGSAYS AED
Bb MKRKAKSILFFLLSTVLEFAQETDGLAEGSKRAEPGELVLDFAELARD PSSTRLDLTNYVDYVYS GAS
*.. ..*..... .. .. ..**.. .. .. .. ..*...

Sh N WVVVWLNESARLTENRRNSYVTNVDSKGNNGAW EAGKVLGVRVHFPLAAWNSYALVKPVYELE
Sa RAL MRSSLAVAQWEVVLNSSARNPVAHAASRVIEAPVSEGA KS FAGERVLGVRVLEPTWDSNANAMIKPAFVIP
Tp RAA MKISLAI PSWEIELASSSQTVENQ TLSLVTAAPVKQDAARYGGETVMGVRIHFPSFGINSFAVIKPPFVIP
Bb GIVKPEDMVVDLGINNWSVLLTPSARLQAYVKN SVVAPAVVKSESKRYAGDTILGVRVLEFPSYSQSS AMIMPPFKIP
.. .. ..*..*..*... ..* * .. .. ..***..** ..*...*...

Sh MYG GADGTYTEGKGV IHNVGEIKS ISSWVYGRNYLISYFVN LQNEFGE LKSYPMGTVY
Sa AYEVM AQVDDQGNVQAPTEEEKASGKGRFEDGYGVVKNVGV LKS IAVNTYGMNYPHGLYVMMRDQDGEVHRYFMGYLL
Tp AYATLGDATAQNAVAG GQF DGFGVLKNVGV IKS IQINILGRNYLNRLS LLEDQNGDEREIVMGYLN
Bb FYS GESGNQFL GKGLIDNIKTMEIKVSVYSLGYEIDLEVL FEFDMNGMEYAYSMTGLK
* .. .. ..*..*..*...*..* .. ..*.. .. ..*.. ..**..

Sh FNGWRQVRWENREYLPNVRDRVLV REPLYPRMIPSVKLD S LGFYRTKDTKGGDFITYVKDVTLEYDVVVVDFEEDID
Sa FDSWKELVWNNPSYISDVRSREVR LYPVYPASTPHVVFEGFMVTRDAAHAGGDYVGYFKDVKI IYDKAVLSTVRDFA
Tp FDGWKSLQWNNPNYQTEVRNRDLQ IVPLYPRSA PLIKLKG I K IHRDGSQEGGDIVSYIKDIKVIYDQAVVDRNSDVD
Bb FKGWADLIWSNPNIYIPNISSRI IKDDVPNYPLASSKMRFKAFRVS KSHSSKVKNFIFVYKDLRVLYDKLSVSDSDID
*..*..*..*..*.....* .. ..*..**.. .. .. .. ..*..**.....** .. ..*..

Sh DEATWQLLKTENDRKQAIESARIREQAE LRDLEQRRIGDGT AADQGAANTGAADTGAAQEQAQ
Sa DEDLWGIQARREAERKRVEVARFGQQV LRYIEQEKLATEVGF T P SGG AQRQEEQQ
Tp DEAIWGI LRQEEQYRNFELAKLGNLQV LRSLEK K KMAKEADFDQ A A P A A A A A R A P A T N
Bb SESVEFKVYETSGTESLRKLKAHETFKRVLK LREKISIAEGSFQNFVEKIESEKPEESSPKN
*.. .. ..*..*..*...*..* .. .. .. .. ..*..*..*...
    
```

FIG. 2. Multiple alignment of FlaA sequences of different spirochetes. Multiple alignment was done with the PepPepSearch Program. Alignment gaps are represented by lines. The asterisks indicate identical residues. The dots indicate conserved amino acid substitutions. The blanked residue is not conserved among the four FlaA proteins. *Sh*, *S. hyodysenteriae*; *Tp*, *T. pallidum*; *Sa*, *S. aurantia*; *Bb*, *B. burgdorferi*.




FIG. 3. RT-PCR and primer extension. (a) Products of RT-PCR examined in a 2% agarose gel with the Eagle Eye system (Stratagene). RT-PCR was done by using the Access RT-PCR System from Promega in accordance with manufacturer's instructions. The amplified products from appropriate primers were loaded in the following lanes: 1, pGEM marker; 2, a pair of primers for the middle of the *ftsZ* gene; 3, same as lane 2 except no reverse transcriptase; 4, a pair of primers from the divergently transcribed *thdF* and *ftgK* regions; 5, primers *ppflaAF* and *pflaARR*; 6, primers *pflaA3* and *pflaA4*; 7, primers *pupcheA* and *pcheA3*. (b) Results of primer extension measured in a 5% polyacrylamide gel. The primer *pflaA6* was used for extension. +1 indicates the transcriptional start site.

Downloaded from <http://j.b.asm.org/> on November 20, 2019 by guest




FIG. 4. Comparison of the *flaA* promoter with consensus σ^{70} and σ^{28} promoters.

pairs were designed to connect the adjacent genes *orfA* and *flaA* (ppflaAF and pflaARR), the middle of *flaA* (pflaA3 and pflaA4), and *flaA* and *cheA* (pupcheA and pcheA3; Fig. 1). RT-PCR was carried out with an Access RT-PCR System from Promega. Positive results were obtained with all three pairs of primers (Fig. 3a, lanes 5, 6, and 7) with the controls working as predicted (lanes 2, 3, and 4). These results indicate that *flaA* is transcribed in cultured cells and that it is transcribed together with *orfA* and *cheA* as one unit. Identical results (data not shown) were obtained with strain HB19.




FIG. 5. Western blotting analysis of *B. burgdorferi* (Bb) FlaA protein. The assay was done with an ECL Western blotting system from Amersham. The anti-*T. pallidum* (Tp) 37.0-kDa FlaA serum was diluted 1:3,000, and the anti-*B. burgdorferi* 41.0-kDa FlaB monoclonal antibody was diluted 1:1,000. Lanes: 1, 2, 4, and 5, whole-cell lysate of *B. burgdorferi* 212; 3, purified *T. denticola* PFs (33). Lane 1 was blotted with normal rabbit serum, lanes 2 and 3 were blotted with anti-*T. pallidum* FlaA serum, lane 4 was blotted with anti-*B. burgdorferi* FlaB monoclonal antibody, and lane 5 was blotted with anti-*T. pallidum* FlaA serum and anti-*B. burgdorferi* FlaB monoclonal antibody.

By using primer extension (Promega AMV Primer Extension System), we determined whether there are any promoters upstream of *flaA* or *cheA*. No detectable signal was identified from the region immediately upstream of the *cheA* gene (data not shown). In contrast, a transcriptional start signal was observed from the region immediately upstream of *flaA* (Fig. 3b). This conclusion was verified by using two different primers (pflaA6 and pflaAR). As illustrated in Fig. 1, the *flaA* promoter has the sequences TAAATT at -10 and TTAAAG at -35, which have weak homology with σ^{70} -like promoters (Fig. 4). To test the strength of this promoter element in *Escherichia coli*, we amplified and cloned the promoter region (the segment between ppflaAF and ppflaAR) into a promoter probe vector (pKK232-8) with a promoterless *cat* gene (6). However, we could not detect chloramphenicol resistance in the transformed *E. coli* cells, indicating that this promoter was not functional in *E. coli*. Although we identified this promoter immediately upstream of *flaA*, we also detected a transcript spanning the region between *orfA* and *flaA* (Fig. 3a, lane 5). These results suggest that the *flaA-cheA* operon is controlled by at least two promoters. A similar transcriptional regulation has been described in *Salmonella typhimurium*, in which a class III gene (*flgK*) operon is controlled by both its own promoter and an upstream class II promoter (23). Together with our previous finding that all five chemotaxis genes and *flaA* are transcribed as one transcript (11), our results suggest that *flaA*, *cheA*, *cheW*, *cheR*, *cheX*, and *cheY* constitute a motility-chemotaxis operon.

We directly tested whether a protein compatible with *flaA* expression is synthesized in cultured cells. Our approach was to examine whether cell lysates of *B. burgdorferi* reacted with an antiserum specific to the *T. pallidum* FlaA protein. This antiserum has been shown to react specifically with the FlaA proteins of *T. pallidum*, *T. phagedenis*, and *T. denticola* (29, 33). Western blot analysis of cell lysates of *B. burgdorferi* indicated that a single band of approximately 38.0 kDa reacted with this antiserum (Fig. 5, lane 2). Three controls were used. No reaction was detected when normal rabbit serum was used (Fig. 5, lane 1). In addition, the antiserum reacted with FlaA in purified *T. denticola* PFs as expected (Fig. 5, lane 3). Finally, monoclonal antibody H9724, specific to *B. burgdorferi* FlaB (3), reacted with a protein with a size (41 kDa) distinct from that of FlaA (Fig. 5, lanes 4 and 5). These results suggest that a FlaA homolog is synthesized in *B. burgdorferi*, that it is distinct from FlaB, and that its size approximates that of the deduced *flaA* gene.

The precise function of *B. burgdorferi* FlaA is unclear. Because of its marked similarity to FlaA proteins of other spirochetes and because it is part of an operon involved in chemotaxis, it is likely to be involved in motility. However, in both *T. pallidum* and *S. aurantia*, *flaA* genes are transcribed from σ^{70} promoters and are monocistronic (18, 31), in contrast to our findings on *B. burgdorferi*. Based on sequence analysis, the *B. burgdorferi* *flaA* homolog likely encodes a flagellar filament sheath protein. It is not clear why *B. burgdorferi* FlaA has not been found to be associated with purified PFs. Perhaps upon purification of the PFs, FlaA readily dissociates from the filaments. Alternatively, *B. burgdorferi* FlaA could conceivably have a function other than being a PF sheath protein. Future experiments using cell fractionation and immunolocalization studies should allow us to determine the location of FlaA in *B. burgdorferi* cells.

Nucleotide sequence accession number. The sequence reported in Fig. 1 has been assigned GenBank accession no. U62900.

We thank J. Ruby for purified *T. denticola* PFs, S. Norris for *T. pallidum* FlaA antiserum, A. Barbour for FlaB monoclonal antibody, and R. C. Johnson, R. Limberger, S. Norris, and I. Saint Girons for sharing unpublished information. We also thank H. Thompson and D. Yelton for suggestions.

This research was supported by U.S. Public Health Service grants AI29743 and DE012046.

REFERENCES

1. Anonymous. 1995. Lyme disease—United States, 1994. *Morbidity and Mortality Weekly Report* **44**:459–462.
2. Assous, M. V., D. Postic, G. Paul, and H. G. Barahona. 1993. Western blot analysis of sera from Lyme borreliosis patients according to the genomic species of the *Borrelia* strains used as antigens. *Eur. J. Clin. Microbiol. Infect. Dis.* **12**:261–268.
3. Barbour, A. G., S. F. Hayes, R. A. Heiland, and M. E. Schrupf. 1986. A *Borrelia*-specific monoclonal antibody binds to a flagellar epitope. *Infect. Immun.* **52**:549–554.
4. Brahmsha, B., and E. P. Greenberg. 1988. A biochemical and cytological analysis of the complex periplasmic flagella from *Spirochaeta aurantia*. *J. Bacteriol.* **170**:4023–4042.
5. Brahmsha, B., and E. P. Greenberg. 1989. Cloning and sequence analysis of *flaA*, a gene encoding a *Spirochaeta aurantia* flagellar filament surface antigen. *J. Bacteriol.* **171**:1692–1697.
6. Brosius, J. 1984. Plasmid vectors for the selection of promoters. *Gene* **27**:151–160.
7. Casjens, S., M. Delange, H. L. Ley III, P. Rosa, and W. M. Huang. 1995. Linear chromosomes of Lyme disease agent spirochetes: genetic diversity and conservation of gene order. *J. Bacteriol.* **177**:2769–2780.
8. Charon, N. W., E. P. Greenberg, M. B. H. Koopman, and R. J. Limberger. 1992. Spirochete chemotaxis, motility, and the structure of the spirochetal periplasmic flagella. *Res. Microbiol.* **143**:597–603.
9. Coleman, J. L., and J. L. Benach. 1989. Identification and characterization of an endoflagellar antigen of *Borrelia burgdorferi*. *J. Clin. Invest.* **84**:322–330.
10. Ge, Y., and N. W. Charon. Identification of a large motility operon in *Borrelia burgdorferi* by semi-random PCR chromosome walking. *Gene*, in press.
11. Ge, Y., and N. W. Charon. 1996. Unpublished data.
12. Ge, Y., I. Old, I. S. Saint Girons, and N. W. Charon. 1996. Unpublished data.
13. Ge, Y., I. Old, I. Saint Girons, and N. W. Charon. The *flgK* motility operon of *Borrelia burgdorferi* is initiated by a σ^{70} like promoter. *Microbiology*, in press.
14. Ge, Y. G., I. Old, I. Saint Girons, D. B. Yelton, and N. W. Charon. 1996. FliH and FliI of *Borrelia burgdorferi* are similar to flagellar and virulence factor export proteins of other bacteria. *Gene* **168**:73–75.
15. Goldstein, S. F., K. F. Buttler, and N. W. Charon. 1996. Structural analysis of the *Leptospiraceae* and *Borrelia burgdorferi* by high-voltage electron microscopy. *J. Bacteriol.* **178**:6539–6545.
16. Goldstein, S. F., N. W. Charon, and J. A. Kreiling. 1994. *Borrelia burgdorferi* swims with a planar waveform similar to that of eukaryotic flagella. *Proc. Natl. Acad. Sci. USA* **91**:3433–3437.
17. Hovind Hougen, K. 1984. Ultrastructure of spirochetes isolated from *Ixodes ricinus* and *Ixodes dammini*. *Yale J. Biol. Med.* **57**:543–548.
18. Isaacs, R. D., and J. D. Radolf. 1990. Expression in *Escherichia coli* of the 37-kilodalton endoflagellar sheath protein of *Treponema pallidum* by use of the polymerase chain reaction and a T7 expression system. *Infect. Immun.* **58**:2025–2034.
19. Johnson, R. C., N. Marek, and C. Kodner. 1984. Infection of Syrian hamsters with Lyme disease spirochetes. *J. Clin. Microbiol.* **20**:1099–1101.
20. Kimsey, R. B., and A. Spielman. 1990. Motility of Lyme disease spirochetes in fluids as viscous as the extracellular matrix. *J. Infect. Dis.* **162**:1205–1208.
21. Koopman, M. B. H., E. Baats, C. J. A. H. V. van Vorstenbosch, B. A. M. van der Zieijst, and J. G. Kusters. 1992. The periplasmic flagella of *Serpulina* (*Treponema*) *hyodysenteriae* are composed of two sheath proteins and three core proteins. *J. Gen. Microbiol.* **138**:2697–2706.
22. Koopman, M. B. H., O. S. de Lee, B. A. M. van der Zieijst, and J. G. Kuster. 1992. Cloning and DNA sequence analysis of a *Serpulina* (*Treponema*) *hyodysenteriae* gene encoding a periplasmic flagellar sheath protein. *Infect. Immun.* **60**:2920–2925.
23. Kutsukake, K., and N. Ide. 1995. Transcriptional analysis of the *flgK* and *fliD* operons of *Salmonella typhimurium* which encode flagellar hook-associated proteins. *Mol. Gen. Genet.* **247**:275–281.
24. Limberger, R. J. 1996. Unpublished data.
25. Limberger, R. J., L. L. Slivenski, M. C. T. El-Afandi, and L. A. Dantuono. 1996. Organization, transcription, and expression of the 5' region of the *fla* operon of *Treponema phagedenis* and *Treponema pallidum*. *J. Bacteriol.* **178**:4628–4634.
26. Lutkenhaus, J., and A. Mukherjee. 1996. Cell division, p. 1615–1626. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology. ASM Press, Washington, D.C.
27. Macnab, R. M. 1996. Flagella and motility, p. 123–145. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology. ASM Press, Washington, D.C.
28. Norris, S. A. 1996. Unpublished data.
29. Norris, S. J., N. W. Charon, R. G. Cook, M. D. Fuentes, and R. J. Limberger. 1988. Antigenic relatedness and N-terminal sequence homology define two classes of major periplasmic flagellar proteins of *Treponema pallidum* subsp. *pallidum* and *Treponema phagedenis*. *J. Bacteriol.* **170**:4072–4082.
30. Norris, S. J., and the *Treponema pallidum* Polypeptide Research Group. 1993. Polypeptides of *Treponema pallidum*: progress toward understanding their structural, function, and immunologic roles. *Microbiol. Rev.* **57**:750–779.
31. Parales, J., and E. P. Greenberg. 1993. Analysis of the *Spirochaeta aurantia* *flaA* gene and transcript. *FEMS Microbiol. Lett.* **106**:245–251.
32. Pugsley, A. P. 1993. The complete general secretory pathway in gram-negative bacteria. *Microbiol. Rev.* **57**:50–108.
33. Ruby, J. D., H. Li, H. Kuramitsu, S. J. Norris, S. F. Goldstein, and N. W. Charon. Relationship of *Treponema denticola* periplasmic flagella and the irregular cell morphology. *J. Bacteriol.*, in press.
34. Sadziene, A., D. D. Thomas, V. G. Bundoc, S. C. Holt, and A. G. Barbour. 1991. A flagella-less mutant of *Borrelia burgdorferi*. Structural, molecular, and in vitro characterization. *J. Clin. Invest.* **88**:82–92.
35. Saint Girons, I., I. G. Old, and B. E. Davidson. 1994. Molecular biology of the *Borrelia* bacteria with linear replicons. *Microbiology* **140**:1803–1816.
36. Suk, K., S. Das, W. Sun, B. Jwang, S. W. Barthold, R. A. Flavell, and E. Fikrig. 1995. *Borrelia burgdorferi* genes selectively expressed in the infected host. *Proc. Natl. Acad. Sci. USA* **92**:4269–4273.
37. Szczepanski, A., and J. L. Benach. 1991. Lyme borreliosis: host responses to *Borrelia burgdorferi*. *Microbiol. Rev.* **55**:21–34.
38. Trueba, G. A., C. A. Bolin, and R. L. Zuerner. 1992. Characterization of the periplasmic flagellum proteins of *Leptospira interrogans*. *J. Bacteriol.* **174**:4761–4768.
39. Trueba, G. A., I. G. Old, I. Saint Girons, and R. C. Johnson. CheA and CheW homologues of the agent of Lyme disease, *Borrelia burgdorferi*. Submitted for publication.
40. Wallich, R., S. E. Moter, M. M. Simon, K. Ebnet, A. Heiberger, and M. D. Kramer. 1990. The *Borrelia burgdorferi* flagellum-associated 41-kilodalton antigen (flagellin): molecular cloning, expression, and amplification of the gene. *Infect. Immun.* **58**:1711–1719.
41. Wilson, D. R., and T. J. Beveridge. 1993. Bacterial flagellar filaments and their component flagellins. *Can. J. Microbiol.* **39**:451–472.