

An ABC Transporter System of *Yersinia pestis* Allows Utilization of Chelated Iron by *Escherichia coli* SAB11

SCOTT W. BEARDEN, TEANNA M. STAGGS,[†] AND ROBERT D. PERRY*

Department of Microbiology and Immunology, University of
Kentucky, Lexington, Kentucky 40536-0084

Received 6 August 1997/Accepted 19 December 1997

The acquisition of iron is an essential component in the pathogenesis of *Yersinia pestis*, the agent of bubonic and pneumonic plague. A cosmid library derived from the genomic DNA of *Y. pestis* KIM6+ was used for transduction of an *Escherichia coli* mutant (SAB11) defective in the biosynthesis of the siderophore enterobactin. Recombinant plasmids which had a common 13-kb *Bam*HI fragment were isolated from SAB11 transductants in which growth but not enterobactin synthesis was restored on media containing the iron chelator EDDA [ethylenediamine-di(*o*-hydroxyphenyl acetic acid)]. Subcloning and transposon mutagenesis revealed a 5.6-kb region, designated *yfe*, essential for SAB11 growth stimulation. In vitro transcription-translation analysis identified polypeptides of 18, 29.5, 32, and 33 kDa encoded by the *yfe* locus. Sequence analysis shows this locus to be comprised of five genes in two separate operons which have potential Fur-binding sequences in both promoters. A putative polycistronic operon, *yfeABCD*, is Fur regulated and responds to iron and manganese. A functional Fur protein is required for the observed manganese repression of this operon. This operon encodes polypeptides which have strong similarity to the ATP-binding cassette (ABC) family of transporters and include a periplasmic binding protein (YfeA), an ATP-binding protein (YfeB), and two integral membrane proteins (YfeC and -D), which likely function in the acquisition of inorganic iron and possibly other ions. The ~21-kDa protein encoded by the separately transcribed *yfeE* gene may be located in the cell envelope, since a *yfeE::TnphoA* fusion is PhoA⁺. Mutations in this gene abrogate growth of SAB11 on iron-chelated media.

Yersinia pestis is the facultative intracellular, gram-negative bacterium that causes bubonic and pneumonic plague. In nature, the organism exists as a zoonotic disease of rodents and their associated fleas and has been responsible for the widespread loss of human life during several catastrophic pandemics (59). Colonization and growth of bacterial pathogens, including *Y. pestis*, are dependent on the ability of the invading organism to acquire iron. Since most of the iron in a mammalian host is complexed with metalloproteins or sequestered by the iron storage protein ferritin, bacteria have evolved elaborate strategies to overcome this iron-deficient environment in order to obtain iron for growth (37, 54). Many utilize a variety of extracellular, low-molecular-weight ferric iron chelators termed siderophores (15, 36, 37). Still others obtain iron directly from hemoproteins or the high-affinity iron-binding glycoproteins transferrin and lactoferrin through specific receptors on the bacterial surface (22, 35, 51, 54).

The relationship between iron availability and virulence was clearly demonstrated in spontaneous nonpigmented (Pgm⁻) mutants of *Y. pestis*, in which virulence in mice was restored when the strains were inoculated via peripheral routes and supplemented with hemin or inorganic iron (49). Iron-independent virulence in mice and formation of dark brown, or pigmented, colonies at 26°C from adsorption of exogenous hemin were the original defining characteristics of the chromosomally encoded pigmentation (Pgm⁺) phenotype of *Y. pestis* (48, 49). Pgm⁻ mutants often arise via spontaneous deletion of the 102-kb pigmentation (*pgm*) locus, which is known to

encode an ~7-kb hemin storage (*hms*) locus required for the Hms⁺ phenotype (pigmented colony formation) and the yersiniabactin-iron transport system (Ybt) (59).

The Ybt transport system is a siderophore-dependent uptake mechanism that is common to the three pathogenic species of *Yersinia*: *Y. pestis*, *Yersinia pseudotuberculosis*, and *Yersinia enterocolitica* (18, 32, 43, 59, 83). An ~23-kb region possesses at least four genes likely encoding enzymes for the biosynthesis of the Ybt siderophore (9, 38) and for the receptor (Psn/FyuA) for Ybt and the bacteriocin pesticin (30, 31, 43, 61). Expression of Psn and the Ybt biosynthetic genes is negatively regulated by the Fur repressor but is also activated by YbtA (an AraC-type transcriptional activator located immediately upstream of the Ybt biosynthetic operon) and by the Ybt siderophore (29, 30, 43, 72, 74). Mutations in the Ybt transport system cause a drastic, or possibly complete, loss of virulence in mice infected subcutaneously (9).

However, *Y. pestis* cells possess additional independent hemoprotein and iron transport systems. A hemin utilization system (Hmu) allows the use of hemin and hemoproteins (46, 58, 69, 73). In addition to the Ybt system, inorganic iron transport is mediated by at least two other separate mechanisms. First, studies which monitored the growth of Pgm⁻ strains in the presence of iron chelators suggest that *Y. pestis* cells possess an iron transport system that is functional at 26°C but not at 37°C (52). Second, Ybt⁻ mutants of *Y. pestis* retain the ability to grow at 37°C under iron-deficient but not iron-chelated conditions (9, 30, 58, 68, 69).

In this study, we have used cloning and transposon mutagenesis to identify a *pgm*-independent transport system (Yfe) encoded in two distinct operons (*yfeABCD* and *yfeE*) and located on the *Y. pestis* chromosome. The data presented here suggest that the *yfe* locus of *Y. pestis* is an ATP-binding cassette (ABC) transport system involved in the acquisition of inorganic iron.

* Corresponding author. Mailing address: Department of Microbiology and Immunology, University of Kentucky, MS415 Med. Ctr., Lexington, KY 40536-0084. Phone: (606) 323-6341. Fax: (606) 257-8994. E-mail: rperry@pop.uky.edu.

[†] Present address: Department of Biology, San Antonio College, San Antonio, TX 78284.

TABLE 1. Bacterial strains and plasmids^a

Strain or plasmid	Relevant characteristics	Source or reference
<i>Y. pestis</i>		
KIM6+	Pgm ⁺ Ybt ⁺ Yfe ⁺ Lcr ⁻	68
KIM6	Pgm ⁻ (Δ pgm) Ybt ⁻ Yfe ⁺ Lcr ⁻	32, 68
KIM6-2030	Pgm ⁻ Fur ⁻ (<i>fur::kan-9</i>) Ybt ⁻ Yfe ⁺ Lcr ⁻ Km ^r	73
<i>Y. enterocolitica</i> WA		
	Serotype O:8, Ybt ⁺ Yfu ⁺ Pst ^s Lcr ⁺	32, 82
<i>Y. pseudotuberculosis</i> PBI/+		
	Serotype 1, Pst ^s Lcr ⁺	32, 82
<i>E. coli</i>		
CC118	Strain for detecting PhoA ⁺ fusion proteins, PhoA ⁻	53
DH5 α	Cloning strain, Yfe ⁻ PhoA ⁻	5
HB101	K-12 \times B hybrid strain, Ent ⁺ Yfe ⁻ Sm ^r	5
LE392	Strain for propagating λ Tn <i>phoA</i>	5
SAB11	Ent ⁻ derivative of HB101	6
Plasmids		
pACYC184	4.2-kb cloning vector, Cm ^r Tc ^r	5
pBluescript II KS ⁺	3.0-kb cloning vector, Ap ^r	Stratagene ^b
pBR322	4.4-kb cloning vector, Ap ^r Tc ^r	5
pUC19	2.7-kb cloning vector, Ap ^r	65
pEUPP1	15.3-kb pEU730-derived reporter plasmid containing the <i>psn</i> promoter region upstream of a promoterless <i>lacZ</i> gene, Sp ^r	29
pSC27.1	Reporter plasmid containing an iron- and Fur-regulated <i>lacZ</i> gene	17
pYHE3	pHC79-derived cosmid clone containing an ~23-kb fragment from <i>Y. pestis</i> KIM6+ genomic DNA, Ap ^r	This study
pYFE1	12.8-kb <i>Bam</i> HI fragment from pYHE3 cloned into pUC19, Yfe ⁺ Ap ^r	This study
pYFE1.1	12.8-kb <i>Bam</i> HI fragment from pYHE3 cloned into pACYC184, Yfe ⁺ Cm ^r	This study
pYFE1.2	pYFE1.1 with <i>Bam</i> HI fragment in the opposite orientation, Yfe ⁺ Cm ^r	This study
pYFE2	5.2-kb <i>Clal/Bam</i> HI fragment of pYFE1 inserted into pBR322, Yfe ⁻ Ap ^r	This study
pYFE3	7.7 kb <i>Bam</i> HI/ <i>Clal</i> fragment of pYFE1 cloned into pBR322, Yfe ⁺ Ap ^r	This study
pYFE3.1	7.7-kb <i>Bam</i> HI/ <i>Clal</i> fragment of pYFE3 cloned into pBluescript, Yfe ⁺ Ap ^r	This study
pYFE4	5.3-kb <i>Bam</i> HI/ <i>Hind</i> III fragment of pYFE1 inserted in pBR322, Yfe ⁻ Ap ^r	This study
pYFE6	3.6-kb <i>Pvu</i> II/ <i>Pst</i> I fragment of pYFE1 cloned into pBR322, Yfe ⁻ Tc ^r	This study
pYFE8	9.7-kb <i>Nco</i> I fragment of pYFE1 cloned into pACYC184, Yfe ⁺ Tc ^r	This study
pYFE9	6.1-kb <i>Pst</i> I fragment of pYFE1 inserted into pBR322, Yfe ⁻ Tc ^r	This study
pYFE10	pYFE1.1 lacking internal 5.3-kb <i>Sph</i> I fragment, Yfe ⁺ Cm ^r	This study
pYFE11	pYFE3 with a <i>cat</i> cassette inserted at the <i>Hind</i> III site, Yfe ^{+/-} Ap ^r Cm ^r	This study
pYFE12	pYFE10 with a <i>kan</i> cassette inserted at the <i>Pst</i> I site, Yfe ⁻ Km ^r Cm ^r	This study
pYFE13	pYFE10 with a <i>kan</i> cassette inserted at the <i>Xho</i> I site, Yfe ⁻ Km ^r Cm ^r	This study
pYFE15	pYFE1 with 4.5-kb <i>Xho</i> I fragment deletion, Yfe ⁻ Ap ^r	This study
pYFE21	4.2-kb <i>Eco</i> RI/ <i>Pst</i> I fragment of pYFE1 cloned into pBluescript, Yfe ⁻ (Δ <i>yfeAB</i>) Ap ^r	This study
pYFE30	~1.8-kb <i>Eco</i> RI fragment deletion of pYFE3.1, Yfe ⁻ (Δ <i>yfeAB</i>) Ap ^r	This study
pYFE34 to -46	Tn <i>phoA</i> insertion mutants of pYFE1.1 (Fig. 1B), Cm ^r Km ^r	This study
pYFE47	14.2-kb <i>Sal</i> I fragment deletion of pYFE34, Yfe ⁻ , PhoA ⁺ reporter for <i>Y. pestis</i> KIM6-2030, Cm ^r Km ^r	This study

^a *Y. pestis* KIM6+ possesses an intact 102-kb *pgm* locus containing the genes for hemin storage (*hms*), *psn*, *ybtA*, and the Ybt biosynthetic genes *irp1*, *irp2*, *ybtT*, and *ybtE*. *Y. pestis* KIM6 is an isolate of KIM6+ with the 102-kb region deleted. *Yersinia* strains producing the siderophore yersiniabactin have been designated Ybt⁺, while those defective in yersiniabactin secretion are Ybt⁻. The Yfe⁺ designation indicates the presence of operons *yfeABCD* and *yfeE*. Mutations in either *yfe* operon are depicted as Yfe⁻. Lcr⁺ and Lcr⁻ refer to the presence or absence, respectively, of the low-calcium response phenotype conferred by the 70- to 75-kb virulence plasmid. Ent⁺ and Ent⁻ indicate production or lack of the siderophore enterobactin, respectively. *E. coli* strains which fail to express alkaline phosphatase are designated PhoA⁻. Ap^r, Cm^r, Km^r, Sm^r, Sp^r, and Tc^r, resistance to ampicillin, chloramphenicol, kanamycin, streptomycin, spectinomycin, and tetracycline, respectively.

^b Stratagene Cloning Systems, La Jolla, Calif.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. The relevant characteristics of all plasmids and bacterial strains are given in Table 1. All bacterial strains were maintained at -20°C in phosphate-buffered glycerol. *Yersinia* strains were grown routinely in heart infusion broth or on tryptose blood agar base (Difco Laboratories) at either 26 or 37°C. For iron-deficient growth, *Y. pestis* strains were grown in the chemically defined medium PMH, which had been deferrated prior to use by extraction with Chelex 100 (Bio-Rad Laboratories); this results in a residual iron content of ~0.3 μ M (73). Iron-deficient growth of *Escherichia coli* strains was accomplished by using Tris-glucose-thymidine medium (TG) prepared as described previously (46). *Y. pestis* KIM6+ and its Δ pgm derivative KIM6 were grown on Congo red agar (78) to determine their hemin storage (Hms) phenotypes (31, 60). *E. coli* strains were cultured in Luria broth (LB) or Terrific broth (79) or on LB solidified with 1.4% (wt/vol) Bacto agar (Difco Laboratories). For growth studies requiring iron-chelated media, *E. coli* SAB11 cells containing

various recombinant plasmids were grown on either LB agar containing 750 μ M EDDA [ethylenediamine-di(*o*-hydroxyphenyl acetic acid)] (LA-EDDA) or TG medium solidified with 1.2% (wt/vol) agarose and supplemented with 75 μ M EDDA (TGE). Where appropriate, media included antibiotics at the following concentrations: ampicillin, 100 μ g/ml; tetracycline, 12.5 μ g/ml; kanamycin, 25, 50, or 200 μ g/ml; and chloramphenicol, 30 μ g/ml.

Recombinant DNA methodology. Plasmids were introduced into *E. coli* strains by standard CaCl₂ transformation (65). Isolation of plasmid and cosmid DNAs was achieved by alkaline lysis (12), and the DNAs were further purified when necessary by polyethylene glycol precipitation (47). DNA probes used in Southern hybridizations (5) were radiolabeled with [α -³²P]dCTP (ICN Biomedicals, Inc.) by nick translation with a commercially available kit (Life Technologies, Inc.). Hybridization to confirm the presence of the *yfe* locus in various recombinant cosmids was carried out as described by Fetherston et al. (32). Radiolabeled blots were exposed to Kodak XAR-5 film at -80°C with intensifying




FIG. 1. (A) Restriction map of pYFE1.1 and various subclones used to test for restoration of growth of *E. coli* SAB11. +, plasmids which promote SAB11 growth; -, no growth stimulation. Arrows indicate the directions of transcription of genes included in the *yfe* locus. Dashed lines and open triangles represent deletions and sites of antibiotic cassette insertion, respectively. (B) Map of relevant *TnphoA* insertions within the *yfe* locus. Filled triangles represent *PhoA*⁺ translational fusions. + and - signs above each insertion site indicate iron-deficient growth and no growth, respectively, of SAB11 carrying that insertion in pYFE1.1. Numbers above each triangle represent plasmids pYFE34 to -46 (pYFE1.1::*TnphoA*34 to -46), respectively (Table 1). Shaded boxes upstream of *yfeA* and *yfeE* represent potential Fur boxes.

screens. Subclones derived from pYFE1.1 (Table 1; Fig. 1) were constructed by using restriction endonucleases and DNA-modifying enzymes according to the manufacturers' instructions. The *Yfe* insertion mutants depicted in Fig. 1 were generated by introduction of antibiotic resistance cassettes (Pharmacia, Inc.) into functional pYFE subclones (Table 1).

DNA sequencing was performed by the dideoxynucleotide chain termination method (66) with Sequenase version 2.0 and ³⁵S-dATP (Amersham Corp.). Resolution of sequencing artifacts and compressions was achieved by using 7-deaza-dGTP (Boehringer Mannheim Biochemicals) and by the inclusion of dimethyl sulfoxide (Sigma Chemical Co.) to 10% (vol/vol) in primer-template annealing reactions. Sequencing reaction products were resolved on 6% polyacrylamide gels containing 8.3 M urea (Sigma Chemical Co.) and cast in Trisborate-EDTA buffer (65). Dried gels were exposed to Kodak BioMax MR film at room temperature. Oligonucleotide primers (Integrated DNA Technologies, Inc.) were used to extend the sequence in the *yfe* region and to determine the sequence of the opposite strand within *yfe* coding regions. DNA sequence analysis was performed with programs from the Intelligenetics software suite.

Isolation of *yfe* clones. Construction of a cosmid library derived from *Y. pestis* KIM6+ and its transduction into *E. coli* SAB11 (Table 1) have been described previously (46). Transduced cells were plated onto LA-EDDA plates supplemented with 5 μ M hemin and ampicillin at 100 μ g/ml. Isolated colonies were subsequently plated onto LA-EDDA containing either 0 or 10 μ M hemin. Transductants requiring hemin for growth have been described elsewhere (46). Isolates exhibiting hemin-independent iron-chelated growth were designated *yfe* clones and are the subject of this study.

***TnphoA* mutagenesis of pYFE1.1.** For transposon mutagenesis, *E. coli* CC118 cells bearing plasmid pYFE1.1 (Table 1; Fig. 1) were transduced with λ *TnphoA* (39) at a multiplicity of infection of 5.0 according to the method of de Bruijn and Lupski (23) with minor modifications. Briefly, transduced cells were spread onto LB agar containing chloramphenicol (30 μ g/ml), kanamycin (200 μ g/ml), and 5-bromo-4-chloro-3-indolyl phosphate (XP) (40 μ g/ml) (Sigma Chemical Co.) and incubated at 30°C for 48 h. Kanamycin-resistant colonies were harvested, and plasmid DNAs from this pool of transformants were isolated by alkaline lysis (12). Pooled plasmid DNA was subsequently used to transform *E. coli* DH5 α (*PhoA*⁻) cells for the selection of individual blue (cells expressing alkaline phosphatase) and white colonies (63). Transposon insertion sites of individual pYFE1.1::*TnphoA* recombinant plasmids were estimated by restriction endonuclease digestion. *TnphoA* insertion sites of mutant plasmids which abrogated growth stimulation of *E. coli* SAB11 cells on TGE medium were verified by DNA sequence analysis with the primer *phoA*.1 (5'-GTGCAGTAATATCGCCCTG AGC-3') derived from the sequence of the *E. coli phoA* gene (19).

Alkaline phosphatase and β -galactosidase assays. Cell lysates were prepared from cultures of *Y. pestis* grown in PMH or in PMH supplemented with either 1.0 μ M MnCl₂, 1.0 μ M FeCl₃, or 10 μ M FeCl₃. Cells were passaged twice for a total of approximately six to eight generations as described previously (72). Alkaline phosphatase and β -galactosidase activities were determined spectrophotometrically by monitoring cleavage, respectively, of the substrates *p*-nitrophenyl phosphate and 4-nitrophenyl- β -D-galactopyranoside (Sigma Chemical Co.). Units of alkaline phosphatase activity were calculated according to the formula described

by Brickman and Beckwith (16), and β -galactosidase activity is expressed in Miller units (55).

Protein analyses. In vitro transcription-translation of plasmid-encoded proteins was performed with an *E. coli* S30 cell extract system (Promega Corp.). Proteins were radiolabeled with ^{35}S -labeled amino acids (DuPont NEN Research Products) according to the manufacturer's recommendations, and equal amounts of trichloroacetic acid-precipitable counts were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Dried gels were exposed to Kodak BioMax MR film at room temperature. Homology searches of protein databases were performed with BLAST (3). Pairwise and multiple sequence alignments were accomplished with the programs Gap (University of Wisconsin Genetics Computer Group software package version 8.1) and CLUSTALW (80), respectively. Determination of the transmembrane-spanning domains in the predicted amino acid sequences of YfeC and YfeD was performed with the program TopPred II (21).

PCR analyses. Amplification of a region of the *yfeA* gene was accomplished by PCR with commercially synthesized oligonucleotides (Integrated DNA Technologies, Inc.) 4151.1 (5'-GAATGGCATGAATCTGGAGCG-3') and 2021.4 (5'-GGCGGGTTTGTGCGAAATAG-3'), derived from the nucleotide sequence of *Y. pestis yfeA*. Similarly, synthetic primers yfuA5.2 (5'-CTGGTGAATCCTGGGTC-3') and yfuA3.3 (5'-TCCAGATCTTCAGCGCAC-3'), derived from the nucleotide sequence of the *Y. enterocolitica yfuA* gene, were used to amplify a fragment of the *yfuA* gene, encoding a putative periplasmic binding protein involved in inorganic iron transport (64). All strains were cultured at 37°C in microtiter wells for 4 to 6 h. For PCR analysis, 1.0 μl of cells (10^4 to 10^6 CFU) from each strain was used. Amplification reactions with mixtures containing either *Taq* DNA polymerase (Promega Corp.) or *Pfu* DNA polymerase (Stratagene), 200 μM deoxynucleoside triphosphates, 1.5 μM MgCl_2 , and 0.4 μM primers were performed for 30 s at 94°C, 55°C, and 72°C for 30 cycles. All PCRs were carried out with a GeneAmp PCR System 2400 thermal cycler (Perkin-Elmer). Products from these reactions were resolved by agarose gel electrophoresis.

Nucleotide sequence accession numbers. The sequences of *yfeABCD* and *yfeE* have been deposited in the GenBank database and assigned accession numbers U50597 and U50903, respectively.

RESULTS

Isolation and subcloning of the *Y. pestis yfe* locus. To identify heme and inorganic iron uptake systems of *Y. pestis*, a cosmid library derived from *Y. pestis* KIM6+ was transduced into *E. coli* SAB11 (Table 1), a mutant defective in the synthesis of the siderophore enterobactin (6). Eleven transductants were isolated from LA-EDDA plates containing 5 μM hemin as described in Materials and Methods and reference 46. Six of the isolates grew on LA-EDDA plates in the absence of added hemin and on TGE medium, precluding the possibility that growth on LA-EDDA was due to contaminating hemin. Southern hybridizations (5) confirmed that these cosmids have a common \sim 13-kb *Bam*HI fragment which is present in both Pgm^+ and Pgm^- strains of *Y. pestis* and in *Y. pseudotuberculosis* (data not shown). This fragment was subcloned from the recombinant cosmid pYHE3 into the vector pACYC184 (5) in both orientations and designated pYFE1.1 and pYFE1.2 (Table 1; Fig. 1). Further subcloning as well as insertion and deletion mutagenesis of this *yfe*-containing recombinant plasmid showed the essential region for SAB11 growth stimulation to be contained within a 7.7-kb *Bam*HI-*Cla*I fragment represented by pYFE3 (Fig. 1A; Table 1). However, restoration of SAB11 growth mediated by pYFE1.1 and various subclones did not result in concomitant biosynthesis of the siderophore enterobactin. *E. coli* SAB11 cells carrying either pACYC184 (Table 1) or the nonfunctional pYFE1.1 subclones pYFE4, pYFE6, pYFE9, pYFE12, pYFE13, pYFE15, and pYFE30 (Fig. 1A) could not be cross-fed by adjacently streaked cells harboring either pYFE1.1 or pYFE3 on iron-deficient media. However, HB101, which synthesizes enterobactin, did cross-feed SAB11 in this assay system. Introduction of antibiotic resistance cassettes into the functional subclones pYFE3 and pYFE10 (Table 1) either abolished (pYFE12 and pYFE13) or inhibited (pYFE11) their growth-promoting properties in SAB11. DNA sequence analysis of the *yfe* locus (see below) places the *Hind*III site into which the chloramphenicol acetyl-

transferase (*cat*) cassette has been ligated in pYFE11 near the carboxy terminus of YfeD. The expression of a truncated YfeD polypeptide which has retained partial function may account for the reduced *E. coli* SAB11 growth for cells containing this plasmid. While similar growth characteristics might be expected for SAB11 cells transformed with pYFE4, it should be noted that this plasmid construct lacks the *yfeE* gene locus, which is an essential component in restoring growth to SAB11 on iron-chelated media (Fig. 1A).

Iron-deficient growth of *E. coli* SAB11. In addition to scoring individual subclones of pYFE1.1 for their ability to promote the growth of *E. coli* SAB11 on iron-chelated media (Fig. 1A), we also examined the growth characteristics of this strain in liquid culture. *E. coli* SAB11 cells containing either pBR322 or the pYFE1.1 subclone pYFE3 (Table 1; Fig. 1A) and HB101 (pBR322) were serially transferred twice (six to eight generations) to acclimate cells to medium conditions prior to monitoring their growth in TG medium with or without 10 μM FeCl_3 for a period of 9 h. As shown in Fig. 2, the iron-limited growth of *E. coli* SAB11(pBR322) is enhanced by providing the *yfe* locus in *trans* (pYFE3). SAB11(pYFE3) shows a slightly increased initial growth rate over those of SAB11(pBR322) and its enterobactin-producing parent strain HB101(pBR322). The relative ineffectiveness of iron acquisition by the enterobactin siderophore system of HB101 is a curious but highly reproducible result. The increased growth of SAB11(pYFE3) over that of the enterobactin-producing HB101(pBR322) may be due to the copy number of the *yfe* operons contained within the pBR322-derived pYFE3 plasmid.

***TnphoA* mutagenesis of plasmid pYFE1.1.** To further define the region necessary for restoring growth to *E. coli* SAB11, we employed a method of random mutagenesis with λTnphoA (39). In mapping transposon insertion sites within pYFE1.1 which abrogated growth of *E. coli* SAB11 on iron-chelated media, we also identified several *TnphoA* insertion mutants which exhibited alkaline phosphatase (PhoA^+) activity by the *phoA* gene product, which requires a foreign signal peptide for export of PhoA to the periplasm where it becomes active (39, 53). PhoA^+ insertions in three separate open reading frames (ORFs) (Fig. 1B), as determined by DNA sequence analysis, indicate that at least three proteins encoded by the *yfe* locus contain sequences which target them for export to the cell envelope. The results in Fig. 1B also show that despite multiple insertions within or upstream of ORF33, SAB11 growth was unaffected, indicating that this gene is not essential for promoting the iron-chelated growth of *E. coli* SAB11. By contrast, insertions in *yfeA*, *yfeD*, and *yfeE* resulted in a loss of the ability of SAB11 to utilize chelated iron. The weak, reduced growth phenotype of the single transposon insertion in *yfeB* cannot be readily explained but likely is aberrant. This is supported by the negative growth phenotype conferred by pYFE13 and by the genetic organization of *yfeABCD* (Fig. 1A), which suggests that interruption of the YfeB-coding region should have polar effects on downstream sequences and thus preclude the ability of such a plasmid to promote the growth of SAB11. Transposon insertions within *yfeE* (Fig. 1B) present apparently conflicting results. DNA sequence analysis of the *TnphoA* insertion proximal to the 5' end of *yfeE* (pYFE45 [Table 1]) shows that the location of the *yfeE-phoA* fusion occurs just 13 nucleotides downstream of the *yfeE* translational start site (*yfeE*::*TnphoA45*). Despite this disruption of the *yfeE* coding region, plasmid pYFE45 is able to restore the growth of SAB11 on iron-chelated media. This is in contrast to the case for pYFE44, in which the transposon insertion site follows nucleotide 139 of the *yfeE* ORF (*yfeE*::*TnphoA44*) and does not restore SAB11 growth (Fig. 1B).




FIG. 2. Growth of *E. coli* HB101 or SAB11 transformed with either pBR322 or pYFE3. Filled symbols indicate strains grown in TG minimal medium supplemented with 10 μ M FeCl₃, while open symbols designate growth in TG without supplementation. OD₆₂₀, optical density at 620 nm.

DNA sequence analysis of the *yfe* locus reveals two distinct operons. The *yfe* locus contains two separate operons (*yfeABCD* and *yfeE*) that are essential for iron acquisition in *E. coli* SAB11. The product of ORF33 (Fig. 1B) is a hypothetical protein, as no evidence for its expression yet exists. The promoter region of each operon has a putative Fur-binding sequence (FBS) (Fig. 3) which has 89% (*yfeABCD*) and 63% (*yfeE*) identity to the *E. coli* FBS consensus sequence (77). Moreover, imperfect inverted repeats can be found downstream of the translational stop codons for *yfeD* and *yfeE* and may serve as transcriptional terminators (Fig. 3). YfeE is an ~21-kDa protein with a pI of 9.3 and significant homology (75.3% similarity) to the product of a 534-bp ORF from *E. coli* designated f178 (13) (GenBank accession number U00096). The PhoA⁺ *yfeE*::*TnphoA* fusions (Fig. 1B) suggest that YfeE is located in the cell envelope despite the absence of an obvious signal sequence. The genetic organization of *yfeABCD* suggests that it is a polycistronic operon. Examination of the intergenic nucleotide sequences indicates that expression of

YfeABCD is translationally coupled (Fig. 3). BLAST homology searches (3) show that the *yfeABCD* operon likely encodes components of a periplasmic binding protein-dependent transport system or ABC transporter (27, 44), with the 35.7-kDa YfeA predicted to be a periplasmic binding protein, the 32.2-kDa YfeB predicted to be an ATP-binding protein, and YfeC and YfeD (31.7 and 32.2 kDa, respectively) predicted to be integral membrane permeases. The predicted translation product of *yfeA* has significant amino acid similarity (82%) to an iron-repressible periplasmic protein of unknown function from *Haemophilus influenzae* (42). More recently, this polypeptide has been shown to be derived from a complex and as-yet-uncharacterized region of the *H. influenzae* genome (HI0359 to HI0362) whose genetic organization is identical to that of *yfeABCD* and whose ORFs have >80% similarity at the amino acid level (33). Additionally, the *yfeABCD* operon has ~70% amino acid similarity with an ABC transporter complex for manganese (*mntCAB*) in the cyanobacterium *Synechocystis* sp. strain PCC 6803 (7). Moreover, a putative ABC transporter




FIG. 3. Partial DNA sequence representation of the *yfeABCD* and *yfeE* operons and ORF33, including single-letter amino acid translations of the 5' and 3' ends of each ORF. Only the coding strand of each gene is shown, and omitted nucleotide sequences are depicted as dots. The FBSs in the promoter regions of *yfeA* and *yfeE* are overlined and underlined, respectively. Nucleotides in boldface represent potential ribosomal binding sites. Nucleotides overlined with arrows, which follow the *yfeD* termination codon and the translational stop of *yfeE*, represent imperfect inverted repeat structures. The translational start sites of the Yfe polypeptides are indicated by arrows. YfeA' indicates an alternate translational start site for the *yfeA* coding region. Underlined amino acids in YfeA depict the region of homology with N-terminal amino acid sequence of a 31-kDa iron-repressible periplasmic binding protein from *H. influenzae* (42). The vertical arrow marks a putative signal sequence cleavage site.

(*troABCD* products) has recently been identified in the syphilis spirochete *Treponema pallidum* (41). This operon encodes proteins TroA to -D, which have amino acid similarities of 51, 61, 54, and 60%, respectively, with YfeA to -D. An amino acid sequence alignment between YfeA, -B, and -C and their *H. influenzae* and Mnt homologs is shown in Fig. 4. For the purpose of clarity, alignments with YfeD were omitted, since the *mnt* operon lacks this additional ORF (7). The deduced amino acid sequence encoded by *yfeB* shows strong similarity to ATP-binding proteins in several systems, including those involved in the uptake of inorganic iron (15, 44). The amino acid sequence alignment between YfeB, HI0361, and MntA (Fig. 4) reveals the high degree of conservation surrounding the ATP-binding motifs (Walker A and B) of these proteins (44). These sites are recognized for their ability to form an ATP-binding pocket and are the distinguishing feature among the family of ABC transporters (44). The integral membrane proteins YfeC and YfeD are strongly hydrophobic, with each containing up to seven membrane-spanning domains and a distinctive domain known as the EAA motif (Fig. 5). This signature motif typically occurs in an hydrophilic loop region and contains an invariant glycine residue positioned about 100 amino acids from the C terminus of the protein (119 and 125 residues for YfeC and YfeD, respectively), and it is highly conserved among cytoplasmic membrane permeases (67). In addition to the YfeC sequence homologies depicted in Fig. 4, the products encoded by *yfeC*

and *yfeD* have homology with a number of hydrophobic integral membrane proteins found in *Streptococcus* spp. (26, 28, 50, 57). Interestingly, despite similarities in their predicted secondary structures, pairwise alignment of the amino acid sequences for YfeC and YfeD (Fig. 5) shows a lesser degree of similarity (60%) than is indicated for YfeC and HI0360 (83%), YfeC and MntB (72%), or YfeD and HI0359 (80%).

Identification of putative *yfe*-encoded proteins. To determine the polypeptides necessary for promoting growth of *E. coli* SAB11 on iron-chelated media, proteins encoded by the plasmid pYFE3 as well as selected subclones were examined by in vitro transcription-translation (Fig. 6). The similar predicted molecular masses of the proteins encoded by the *yfeABCD* operon have made specific identification of these gene products problematic due to possible comigration of several polypeptides in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (Fig. 6). We have identified at least four polypeptides, with molecular masses of 18, 29.5, 32, and 33 kDa (Fig. 6A), encoded by the 7.7-kb insert of pYFE3, which had previously been designated the essential region for promoting the iron-deficient growth of SAB11. The most readily identifiable gene product, YfeE, migrates at ~18 kDa, compared to its predicted mass of 21 kDa, and is evident in the protein profiles of pYFE3, pYFE8, pYFE9, pYFE12, pYFE13, and pYFE21 (Fig. 6). YfeE is absent from the profile of pYFE15 (Fig. 6A). In addition to the loss of YfeE expression




FIG. 4. CLUSTALW amino acid sequence alignments of *Y. pestis* proteins YfeA, YfeB, and YfeC with components of a manganese transporter from *Synechocystis* sp. strain PCC 6803 (MntCAB) and with polypeptides from a putative ABC transporter from *H. influenzae* (HI0360, HI0361, and HI0362). Amino acids in boldface represent ATP-binding motifs (Walker A and Walker B). The consensus line displayed below the aligned sequences depicts identical amino acids as asterisks, with conserved residues shown as dots.

from pYFE15, it was expected that at least two other proteins (YfeC and YfeD), whose genes are contained within the deleted region of this plasmid (Fig. 1A), would be lost. While the protein band migrating at ~32 kDa is no longer evident, the persistence of a 33-kDa band may be explained by the presence of an additional 5.1 kb of *Y. pestis* DNA downstream of the *XhoI* deletion in this plasmid. Products potentially encoded by this region of pYFE1.1 have not been extensively examined. While *yfeA* is predicted to encode a 35.7-kDa protein, a polypeptide migrating at this position was not detected (Fig. 6). To determine the relative migration of YfeA, we examined the protein profile of pYFE9 (Fig. 6A). Construction of this plasmid resulted in a fortuitous translational fusion between the pBR322-encoded β-lactamase protein (Bla) and YfeA, resulting in a fusion protein with a predicted molecular mass of ~53 kDa. The appearance of the Bla-YfeA fusion in pYFE9 and the concomitant loss of the species migrating at 29.5 kDa (Fig. 6A) suggest that this band corresponds to YfeA. The protein bands migrating at 32 and 33 kDa account for the remaining proteins, YfeB to -D. Since there are only two bands present to account for three distinct proteins, two-dimensional gel electrophoresis will be required to resolve individual YfeB-

-C, and -D polypeptides. Finally, since the organization of *yfeABCD* and supporting DNA sequence data suggest that it is a polycistronic operon, we examined the in vitro transcription-translation protein profiles of plasmids that either lacked the *yfeA* promoter region or contained an insert which interrupted the *yfeA* coding region. The profile of plasmid pYFE12 shows that insertion of a *kan* cassette near the 5' end of *yfeA* causes polar effects on downstream sequences, resulting in the apparent loss of polypeptides YfeA to -D (Fig. 6B). Similarly, plasmid pYFE21 (Table 1), which lacks the *yfeA* promoter region as well as *yfeA* and a portion of *yfeB*, fails to express downstream sequences. However, both pYFE12 and pYFE21 maintain expression of the 18-kDa protein, confirming that YfeE is derived from a separate transcriptional unit (Fig. 6).

Regulation of the *yfeA* promoter by iron, manganese, and Fur. We introduced plasmid pYFE34 (*yfeA*::*TnphoA34*) into *Y. pestis* KIM6 and KIM6+ by electroporation to examine expression of the plasmid-encoded YfeA-PhoA fusion protein under repressive or nonrepressive conditions. The alkaline phosphatase activities of these strains grown in the presence or absence of iron or manganese are reported in Table 2. *Y. pestis* KIM6(pYFE34) cells grown in PMH medium under iron-de-




FIG. 6. Autoradiograms of plasmid-encoded proteins labeled with ³⁵S-amino acids by in vitro transcription-translation. Lanes MW, molecular mass markers. Other lane designations indicate the plasmids used. Maps of pYFE plasmids are shown in Fig. 1A. Relevant proteins are indicated by arrows. Bla/YfeA, β -lactamase-YfeA translational fusion; CAT, chloramphenicol acetyltransferase. Numbers on the left and right are molecular masses in kilodaltons.

larger of these, *yfeABCD*, encodes polypeptides which are translationally coupled (Fig. 3) and collectively resemble a periplasmic binding protein-dependent transport system (14, 45) belonging to the superfamily of ABC transporters (27, 44). The

yfeA gene encodes the putative substrate-binding protein YfeA, which likely functions within the periplasmic space of *Y. pestis* cells. YfeB has signature ATP-binding motifs (44), while YfeC and YfeD possess characteristics of integral mem-

TABLE 2. β -galactosidase and alkaline phosphatase activities of *Y. pestis* KIM6+ and KIM6 derivatives grown to mid-log phase in defined PMH medium

Strain	Alkaline phosphatase or β -galactosidase activity ^a of cells grown in medium with:				Induction ratio ^b				
	10 μ M FeCl ₃ (37°C)	1.0 μ M FeCl ₃ (37°C)	1.0 μ M MnCl ₂ (37°C)	No added FeCl ₃ or MnCl ₂		-Fe/+10 μ M Fe	-Fe/+1.0 μ M Fe	-Fe/+Mn	26°C/37°C
				37°C	26°C				
PhoA strains									
KIM6(pYFE34)+ ^c	229	369	576	1,532		5.1 (1.6)	4.7 (1.4)	2.6 (0.2)	
KIM6(pYFE34)	326	436	647	2,071	3,295	7.2 (2.4)	5.7 (0.9)	3.5 (0.2)	1.8 (0.3)
KIM6(pYFE47)	239	317	537	1,306		5.4 (1.4)	4.8 (2.1)	2.4 (0.3)	
KIM6-2030(pYFE47)	1,830	1,865	1,940	2,046		1.2 (0.3)	0.9 (0.4)	0.9 (0.2)	
LacZ strains									
KIM6(pEUAPP1)+	196	623	1,102	1,403		7.3 (2.7)	2.3 (0.05)	1.3 (0.04)	
KIM6(pSC27.1)+	1,060	1,412	2,265	2,648		2.6 (0.6)	2.2 (0.2)	1.2 (0.04)	
KIM6(pSC27.1)				1,726	4,307				2.5 (0.1)

^a Units of phosphatase activity were calculated as $1,000 \times [(OD_{420} - 1.75 \times OD_{550}) / \text{time} \times OD_{620} \times \text{volume}]$, where OD_{420} is optical density at 420 nm (16). β -Galactosidase activity is expressed in Miller units (55). Values are the means for samples taken from at least two independent experiments. In the absence of reporter genes, *Y. pestis* cells are phenotypically PhoA⁻ and β -galactosidase negative.

^b Values in parentheses represent standard deviations.

^c Names followed by a + are those of *Y. pestis* strains containing an intact 102-kb pigmentation (*pgm*) locus (59).




FIG. 7. PCR of DNAs derived from whole cells of *Yersinia* spp. or *E. coli* DH5α. (A) Oligonucleotide primers derived from a region of *Y. enterocolitica* *yfuA* were used to PCR amplify genomic DNAs from the indicated strains. The predicted product is indicated by the arrow. (B) Oligonucleotide primers were derived from a region of *Y. pestis* *yfeA*. The predicted amplicon is designated with an arrow. Reactions were performed with *Taq* DNA polymerase for 25 cycles of 94°C for 30 s, 50°C for 30 s, and 72°C for 30 s.

brane permeases (67). YfeE, encoded by the second operon of the *yfe* locus, is apparently essential for restoring growth to *E. coli* SAB11 on iron-chelated media. The mechanism which enables pYFE46 (*yfeE::TnphoA46*) to restore iron-chelated growth to SAB11 is unknown; however, it is plausible that a spurious promoter generated at the *TnphoA* insertion site in pYFE46 facilitated some level of YfeE (lacking only the first five amino acids) expression. This rationale might also explain the weak growth observed with SAB11(pYFE36), which has a *TnphoA* insertion in *yfeB* (Fig. 1B). In most instances insertions of transposon Tn5, and derivatives such as *TnphoA*, have strongly polar effects on the expression of distal genes. However, other studies have shown that low levels of distal gene expression can occur in some cases with transcription originating from within the end of Tn5 (10). While the function of YfeE is unknown, *TnphoA* mutagenesis of *yfeE* suggests that YfeE is transported to the cell envelope (Fig. 1B). Although we have demonstrated, by subcloning and mutagenesis, only that YfeD and YfeE are essential for this system to function in *E. coli* SAB11, sequence information makes it likely that all five *yfe* gene products are necessary for transport activity.

Several well-characterized ABC transporters involved in inorganic iron uptake have been described for *E. coli*. However, these systems are largely defined by their iron-chelating siderophores and associated outer membrane receptors (15). Re-

cently, other ABC transporter systems for inorganic iron that are more readily characterized based on their periplasmic binding proteins rather than an associated siderophore or outer membrane protein receptor have been described. The *fbpABC* locus of *N. gonorrhoeae*, *hitABC* of *H. influenzae*, and *sfuABC* of *S. marcescens* all encode systems which transport iron from the periplasm to the cytosol in a siderophore-independent manner. These transport systems may utilize several different specific outer membrane receptors (1, 2, 4, 20, 34, 85). Amino acid sequence homologies indicate that the *yfuABC* locus of *Y. enterocolitica* also belongs to this family of iron transporters and encodes polypeptides which facilitate iron uptake (62, 64). The Yfe system of *Y. pestis* is functionally and organizationally similar to the Yfu system of *Y. enterocolitica*. However, except for ATP-binding motifs found in YfeB, it shows no significant amino acid homology to Yfu or to the related ABC iron transport systems of *H. influenzae* (HitABC), *N. gonorrhoeae* (FbpABC), and *S. marcescens* (SfuABC) (1, 2, 4, 20, 34, 85). Moreover, PCR analysis indicates that the *yfu* locus is restricted to *Y. enterocolitica* (Fig. 7A), whereas the *yfeA* gene and presumably *yfeBCD* were detected in all three pathogenic yersiniae, including *Y. pestis* KIM6, the Pgm⁻ derivative of KIM6+ (Fig. 7B). Recent database searches and the alignments depicted in Fig. 4 show that the Yfe system has a high degree of similarity with a manganese-specific periplasmic binding protein-dependent transporter of the cyanobacterium *Synechocystis* sp. strain PCC 6803 (MntCAB) as well as with putative periplasmic permeases found in *H. influenzae* (HI0359 to HI0362) and *T. pallidum* (TroA to -D) (7, 33, 41). It has been suggested that the *Y. pestis* Yfe transport system and the HI0359 to HI0362, TroA to -D, and MntCAB systems are members of a new subfamily of ABC transporters (26). This subfamily also includes a number of potentially metal-binding streptococcal adhesins and two ABC metal permease systems from *Streptococcus pneumoniae* involved in the transport of zinc (Adc) and manganese (Psa). These two cation transport systems are important for the competence and virulence of this human pathogen (25, 26).

A recent study using the manganese ABC transporter, Mnt, showed that the uptake of ⁵⁴Mn²⁺ was uninhibited in the presence of FeCl₃ (8). However, the accumulation of manganese by the Mnt system was competitively inhibited by Cd²⁺, Co²⁺, and Zn²⁺ (8). Although it is clear that the *yfeABCDE* locus functions to provide iron to *E. coli* SAB11 in our system, the similarities between the Mnt and Yfe ABC transport systems allow for reasonable speculation that the Yfe system is involved not only in the transport of iron but also in that of other metal ions. The promoter regions of both *yfe* operons (Fig. 3) contain FBSs, and we have used transcriptional reporter gene studies to show that the *yfeABCD* promoter is iron and Fur regulated. However, the high degree of similarity to the manganese transporter of *Synechocystis* sp. strain PCC 6803 (7, 8) suggests that the Yfe system may also function in manganese transport.

In *E. coli*, manganese is transported by a specific, high-affinity system (71). Although the genetic components of this system have not been isolated, studies using *E. coli* right-side-out membrane vesicles demonstrated functional manganese transport in the absence of ATP, indicating that the *E. coli* system derives its energy from the chemiosmotic membrane potential of the cell (11). Since the manganese transporter of *E. coli* is presumed to be encoded by a single gene (70) and there is no evidence to indicate that *E. coli* SAB11 or its parent strain, HB101, is defective in manganese transport, it is unlikely that the Yfe ABC transporter would restore a specific manganese transport defect in *E. coli* SAB11. Still, it is interesting that

PhoA expression driven by the *yfeA* promoter was significantly repressed in *Y. pestis* cells grown in the presence of manganese. In *E. coli*, binding to Fur-regulated promoters has been demonstrated in DNase I footprinting and gel retardation assays using the Fur repressor complexed with Mn^{2+} (24, 84). In addition, resistance to high concentrations of Mn^{2+} was used to select an *E. coli* Fur⁻ mutant (40). Therefore, we examined the possibility that manganese repression of the *yfeABCD* promoter is mediated in conjunction with Fur by examining regulation in other Fur-regulated promoters and in a *Y. pestis* Fur⁻ mutant. We showed that manganese repression of transcription from the *yfeABCD* promoter was Fur dependent; however, β -galactosidase expression from two separate iron- and Fur-regulated promoters was not significantly repressed by the presence of manganese in the growth medium (Table 2). This indicates that the manganese regulation shown by the *yfeABCD* promoter is not common to all Fur-regulated promoters. Moreover, it is possible that manganese regulation of *yfeABCD* occurs through an altered FBS which enhances binding of the Fur- Mn^{2+} complex. Whether Fur proteins from other bacteria are capable of manganese regulation remains to be determined.

In the *Y. pestis* and *Y. enterocolitica* siderophore-mediated iron uptake system, many of the key elements involved in the biosynthesis, uptake, and regulation of the Ybt siderophore have been established (9, 29–31, 43, 59, 61). However, only the outer membrane receptor for the Ybt transport system has been identified (30, 31, 59, 61). In recent cell fractionation studies of *Y. pestis*, a number of iron-repressible periplasmic and cytoplasmic membrane proteins were detected, several of which were unrelated to the *pgm* locus (52, 59). The presence of the *yfe* locus in *Y. pestis* KIM6+ and KIM6, as determined by PCR (Fig. 7B) and Southern blot analyses (data not shown), suggests that the Yfe system functions independently of the *pgm*-linked Ybt siderophore-mediated uptake system of *Y. pestis* (9, 30). Furthermore, the presence of an uptake system which utilizes heme and hemoproteins (46, 58, 69, 73), an iron transport system that functions at 26°C but not at 37°C, and a Ybt-independent iron transport system functioning at 37°C (30, 52, 69) supports the hypothesis that multiple mechanisms for heme and iron acquisition in *Y. pestis* are fundamental to the success of plague pathogenesis. The lack of specific temperature-regulated expression from the *yfeABCD* promoter and the ability of the transport system to function at 37°C in *E. coli* SAB11 suggest that the *yfe* locus does not represent the 26°C iron transport system of *Y. pestis*. However, it may correspond to an iron transport system that, in the absence of a functional Ybt system, allows growth of *Y. pestis* cells in iron-deficient but not iron-chelated media at 37°C (30, 52, 69).

With at least four independent iron/heme transport systems, *Y. pestis* is in the company of a number of bacterial pathogens that have evolved several strategies for acquiring iron. *Vibrio cholerae*, for example, utilizes ferric citrate, heme-containing compounds, and a siderophore-mediated uptake system and produces an iron-regulated hemolysin which may facilitate the acquisition of iron or heme compounds liberated from lysed cells (75, 76). Similarly, *Shigella flexneri* uses siderophores and heme compounds to obtain iron but is also able to bind lactoferrin (56, 81). Receptors for both lactoferrin and transferrin, as well as heme and hemoprotein transport systems, are expressed by the pathogenic neisseriae and by *H. influenzae* (51, 54). The multiple systems for obtaining iron possessed by numerous bacterial pathogens emphasize the importance of iron acquisition to the pathogenic process. In addition, it is feasible that different iron/heme transport systems are required for the variety of environmental niches that pathogens inhabit—dif-

ferent organ systems within the host as well as nonhost environments. In the plague bacillus, *Y. pestis*, the role that the Yfe transport system plays in establishing or maintaining an infection and the conditions under which it is optimally used will likely provide new insights into the relationship between virulence and iron acquisition.

ACKNOWLEDGMENTS

This work was supported by Public Health Service grant AI33481 from the National Institutes of Health.

We thank B. R. Byers for providing *E. coli* SAB11.

REFERENCES

- Adhikari, P., S. A. Berish, A. J. Nowalk, K. L. Veraldi, S. A. Morse, and T. A. Mietzner. 1996. The *fbpABC* locus of *Neisseria gonorrhoeae* functions in the periplasm-to-cytosol transport of iron. *J. Bacteriol.* **178**:2145–2149.
- Adhikari, P., S. D. Kirby, A. J. Nowalk, K. L. Veraldi, A. B. Schryvers, and T. A. Mietzner. 1995. Biochemical characterization of a *Haemophilus influenzae* periplasmic iron transport operon. *J. Biol. Chem.* **270**:25142–25149.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Angerer, A., B. Klupp, and V. Braun. 1992. Iron transport systems of *Serratia marcescens*. *J. Bacteriol.* **174**:1378–1387.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1987. Current protocols in molecular biology. John Wiley & Sons, New York, N.Y.
- Barghouthi, S., S. M. Payne, J. E. L. Arceneaux, and B. R. Byers. 1991. Cloning, mutagenesis, and nucleotide sequence of a siderophore biosynthetic gene (*amoA*) from *Aeromonas hydrophila*. *J. Bacteriol.* **173**:5121–5128.
- Bartsevich, V. V., and H. B. Pakrasi. 1995. Molecular identification of an ABC transporter complex for manganese: analysis of a cyanobacterial mutant strain impaired in the photosynthetic oxygen evolution process. *EMBO J.* **14**:1845–1853.
- Bartsevich, V. V., and H. B. Pakrasi. 1996. Manganese transport in the cyanobacterium *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **271**:26057–26061.
- Bearden, S. W., J. D. Fetherston, and R. D. Perry. 1997. Genetic organization of the yersiniabactin biosynthetic region and construction of avirulent mutants in *Yersinia pestis*. *Infect. Immun.* **65**:1659–1668.
- Berg, D. E., A. Weiss, and L. Crossland. 1980. Polarity of Tn5 insertion mutations in *Escherichia coli*. *J. Bacteriol.* **142**:439–446.
- Bhattacharyya, P. 1970. Active transport of manganese in isolated membranes of *Escherichia coli*. *J. Bacteriol.* **104**:1307–1311.
- Birnboim, H. C., and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* **7**:1513–1523.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collada-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453–1474.
- Boos, W., and J. M. Lucht. 1996. Periplasmic binding protein-dependent ABC transporters, p. 1175–1209. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C.
- Braun, V., and K. Hantke. 1991. Genetics of bacterial iron transport, p. 107–138. *In* G. Winkelman (ed.), *Handbook of microbial iron chelates*. CRC Press, Inc., Boca Raton, Fla.
- Brickman, E., and J. Beckwith. 1975. Analysis of the regulation of *Escherichia coli* alkaline phosphatase synthesis using deletions and $\phi 80$ transducing phages. *J. Mol. Biol.* **96**:307–316.
- Calderwood, S. B., and J. J. Mekalanos. 1988. Confirmation of the Fur operator site by insertion of a synthetic oligonucleotide into an operon fusion plasmid. *J. Bacteriol.* **170**:1015–1017.
- Carniel, E., D. Mazigh, and H. H. Mollaret. 1987. Expression of iron-regulated proteins in *Yersinia* species and their relation to virulence. *Infect. Immun.* **55**:277–280.
- Chang, C. N., W.-J. Kuang, and E. Y. Chen. 1986. Nucleotide sequence of the alkaline phosphatase gene of *Escherichia coli*. *Gene* **44**:121–125.
- Chen, C.-Y., S. A. Berish, S. A. Morse, and T. A. Mietzner. 1993. The ferric iron-binding protein of pathogenic *Neisseria* spp. functions as a periplasmic transport protein in iron acquisition from human transferrin. *Mol. Microbiol.* **10**:311–318.
- Claros, M. G., and G. von Heijne. 1994. TopPred II: an improved software for membrane protein structure predictions. *CABIOS* **10**:685–686.
- Cornelissen, C. N., and P. F. Sparling. 1994. Iron piracy: acquisition of transferrin-bound iron by bacterial pathogens. *Mol. Microbiol.* **14**:843–850.
- de Bruijn, F. J., and J. R. Lupski. 1984. The use of Tn5 mutagenesis in the rapid generation of correlated physical and genetic maps of DNA segments

- cloned into multicopy plasmids—a review. *Gene* **27**:131–149.
24. De Lorenzo, V., F. Giovannini, M. Herrero, and J. B. Neilands. 1988. Metal ion regulation of gene expression. Fur repressor-operator interaction at the promoter region of the aerobactin system of pColV-K30. *J. Mol. Biol.* **203**: 875–884.
 25. Dintilhac, A., G. Alloing, C. Granadel, and J.-P. Claverys. 1997. Competence and virulence of *Streptococcus pneumoniae*: Adc and PsaA mutants exhibit a requirement for Zn and Mn resulting from inactivation of putative ABC metal permeases. *Mol. Microbiol.* **25**:727–939.
 26. Dintilhac, A., and J.-P. Claverys. 1997. The *adc* locus, which affects competence for genetic transformation in *Streptococcus pneumoniae*, encodes an ABC transporter with a putative lipoprotein homologous to a family of streptococcal adhesins. *Res. Microbiol.* **148**:119–131.
 27. Doige, C. A., and G. F.-L. Ames. 1993. ATP-dependent transport systems in bacteria and humans: relevance to cystic fibrosis and multidrug resistance. *Annu. Rev. Microbiol.* **47**:291–319.
 28. Fenno, J. C., A. Shaikh, G. Spatafora, and P. Fives-Taylor. 1995. The *fimA* locus of *Streptococcus parasanguis* encodes an ATP-binding membrane transport system. *Mol. Microbiol.* **15**:849–863.
 29. Fetherston, J. D., S. W. Bearden, and R. D. Perry. 1996. YbtA, an AraC-type regulator of the *Yersinia pestis* pesticin/yersiniabactin receptor. *Mol. Microbiol.* **22**:315–325.
 30. Fetherston, J. D., J. W. Lillard, Jr., and R. D. Perry. 1995. Analysis of the pesticin receptor from *Yersinia pestis*: role in iron-deficient growth and possible regulation by its siderophore. *J. Bacteriol.* **177**:1824–1833.
 31. Fetherston, J. D., and R. D. Perry. 1994. The pigmentation locus of *Yersinia pestis* KIM6+ is flanked by an insertion sequence and includes the structural genes for pesticin sensitivity and HMW2. *Mol. Microbiol.* **13**:697–708.
 32. Fetherston, J. D., P. Schuetze, and R. D. Perry. 1992. Loss of the pigmentation phenotype in *Yersinia pestis* is due to the spontaneous deletion of 102 kb of chromosomal DNA which is flanked by a repetitive element. *Mol. Microbiol.* **6**:2693–2704.
 33. Fleischmann, R. D., M. D. Adams, O. White, R. A. Clayton, E. F. Kirkness, A. R. Kerlavage, C. J. Bult, J.-F. Tomb, B. A. Dougherty, J. M. Merrick, K. McKenney, G. Sutton, W. FitzHugh, C. A. Fields, J. D. Gocayne, J. D. Scott, R. Shirley, L.-I. Liu, A. Glodek, J. M. Kelley, J. F. Weidman, C. A. Phillips, T. Spriggs, E. Hedblom, M. D. Cotton, T. R. Utterback, M. C. Hanna, D. T. Nguyen, D. M. Saudek, R. C. Brandon, L. D. Fine, J. L. Fritchman, J. L. Fuhrmann, N. S. M. Geoghagen, C. L. Gnehm, L. A. McDonald, K. V. Small, C. M. Fraser, H. O. Smith, and J. C. Ventor. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496–512.
 34. Genco, C. A., and P. J. Desai. 1996. Iron acquisition in the pathogenic *Neisseria*. *Trends Microbiol.* **4**:179–184.
 35. Gray-Owen, S. D., and A. B. Schryvers. 1996. Bacterial transferrin and lactoferrin receptors. *Trends Microbiol.* **4**:185–191.
 36. Griffiths, E. 1987. The iron-uptake systems of pathogenic bacteria, p. 69–137. In J. J. Bullen and E. Griffiths (ed.), *Iron and infection: molecular, physiological, and clinical aspects*. John Wiley & Sons, Inc., New York, N.Y.
 37. Guerinot, M. L. 1994. Microbial iron transport. *Annu. Rev. Microbiol.* **48**: 743–772.
 38. Guilvout, I., O. Mercereau-Pujalon, S. Bonnefey, A. P. Pugsley, and E. Carniel. 1993. High-molecular-weight protein 2 of *Yersinia enterocolitica* is homologous to AngR of *Vibrio anguillarum* and belongs to a family of proteins involved in nonribosomal peptide synthesis. *J. Bacteriol.* **175**:5488–5504.
 39. Gutierrez, C., J. Barondess, C. Manoil, and J. Beckwith. 1987. The use of transposon *TnphoA* to detect genes for cell envelope proteins subject to a common regulatory stimulus. *J. Mol. Biol.* **195**:289–297.
 40. Hantke, K. 1987. Selection procedure for deregulated iron transport mutants (*fur*) in *Escherichia coli* K12: *fur* not only affects iron metabolism. *Mol. Gen. Genet.* **210**:135–139.
 41. Hardham, J. M., L. V. Stamm, S. F. Porcella, J. G. Frye, N. Y. Barnes, J. K. Howell, S. L. Mueller, J. D. Radolf, G. M. Weinstock, and S. J. Norris. 1997. Identification and transcriptional analysis of a *Treponema pallidum* operon encoding a putative ABC transport system, an iron-activated repressor protein homolog, and a glycolytic pathway enzyme homolog. *Gene* **197**:47–64.
 42. Harkness, R. E., P. Chong, and M. H. Klein. 1992. Identification of two iron-repressed periplasmic proteins in *Haemophilus influenzae*. *J. Bacteriol.* **174**:2425–2430.
 43. Heesemann, J., K. Hantke, T. Vocke, E. Saken, A. Rakin, I. Stojiljkovic, and R. Berner. 1993. Virulence of *Yersinia enterocolitica* is closely associated with siderophore production, expression of an iron-repressible outer membrane polypeptide of 65,000 Da and pesticin sensitivity. *Mol. Microbiol.* **8**:397–408.
 44. Higgins, C. F. 1992. ABC transporters: from microorganisms to man. *Annu. Rev. Cell Biol.* **8**:67–113.
 45. Higgins, C. F., S. C. Hyde, M. M. Mimmack, U. Gileadi, D. R. Gill, and M. P. Gallagher. 1990. Binding protein-dependent transport systems. *J. Bioenerg. Biomembr.* **22**:571–592.
 46. Hornung, J. M., H. A. Jones, and R. D. Perry. 1996. The *hmu* locus of *Yersinia pestis* is essential for utilization of free haemin and haem-protein complexes as iron sources. *Mol. Microbiol.* **20**:725–739.
 47. Humphreys, G. O., G. A. Willshaw, and E. S. Anderson. 1975. A simple method for the preparation of large quantities of pure plasmid DNA. *Biochim. Biophys. Acta* **383**:457–463.
 48. Jackson, S., and T. W. Burrows. 1956. The pigmentation of *Pasteurella pestis* on a defined medium containing haemin. *Br. J. Exp. Pathol.* **37**:570–576.
 49. Jackson, S., and T. W. Burrows. 1956. The virulence-enhancing effect of iron on nonpigmented mutants of virulent strains of *Pasteurella pestis*. *Br. J. Exp. Pathol.* **37**:577–583.
 50. Kolenbrander, P. E., R. N. Andersen, and N. Ganeshkumar. 1994. Nucleotide sequence of the *Streptococcus gordonii* PK488 coaggregation adhesin gene, *scaA*, and ATP-binding cassette. *Infect. Immun.* **62**:4469–4480.
 51. Lee, B. C. 1995. Quelling the red menace: haem capture by bacteria. *Mol. Microbiol.* **18**:383–390.
 52. Lucier, T. S., J. D. Fetherston, R. R. Brubaker, and R. D. Perry. 1996. Iron uptake and iron-repressible polypeptides in *Yersinia pestis*. *Infect. Immun.* **64**:3023–3031.
 53. Manoil, C., and J. Beckwith. 1985. *TnphoA*: a transposon probe for protein export signals. *Proc. Natl. Acad. Sci. USA* **82**:8129–8133.
 54. Mietzner, T. A., and S. A. Morse. 1994. The role of iron-binding proteins in the survival of pathogenic bacteria. *Annu. Rev. Nutr.* **14**:471–493.
 55. Miller, J. F. 1992. A short course in bacterial genetics. A laboratory manual and handbook for *Escherichia coli* and related bacteria. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 56. Payne, S. M. 1988. Iron and virulence in *Shigella*. *Mol. Microbiol.* **3**:1301–1306.
 57. Pearce, B. J., Y. B. Yin, and H. R. Masure. 1993. Genetic identification of exported proteins in *Streptococcus pneumoniae*. *Mol. Microbiol.* **9**:1037–1050.
 58. Perry, R. D., and R. R. Brubaker. 1979. Accumulation of iron by yersiniae. *J. Bacteriol.* **137**:1290–1298.
 59. Perry, R. D., and J. D. Fetherston. 1997. *Yersinia pestis*—etiologic agent of plague. *Clin. Microbiol. Rev.* **10**:35–66.
 60. Perry, R. D., M. L. Pendrak, and P. Schuetze. 1990. Identification and cloning of the hemin storage locus involved in the pigmentation phenotype of *Yersinia pestis*. *J. Bacteriol.* **172**:5929–5937.
 61. Rakin, A., E. Saken, D. Harmsen, and J. Heesemann. 1994. The pesticin receptor of *Yersinia enterocolitica*: a novel virulence factor with dual function. *Mol. Microbiol.* **13**:253–263.
 62. Rakin, A. V., E. Saken, and J. Heesemann. 1997. Variability of iron acquisition systems in pathogenic *Yersinia*, abstr. B-377, p. 93. In Abstracts of the 97th General Meeting of the American Society for Microbiology 1997. American Society for Microbiology, Washington, D.C.
 63. Rodriguez-Quinones, F., and V. J. Benedi. 1993. *Escherichia coli* strain DH5 α is a suitable host for the study of *phoA* insertions. *Focus* **15**:110–112.
 64. Saken, E. M., and J. Heesemann. 1995. GenBank accession number Z47200.
 65. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 66. Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
 67. Saurin, W., W. Köster, and E. Dassa. 1994. Bacterial binding protein-dependent permeases: characterization of distinctive signatures for functionally related integral cytoplasmic membrane proteins. *Mol. Microbiol.* **12**:993–1004.
 68. Sikkema, D. J., and R. R. Brubaker. 1987. Resistance to pesticin, storage of iron, and invasion of HeLa cells by yersiniae. *Infect. Immun.* **55**:572–578.
 69. Sikkema, D. J., and R. R. Brubaker. 1989. Outer membrane peptides of *Yersinia pestis* mediating siderophore-independent assimilation of iron. *Biol. Metals* **2**:174–184.
 70. Silver, S. 1996. Transport of inorganic cations, p. 1091–1102. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C.
 71. Silver, S., and J. Lusk. 1987. Magnesium, manganese and zinc transport, p. 165–180. In B. P. Rosen and S. Silver (ed.), *Ion transport in prokaryotes*. Academic Press, Inc., New York, N.Y.
 72. Staggs, T. M., J. D. Fetherston, and R. D. Perry. 1994. Pleiotropic effects of a *Yersinia pestis fur* mutation. *J. Bacteriol.* **176**:7614–7624.
 73. Staggs, T. M., and R. D. Perry. 1991. Identification and cloning of a *fur* regulatory gene in *Yersinia pestis*. *J. Bacteriol.* **173**:417–425.
 74. Staggs, T. M., and R. D. Perry. 1992. Fur regulation in *Yersinia* species. *Mol. Microbiol.* **6**:2507–2516.
 75. Stoebner, J. A., J. R. Butterton, S. B. Calderwood, and S. M. Payne. 1992. Identification of the vibriobactin receptor of *Vibrio cholerae*. *J. Bacteriol.* **174**:3270–3274.
 76. Stoebner, J. A., and S. M. Payne. 1988. Iron-regulated hemolysin production and utilization of heme and hemoglobin by *Vibrio cholerae*. *Infect. Immun.* **56**:2891–2895.
 77. Stojiljkovic, I., A. J. Bäuml, and K. Hantke. 1994. Fur regulon in gram-negative bacteria. Identification and characterization of new iron-regulated *Escherichia coli* genes by a Fur titration assay. *J. Mol. Biol.* **236**:531–545.

78. **Surgalla, M. J., and E. D. Beesley.** 1969. Congo red-agar plating medium for detecting pigmentation in *Pasteurella pestis*. *Appl. Microbiol.* **18**:834–837.
79. **Tartof, K. D., and C. A. Hobbs.** 1987. Improved media for growing plasmid and cosmid clones. *Focus* **9**:12.
80. **Thompson, J. D., D. G. Higgins, and T. J. Gibson.** 1994. CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
81. **Tigyi, Z., A. R. Kishore, J. A. Mæland, A. Forsgren, and A. S. Naidu.** 1992. Lactoferrin-binding proteins in *Shigella flexneri*. *Infect. Immun.* **60**:2619–2626.
82. **Une, T., and R. R. Brubaker.** 1984. In vivo comparison of avirulent Vwa⁻ and Pgm⁻ or Pst⁺ phenotypes of yersiniae. *Infect. Immun.* **43**:895–900.
83. **Wake, A., M. Misawa, and A. Matsui.** 1975. Siderochrome production by *Yersinia pestis* and its relation to virulence. *Infect. Immun.* **12**:1211–1213.
84. **Wee, S., J. B. Neilands, M. L. Bittner, B. C. Hemming, B. L. Haymore, and R. Seetharam.** 1988. Expression, isolation and properties of Fur (ferric uptake regulation) protein of *Escherichia coli* K 12. *Biol. Metals* **1**:62–68.
85. **Zimmermann, L., A. Angerer, and V. Braun.** 1989. Mechanistically novel iron(III) transport system in *Serratia marcescens*. *J. Bacteriol.* **171**:238–243.