

Melanin Biosynthesis in *Cryptococcus neoformans*

PETER R. WILLIAMSON,^{1*} KAZUMASA WAKAMATSU,² AND SHOSUKE ITO²

*Division of Infectious Diseases, University of Illinois at Chicago College of Medicine, Chicago, Illinois 60612,¹
and Fujita Health University School of Health Sciences, Toyoake, Aichi 470-1192, Japan²*

Received 17 November 1997/Accepted 13 January 1998

Pigment production by *Cryptococcus neoformans* is virulence associated. Dopamine- and 3,4-dihydroxyphenylalanine–melanin products were identified after acidic permanganate oxidation, alkaline hydrogen peroxide oxidation, or hydrolysis with hydriodic acid. These data provide direct chemical evidence for the formation of eumelanin polymers by catecholamine oxidation by laccase alone followed by oxidative coupling of dihydroxyindole.

Melanin, produced by the enzyme laccase, has been thought to be a major virulence factor in *Cryptococcus neoformans* since its first description by Staib (17). However, the chemical structure of *C. neoformans* pigment has not been established; only indirect studies implicate the presence of eumelanin-type pigments. Pigment production is dependent on the presence of *o*- or *p*-dihydroxybenzenes or diaminobenzenes (2), and the intermediate dopaminechrome involved in eumelanin formation is produced from purified laccase (20). The metal environment is similar to that of eumelanins as determined by electron paramagnetic resonance spectroscopy, although an unusually high carbon-to-nitrogen ratio in melanin particles prompted Wang et al. to conclude that the structure of *Cryptococcus* melanin differs from those of mammalian and other fungal eumelanins (19). Knowledge of the structure of *Cryptococcus* melanin is important not only for the accurate simulation of pigments for immunological experiments but also for the identification of possible ancillary enzymatic activities that may have major impacts on virulence mechanisms and their regulation in this important pathogen.

To understand the biosynthesis of eumelanin pigments in *C. neoformans* more fully, we used recently developed microanalytical high-performance liquid chromatography methods to chemically quantify and characterize cryptococcal pigment, based on the formation of specific degradative products (7, 8). Representative isolates of *C. neoformans*, namely, serotype A (H99) (14), serotype D (ATCC 34873) (9), and a set of isogenic *CNLAC1* and *cnlac1* strains (2E-TUC4 and 2E-TU4, respectively) (16), were each incubated on asparagine agar (pH 6.5) containing 100 mg of either dopamine (DA) or 3,4-dihydroxyphenylalanine (DOPA) per liter for 2 days at 30°C. Recombinant laccase was expressed with plasmid pPIC93 (Invitrogen, San Diego, Calif.) in *Pichia pastoris* (4) and used to generate pigment (isolate PIC93). (Recombinant laccase was purified by methods used to purify the native enzyme [20] and characterized by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, which showed a single band of approximately 100 kDa, consistent with a glycosylated laccase-*Pichia* leader sequence fusion protein.) Laccase activities were determined (20) on cells induced on asparagine agar as described above and were as follows: 70.0 U/10⁷ cells with H99, 31.3 U/10⁷ cells with ATCC 34873, 82.0 U/10⁷ cells with PIC93, 15.0 U/10⁷ cells with 2E-TUC4, and <1 U/10⁷ cells with 2E-TU4.

Solubilization of melanin from 10 million cells by Soluene-350 as described previously (8) produced an A_{500} proportional to the amount of pigment observed in cryptococcal cells and proportional to laccase activity (Table 1). Notable was the low amount of solubilized pigment from PIC93 relative to the cryptococcal strains in spite of high laccase activity (82.0 U/10⁷ cells). This may be due to the ability of the anionic cell wall and capsule of *C. neoformans* (13) to adsorb greater amounts of melanin pigments similar to the property of anionic carboxymethylcellulose beads (3). Laccase-negative 2E-TU4 produced a minimum of pigment at pH 6.5, but it did not produce pigment at a lower pH of 5.2, consistent with auto-oxidation of the catecholamine at the higher pH (10).

Acidic permanganate oxidation of *Cryptococcus* pigment was performed as described previously (7) on 5 million cells grown on DA agar. Pyrrole-2,3,5-tricarboxylic acid (PTCA), a specific breakdown product of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) polymeric units, was detected (7). Relative yields were less from strains having less laccase activity, with the laccase-negative strain 2E-TU4 having a minimal level of PTCA due to auto-oxidation (Table 1).

In order to assess for possible copolymerization of cysteine to form pheomelanins, reductive hydrolysis of *Cryptococcus* pigment with hydriodic acid (HI) from 5 million cells was performed (8). Aminohydroxyphenylalanine (AHP) is a specific indicator of pheomelanins (7), and its presence was confirmed in this study by subjecting various synthetic pheomelanins (obtained as described in reference 8) to reductive hydrolysis with HI (Table 1). AHP was not found in significant quantities from cryptococcal pigment, showing that cysteine is not incorporated in the form of pheomelanin (Table 1). Detectable amounts of aminohydroxyphenylethylamine (AHPEA) were found with cells grown on DA but not on DOPA. AHPEA has been proposed to be an indicator of cysteinyl-DA-derived units in melanin (1, 18) but has recently been found in DA-melanin as well (8). AHPEA found in this study is most likely due to DA-melanin formed from cells grown on DA, since (i) the ratio of AHPEA to A_{500} from strain H99 (0.50) was similar to that of synthetic DA-melanin (0.40) and (ii) the ratio of the 3-amino isomer of AHPEA to the 4-amino isomer (0.80) from strain H99 was similar to that of synthetic DA-melanin (0.95) (8).

In order to clarify this issue further, additional studies which showed that AHPEA is formed from DA-melanin and that oxidation by alkaline H₂O₂ could be used to identify DA-melanin were undertaken (8). Oxidation by alkaline H₂O₂ was performed on 5 million cells by the method of Napolitano (11)

* Corresponding author. Mailing address: Rm. 888, m/c 735, 808 S. Wood St., Chicago, IL 60612. Phone: (312) 996-6070. Fax: (312) 996-5704. E-mail: prw@uic.edu.

TABLE 1. Chemical characterization of melanins in *C. neoformans* in comparison with that of synthetic melanins

Melanin (quantity)	A_{500} (1,000) after solubilization ^a	H_2O_2/OH^- production of ^b :		KMnO ₄ /H ⁺ production of PTCA ^b	HI hydrolysis production of ^b :		
		PTCA	PDCA		AHPEA	AHP	
<i>C. neoformans</i> in DA medium (5×10^6 cells)							
2E-TU4	2.8	2.3	0.4	0.3	0.4	1.1	
2E-TUC4	6.8	3.1	1.9	1.3	1.9	0.9	
H99	133	9.4	35.2	4.5	66.3	1.1	
ATCC 34873	10.8	2.0	3.4	1.5	2.3	0.5	
PIC93	4.5	1.4	1.0	0.9	1.0	1.1	
<i>C. neoformans</i> in DOPA medium (5×10^6 cells)							
2E-TU4	6.0	9.8	1.5	0.6	<0.2	1.1	
2E-TUC4	17.5	11.4	3.6	3.1	<0.2	1.6	
H99	71.0	32.9	9.3	17.0	<0.2	1.5	
ATCC 34873	9.8	6.6	1.4	3.9	<0.2	1.3	
PIC93	3.0	1.3	0.0	1.9	<0.2	1.1	
Synthetic (0.1 mg) ^c							
DA-melanin	769	283	377	186	304	0	
DOPA-melanin	954	429	167	255	0	6	
DHI-melanin	941	329	166	206	ND ^d	ND	
DHICA-melanin	467	5,350	<4	2,490	ND	ND	
DA + Cys-melanin ^e	435	41	534	18	12,200	26	
5-S-Cysteiny-DOPA-melanin	437	708	430	38	<20	17,500	

^a Absorbance unit times 1,000 per 5×10^6 cells or per 0.1 mg of melanin dissolved in 1 ml of Soluene-350 plus water (9:1). Values are averages of two determinations.

^b Values are expressed in nanograms per 5×10^6 cells or per 0.1 mg of melanin. Except for AHP values (single determinations), values are averages of two determinations.

^c Data cited are from reference 8.

^d ND, not determined.

^e Prepared from a mixture of DA (10 mM) and cysteine (5 mM). The N/S ratio indicates a DA/Cys-DA molar ratio of ca. 1:3 (18).

as modified by Ito and Wakamatsu (8). H_2O_2 oxidation of *Cryptococcus* cells produced, in addition to PTCA, pyrrole-2,3-dicarboxylic acid (PDCA), a specific degradation product of 5,6-dihydroxyindole (DHI) units. High PDCA/PTCA ratios from pigment obtained from growth on DA agar (Table 1) indicate that AHPEA derived from *Cryptococcus* is from DA-melanin and not cysteinyl-DA-melanin.

To assess DOPA decarboxylase and dopachrome tautomerase activities, strains of *Cryptococcus* were incubated in the

presence of DOPA as described above. High PTCA/PDCA ratios obtained after oxidation by alkaline H_2O_2 (Table 1) showed that DOPA-melanin was produced without DOPA decarboxylation. This result differentiates melanogenesis in *Cryptococcus* from that in other systems, such as *Drosophila melanogaster*, where DOPA decarboxylase activity is required for effective melanogenesis and protection from parasites (12). In addition, the similarity of DOPA-melanin of *C. neoformans* to synthetic DOPA-melanin argues against ancillary dopachrome tautomerase activity in the fungus. This supports a Mason-Raper-derived scheme of melanogenesis (Fig. 1) wholly dependent on laccase (15). Finally, lack of AHP after HI hydrolysis of cells grown on DOPA medium shows that, unlike in mammalian systems, these *Cryptococcus* strains do not incorporate cysteine from intrinsic sources into melanin. We found nothing to suggest any unusual structure in the melanin polymer, in contrast to the structure suggested by a high C/N ratio found in an elemental analysis of melanin "ghosts" by Wang et al. (19). This may be a strain-related phenomenon, or the previously reported high C/N ratio may have resulted from impurities which are difficult to remove from melanin (5) or from the introduction of impurities during isolation procedures (6).

P.R.W. was supported by NIH grant AI38258.

We are grateful for the gifts of *Cryptococcus* strains H99 (J. Perfect) and ATCC 34873/B-3501 (K. J. Kwon-Chung).

REFERENCES

- Carstam, R., A. Brinck, A. Hindemith-Augustsson, H. Rorsman, and E. Rosengren. 1991. The neuromelanin of the human substantia nigra. *Biochim. Biophys. Acta* **1097**:152-160.
- Chaskes, S., and R. L. Tyndall. 1975. Pigment production by *Cryptococcus neoformans* from *para*- and *ortho*-diphenols: effect of the nitrogen source. *J. Clin. Microbiol.* **1**:509-514.
- Gorman, M. J., and S. M. Paskewitz. 1997. A genetic study of a melanization




FIG. 1. Melanogenesis from DA in *C. neoformans*.

- response to Sephadex beads in *Plasmodium*-refractory and -susceptible strains of *Anopheles gambiae*. *Am. J. Trop. Med. Hyg.* **56**:446–451.
4. **Grinna, L. S., and J. F. Tschopp.** 1989. Size distribution and general structural features of N-linked oligosaccharides from the methylotrophic yeast *Pichia pastoris*. *Yeast* **5**:107–115.
 5. **Ito, S.** 1993. Biochemistry and physiology of melanin, p. 33–59. *In* N. Levine (ed.), *Pigmentation and pigmentary disorders*. CRC Press, Boca Raton, Fla.
 6. **Ito, S.** 1986. Reexamination of the structure of eumelanin. *Biochim. Biophys. Acta* **883**:155–161.
 7. **Ito, S., and K. Fujita.** 1985. Microanalysis of eumelanin and pheomelanin in hair and melanomas by chemical degradation and liquid chromatography. *Anal. Biochem.* **144**:527–536.
 8. **Ito, S., and K. Wakamatsu.** Chemical degradation of melanins. Application to identification of dopamine-melanin. *Pigm. Cell Res.*, in press.
 9. **Kwon-Chung, K. J., J. C. Edman, and B. L. Wickes.** 1992. Genetic association of mating types and virulence in *Cryptococcus neoformans*. *Infect. Immun.* **60**:602–605.
 10. **Lin, J., and B. M. Christensen.** 1994. Effect of pH on the oxidation pathway of dopamine and DOPA. *J. Electroanal. Chem.* **375**:219–231.
 11. **Napolitano, A., A. Pezella, M. R. Vincensi, and G. Protta.** 1995. Oxidative degradation of melanins to pyrrole acids: a model study. *Tetrahedron* **51**:5913–5920.
 12. **Nappi, A. J., Y. Carton, and E. Vass.** 1992. Reduced cellular immune competence of a temperature-sensitive dopa decarboxylase mutant strain of *Drosophila melanogaster* against the parasite *Leptopilina boulardi*. *Comp. Biochem. Physiol. B.* **101**:453–460.
 13. **Nosanchuk, J. D., and A. Casadevall.** 1997. Cellular charge of *Cryptococcus neoformans*: contributions from the capsular polysaccharide, melanin, and monoclonal antibody binding. *Infect. Immun.* **65**:1836–1841.
 14. **Perfect, J. R., S. D. Lang, and D. T. Durack.** 1980. Chronic cryptococcal meningitis: a new experimental model in rabbits. *Am. J. Pathol.* **101**:177–194.
 15. **Polacheck, L., and K. J. Kwon-Chung.** 1988. Melanogenesis in *Cryptococcus neoformans*. *J. Gen. Microbiol.* **134**:1037–1041.
 16. **Salas, S. D., J. E. Bennett, K. J. Kwon-Chung, J. R. Perfect, and P. R. Williamson.** 1996. Effect of the laccase gene *CNLAC1* on virulence of *Cryptococcus neoformans*. *J. Exp. Med.* **184**:377–386.
 17. **Staib, F.** 1962. *Cryptococcus neoformans* and *Guizotia abyssinica* (syn. *G. oleifera* D.C.) (farbreadaktion für *C. neoformans*). *Z. Hyg. Infektionskr.* **148**:466–475.
 18. **Wakamatsu, K., S. Ito, and T. Nagatsu.** 1991. Cysteinyldopamine is not incorporated into neuromelanin. *Neurosci. Lett.* **131**:57–60.
 19. **Wang, Y., P. Aisen, and A. Casadevall.** 1996. Melanin, melanin “ghosts,” and melanin composition in *Cryptococcus neoformans*. *Infect. Immun.* **64**:2420–2424.
 20. **Williamson, P. R.** 1994. Biochemical and molecular characterization of the diphenol oxidase of *Cryptococcus neoformans*: identification as a laccase. *J. Bacteriol.* **176**:656–664.