

Cloning and Sequencing of a Form II Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase from the Bacterial Symbiont of the Hydrothermal Vent Tubeworm *Riftia pachyptila*

JONATHAN J. ROBINSON,¹ JEFFREY L. STEIN,² AND COLLEEN M. CAVANAUGH^{1*}

Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts 02138,¹ and Diversa Corp., San Diego, California 92121²

Received 4 August 1997/Accepted 13 January 1998

The bacterial symbiont of the hydrothermal vent tubeworm fixes carbon via the Calvin-Benson cycle and has been shown previously to express a form II ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO). The gene *cbbM*, which encodes this enzyme, has been cloned and sequenced. The gene has the highest identity with the *cbbM* gene from *Rhodospirillum rubrum*, and analysis of the inferred amino acid sequence reveals that all active-site residues are conserved. This is the first form II RubisCO cloned and sequenced from a chemoautotrophic symbiont and from a deep-sea organism.

Hydrothermal vent environments are dominated by dense assemblages of invertebrates which harbor chemoautotrophic sulfur-oxidizing bacteria within their tissues. This nutritional interaction between prokaryotic symbionts and various animal hosts is dependent upon the biological fixation of inorganic carbon by the symbionts and the subsequent supply of organic carbon to the host in a manner analogous to the chloroplasts of green plants and algae (reviewed in reference 2). Fundamental to the initial and subsequent characterization of these symbioses has been the detection of the key Calvin-Benson cycle enzyme, ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO) (1, 8). The hydrothermal vent tubeworm *Riftia pachyptila* is of particular interest with regard to its carbon fixation abilities, as this animal completely lacks a mouth, gut, or anus (13) but is capable of extreme size and high growth rates due to its symbiotic association (16).

The primary carbon fixation step in the Calvin-Benson cycle is catalyzed by RubisCO, which carboxylates ribulose-1,5-bisphosphate with CO₂ to yield two molecules of 3-phosphoglyceric acid. The enzyme is found in two forms, called form I and form II (30), which are distinct in primary and quaternary structure (26), reaction mechanism, and kinetic isotope effect (KIE) (22, 23). The form I RubisCO, found in the vast majority of eukaryotic and prokaryotic autotrophs, consists of eight large subunits and eight small subunits, with the holoenzyme having a molecular weight between 500 and 560 kDa (26). The form II enzyme is structurally less complicated, consisting of a dimer of only two large (L) subunits found in either an L₂ configuration (26), as in *Rhodospirillum rubrum*, or an L₂-to-L₆ configuration, as reported for other species (30). The two forms are ~25% identical to each other at the amino acid level (18). To date, the form II enzyme has only been characterized at the nucleic acid sequence level from five prokaryotes (31) and two dinoflagellates (17, 24).

The bacterial symbiont of the vestimentiferan *R. pachyptila* has been shown to express a form II RubisCO (21). In this work we report the cloning and sequencing of the *cbbM* gene,

which encodes a form II enzyme, from the *R. pachyptila* symbiont.

Bacterial strains, plasmids, and polyclonal antisera. The *Escherichia coli* construct pRR2119 (ATCC 37846) was used to generate probes for hybridization during library screening. This clone harbors the plasmid pXG9 containing the cloned form II RubisCO from *Rhodospirillum rubrum* (28). *E. coli* INF^aF' (Invitrogen) was used for cloning steps and grown in Luria broth supplemented with ampicillin (40 mg liter⁻¹). Plasmid pCRII (Invitrogen) was used for subcloning, DNA sequence analysis, and protein expression studies.

The lambda DNA library (see below) was screened with polyclonal antiserum directed against the *R. rubrum* form II RubisCO (anti-RrFII) (antiserum generously provided by George Lorimer [DuPont]), which has been shown to be specific to form II RubisCOs and to cross-react with the *R. pachyptila* enzyme (21). In all cases antiserum was used at a 1:3,000 dilution.

***R. pachyptila* genomic DNA library construction.** Tubeworm specimens used for genomic DNA library construction were collected from a depth of 2,600 m using the DSV *Alvin* from hydrothermal vents on the East Pacific Rise at the 13°N site (12°48'N, 103°56'W; November 1987). The worms were transported to the surface in a thermally insulated container and the symbiont-containing trophosome tissue was immediately dissected on board ship. Tissue was homogenized in a 1:1 (wt/vol) solution of ice-cold *Riftia* saline (46 mM imidazole, 0.46 M NaCl, 30 mM MgSO₄, 2.5 mM KCl, 10 mM CaCl₂; pH 7.1) at 30 to 40% speed in an Ultraturrax homogenizer for 2 min on ice. Symbionts in this solution were separated from host cells on 80% Percoll density gradients according to the method of Distel and Felbeck (6) with modifications.

DNA was extracted from the symbiont preparation by using a 5 M guanidinium isothiocyanate solution (15). DNA (75 μg) was sheared to an average size of 3 to 6 kbp by vigorous passage through a 25-gauge needle in a 1-ml syringe. The sheared DNA was blunt ended with mung bean nuclease and ligated to *Eco*RI linkers, and 3- to 6-kbp fragments were cloned into lambda gt11 (27). The library titer was estimated to be 1.5 × 10¹⁰ PFU (25).

Library screening. Phage were plated and screened by standard methods on a lawn of *E. coli* Y1090 (25). Plaques were screened for the expression of the form II RubisCO by incu-

* Corresponding author. Mailing address: Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Ave., Cambridge, MA 02138. Phone: 617-495-2177. Fax: 617-496-6933. E-mail: cavanaugh@hump.harvard.edu.

bation with anti-RrFII antiserum (25). Plaques which were immunologically positive were rescreened with a ³²P-labelled *Bgl*III/*Sma*I fragment of the *R. rubrum* form II RubisCO derived from plasmid pXG9 (28).

Two lambda clones were isolated. Inserts were amplified from purified lambda DNA by PCR with primers specific to the *lacZ* cloning region (Promega) and subcloned into the pCR II vector for transformation into *E. coli* host strain INF^αF' and subsequent DNA sequencing. The two clones are different sizes, with insert sizes of ~3,300 bp for pRpR-1 and ~2,200 bp for pRpR-2, and are oriented in opposite directions.

DNA sequence analysis. Oligonucleotide primer walking was used to generate a double-stranded sequence for the region encoding the form II RubisCO and immediate flanking regions for both clones. Sequencing was conducted with the Applied Biosystems Inc. (ABI) Dye Terminator Cycle Sequencing kit under standard conditions, an Ericomp thermal cycler, and an ABI model 373 automated sequencer. Sequences of regions flanking the *Eco*RI cloning sites were also determined, by using the M13 universal primers designated reverse and -40 forward.

Sequencing of a 1,678-bp region from both clones revealed open reading frames with high identity to previously sequenced *cbmM* genes (Fig. 1 and 2). *cbmM* is preceded by an in-frame TAG stop codon at position -9 and then begins with an ATG and proceeds 1,383 bp to a TAA stop codon, followed by a putative hairpin loop beginning 27 bp downstream (Fig. 1). The *cbmM* coding region is composed of 57.5 mol% G+C, and a 461-amino-acid protein with a calculated molecular weight of 50,552 Da is predicted. Efforts to express active recombinant form II RubisCO failed to yield enzyme with significant activity, suggesting that the recombinant does not fold properly in *E. coli* or is posttranslationally modified by the bacterial symbiont. Therefore, biochemical characterization of this RubisCO is currently being conducted on native enzyme.

Analysis of sequence flanking the *Eco*RI cloning sites revealed the presence of an open reading frame sharing identity to the LysR type regulator *cbmR* (not shown). This gene is upstream of *cbmM* and in the opposite orientation. The deduced amino acid sequence of the *cbmR* element has 61% identity with the *cbmR* of *Chromatium vinosum* (31) over the region sequenced, which spans 71 residues at the 5' end.

Translation of the open reading frame and alignment with other form II enzymes and a representative form I RubisCO (Fig. 2) revealed strict conservation of residues known to form the enzyme active site (11, 26), e.g., the specific lysine residue which is carbamylated during enzyme activation and corresponds to position 191 of the *R. rubrum* sequence. N-terminal sequence analysis indicates that the first-position methionine residue is posttranslationally cleaved (3), a situation encountered in plant RubisCO enzymes (12). Amino acid identity with other form II RubisCOs ranges from a high of 76.2% with *R. rubrum* to a low of 69.1% with the dinoflagellate *Gonyaulax polyedra*. With regard to amino acid similarity, i.e., by comparison of major amino acid biochemical groupings, the *R. pachyptila* enzyme is most similar (89.2%) to the *Rhodobacter sphaeroides* form II enzyme and shows 78 to 89% similarity with all the other form II enzymes. In contrast, the *R. pachyptila* RubisCO shows only 22 to 32% amino acid identity with the gene encoding the large subunit of representative form I RubisCOs, including that of *Spinacia oleracea*.

The discovery of a form II RubisCO in a deep-sea organism indicates that this enzyme is found in diverse settings and is not as rare as once thought. Indeed, six other deep-sea symbionts and two bacterial mats have recently been shown to express this form of RubisCO (3, 20, 21). While both forms of

```

1  AACCATAGATTTACCTTATCATGCTGATCAAACATTTTAGTTTATTTATATGTGCACC
61  GAGGGTAAGGTCACACCCAGTCCCACCATGATGTGGAATCGATAGGCAGACCTCCG
121  GGTTCGCCGACAATCAACACAATCGACTCGATATGGAGTAGATCAACATGGCATTAGAC
      M A L D
181  CAAACCAATCGTTATTCAGACCTGAGTTTGAAGAAGAGAGCTGATCGCAAGCGGTGAC
5  Q T N R Y S D L S L K E D E L I A S G D
241  TAGCTACTTTGGCCTACCTGATGAAGCCAAAGTCCGGCTACGGTTACCTGGAAGCTGCT
25  Y V L C A Y L M K N P K S G M Y G Y L E A A
301  GCTCACITTCGAGCTGAGTCCCTCCACCGCACCACCGTGGAAAGTCTCTACCACATGATGAC
45  A H F A A E S S T G T N V E V S T T D D
361  TTCACCAAGGGTGTGACCGCTGGTCTATGAGATCGATGAAGCCAAAGAGCTGATGAAG
65  F T K G V D A L V Y E I D E N G I K E L M K
421  ATCGCCTACCCCGTGTGACTTATTCGATATCAACATCATCGAGGGCGTCCCATGCTGGCA
85  I A Y P V D L F D I N I I D G R A M L A
481  TCGTTCCTGACCCCTGACCATCGGTAAACACCGAGGTATGGGGGATATCGAATACGCCAAG
105  S F L T L T I G N N P I E N G I G Y I A K
541  ATGCTCGACTTCTACATGCCACCCAAATACCTGCGCGTGTACGACGGTCCGCGGTCAAC
125  M L D F Y M P P K Y L R L Y D G P A V N
601  ATCCAGGATATGTGGCCATTCCTGGGCGTCCAATGGAAGATGGTGGTTACATCGCCGGC
145  I Q D M W R I L G R P I E N G I G Y I A K
661  ACCATCATCAAACCAAACTGGGCGTCCGCTCCCGAGCCCTTTGCTGAAGCGGCATACCAG
165  T I I K P K L G L R P E P F A E A A Y Q
721  TTCTGGCTGGGGGTGACTTCATTAAGAAGCAGCCGCCAGGGTAACCGACCCCTTCTCC
185  F W L G G D F I K N D E P A E M I A
781  CCCATGAAGAAGACCATCCCGCTGGTGGCAGACGCCATCGCTCGCGCCAGGATGAGACT
205  P M K K T I P L V A D A M R R A Q D E T
841  GCGAAGCCAGCTCTCTCCGCCAACATCACCCTGATGATCCAGCCGAGATGATTGCC
225  G E A K L F S A N I T A D D P A E M I A
901  CGTGTGAGTTCGCTCCGAGACCTCCGCTTTTGGGCTCTCAGGTTGCCTTCTCGGTT
245  R G E F V L E T F G F E A S Q V A F L V
961  GATGTAGCTAGCCGCTTACCGCTGTTCGCCACCGCTCCGCACTCCCAACCCAG
265  D G Y V A G P T A V A T A R R N P P N Q
1021  TTCTGCACTTCCATCGTCCGGTCCAGTTCGCTCCCTCCCGAGGACCGCTGGT
285  F L H F H R A G H G A V T S P Q S K R G
1081  TACACCGCTTTCGTACACATCAAGATGACCCGCTCTGCTGGTGCATCCGGTATGCACGTC
305  Y T A F V H I K M T R L L G A S G M H V
1141  GGCACCATGGCTACGGCAAGATGGAAGCGCAGCCAGGACAACTGATTGCCCTACATG
325  G T M G Y G K M E G E A S D K L I A Y M
1201  ATCGAGGCTGACAGCGCGATGGTCTTTCTACATCGAGGAGTGGGTGGCATGAAACCC
345  I E R D S A D G P F Y H Q E W A G M K P
1261  ACCACCCGATCATCTCCGGCGCATGAACCGCTTCGGTCTACCCGGCTTCTCGAGAAC
365  T P T I I S G G M N A L R L P G F P E N
1321  CTGGGCCACGGCAACGATCAACACCGAGTGGTGGTACTTACGGTACATCGACAGC
385  L G H G N V I N T A G G G T Y G H I D S
1381  CCGCGCGTGGTGTCTCCCTGCTGACGCTACGAGTGTCTGGAAGAGGGTGCAGAC
405  P A A G A V S L R Q A Y E C W K E G A D
1441  CCGTTCGAGTACGCGAAAGAGCAAAAGTTCGCTCCCGCTTCGAGTCTTTCGCGCAC
425  P V E Y A K E H K E F A R A F E S F P H
1501  GATCGGACGCTATCTCCCGGATGGCCGACAAAGCTGGCGTACACAA*TAATCGAAC
445  D A D A I F P G W R D K L G V H K
1561  GGTTACTGTACATCCAGTGCAAAAAAGCCCGCTTCGGCGGGGCTTTTTTTTATCCCTT
1621  GACTTTACTGCGCGCCCGCCAGCCACACATCTGTACCCCTTCTCCCAACACAG
    
```

FIG. 1. Nucleic acid sequence of the *R. pachyptila* *cbmM* gene. The deduced amino acid sequence of the form II RubisCO is shown, with the putative Shine-Dalgarno sequence and hairpin loop underlined. An asterisk marks the TAA stop codon.

RubisCO are expressed in some free-living bacteria (30, 31), the *R. pachyptila* symbiont appears to encode and express only a form II enzyme. In the *R. pachyptila* symbiosis, hybridization to a form I heterologous gene probe was not detected during library screening or Southern analysis of trophosome DNA, in contrast to earlier reports (29, 32), nor was a form I gene product detected (21). Indeed, other researchers have also failed to detect the gene encoding the form I enzyme in the *R. pachyptila* symbiont, detecting only the *cbmM* gene (14). Physiologically, the use of a form II RubisCO in this symbiosis is

<i>R. pachyptila</i>	MALDQ	TWRYSDLSIK	EBELIASGDY	VLCAYLMLPK	35	
<i>R. rubrum</i>	M	SS-VN-A-K	ED-V-G-EHI.....	33	
<i>R. sphaeroides</i>	M	S-AR-D-Q	AD-G-RHV.....	33	
<i>T. denitrificans</i>	M	SA-A-.....	ED-KG-RH	L-V-K-.....	33	
<i>H. marinus</i>	M	S-A-T-T	EK-V-D-NH	L-V-RL-A	33	
<i>G. polyedra</i>	...	SS-A-.....	ED-VKN-KHV-YI-NQ	32	
Symbiodinium spp.	...	SS-A-.....	T-RN-KHV-YI-NQ	33	
<i>S. oleracea</i>	MSPQTEKAS	VEFKAGVKDY	KLT-YTPEYE	-----TLDTD	I-A-PRVS-Q	45
* * *						
<i>R. pachyptila</i>	SGVGLLEAA	HFAESSVGT	NVEVST-TDD	PTKGDVALVY	EIDEA-----	79
<i>R. rubrum</i>	A-...VAT...C.....R.....V.....H....	77
<i>R. sphaeroides</i>	A-...T...R.....SPE...	77
<i>T. denitrificans</i>	A-...F-V...R.....AFGDK	82
<i>H. marinus</i>	G-D-AT...N-C.....T.....Y-PE...	76
<i>G. polyedra</i>	A-D-AT...P.....N-C.....S.....Y-PD...	77
Symbiodinium spp.	P-VPPE-G	AV.....	WTPVW-DGLT	NLDRYKRC	H-EPVAGEEN	95
<i>S. oleracea</i>
* * *						
<i>R. pachyptila</i>	KELMKIAYPV	DEPDINLDG	RAMLASPLTL	TIGNNQMGD	IEYARMLDFY	129
<i>R. rubrum</i>	R-T-.....	A-R-T-K	I-.....	M-.....	V-.....H...	127
<i>R. sphaeroides</i>I-.....	E-ER.....C.....V.....H...	127
<i>T. denitrificans</i>	S-D-RV-L	E-ER-VT...	F-V-.....	A-.....H-I...	127
<i>H. marinus</i>	GG-.....P-L...	HYNVSRMWS	IL-.....	H-GLR-...L	132
<i>G. polyedra</i>	V-ERICQMP	-TD-SMLV-V...S.....	V-G-IT-T	126
Symbiodinium spp.	N-E-.....	P-R-T-...	G-MC-...	A-.....	V-G-IV...	127
<i>S. oleracea</i>	QYICV-...LE-.....	RGSVTNMF-S	IV-VF-FKA	LRLALR-LR	139
* * *						
<i>R. pachyptila</i>	MPPKYLRLYD	GPVNIQDMW	RILGRPIENG	GVIAGTIIPK	KLGLRPEPPA	179
<i>R. rubrum</i>	V-ER-RA-PS-SAL	KV-.....EVD	LAV-.....K...	177
<i>R. sphaeroides</i>	V-C-...P...	SM-A-.....	V-DVR-...	MVV-.....K...	177
<i>T. denitrificans</i>	V-E-MVRRP	-TD-S-L	KV-.....EVD	182
<i>H. marinus</i>	F-S-...FF...	CSIL-...	DMTD-...	LUV-.....Q-K-G	176
<i>G. polyedra</i>	L-AF-L-...	S-V-VE...	GTT-...	LUV-.....Q-K-G	177
Symbiodinium spp.	I-VA-VKTEQPHG-VF	DK-NKY-...	RPL-CT-...SAKNYG	186
<i>S. oleracea</i>
* * *						
<i>R. pachyptila</i>	EAAYQFVLWG	-DFIKNDPQ	GNQFSPMKK	TIPLVADAMR	RAQDETEGAK	228
<i>R. rubrum</i>	-CHE-.....	A-LD-...	226
<i>R. sphaeroides</i>	D-CHE-.....	T-A-L-E	-R-.....K...	226
<i>T. denitrificans</i>	K-.....	V-C-L-K	V-.....Y-...KD-Q...	226
<i>H. marinus</i>	K-C-D-.....	A-N-C-EV	V-K-E-E-DQA-Q...	231
<i>G. polyedra</i>	-C-A-GQ...	V-CQ-NE	C-E-VT-K	ACIK-...SE	225
Symbiodinium spp.	-C-S-WQ...	V-CQ-NE	C-E-VK-...	ACVK-...SS	226
<i>S. oleracea</i>	R-V-ECLR...	L-T-D-NV	NS-...MRWRD	RFLFC-E-LY	K-A-...BT	236
* * *						
<i>R. pachyptila</i>	LFSANIADAD	PAEMIARGE	VLETFPFAS	--QVAFIVDG	VYAGPTAVAT	276
<i>R. rubrum</i>F-I-...YEN-...	-H-L-...AA-IP	274
<i>R. sphaeroides</i>	HY-V-...Y	I-...EN-D	-H-...T-A-IP	274
<i>T. denitrificans</i>	M-...ADY	A-V-PD-D	-KL-...G-GM-T	274
<i>H. marinus</i>	V-F-HE	K-Y-...GE	AKYGN	EKH-...F-T-AG-T	286
<i>G. polyedra</i>KY	I-G-Q-PM-E	-NC-.....G-G-TV	273
Symbiodinium spp.	IMSQ-PLSE	-NC-.....G-G-TV	274
<i>S. oleracea</i>	GHLV-A-GT	CED-MK-AVF	AR-LG-...	--VPIVMH-Y	LIG-F-NP-	279
* * *						
<i>R. pachyptila</i>	ARRNFP--NQ	FLHFHRAGH	AVTSPQSKRG	YTAHVHIKMT	RLLAGSMMIV	324
<i>R. rubrum</i>	...R-...DN	...Y-.....C-AQ-...I-T	322
<i>R. sphaeroides</i>	...Q-...R	...Y-.....LS-AQ-...I-T	322
<i>T. denitrificans</i>	...QV-...G	Y-Y-.....I-L-A-SVQ-...D-T	322
<i>H. marinus</i>	S-A-...DT	Y-P-.....YK-PM	MDPLCYM-LAM-...I-T	334
<i>G. polyedra</i>	...C-...K	...P-...Y	...TQ-...	...T-IS	...V-I-...I	321
Symbiodinium spp.	C-...K	...Y-.....	...S-...TQ	...T-IS	...V-I-...I	322
<i>S. oleracea</i>	LSHYCRDGL	L-I-...M-A	VIDR--KNH	MHFR-LA-AL	--S-GHI-S	328
* * *						
<i>R. pachyptila</i>	GTMGYGRMEG	EA-SDKLLAY	MIERDSAD--	--GPPYHQ	WAGMKPTPI	368
<i>R. rubrum</i>	...F-...S	...RA-...	...LTQ-E-Q	...R-S	...G-AC-...	366
<i>R. sphaeroides</i>	...F-...L	...IM-...	...LIDEA-E	...R-T	...GW-S-A-...	366
<i>T. denitrificans</i>	...F-...L	...EPT-...	...LIFE-OBQ	...P-F-S	...RDTACSAM	366
<i>H. marinus</i>	...S-...V	...D-...G	...LQDDA-G	...Y-K	...E-VVQ-...	378
<i>G. polyedra</i>	...SF-...V	...D-...G	...LQDDA-G	...Y-K	...E-VVQ-...	365
Symbiodinium spp.	...SF-...D	...N-...G	...LQDDA-G	...Y-R	...K-KE-...	366
<i>S. oleracea</i>	...VV-...L	...RDITLGFVD	LLRD-YTEKD	RSR-IYFT-S	-VSTPGVL-V	377
* * *						
<i>R. pachyptila</i>	ISGQNALRL	PGFFELHNG	NVINTAGGGT	YGHIDSPAG	AVSLRQAYBC	418
<i>R. rubrum</i>MNRL-...A	F-...G-V-R-...WQA	416
<i>R. sphaeroides</i>MHSQ-S-A	F-L-GCT-K-...SH-A	416
<i>T. denitrificans</i>	A-...HG-	M-S-...EP	D-VL-...	F-...G-VDA	RAN-H-W-A	416
<i>H. marinus</i>DH-.....C-S	F-.....	GI-...G-A	428
<i>G. polyedra</i>AS	Y-L-...	F-K-G-KQ	T-C-DE-A	415
Symbiodinium spp.AS	K-L-...	F-K-G-KI	I-C-GE-A	416
<i>S. oleracea</i>	A-...IHWVM	-ALT-IF--D	DSVLQF-...	L-...PWGN-P	-VAN-V-L-A	426
* * *						
<i>R. pachyptila</i>	MKEGAD----	-----PVE	YAKEH----	EPARAFESFP	HDADAIFPGW	453
<i>R. rubrum</i>	RD-VP----	-----VLD	-R-.....	L-.....	G-Q-Y-...	451
<i>R. sphaeroides</i>	MA-V-....	-----VLD	-R-.....	L-.....	A-KFY-...	451
<i>T. denitrificans</i>	RD-VR----	-----VLD	-R-.....	VL-D-K-WA	G-E-Y-R	451
<i>H. marinus</i>	T-E-.....	-----I	APR-F----	G-K-.....	460
<i>G. polyedra</i>	LWKAGTYG	DVLSLSDG-IT-.....	E-IGK-LTFQ	K-S-Q-Y-...	461
Symbiodinium spp.	QWKAQZG	NLSLSDG-IT-.....	E-IGK-LTFQ	K-A-Q-Y-...	462
<i>S. oleracea</i>	CVQARNEGHD	LARE--GWT	IIR-AYKWSF	EL-A-C-VWK	EIKFEFPAMD	473
* * *						
<i>R. pachyptila</i>	RDKLGVHK	461
<i>R. rubrum</i>	KA-...EDTRS	ALPA	466
<i>R. sphaeroides</i>	DR-HRAA	459
<i>T. denitrificans</i>	KSM-...	457
<i>H. marinus</i>	DE-.....	468
<i>G. polyedra</i>	KE-...YTGESS	QVQASPDWKG	KAA	464
Symbiodinium spp.	KE-...YTGESS	QVQASPDWAK	RA	464
<i>S. oleracea</i>	TV	475

FIG. 2. Alignment of *R. pachyptila* RubisCO with other form II enzymes and a representative form I from *S. oleracea* (shown for comparison). Asterisks refer to residues known to be involved in the active site and activating CO₂ binding site. Dots indicate residues identical to those in the *R. pachyptila* sequence, and dashes are added to preserve the alignment. The alignment of form I and II RubisCOs is based on the three-dimensional structure (26).

not surprising, given that form II enzymes typically have a low affinity for CO₂ and that concentrations of CO₂ are extremely high in the blood of the tubeworm, where concentrations of total dissolved inorganic carbon can be greater than 30 mM (5).

The expression of form I and II RubisCO has recently been suggested to account for the difference observed in stable carbon isotope ratios ($\delta^{13}\text{C}$) of hydrothermal vent invertebrate-chemoautotrophic bacterial symbioses (3, 21). These symbioses fall into two groups based upon their $\delta^{13}\text{C}$ values, with $\delta^{13}\text{C} = -27$ to -35% for mollusc symbioses and $\delta^{13}\text{C} = -9$ to 16% for tubeworm and shrimp symbioses (references 21 and references within). Several hypotheses, such as carbon limitation (9, 19), a C₄-type pathway in the tubeworms (7), or the use of isotopically different source CO₂ utilized by the two groups (4), have been proposed to explain the differences in $\delta^{13}\text{C}$ values but have failed to be corroborated by experimental data. The KIEs of the few form I (for *S. oleracea*, 29%; for *Anacystis nidulans*, 22% [10, 22]) and form II (for *R. rubrum*, 17.8 to 23% [10, 23]) RubisCOs examined by high-precision methods indicate that the two forms fractionate carbon isotopes to differing degrees. Given the high identity between the *R. rubrum* and *R. pachyptila cbbM* sequences, the expression of a form II RubisCO in *R. pachyptila* could account for the heavier isotopic composition if the extreme values for the KIE of the form II RubisCO are considered.

Chemoautotrophic symbioses and free-living chemoautotrophs represent a vast resource for examining different adaptations that have occurred in RubisCO biochemistry and evolution. These organisms promise to yield important new information regarding enzymological adaptation, regulation, and genetic diversity, as they inhabit many niches which are too inhospitable for photoautotrophs. The examination of a greater diversity of species for the form II RubisCO is necessary to determine the distribution of this enzyme among autotrophs.

Nucleotide sequence accession number. The *R. pachyptila* symbiont *cbbM* gene sequence has been deposited in GenBank under accession no. AF047688.

We thank the chief scientists and captains and crew of the RV *Atlantis II* and DSV *Alvin* for their excellent assistance in sample collection, George Lorimer for the generous gift of antiserum, and Marjory Snead for the *R. pachyptila* DNA library construction.

This work was supported in part by NSF grants OCE-9317734 (J.L.S.) and OCE-9504257 (C.M.C.). J.J.R. was also supported by a Graduate Assistance in Areas of National Need fellowship from the Department of Education.

REFERENCES

- Cavanaugh, C. M. 1983. Symbiotic chemoautotrophic bacteria in marine invertebrates from sulfide-rich habitats. *Nature (London)* **302**:58-61.
- Cavanaugh, C. M. 1994. Microbial symbiosis: patterns of diversity in the marine environment. *Am. Zool.* **34**:79-89.
- Cavanaugh, C. M., and J. J. Robinson. 1996. CO₂ fixation in chemoautotrophic-invertebrate symbioses: expression of form I and form II RubisCO. *In* M. E. Lidstrom and F. R. Tabita (ed.), *Microbial growth on C₁ compounds*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Childress, J. J., and C. R. Fisher. 1992. The biology of hydrothermal vent animals: physiology, biochemistry, and autotrophic symbioses. *Oceanogr. Mar. Biol. Annu. Rev.* **30**:337-441.
- Childress, J. J., R. W. Lee, N. K. Sanders, H. Felbeck, D. R. Oros, A. Toulmond, D. Desbruyeres, M. C. Kennicutt, and J. Brooks. 1993. Inorganic carbon uptake in hydrothermal vent tubeworms facilitated by high environmental pCO₂. *Nature* **362**:147-149.
- Distel, D. L., and H. Felbeck. 1988. Pathways of inorganic carbon fixation in the endosymbiont-bearing lucinid clam *Lucinoma aequizonata*. Part 1: purification and characterization of endosymbiotic bacteria. *J. Exp. Zool.* **247**: 1-10.
- Felbeck, H. 1985. CO₂ fixation in the hydrothermal vent tube worm *Riftia pachyptila* (Jones). *Physiol. Zool.* **58**:272-281.

Downloaded from http://jlb.asm.org/ on November 14, 2019 by guest

8. **Felbeck, H., J. J. Childress, and G. N. Somero.** 1981. Calvin-Benson cycle and sulfide oxidation enzymes in animals from sulfide-rich habitats. *Nature (London)* **293**:291–293.
9. **Fisher, C. R., M. C. Kennicutt, and J. M. Brooks.** 1990. Stable carbon isotope evidence for carbon limitation in hydrothermal vent vestimentiferans. *Science* **247**:1094–1096.
10. **Guy, R. D., M. L. Fogel, and J. A. Berry.** 1993. Photosynthetic fractionation of the stable isotopes of oxygen and carbon. *Plant Physiol.* **101**:37–47.
11. **Hartman, F. C., and M. R. Harpel.** 1993. Chemical and genetic probes of the active site of D-ribulose-1,5-bisphosphate carboxylase/oxygenase: a retrospective based on the three-dimensional structure. *Adv. Enzymol.* **63**:1–75.
12. **Houtz, R. L., J. T. Stults, R. M. Mulligan, and N. E. Tolbert.** 1989. Post-translational modifications in the large subunit of ribulose bisphosphate carboxylase/oxygenase. *Proc. Natl. Acad. Sci. USA* **86**:1855–1859.
13. **Jones, M. L.** 1981. *Riftia pachyptila* Jones: observations on the vestimentiferan worm from the Galapagos Rift. *Science* **213**:333–336.
14. **Laue, B. E., and D. C. Nelson.** 1997. Sulfur-oxidizing symbionts have not co-evolved with their hydrothermal vent tubeworm hosts: an RFLP analysis. *Mol. Mar. Biol. Biotechnol.* **6**:180–188.
15. **Lippke, J. A., M. N. Strepko, F. F. Raia, S. L. Simon, and C. K. French.** 1987. Isolation of intact high-molecular-weight DNA by using guanidine isothiocyanate. *Appl. Environ. Microbiol.* **53**:2588–2589.
16. **Lutz, R. A., T. M. Shank, D. J. Fornari, R. M. Haymon, M. D. Lilley, K. L. Von Damm, and D. Desbruyers.** 1994. Rapid growth rates at deep-sea vents. *Nature (London)* **371**:663–664.
17. **Morse, D., P. Salois, P. Markovic, and J. W. Hastings.** 1995. A nuclear encoded form II rubisco in dinoflagellates. *Science* **268**:1662–1674.
18. **Nargang, F., L. McIntosh, and C. Somerville.** 1984. Nucleotide sequence of the ribulosebisphosphate carboxylase gene from *Rhodospirillum rubrum*. *Mol. Gen. Genet.* **193**:220–224.
19. **Rau, G. H.** 1981. Hydrothermal vent clam and tube worm $^{13}\text{C}/^{12}\text{C}$: further evidence of non-photosynthetic food sources. *Science* **213**:338–340.
20. **Robinson, J. J.** 1997. Biochemistry and molecular biology of RubisCO expression and stable carbon isotope fractionation in chemoautotrophic symbioses. Ph.D. thesis. Harvard University, Cambridge, Mass.
21. **Robinson, J. J., and C. M. Cavanaugh.** 1995. Expression of form I and form II ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) in chemoautotrophic symbioses: implications for the interpretation of stable carbon isotope ratios. *Limnol. Oceanogr.* **40**:1496–1502.
22. **Roeske, C. A., and M. H. O'Leary.** 1984. Carbon isotope effects on the enzyme-catalyzed carboxylation of ribulose biphosphate. *Biochemistry* **23**:6275–6284.
23. **Roeske, C. A., and M. H. O'Leary.** 1985. Carbon isotope effect on carboxylation of ribulose bisphosphate catalyzed by ribulosebisphosphate carboxylase from *Rhodospirillum rubrum*. *Biochemistry* **24**:1603–1607.
24. **Rowan, R., S. M. Whitney, A. Fowler, and D. Yellowlees.** 1996. Rubisco in marine symbiotic dinoflagellates: form II enzymes in eukaryotic oxygenic phototrophs encoded by a nuclear multigene family. *Plant Cell* **8**:539–553.
25. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
26. **Schneider, G., Y. Lindqvist, and C. Branden.** 1992. RubisCO: structure and mechanism. *Annu. Rev. Biophys. Biomol. Struct.* **21**:119–143.
27. **Snead, M., P. Kretz, and J. Short.** 1995. Methods for generating plant genomic libraries. In S. B. Gelvin and R. A. Schilperoort (ed.), *Plant molecular biology manual*, 2nd ed. Kluwer Academic Publishers, Dordrecht, The Netherlands.
28. **Somerville, C. R., and S. C. Somerville.** 1984. Cloning and expression of the *Rhodospirillum rubrum* ribulosebisphosphate carboxylase gene in *E. coli*. *Mol. Gen. Genet.* **193**:214–219.
29. **Stein, J., M. Haygood, and H. Felbeck.** 1990. Diversity of ribulose 1,5 bisphosphate carboxylase genes in thiotrophic symbioses. In P. Nardon et al. (ed.), *Endocytobiology IV*. INRA, Paris, France.
30. **Tabita, F. R.** 1988. Molecular and cellular regulation of autotrophic carbon dioxide fixation in microorganisms. *Microbiol. Rev.* **52**:155–189.
31. **Tabita, F. R.** 1995. The biochemistry and metabolic regulation of carbon metabolism and CO₂ fixation in purple bacteria. In R. E. Blankenship, M. T. Madigan, and C. E. Bauer (ed.), *Anoxygenic photosynthetic bacteria*, Kluwer Academic Publishers, Dordrecht, The Netherlands.
32. **Williams, C. A., D. C. Nelson, B. A. Farah, H. W. Jannasch, and J. M. Shively.** 1988. Ribulose bisphosphate carboxylase of the prokaryotic symbiont of a hydrothermal vent tube worm: kinetics, activity and gene hybridization. *FEMS Microbiol. Lett.* **50**:107–112.