

Effect of *rpoS* Mutation on the Stress Response and Expression of Virulence Factors in *Pseudomonas aeruginosa*

SANG-JIN SUH,^{1,2*} LAURA SILO-SUH,² DONALD E. WOODS,³ DANIEL J. HASSETT,⁴
SUSAN E. H. WEST,⁵ AND DENNIS E. OHMAN²

Department of Microbiology and Immunology, University of Tennessee, and Veterans Affairs Medical Center, Memphis, Tennessee 38104¹; Department of Microbiology and Immunology, Medical College of Virginia campus of Virginia Commonwealth University, and McGuire Veterans Affairs Medical Center, Richmond, Virginia 23298-0678²; Department of Microbiology and Infectious Diseases, University of Calgary, Calgary, Alberta, Canada T2N 4N1³; Department of Molecular Genetics, Biochemistry and Microbiology, University of Cincinnati College of Medicine, Cincinnati, Ohio 45257⁴; and Department of Pathobiological Sciences, School of Veterinary Medicine, University of Wisconsin—Madison, Madison, Wisconsin 53706⁵

Received 7 December 1998/Accepted 20 April 1999

The sigma factor RpoS (σ^S) has been described as a general stress response regulator that controls the expression of genes which confer increased resistance to various stresses in some gram-negative bacteria. To elucidate the role of RpoS in *Pseudomonas aeruginosa* physiology and pathogenesis, we constructed *rpoS* mutants in several strains of *P. aeruginosa*, including PAO1. The PAO1 *rpoS* mutant was subjected to various environmental stresses, and we compared the resistance phenotype of the mutant to that of the parent. The PAO1 *rpoS* mutant was slightly more sensitive to carbon starvation than the wild-type strain, but this phenotype was obvious only when the cells were grown in a medium supplemented with glucose as the sole carbon source. In addition, the PAO1 *rpoS* mutant was hypersensitive to heat shock at 50°C, increased osmolarity, and prolonged exposure to high concentrations of H₂O₂. In accordance with the hypersensitivity to H₂O₂, catalase production was 60% lower in the *rpoS* mutant than in the parent strain. We also assessed the role of RpoS in the production of several exoproducts known to be important for virulence of *P. aeruginosa*. The *rpoS* mutant produced 50% less exotoxin A, but it produced only slightly smaller amounts of elastase and LasA protease than the parent strain. The levels of phospholipase C and casein-degrading proteases were unaffected by a mutation in *rpoS* in PAO1. The *rpoS* mutation resulted in the increased production of the phenazine antibiotic pyocyanin and the siderophore pyoverdine. This increased pyocyanin production may be responsible for the enhanced virulence of the PAO1 *rpoS* mutant that was observed in a rat chronic-lung-infection model. In addition, the *rpoS* mutant displayed an altered twitching-motility phenotype, suggesting that the colonization factors, type IV fimbriae, were affected. Finally, in an alginate-overproducing cystic fibrosis (CF) isolate, FRD1, the *rpoS101::aacCI* mutation almost completely abolished the production of alginate when the bacterium was grown in a liquid medium. On a solid medium, the FRD1 *rpoS* mutant produced approximately 70% less alginate than did the wild-type strain. Thus, our data indicate that although some of the functions of RpoS in *P. aeruginosa* physiology are similar to RpoS functions in other gram-negative bacteria, it also has some functions unique to this bacterium.

Microorganisms continuously sense and respond to environmental stimuli, such as starvation, desiccation, osmotic stress, oxidative stress, and changes in temperature. In many instances, an appropriate response to an environmental stimulus requires changes in the levels of specific gene products. One way to accomplish this is by modulating gene expression through the replacement of the main sigma factor of RNA polymerase with an alternative sigma factor that recognizes a specific promoter of the stimulus response gene (19, 39). The sigma factor RpoS (σ^S) was originally identified in *Escherichia coli* and *Salmonella typhimurium* as an alternative sigma factor that activates the expression of numerous genes required to maintain cell viability during the stationary phase of growth when cells are experiencing nutrient starvation (for reviews, see references 30 and 38). Activation of these genes makes the bacterium more resistant to environmental stresses, such as

prolonged starvation, osmotic stress, and oxidative stress. RpoS also plays an important role in exponentially growing cells that are exposed to increased osmolarity (22). In *E. coli* and *S. typhimurium*, RpoS induces the expression of more than 30 genes in response to various stresses (33, 42). RpoS also activates the genes necessary for virulence in several bacteria, including *E. coli*, *S. typhimurium*, and *Shigella flexneri* (3, 31, 48, 55). Among pseudomonads, the role of RpoS as a general stress response regulator has been described for *Pseudomonas putida* (43, 51) and *Pseudomonas fluorescens* (52). In *P. fluorescens*, RpoS also modulates the production of several antibiotics and affects the biological control activity of the bacterium.

Pseudomonas aeruginosa is a gram-negative bacterium that is found in various environmental niches, including soil, water, plants, hospital environments, and human infections. As an opportunistic pathogen, this bacterium primarily infects patients who are immunocompromised (11). During infection, *P. aeruginosa* must survive major environmental changes, including changes in temperature, pH, ionic strength, and the presence of reactive oxygen intermediates and antibodies. With the exception of several genes that have been identified as being involved in the oxidative stress response, such as *katA*, *katB*,

* Corresponding author. Mailing address: Department of Microbiology and Immunology, Medical College of Virginia campus of Virginia Commonwealth University, 1101 East Marshall St., P.O. Box 980678, Richmond, VA 23298-0678. Phone: (804) 628-0247. Fax: (804) 828-9946. E-mail: ssuh@hsc.vcu.edu.

TABLE 1. Bacterial strains and plasmids

Strain or plasmid	Genotype or relevant characteristic ^a	Source or reference
<i>P. aeruginosa</i>		
PAO1	Prototroph	23
PA103	Exotoxin A-overproducing strain	36
FRD1	Alginate-overproducing strain	45
SS6	PA103 <i>rpoS101::aacCI</i>	This study
SS24	PAO1 <i>rpoS101::aacCI</i>	This study
SS138	FRD1 <i>rpoS101::aacCI</i>	This study
<i>E. coli</i>		
DH10B	F ⁻ <i>mcrA</i> Δ(<i>mrr-hsdRMS-mcrBC</i>) φ80 <i>dlacZ</i> Δ <i>M15</i> Δ <i>lacX74</i> <i>deoR</i> <i>recA1</i> <i>endA1</i> <i>araD139</i> Δ(<i>ara</i> , <i>leu</i>)7697 <i>galU</i> <i>galK</i> <i>l-rpsL</i> <i>nupG</i>	Gibco BRL
HB101	F ⁻ Δ(<i>gpt-proA</i>)62 <i>leuB6</i> <i>supE44</i> <i>ara-14</i> <i>galK2</i> <i>lacY1</i> Δ(<i>mcrC-mrr</i>) <i>rpsL20</i> <i>xyl-5</i> <i>mtl-1</i> <i>recA13</i>	Lab collection
Plasmids		
pDB18R	1.8-kb <i>KpnI-HindIII</i> fragment of <i>P. aeruginosa rpoS</i> in pTZ18R	58
pRK2013	Tra1 (RK2), ColE1; Km ^r	15
pSF1	pMB1 replicon, 1.3-kb <i>oriT</i> from (RK2); Ap ^r	54
pSS1	<i>rpoS101::aacCI</i> in pTZ18R; Ap ^r /Cb ^r Gm ^r	This study
pSS6	<i>rpoS101::aacCI</i> in pTZ18R plus 1.2-kb <i>EcoRI</i> fragment of <i>oriT</i> from pSF1	This study
pTZ18R	pUC18-derived general cloning vector; Ap ^r /Cb ^r	Pharmacia
pUCGM1	pUC1918 carrying the <i>aacCI</i> (Gm ^r) cassette; Ap ^r /Cb ^r Gm ^r	53

^a Km^r, kanamycin resistant; Ap^r, ampicillin resistant; Cb^r, carbenicillin resistant; Gm^r, gentamicin resistant.

sodA, *sodB*, and alginate genes (6, 8, 21, 40, 41, 62), little is known about the stress responses of *P. aeruginosa*. However, some information regarding the stress responses of this bacterium can be obtained from studies on the regulation of its various exoproducts. *P. aeruginosa* secretes a battery of extracellular products that are important for virulence: exotoxin A, elastase, LasA protease, phospholipase C, alkaline protease, rhamnolipid, pyocyanin, and alginate. Many of these toxic exoproducts are involved in procuring nutrients and/or protecting the bacterium from the host immune system (14). Furthermore, some of these exoproducts are synthesized in response to the scarcity of specific nutrients (37). For example, exotoxin A and phospholipase C are produced when iron and phosphate are limiting, respectively. Thus, the synthesis of some of these exoproducts may be regulated by various stresses that *P. aeruginosa* encounters in specific niches.

The *P. aeruginosa rpoS* gene, which encodes RpoS, or σ^S, has been cloned and sequenced by Tanaka and Takahashi (58). The *P. aeruginosa rpoS* gene, like the *E. coli rpoS* gene, is growth phase regulated (16). In both *E. coli* and *P. aeruginosa*, the level of *rpoS* mRNA is low during exponential growth but increases severalfold as cells enter the stationary phase. In *P. aeruginosa*, the induction of *rpoS* gene expression has been reported to be regulated by the quorum-sensing cascade involving Vfr, the *las* system, and the *rhl* system (1, 34, 49). The quorum-sensing systems of *P. aeruginosa* regulate the synthesis of many virulence factors, including exotoxin A, elastase, LasA protease, and pyocyanin (5, 35). Thus, the synthesis of these virulence factors may be regulated through RpoS.

Because stress responses may play important roles in pathogenesis, we examined the role of RpoS when *P. aeruginosa* responds to various environmental changes. In this study, we present evidence that RpoS mediates the general stress response and modulates synthesis of several exoproducts, including some secondary metabolites.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. Bacterial strains and plasmids used in this study are listed in Table 1. Bacteria were grown in L broth (LB) or LB supplemented with appropriate antibiotics at 37°C with aeration, unless other-

wise indicated. For minimal salts medium, no-carbon-E minimal medium (NCE) (12) supplemented with either glucose (10 mM) or succinate (20 mM) as a carbon source was used. For exotoxin A assays, iron-depleted Trypticase soy broth dialysate was used as previously described (47). For phospholipase C assays and pyocyanin extractions, 1% peptone broth supplemented with 1% glycerol and 1% NaCl was used as the growth medium. For pyoverdine assays, *P. aeruginosa* was grown in King B medium (28). Following conjugations, pseudomonas isolation agar was used to counterselect against *E. coli*. Media were solidified with 1.5% Bacto Agar (Difco). The following antibiotic concentrations (per milliliter) were used in this study: ampicillin, 100 μg for *E. coli*; carbenicillin, 300 μg for *P. aeruginosa*; gentamicin, 20 μg for *E. coli* and 300 μg for *P. aeruginosa*; kanamycin, 50 μg for *E. coli*. Antibiotics were purchased from Sigma (St. Louis, Mo.).

DNA manipulations, transformations, and conjugations. *E. coli* DH10B or HB101 was routinely used as a host strain for cloning. Plasmids were purified with QIAprep spin miniprep columns (Qiagen, Santa Clarita, Calif.). DNA fragments were excised from agarose gels and purified with the QIAx II DNA gel extraction system (Qiagen), according to the manufacturer's instructions. Restriction enzymes and DNA modification enzymes were purchased from New England Biolabs (Beverly, Mass.), Gibco BRL (Gaithersburg, Md.), or Boehringer Mannheim (Indianapolis, Ind.). For DNA amplifications with a GeneAmp 2400 (Perkin-Elmer, Foster City, Calif.), either *Pfu* (Stratagene, La Jolla, Calif.) or Amplitaq (Perkin-Elmer) was used. Oligonucleotides were purchased from Operon, Inc. (Alameda, Calif.). DNA was introduced into *E. coli* or *P. aeruginosa* either by electroporation or by conjugation. A standard electroporation procedure was used for *E. coli* with the *E. coli* gene pulser (Bio-Rad, Hercules, Calif.). For *P. aeruginosa*, electrocompetent cells were prepared as follows. Cells were grown to mid-log phase (optical density at 600 nm [OD₆₀₀], 0.5 to 0.7) in 100 ml of LB and harvested by centrifugation. Following two washes with 10 ml of 0.3 M sucrose, the cells were resuspended in 1 ml of 0.3 M sucrose, and 40 μl of this cell suspension was used for electroporation. The conditions used to electroporate *P. aeruginosa* were 200 Ω, 200 μF, and 1.6 kV. For conjugations, triparental matings were performed with the helper plasmid pRK2013 as previously described (17), with the following modifications. Briefly, 50 to 100 μl of overnight cultures of donor, helper, and recipient strains were mixed in 1 ml of saline, centrifuged, washed with 1 ml of saline, and resuspended in 50 μl of saline. The cell suspension was spotted on an LB plate and incubated overnight at 30°C. Following incubation, the cells were scraped with a sterile cotton swab, resuspended in 2 ml of saline, and plated on selective plates. Introduction of plasmid DNA into *P. aeruginosa* FRD1 was done exclusively by conjugation.

Construction of *P. aeruginosa rpoS* mutants. To generate *P. aeruginosa rpoS* mutants, the suicide plasmid pSS1, which carries the *rpoS101::aacCI* null allele, was constructed. Briefly, pDB18R, which carries *P. aeruginosa rpoS* on a 1.8-kb *KpnI-HindIII* fragment, was digested with *SphI* to cut once within *rpoS*. Following T4 DNA polymerase digestion to blunt the protruding ends, the linearized plasmid was digested with *HincII* to delete approximately 420 nucleotides within the *rpoS* gene. The *aacCI* gene, which encodes gentamicin resistance (Gm^r) and which was obtained from pUCGM1 as a *SmaI* fragment, was inserted to generate the *rpoS101* allele. pSS1 was introduced into *P. aeruginosa* PAO1 and PA103 by electroporation, and the potential *rpoS* mutants were isolated as having Gm^r

(encoded by the *rpoS101::aacCI* allele) but lacking carbenicillin resistance (Cb^r) (encoded by the plasmid vector), indicating a double crossover event. Introducing the suicide plasmid pSS6 by conjugation and isolating mutants that had undergone allelic exchange generated an *rpoS* mutation in FRD1. pSS6 carries a 1.2-kb *oriT* fragment isolated from pSF1 as an *EcoRI* fragment in the unique *EcoRI* site of pSS1. The presence of the *rpoS101::aacCI* allele in these mutants was verified by two methods, Southern blot hybridization (57) and PCR analysis, to demonstrate the replacement of the wild-type *rpoS* gene with the *rpoS101* allele (data not shown).

For Southern blot analysis, chromosomal DNAs from the putative *rpoS* mutants and their respective wild-type strains were digested independently with either *EcoRI* or *EcoRV*, electrophoresed on agarose gels, and blotted to nitrocellulose. The blots were probed with *rpoS*, a 420-bp *HincII-SphI* internal fragment of *rpoS*, the 820-bp *aacCI* gene, or the plasmid vector. The *rpoS* gene and the 820-bp *aacCI* gene both hybridized to the same-sized *EcoRI* or *EcoRV* fragments of chromosomal DNA isolated from putative mutants. Neither the internal *rpoS* fragment nor the plasmid vector hybridized to chromosomal DNA from the putative mutants. For PCR analysis, oligonucleotides that hybridized to the *rpoS* flanking sequences were used. In *rpoS* mutants, we observed the disappearance of the DNA fragment corresponding to the wild-type *rpoS* gene and the appearance of the larger fragment corresponding to the *rpoS101::aacCI* allele.

Stress response assays. The ability of the wild-type cells (PAO1) and the *rpoS* mutant cells (SS24) to survive during prolonged starvation was determined with cells grown at 37°C with aeration in LB, NCE with 20 mM succinate, or NCE with 10 mM glucose. After cells entered the stationary phase of growth, incubation was continued for several days; cell viability was determined by taking periodic aliquots and measuring the CFU on LB plates. The time at which the cells entered the stationary phase of growth was taken as day 0, and aliquots were taken in periodic increments of 24 h from that initial point. The viability was measured as a percentage of the highest number of CFU for PAO1 and SS24 in each experiment.

To measure cells' ability to survive heat shock, cells grown overnight in LB at 37°C with aeration were washed twice and diluted in NCE to a density of approximately 7,000 CFU/ml. One milliliter of the diluted culture was placed in a prewarmed tube at 50°C, and the viability was determined by taking periodic aliquots and plating them directly on LB plates to measure the CFU. To measure sensitivity to osmotic stress, cells were grown overnight in LB at 37°C with aeration, washed twice in NCE, and resuspended in NCE, NCE with NaCl, or NCE with sucrose. Resuspended cells were incubated at 37°C with aeration, periodic aliquots were taken, and serial dilutions of the samples were plated on LB plates to determine the CFU. NaCl concentrations of 2, 2.5, and 3 M and sucrose concentrations of 1, 1.5, and 2 M were tested. Sensitivity to hydrogen peroxide (H_2O_2) was measured on cells grown for 17 h in LB at 37°C as previously described (21). Cells from cultures in mid-log (OD_{600} , 0.3) and stationary phases of growth were plated on LB plates. Sterile Whatman no. 1 filter disks impregnated with 10 μ l of 30% H_2O_2 were placed on top of bacterium-seeded plates and incubated at 37°C for 24 h, and the zones of inhibition were measured.

Enzyme assays. The catalase assays were performed as described by Beers and Sizer (4) with 19.5 mM H_2O_2 with cells grown in LB. Cell extracts for the catalase assays were prepared as previously described (6). For the exotoxin A assays, cells were grown in Trypticase soy broth dialysate at 32°C with aeration for 18 h. The cells were harvested by centrifugation, and 10 μ l of the supernatant was used to measure the exotoxin A activity, as previously described (47). The values for the ADP-ribosyltransferase activity were measured as counts per minute per 10 μ l of supernatant. To measure elastase, LasA protease, and casein-degrading protease activities, cells were grown in LB at 37°C for 14 to 17 h with aeration. Following centrifugation to harvest the cells, the quantity of total proteins in the culture supernatants was determined by the Bradford method (with a kit from Bio-Rad). Elastase activity was determined by the elastin Congo red hydrolysis assay with 20 to 30 μ g of supernatant protein, as previously described (46). The elastase activity was calculated as the increase in A_{495} $minute^{-1}$ gram of protein $^{-1}$. The staphylococcal activity of LasA protease was determined with approximately 25 μ g of total supernatant protein, as previously described (26). The LasA protease activity was calculated as the rate of decrease in A_{595} $minute^{-1}$ milligram of protein $^{-1}$. To determine the phospholipase C activity, the cells were grown at 30°C for 20 to 24 h in 1% peptone broth supplemented with 1% glycerol and 1% NaCl. The ability of phospholipase C to degrade *p*-nitrophenylphosphorylcholine was determined as previously described (32) with approximately 10 μ g of supernatant protein.

Secondary metabolite assays. To isolate pyocyanin, cells were grown in peptone broth (1% peptone, 1% NaCl, 1% glycerol) for 17 h at 37°C with aeration. Pyocyanin from 4 ml of supernatant was extracted by repeated extractions (a total of four times) with 1 ml of chloroform (9). The organic layer containing pyocyanin was collected, and the relative amount of pyocyanin isolated was determined by measuring the OD_{695} . The amount of pyoverdine produced and secreted by *P. aeruginosa* was determined by measuring the fluorescence of pyoverdine when excited at 400 nm, as previously described (50). Briefly, the cells were grown in King B medium for 16 to 17 h at 37°C with aeration. The culture supernatants were serially diluted in 10 mM Tris-HCl (pH 7.5) and excited at 400 nm, and the emission at 460 nm was recorded. The optical density measurements were normalized to the OD_{600} of the cultures used for the assay.




FIG. 1. Construction of *P. aeruginosa rpoS* mutants. To generate a null allele of the *P. aeruginosa rpoS* gene, approximately 420 bp of the *rpoS* coding sequence on a 1.8-kb fragment was replaced with the *aacCI* gene, which encodes Gm^r , to form pSS1. The wild-type *rpoS* was replaced with the *rpoS101::aacCI* allele carried on the suicide plasmid pSS1, as described in Materials and Methods, to generate *P. aeruginosa rpoS* mutants.

Alginate assays. To quantitate the amount of alginate produced when it was grown in liquid medium, cells were grown in LB for 20 to 24 h at 37°C with aeration. Alginate in the culture supernatant was separated from the cells by centrifugation (17,000 $\times g$ for 10 min) and dialyzed at 4°C against deionized water for 24 h with three changes of water. Dialysis tubing with a molecular weight cutoff of 10,000 (Spectrum, Houston, Tex.) was used. To quantitate the alginate produced on solid medium, overnight cultures of FRD1 or SS138 were serially diluted in saline (0.85% NaCl), and 0.1 ml of diluted cultures was plated on LB plates. The plates were incubated at 37°C for 25 to 30 h; alginate was collected by scraping the cells and alginate with saline, followed by centrifugation to separate the alginate from the cells. The alginate thus collected was subjected to 24 h of dialysis at 4°C against three changes of deionized water. The amount of alginate present in the samples was quantitated by the carbazole method of Knutson and Jeanes (29), with *Macrocystis pyrifera* alginate (Sigma) as a standard. The values for alginate production were obtained by normalizing the total amount of alginate produced to the CFU of the culture or the number of colonies on plates.

Twitching-motility assay. A twitching-motility assay was performed by stabbing a colony of bacteria into the bottom of a petri plate containing 10 ml of L agar (1% agar content). Following incubation at 37°C for 17 h, the volume of agar was reduced with a thick stack of circular paper towels to absorb the moisture. The zone of motility was visualized by staining the agar with Coomassie blue, as described by Alm and Mattick (2).

Chronic-lung-infection assay. The rat chronic-lung-infection assay was performed as previously described (60). Briefly, agar beads embedded with approximately 3×10^4 cells of either the wild-type bacteria or the *rpoS* mutant were placed in the left lungs of rats. The lungs of three animals from each group were isolated at days 7 and 14 for bacterial quantification. The lungs of two additional animals from each group were isolated at the same times for histopathological examination as previously described (56). Briefly, sagittal slices of the entire left lobes of the fixed lungs were dehydrated in graded alcohols, embedded in paraffin, and cut into 6- μ m-thick sections. Mounted sections were stained with hematoxylin and eosin. Infiltration of the lung by inflammatory cells and exudate was measured microscopically with a Zeiss integrating eyepiece. The number of points overlying the surface area of the infiltrate was divided by the total number of points counted over the entire surface area of the left lobe to measure the percent infiltration. This procedure was repeated with left lobe slices from each animal's lung.

RESULTS

Construction of *P. aeruginosa rpoS* mutants. *P. aeruginosa rpoS* mutants were isolated in order to better understand the potential role of RpoS (σ^S) in *P. aeruginosa* physiology and pathogenesis. Figure 1 shows the schematic diagram for constructing the suicide plasmid, pSS1, which carries the null *P. aeruginosa rpoS101::aacCI* allele, and the subsequent generation of *P. aeruginosa rpoS* mutants with pSS1. We generated a




FIG. 2. Effect of *rpoS* mutation on stress responses of *P. aeruginosa*. To assay cell survival after prolonged starvation (A to C), PAO1 (□) and SS24 (◆) cells were grown in the indicated growth media at 37°C with aeration. Day 0 is the time point at which cells entered stationary phase of growth, and then samples were taken at 24-h intervals. Viability was measured as the number of CFU and was expressed as a percentage of the number of CFU at day 0. Minimal salt medium was supplemented with either 10 mM glucose (A) or 20 mM succinate (B). LB (C) was used as a rich medium. To assay cell survival of heat shock at 50°C (D), stationary-phase cultures of PAO1 (□) and SS24 (◆) were grown in LB, washed, and transferred to prewarmed tubes. The number of viable cells was measured by plating aliquots on LB plates at each time point and determining the number of CFU. Viability is expressed as a percentage of the number of CFU at time 0. To assay survival of exposure to 2 M NaCl (E), stationary-phase cultures of PAO1 (□) and SS24 (◆) were grown in LB, washed, resuspended either in minimal salts or in minimal salts supplemented with 2 M NaCl, and incubated at 37°C with aeration. Cell viability was determined at 2 and 6 h. One hundred percent viability corresponds to the CFU present immediately following resuspension in 2 M NaCl. The carbon starvation experiments were performed at least four times, and the other stress response assays were performed at least twice. The data shown are representative of those experiments.

null allele of *rpoS* by deleting the 420-bp *HincII-SphI* fragment of the *rpoS* coding sequence and replacing it with an 820-bp *SmaI* fragment that confers the Gm^r encoded by *aacCI*. We introduced pSS1 into *P. aeruginosa* and isolated colonies that had undergone the allelic exchange reaction replacing the wild-type *rpoS* with the *rpoS101::aacCI* allele. The inheritance of the *rpoS101::aacCI* allele in *P. aeruginosa* was confirmed by Southern blot hybridization analysis and/or PCR analysis (data not shown). The *rpoS* mutants of PAO1, PA103, and FRD1 were designated SS24, SS6, and SS138, respectively (Table 1).

Response of *rpoS* mutants to environmental stresses. To determine whether RpoS confers cross-protection for *P. aeruginosa* against various stresses, the resistance of the parent (PAO1) and *rpoS* mutant (SS24) against prolonged starvation, heat shock, increased osmolarity, and H₂O₂ was determined. To study its response to prolonged carbon starvation, we grew *P. aeruginosa* on three different growth media to determine whether growth on a specific carbon source affected the survival of *P. aeruginosa*. The media used were minimal medium with glucose, minimal medium with succinate, and LB. For *P. aeruginosa*, glucose is a less preferred carbon source than succinate. There was no difference in growth rate between the parent and *rpoS* mutant in each of three growth media (data not shown). When bacteria were grown with glucose as the sole carbon source, we observed increased sensitivity by the *rpoS* mutant to prolonged starvation (Fig. 2A). However, the difference in survival between the parent and the *rpoS* mutant was not as pronounced as that observed in other bacteria. For example, an *E. coli rpoS* mutant is almost 50-fold more sensi-

tive to starvation than the parent strain after 9 days of incubation (33). In *P. aeruginosa*, we consistently observed only a four- to eightfold difference between the parent and *rpoS* mutant after 12 to 14 days of incubation. In contrast, when bacteria were grown with succinate as a sole carbon source (Fig. 2B) or in LB (Fig. 2C), we observed little difference in survival between the *rpoS* mutant and the parent. The most striking difference observed among the three growth conditions was the relative resistance of *P. aeruginosa* to prolonged starvation when the cells were grown with glucose as the carbon source. When cells were grown on glucose as the sole carbon source, approximately 12% of the parent cells and 4% of the *rpoS* mutant cells were viable after 15 days of starvation. In contrast, when cells were grown with succinate, approximately 1% of the parent cells and 0.8% of the *rpoS* mutant cells were viable after 13 days of incubation. When cells were grown in LB, no viable cells were recovered after 10 to 16 days of continued incubation (data not shown). Thus, *P. aeruginosa* is sensitive to prolonged starvation, and RpoS had little effect in protecting the bacterium from prolonged starvation. In addition, the growth of *P. aeruginosa* on glucose appeared to decrease the sensitivity of the bacterium to prolonged starvation. Although only three substrates were tested, it appeared that survival of *P. aeruginosa* was enhanced when a poorer substrate was used.

When stationary-phase cultures of the parent (PAO1) and the *rpoS* mutant (SS24) were exposed to a sudden shift in temperature from 24° to 50°C, the *rpoS* mutant was more sensitive to heat shock at 50°C. After 8 min of incubation at 50°C, there was a 50-fold difference in survival between the

TABLE 2. Effect of *rpoS* mutation on oxidative stress in *P. aeruginosa*^a

Growth phase and phenotype	H ₂ O ₂ sensitivity ^b	Catalase activity (U/mg of protein)
Mid-log		
RpoS ⁺	25.8 ± 0.7	116 ± 8
RpoS ⁻	25.5 ± 2.8	71 ± 7
Stationary		
RpoS ⁺	29.8 ± 0.2	1,206 ± 41
RpoS ⁻	36.3 ± 0.9	476 ± 21

^a Values are means ± standard deviations.^b Zone of inhibition was measured in millimeters.

parent and the *rpoS* mutant (Fig. 2D). When the cells were subjected to an increase in osmotic pressure caused by the addition of a high concentration of salt, the *rpoS* mutant was more sensitive to a 2 M concentration of sodium chloride (Fig. 2E). After 6 h of exposure, there was approximately a 37-fold difference in relative viability between the parent and the *rpoS* mutant. Thus, RpoS was essential for resistance against osmotic stress caused by exposure to salt. Against nonionic osmotic stress, the wild type was only slightly more resistant (an approximately threefold difference after 6 h of exposure) to 2 M sucrose than the *rpoS* mutant (data not shown). This indicated that the role of RpoS in protecting *P. aeruginosa* from nonionic osmotic stress may be small.

We also assayed whether RpoS mediated protection against H₂O₂ in *P. aeruginosa* (Table 2). In exponentially growing cells, there was no difference in sensitivity to H₂O₂ between the parent and the *rpoS* mutant. However, when stationary-phase cells were tested, the zone of inhibition caused by H₂O₂ was approximately 20% larger for the *rpoS* mutant than for the parent strain. We tested whether this hypersensitivity of the *rpoS* mutant to H₂O₂ was due to a decreased production of catalases in the mutant. In stationary-phase cultures, the *rpoS* mutant produced 60% less catalase activity than the parent strain. Thus, RpoS was required for maximum protection against H₂O₂ by mediating the production of catalase in *P. aeruginosa*.

Effect of *rpoS* mutation on production of exoproducts. *P. aeruginosa* produces and secretes numerous toxins, proteases, and secondary metabolites that contribute to the pathogenesis of the bacterium. We investigated whether RpoS controls the production and secretion of exotoxin A, elastase, LasA protease, casein-degrading proteases, phospholipase C, pyocyanin, and pyoverdine. The data are shown in Table 3.

Exotoxin A is an ADP-ribosyltransferase that inhibits eukaryotic protein synthesis (36). Exotoxin A activity in the culture supernatants was decreased by 50% in the PAO1 *rpoS* mutant (SS24). In the exotoxin A-overproducing PA103 and its respective *rpoS* mutant (SS6), the RpoS⁻ defect resulted in a 70% reduction in exotoxin A activity (18,502 ± 1,276 for the parent versus 5,809 ± 407 for the *rpoS* mutant). We also determined the effect of the *rpoS* mutation on the extracellular accumulation of proteases and phospholipase C in the PAO1 background. Culture supernatants of the PAO1 *rpoS* mutant consistently had 20% less elastase activity and 10% less LasA protease activity. The decreased accumulation of these extracellular proteins in the culture supernatant of *rpoS* mutant was fully restored to the level of the parental strain when RpoS was provided in *trans* from a plasmid (data not shown). A measure of casein degradation detects many proteases produced by the

TABLE 3. Effect of *rpoS* mutation on *P. aeruginosa* PAO1 exoproducts^a

Exoproduct	Activity in strain:		% Relative activity
	PAO1 (RpoS ⁺)	SS24 (RpoS ⁻)	
Exotoxin A	1,302 ± 46	660 ± 20	50.7
Elastase	161 ± 6	126 ± 2	78.6
LasA protease	0.587 ± 0.002	0.518 ± 0.014	88.3
Phospholipase C	33.45 ± 0.65	32.97 ± 0.82	98.6
Pyocyanin	0.347	0.716	206.3
Pyoverdine	0.511	0.755	147.7

^a Exotoxin A activity is measured in counts per minute per 10 μl of supernatant. Elastase activity is measured as the increase in A₄₉₅ minute⁻¹ gram of protein⁻¹. LasA protease activity is measured as the rate of decrease in A₅₉₅ minute⁻¹ milligram of protein⁻¹. Phospholipase C activity is measured as the increase in OD₄₀₅ minute⁻¹ milligram of protein⁻¹. The pyocyanin level was measured as OD₆₉₅ and the pyoverdine level was measured as OD₄₆₀ following excitation at OD₄₀₀ (see Materials and Methods for details). The relative activity was calculated as follows: (activity of the *rpoS* mutant/activity of the parent) × 100. Values are means ± standard deviations.

bacterium (i.e., elastase and alkaline protease). The difference in the production of casein-degrading protease between the parent strain and the *rpoS* mutant was negligible (data not shown). Similarly, little difference in the production and secretion of total phospholipase C (Table 3) or hemolytic phospholipase C (data not shown) between the parent and the *rpoS* mutant was observed. Based on these data, we concluded that RpoS was required for the maximum production of exotoxin A but that it was not required for the maximum production of casein-degrading proteases or phospholipase C.

Pyocyanin and pyoverdine are secondary metabolites produced by *P. aeruginosa*. Pyocyanin is a blue compound that plays an important role in *P. aeruginosa* pathogenesis (59). Pyocyanin kills bacteria and inhibits lymphocyte proliferation and ciliary function. The color of *P. aeruginosa rpoS* mutant culture supernatants was an intense blue relative to parent strains, suggesting that *rpoS* mutants were producing and secreting more pyocyanin. Measurements showed that culture supernatants from the *rpoS* mutant contained twice as much pyocyanin as the parent strain PAO1. Conversely, the overproduction of RpoS from a high-copy-number plasmid in PAO1 resulted in a decrease in pyocyanin levels in culture supernatants (data not shown). We also determined the effect of *rpoS* mutation on the accumulation of pyoverdine, the primary siderophore produced by *P. aeruginosa*, in culture supernatant (10). Culture supernatant from the *rpoS* mutant contained 1.5-fold more pyoverdine than the parent strain. These data indicated that RpoS modulates the level of pyocyanin and pyoverdine production in *P. aeruginosa*.

Effect of *rpoS* mutation on alginate production. Alginate is an exopolysaccharide that is overproduced by most of the *P. aeruginosa* strains isolated from CF patients suffering from chronic lung infections (18). The overproduction of alginate contributes to the persistence of the bacterium in the CF lung. To determine whether RpoS is involved in the production of alginate, we examined SS138, the *rpoS101::aacCI* mutant of FRD1 (an alginate-overproducing CF isolate). As shown in Fig. 3, when the cells were grown in a liquid medium, alginate production was almost completely abolished in the FRD1 *rpoS* mutant. In contrast, when grown on plates (seeded with 10² CFU/plate), the *rpoS* mutant produced approximately 35% of the alginate produced by the wild-type strain. Interestingly, when agar plates were seeded with a high cell density (i.e., ≥10⁴ CFU), the *rpoS* mutant accumulated as much alginate as




FIG. 3. The *rpoS* mutation affects alginate accumulation in mucoid strain FRD1. To evaluate growth in liquid, FRD1 and its *rpoS* mutant derivative (SS138) were grown in LB at 37°C with aeration for 20 to 24 h. For growth on plates, approximately 1,000 cells of the parent or the *rpoS* mutant were spread on L agar plates and incubated at 37°C for 25 to 30 h. Alginate was separated from the cells and quantitated as described in Materials and Methods. The amount of alginate produced was normalized to the number of cells in the culture or to the number of colonies on plates to obtain the amount of alginate produced per CFU. One hundred percent of alginate production corresponds to the level produced by the wild-type FRD1 strain. The values obtained for alginate were as follows: FRD1 (liquid), 1.23×10^{-6} $\mu\text{g CFU}^{-1}$; SS138 (liquid), 1.39×10^{-8} $\mu\text{g CFU}^{-1}$; FRD1 (plate), $0.77 \mu\text{g colony}^{-1}$; SS138 (plate), $0.27 \mu\text{g colony}^{-1}$.

the wild-type FRD1 (data not shown). Because FRD1 and the *rpoS* mutant grew at similar rates (data not shown), the difference in alginate production could not be attributed to a growth defect. Thus, RpoS, in combination with cell density, was required for maximum alginate synthesis in *P. aeruginosa*. Providing RpoS in *trans* on a multicopy plasmid restored the alginate production of the *rpoS* mutant to that of the parental strain FRD1 (data not shown).

Effect of *rpoS* mutation on the rat chronic-lung-infection model. We assessed the role of RpoS in the virulence of *P. aeruginosa* using the rat chronic-lung-infection model. In this model, the bacteria are embedded in agar beads which protect them from the host immune response (60). The results showed that the *rpoS* mutant survived as well as the wild-type bacteria when the mutant was embedded in agar beads and placed in rat lungs. Interestingly, the *rpoS* mutant appeared to cause greater damage to lung tissues than the wild-type strain (Table 4). The increased damage inflicted on the lungs of rats by the *rpoS* mutant may be due to the overproduction of pyocyanin.

Although the rat chronic-lung-infection model is a good model for assessing the roles of extracellular products in infection, it does not provide information about colonization functions and other initial stages of infection. The colonization of *P. aeruginosa* is mediated by type IV fimbriae (13), which also mediate the twitching motility. We examined the twitching




FIG. 4. Effect of *rpoS* mutation on twitching motility. A colony of wild-type (PAO1) or *rpoS* mutant (SS24) cells was stabbed into the bottom of a petri plate containing 10 ml of L agar (1% agar content). Following incubation at 37°C for 17 h, the volume of agar was reduced by absorbing the moisture with a stack of paper towels, and the zone of motility was visualized by staining the agar with Coomassie brilliant blue.

motility of the *rpoS* mutant to determine whether RpoS may be involved in *P. aeruginosa* colonization. As shown in Fig. 4, the zone of motility of the *rpoS* mutant (SS24) was approximately 60 to 70% of that of the parent (PAO1). In addition, the shape of the zone observed for the *rpoS* mutant was typically elliptical, rather than the circular shape that was observed for the parent strain. Finally, the *rpoS* mutant demonstrated altered swarming motility on the top of the agar. These data suggest that RpoS may be involved in the colonization of *P. aeruginosa* as mediated by the type IV fimbriae.

DISCUSSION

The ability of *P. aeruginosa* to resist, adapt, and survive in a wide variety of environments makes it a ubiquitous bacterium in nature and likely contributes to its opportunistic pathogenic behavior. As an initial effort to elucidate and understand the stress responses of this bacterium, we isolated and characterized *P. aeruginosa rpoS* mutants to determine the effect of an RpoS⁻ defect on the stress responses and the synthesis of virulence factors in this bacterium.

Our data demonstrate both similarities and differences between the role of RpoS in *P. aeruginosa* and its role in other bacteria. Some of the similarities include the RpoS-mediated resistance of *P. aeruginosa* to heat-shock, H₂O₂, and osmotic stress. These data indicate that RpoS is a general stress response regulator in *P. aeruginosa*. However, because the physiology and metabolism of *P. aeruginosa* are different from those of *E. coli*, we do not know whether RpoS-mediated resistance to various stresses in *P. aeruginosa* is similar or even proceeds via analogous genes. For example, the *E. coli rpoS* mutant is hypersensitive to 15 mM H₂O₂ (33), while the *P. aeruginosa rpoS* mutant is resistant to 100 mM H₂O₂ (data not shown), despite reduced catalase production in the *P. aeruginosa rpoS* mutant. In *E. coli*, RpoS regulates the expression of two catalase biosynthetic genes, *katE* and *katG*, in order to detoxify

TABLE 4. Effect of *rpoS* mutation on chronic lung infection caused by *P. aeruginosa*

Parent or mutant	Initial inoculum (CFU/ml)	Quantitative bacteriology (CFU/ml) on day ^a :		Quantitative pathology (% damage) on day:	
		7	14	7 ^b	14
RpoS ⁺ (PAO1)	3.5×10^4	$(4.4 \pm 4.3) \times 10^5$	$(1.3 \pm 1.2) \times 10^5$	18.5 ± 4.9	20.9 ± 8.8
RpoS ⁻ (SS24)	2.8×10^4	$(2.3 \pm 2.2) \times 10^6$	$1.2 \times 10^6 \pm 9.2 \times 10^5$	32.7 ± 3.8^c	56.2 ± 19.3^c

^a Values are means \pm standard deviations.

^b Infiltration index.

^c Significantly different from RpoS⁺ value ($P < 0.05$) by Student's *t* test.

H₂O₂ (25, 44). *P. aeruginosa* also possesses two catalases, KatA and KatB (6, 40). It was shown that KatA activity is under quorum-sensing control and thus is maximal in stationary phase (20). Because the *rpoS* gene is also under quorum-sensing control (34), it may be that RpoS directly controls *kata* expression in stationary phase.

We also found several differences in the role of RpoS between *E. coli* and *P. aeruginosa*. One of the differences was the minor role of RpoS in protecting *P. aeruginosa* from prolonged carbon starvation. *P. aeruginosa* was quite sensitive to prolonged starvation (Figure 2A to C), despite the fact that *P. aeruginosa* is almost ubiquitous in nature, including in low-nutrient environments. One possible explanation is that because *P. aeruginosa*, like other members of the pseudomonads, can metabolize so many compounds as nutrients, it has not developed resistance against prolonged carbon starvation. Alternatively, carbon starvation resistance in *P. aeruginosa* may be regulated via a mechanism not addressed by our methods and/or by a different regulator than RpoS. In *P. putida*, protection against carbon starvation has been shown to be regulated by both RpoS (43, 51) and RpoN (27), the sigma factor that regulates nitrogen metabolism in several bacteria.

Another interesting result was the differential effects that various carbon sources had on the subsequent resistance of *P. aeruginosa* to prolonged carbon starvation. *P. aeruginosa* grown on glucose was approximately 10- to 15-fold more resistant to prolonged starvation than when grown on succinate and at least 500-fold more resistant than when grown in LB. There are several possibilities for these differences. *P. aeruginosa* growing in LB or on succinate may accumulate compounds that are toxic to the cells, or growth on glucose may induce the expression of genes under RpoS control that confer enhanced resistance to prolonged starvation. It is intriguing to consider the possibility that growth on a less preferred carbon source (such as glucose) may better prepare the bacterium for the onset of starvation.

RpoS has been demonstrated to be required for production of extracellular toxins or proteases in three other organisms: *Yersinia enterocolitica*, *Vibrio cholerae*, and *P. fluorescens* (24, 52, 61). Of a subset of extracellular products secreted by *P. aeruginosa*, the accumulation of exotoxin A, pyocyanin, and pyoverdine was most dramatically affected by the RpoS⁻ defect. However, it is unclear whether regulation of these exoproducts by RpoS occurs at the level of synthesis or secretion. The expression of *P. aeruginosa* *xcp* operons, which encode the type II secretion apparatus for exotoxin A, elastase, LasA protease, and phospholipase C, among other proteins, is regulated by the quorum-sensing systems (7), which in turn regulate the expression of *rpoS* (34). This led Chapon-Hervé et al. (7) to propose the possibility that regulation of *xcp* operons may proceed via RpoS in *P. aeruginosa*. This proposal was also based on the presence of a potential RpoS consensus promoter upstream of *xcp*. If the *xcp* operons are regulated by RpoS, then the *rpoS* mutant should be deficient for the accumulation of many exoproteins. However, only the accumulation of exotoxin A was significantly affected by the *rpoS* mutation in our study. Further data are needed to clarify whether RpoS regulates the expression of genes encoding exotoxin A, elastase, and LasA protease directly or regulates the secretion of these proteins via Xcp.

RpoS plays a major role in the production of alginate, and this effect is dependent on whether cells are grown in liquid or on a solid surface. When bacteria are grown on a solid surface, a higher cell density can be reached. Thus, RpoS-mediated alginate production may be cell density dependent. Our data suggest that a high cell density can minimize or even negate the

RpoS requirement for the maximum accumulation of alginate and that RpoS is essential for alginate accumulation only when cell density is relatively low. These results pose an interesting question. How can RpoS induce the synthesis or secretion of alginate under low-cell-density conditions when the expression of *rpoS* itself is induced by high cell density? Perhaps the basal level of *rpoS* expressed in FRD1 at a low cell density may be sufficient for the induction of alginate production and/or secretion. Alternatively, due to the close proximity of the cells growing on a solid surface, a few cells may be sufficient to induce the quorum-sensing system.

In conclusion, the mutation in *rpoS* affects a variety of functions in *P. aeruginosa*. These include the general stress response, the accumulation of virulence factors, and twitching motility. Our data suggest that one of the functions of RpoS in *P. aeruginosa* is to fine-tune the cell metabolism, which includes modulating the production of several secondary metabolites, in order to maintain the optimal conditions for cells in stationary phase. The two-dimensional gel analysis on PAO1 and SS24 indicated that the *rpoS* mutation results in an alteration in the production and accumulation of at least 37 proteins (data not shown). Studies to identify the genes regulated by RpoS in *P. aeruginosa* are in progress.

ACKNOWLEDGMENTS

We thank Kan Tanaka for providing us with pDB18R and pDB19R. We also thank Tricia Maleniak, Robert Kazmierczak, Laura Regassa, Laura Runyen-Janecky, and Jorge Escalante-Semerena for help and encouragement.

This research was supported by Public Health Service grant AI-19146 from the National Institute of Allergy and Infectious Diseases and Veterans Administration Medical Research Funds, awarded to D.E.O. Studies at the University of Calgary were supported by a grant from the Canadian Cystic Fibrosis Foundation, awarded to D.E.W. Studies at the University of Cincinnati were supported by Public Health Service grant AI-40541 from the National Institute of Allergy and Infectious Diseases and Cystic Fibrosis Foundation grant Has-setPO97, awarded to D.J.H. Studies at the University of Wisconsin were supported by Public Health Service grant AI-31477 from the National Institute of Allergy and Infectious Diseases, awarded to S.E.H.W. S.-J.S. was supported by Cystic Fibrosis Foundation post-doctoral fellowship SUH96F0.

REFERENCES

1. Albus, A. M., E. C. Pesci, L. J. Runyen-Janecky, S. E. H. West, and B. H. Iglewski. 1997. Vfr controls quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* **179**:3928–3935.
2. Alm, R. A., and J. S. Mattick. 1995. Identification of a gene, *pilV*, required for type 4 fimbrial biogenesis in *Pseudomonas aeruginosa*, whose product possesses a pre-pilin-like leader sequence. *Mol. Microbiol.* **16**:485–496.
3. Arnold, K. W., and C. W. Kaspar. 1995. Starvation- and stationary-phase-induced acid tolerance in *Escherichia coli* O157:H7. *Appl. Environ. Microbiol.* **61**:2037–2039.
4. Beers, R. F., Jr., and I. W. Sizer. 1952. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.* **195**:133–140.
5. Brint, J. M., and D. E. Ohman. 1995. Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhII, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *J. Bacteriol.* **177**:7155–7163.
6. Brown, S. M., M. L. Howell, M. L. Vasil, A. J. Anderson, and D. J. Hassett. 1995. Cloning and characterization of the *katB* gene of *Pseudomonas aeruginosa* encoding a hydrogen peroxide-inducible catalase: purification of KatB, cellular localization, and demonstration that it is essential for optimal resistance to hydrogen peroxide. *J. Bacteriol.* **177**:6536–6544.
7. Chapon-Hervé, V., M. Akrim, A. Latifi, P. Williams, A. Lazdunski, and M. Bally. 1997. Regulation of the *xcp* secretion pathway by multiple quorum-sensing modulons in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **24**:1169–1178.
8. Cochran, W. L., S.-J. Suh, G. A. McFeters, and P. S. Stewart. Unpublished data.
9. Cox, C. D. 1986. Role of pyocyanin in the acquisition of iron from transferrin. *Infect. Immun.* **52**:263–270.

10. Cox, C. D., and R. Graham. 1979. Isolation of an iron-binding compound from *Pseudomonas aeruginosa*. J. Bacteriol. 137:357–364.
11. Cross, A., J. R. Allen, J. Burke, G. Ducloux, A. Harris, J. John, D. Johnson, M. Lew, B. MacMillan, R. Skalova, R. Wenzel, and J. Tenney. 1983. Nosocomial infections due to *Pseudomonas aeruginosa*: review of recent trends. Rev. Infect. Dis. 5(Suppl.):S837–S845.
12. Davis, R. W., D. Botstein, and J. R. Roth. 1980. Advanced bacterial genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
13. Doig, P., T. Todd, P. A. Sastry, K. K. Lee, R. S. Hodges, W. Paranchych, and R. T. Irvin. 1988. Role of pili in adhesion of *Pseudomonas aeruginosa* to human respiratory epithelial cells. Infect. Immun. 56:1641–1646.
14. Doring, G., M. Maier, E. Muller, Z. Bibi, B. Tummeler, and A. Kharazmi. 1987. Virulence factors of *Pseudomonas aeruginosa*. Antibiot. Chemother. 39:136–148.
15. Figurski, D., and D. R. Helinski. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. Proc. Natl. Acad. Sci. USA 76:1648–1652.
16. Fujita, M., K. Tanaka, H. Takahashi, and A. Amemura. 1994. Transcription of the principal sigma-factor genes, *rpoD* and *rpoS*, in *Pseudomonas aeruginosa* is controlled according to the growth phase. Mol. Microbiol. 13:1071–1077.
17. Goldberg, J. B., and D. E. Ohman. 1984. Cloning and expression in *Pseudomonas aeruginosa* of a gene involved in the production of alginate. J. Bacteriol. 158:1115–1121.
18. Govan, J. R. W., and G. S. Harris. 1986. *Pseudomonas aeruginosa* and cystic fibrosis: unusual bacterial adaptation and pathogenesis. Microbiol. Sci. 3:302–308.
19. Gross, C. A., M. Lonetto, and R. Losick. 1992. Bacterial sigma factors, p. 129–176. In S. McKnight (ed.), Transcriptional regulation. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
20. Hassett, D. J., J.-F. Ma, J. G. Elkins, U. A. Ochsner, S. E. H. West, C. T. Huang, J. Fredericks, S. Burnett, P. S. Stewart, G. McFeters, and T. R. McDermott. Quorum sensing in *Pseudomonas aeruginosa* controls catalase and superoxide dismutase genes and mediates biofilm susceptibility to hydrogen peroxide. Submitted for publication.
21. Hassett, D. J., H. P. Schweizer, and D. E. Ohman. 1995. *Pseudomonas aeruginosa* *sodA* and *sodB* mutants defective in manganese- and iron-cofactored superoxide dismutase activity demonstrate the importance of the iron-cofactored form in aerobic metabolism. J. Bacteriol. 177:6330–6337.
22. Hengge-Aronis, R. 1996. Back to log phase: σ^S as a global regulator in the osmotic control of gene expression in *Escherichia coli*. Mol. Microbiol. 21:887–893.
23. Holloway, B. W., V. Krishnapillai, and A. F. Morgan. 1979. Chromosomal genetics of *Pseudomonas*. Microbiol. Rev. 43:73–102.
24. Iriarte, M., I. Stainier, and G. R. Cornelis. 1995. The *rpoS* gene from *Yersinia enterocolitica* and its influence on expression of virulence factors. Infect. Immun. 63:1840–1847.
25. Ivanova, A., C. Miller, G. Glinsky, and A. Eisenstark. 1994. Role of *rpoS* (*katF*) in *oxyR*-independent regulation of hydroperoxidase I in *Escherichia coli*. Mol. Microbiol. 12:571–578.
26. Kessler, E., M. Safrin, J. C. Olson, and D. E. Ohman. 1993. Secreted LasA of *Pseudomonas aeruginosa* is a staphylolytic protease. J. Biol. Chem. 268:7503–7508.
27. Kim, Y., L. S. Watrud, and A. Matin. 1995. A carbon starvation survival gene of *Pseudomonas putida* is regulated by σ^{54} . J. Bacteriol. 177:1850–1859.
28. King, E. O., M. K. Ward, and D. E. Raney. 1954. Two simple media for the demonstration of pyocyanin and fluorescin. J. Lab. Clin. Med. 44:301–307.
29. Knutson, C. A., and A. Jeanes. 1968. A new modification of the carbazole analysis: application to heteropolysaccharides. Anal. Biochem. 24:470–481.
30. Kolter, R., D. A. Siegle, and A. Tormo. 1993. The stationary phase of the bacterial life cycle. Annu. Rev. Microbiol. 47:855–874.
31. Kowarz, L., C. Coyault, V. Robbe-Saule, and F. Norel. 1994. The *Salmonella typhimurium katF* (*rpoS*) gene: cloning, nucleotide sequence, and regulation of *spvR* and *spvABCD* virulence plasmid genes. J. Bacteriol. 176:6852–6860.
32. Kurioka, S., and M. Matsuda. 1976. Phospholipase C activity using *p*-nitrophenylphosphorylcholine together with sorbitol and its application to studying the metal and detergent requirement of the enzyme. Anal. Biochem. 75:281–289.
33. Lange, R., and R. Hengge-Aronis. 1991. Identification of a central regulator of stationary-phase gene expression in *Escherichia coli*. Mol. Microbiol. 5:49–59.
34. Latifi, A., M. Foglino, K. Tanaka, P. Williams, and A. Lazdunski. 1996. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhlR (VsmR) to expression of the stationary-phase sigma factor RpoS. Mol. Microbiol. 21:1137–1146.
35. Latifi, A., M. K. Winson, M. Foglino, B. W. Bycroft, G. S. A. B. Stewart, A. Lazdunski, and P. Williams. 1995. Multiple homologues of LuxR and LuxI control the expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. Mol. Microbiol. 17:333–343.
36. Liu, P. V. 1973. Exotoxins of *P. aeruginosa*. I. Factors that influence the production of exotoxin A. J. Infect. Dis. 128:506–513.
37. Liu, P. V. 1974. Extracellular toxins of *Pseudomonas aeruginosa*. J. Infect. Dis. 130(Suppl.):S94–S99.
38. Loewen, P. C., and R. Hengge-Aronis. 1994. The role of the sigma factor σ^S (KatF) in bacterial global regulation. Annu. Rev. Microbiol. 48:53–80.
39. Losick, R., and P. Stragier. 1992. Crisscross regulation of cell-type-specific gene expression during development in *B. subtilis*. Nature 355:601–604.
40. Ma, J.-F., U. A. Ochsner, M. G. Klotz, V. K. Nanayakkara, M. L. Howell, Z. Johnson, J. E. Posey, M. L. Vasil, J. J. Monaco, and D. J. Hassett. Bacterioferritin A modulates catalase (KatA) activity and resistance to hydrogen peroxide in *Pseudomonas aeruginosa*. J. Bacteriol. 181:3730–3742.
41. Mathee, K., O. Ciofu, C. K. Sternberg, P. Lindum, J. Campbell, P. Jensen, A. Jøhnsen, M. Givskov, D. Ohman, S. Molin, N. Høiby, and A. Kharazmi. 1999. Mucoid conversion of *Pseudomonas aeruginosa* by hydrogen peroxide: a mechanism for virulence activation in the cystic fibrosis lung. Microbiology 145:1349–1357.
42. McCann, M. P., J. P. Kidwell, and A. Matin. 1991. The putative σ factor KatF has a central role in development of starvation-mediated general resistance in *Escherichia coli*. J. Bacteriol. 173:4188–4194.
43. Miura, K., S. Inouye, and A. Nakazawa. 1998. The *rpoS* gene regulates OP2, an operon for the lower pathway of xylene catabolism on the TOL plasmid, and the stress response in *Pseudomonas putida* mt-2. Mol. Gen. Genet. 259:72–78.
44. Mulvey, M. R., J. Switala, A. Borys, and P. C. Loewen. 1990. Regulation of transcription of *katE* and *katF* in *Escherichia coli*. J. Bacteriol. 172:6713–6720.
45. Ohman, D. E., and A. M. Chakrabarty. 1981. Genetic mapping of chromosomal determinants for the production of the exopolysaccharide alginate in a *Pseudomonas aeruginosa* cystic fibrosis isolate. Infect. Immun. 33:142–148.
46. Ohman, D. E., S. J. Cryz, and B. H. Iglewski. 1980. Isolation and characterization of a *Pseudomonas aeruginosa* PAO mutant that produces altered elastase. J. Bacteriol. 142:836–842.
47. Ohman, D. E., J. C. Sadoff, and B. H. Iglewski. 1980. Toxin A-deficient mutants of *Pseudomonas aeruginosa* PA103: isolation and characterization. Infect. Immun. 28:899–908.
48. Olsen, A., A. Arnqvist, M. Hammar, S. Sukupolvi, and S. Normark. 1993. The RpoS sigma factor relieves H-NS-mediated transcriptional repression of *csgA*, the subunit gene of fibronectin-binding curli. Mol. Microbiol. 7:523–536.
49. Pesci, E. C., J. P. Pearson, P. C. Seed, and B. H. Iglewski. 1997. Regulation of *las* and *rhl* quorum sensing in *Pseudomonas aeruginosa*. J. Bacteriol. 179:3127–3132.
50. Prince, R. W., C. D. Cox, and M. L. Vasil. 1993. Coordinate regulation of siderophore and exotoxin A production: molecular cloning and sequencing of the *Pseudomonas aeruginosa fur* gene. J. Bacteriol. 175:2589–2598.
51. Ramos-González, M. I., and S. Molin. 1998. Cloning, sequencing, and phenotypic characterization of the *rpoS* gene from *Pseudomonas putida* KT2440. J. Bacteriol. 180:3421–3431.
52. Sarniguet, A., J. Kraus, M. Henkels, A. M. Muehchen, and J. E. Loper. 1995. The sigma factor σ^S affects antibiotic production and biological control activity of *Pseudomonas fluorescens* Pf-5. Proc. Natl. Acad. Sci. USA 92:12255–12259.
53. Schweizer, H. D. 1993. Small broad-host-range gentamicin resistance gene cassettes for site-specific insertion and deletion mutagenesis. BioTechniques 15:831–834.
54. Selvaraj, G., Y. C. Fong, and V. N. Iyer. 1984. A portable DNA sequence carrying the cohesive site (*cos*) of bacteriophage λ and the *mob* (mobilization) region of the broad-host-range plasmid RK2: a module for the construction of new cosmids. Gene 32:235–241.
55. Small, P., D. Blankenhorn, D. Welty, E. Zinser, and J. L. Slonczewski. 1994. Acid and base resistance in *Escherichia coli* and *Shigella flexneri*: role of *rpoS* and growth pH. J. Bacteriol. 176:1729–1737.
56. Sokol, P. A., and D. E. Woods. 1988. Effect of pyochelin on *Pseudomonas cepacia* respiratory infections. Microb. Pathog. 5:197–205.
57. Southern, E. M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. J. Mol. Biol. 98:503–517.
58. Tanaka, K., and H. Takahashi. 1994. Cloning, analysis and expression of an *rpoS* homologue gene from *Pseudomonas aeruginosa* PAO1. Gene 150:81–85.
59. Ulmer, A. J., J. Pryjma, Z. Tarnok, M. Ernst, and H.-D. Flad. 1990. Inhibitory and stimulatory effects of *Pseudomonas aeruginosa* pyocyanin on human T and B lymphocytes and human monocytes. Infect. Immun. 58:808–815.
60. Woods, D. E., S. J. Cryz, R. L. Friedman, and B. H. Iglewski. 1982. Contribution of toxin A and elastase to virulence of *Pseudomonas aeruginosa* in chronic lung infections of rats. Infect. Immun. 36:1223–1228.
61. Yildiz, F. H., and G. K. Schoolnik. 1998. Role of *rpoS* in stress survival and virulence of *Vibrio cholerae*. J. Bacteriol. 180:773–784.
62. Yu, H., M. J. Schurr, and V. Deretic. 1995. Functional equivalence of *Escherichia coli* σ^E and *Pseudomonas aeruginosa* AlgU: *E. coli* *rpoE* restores mucoidy and reduces sensitivity to reactive oxygen intermediates in *algU* mutants of *P. aeruginosa*. J. Bacteriol. 177:3259–3268.