

Expression of the σ^B -Dependent General Stress Regulon Confers Multiple Stress Resistance in *Bacillus subtilis*

UWE VÖLKER,¹ BJÖRN MAUL,² AND MICHAEL HECKER^{2*}

Laboratorium für Mikrobiologie und MPI für terrestrische Mikrobiologie, Philipps-Universität, 35043 Marburg,¹ and
Institut für Mikrobiologie und Molekularbiologie, Ernst-Moritz-Arndt-Universität, 17487 Greifswald,² Germany

Received 16 February 1999/Accepted 19 April 1999

The alternative sigma factor σ^B of *Bacillus subtilis* is required for the induction of approximately 100 genes after the imposition of a whole range of stresses and energy limitation. In this study, we investigated the impact of a null mutation in *sigB* on the stress and starvation survival of *B. subtilis*. *sigB* mutants which failed to induce the regulon following stress displayed an at least 50- to 100-fold decrease in survival of severe heat (54°C) or ethanol (9%) shock, salt (10%) stress, and acid (pH 4.3) stress, as well as freezing and desiccation, compared to the wild type. Preloading cells with σ^B -dependent general stress proteins prior to growth-inhibiting stress conferred considerable protection against heat and salt. Exhaustion of glucose or phosphate induced the σ^B response, but surprisingly, σ^B did not seem to be required for starvation survival. Starved wild-type cells exhibited about 10-fold greater resistance to salt stress than exponentially growing cells. The data argue that the expression of σ^B -dependent genes provides nonsporulated *B. subtilis* cells with a nonspecific multiple stress resistance that may be relevant for stress survival in the natural ecosystem.

One of the earliest responses of a *Bacillus subtilis* cell population to different stressful environmental conditions is the striking and immediate induction of a large number of general stress proteins. Heat, acid, ethanol, and salt stress or starvation for glucose, oxygen, or phosphate induces probably more than 100 stress genes that form the σ^B -dependent general stress regulon (4, 11, 14, 45). The mechanism of this induction has been elucidated: the *sigB* operon itself is induced by all of the stress and starvation signals (9, 10, 13, 25, 28, 45). The complex signal transduction network conveying the different stress and starvation stimuli to *sigB* has been elucidated by the laboratories of Price, Haldenwang, and Losick: all of these stimuli enhance the activity of σ^B , thereby inducing the *sigB* operon itself and consequently the entire σ^B regulon (1–3, 8, 9, 15, 19, 29, 47, 48, 51).

Contrary to the wealth of information on the regulation of the activity of σ^B , the physiological function of σ^B remained obscure despite the increasing list of newly described genes subject to σ^B control (24). Based on the induction profile, we suggested in 1986 that these proteins may have a rather nonspecific but nevertheless essential protective function under stress regardless of the specific growth-limiting factor (38). Later we found that the translational capacity used for the production of general stress proteins increases from about 1% in growing cells to 20 or even 30% in starved or stressed cells, suggesting a crucial physiological role of the proteins in stress adaptation (11). Elucidation of the structure and function of this stress and starvation regulon seemed to be essential for understanding the physiology of growth-restricted bacterial cells in general.

Two principal approaches should help to elucidate the function of the general stress regulon: (i) identification of the function of stress proteins belonging to the σ^B regulon may allow predictions on the function of the entire regulon, and (ii)

comparison of the stress tolerance of wild-type cells with that of strains carrying mutations in *sigB* itself or in σ^B -dependent general stress genes will provide experimental evidence for the function of the entire regulon.

Some of the general stress proteins identified so far seem to be involved in the nonspecific protection from oxidative stress, from osmotic and water stress, or from heat stress. Therefore, we recently concluded that the products of σ^B -dependent stress genes most likely provide *B. subtilis* cells in transition from a growing to a nongrowing state with a comprehensive multiple stress resistance machine in anticipation of future stress during long-term survival as nongrowing vegetative cells. Furthermore, we suggested that this stress-resistant state of growth-restricted cells could be an essential part of a survival strategy alternative to sporulation (for a review, see reference 26).

However, the phenotypic studies of *sigB* mutants available up to 1995 did not indicate any essential role in stress adaptation. σ^B mutants were impaired in neither growth nor sporulation (12, 20). Furthermore, σ^B seemed to be nonessential for the survival of *B. subtilis* following heat shock, since a *sigB* null mutant was as temperature resistant as the wild type (10). First hints for a role of σ^B -dependent genes emerged when an insertional inactivation of the σ^B -dependent *ctc* gene resulted in a sporulation frequency at 48°C lower than that in the wild type (43).

Considering the discrepancies between the growing list of newly described σ^B -dependent genes on the one hand and the lack of a distinct stress-sensitive phenotype of *sigB* mutants on the other, a systematic reinvestigation of the function of the σ^B -dependent stress response became necessary. Recently, a *sigB* mutant was found to be extremely sensitive to the radical-producing agent cumene hydroperoxide both during growth and in stationary phase (5). Furthermore, the *sigB* mutant was impaired in the development of the resistance against hydrogen peroxide nonspecifically acquired in glucose-starved cells (21) as well as against acid or alkaline stress (22). In this report, we present the results of a systematic study on the stress resistance of *sigB* mutants compared to the wild type, underlining

* Corresponding author. Mailing address: Ernst-Moritz-Arndt-Universität, Institut für Mikrobiologie und Molekularbiologie, Friedrich-Ludwig-Jahn-Str. 15, 17487 Greifswald, Germany. Phone: 0049-3834-864200. Fax: 0049-3834-864202. E-mail: hecker@microbio7.biologie.uni-greifswald.de.




FIG. 1. Growth and survival of *B. subtilis* strains during heat shock and ethanol stress. The wild-type strain 168 (squares) and its isogenic *sigB* mutant ML6 (triangles) were grown in LB and exposed to either heat shock (A and B) or ethanol stress (C). (A) The growth of cultures was monitored by measuring the optical density at 540 nm. The strains were cultivated at 37°C (open symbols) or transferred at time zero to 52°C (black symbols) or 54°C (gray symbols). (B) Survival was determined by plating appropriate dilutions of samples taken at the time indicated. The number of cells of each strain present at the time zero immediately before treatment was set at 100%. Open symbols, cultures shifted to 54°C at time zero; filled symbols, cultures incubated for 30 min at 48°C prior to transfer to the final temperature of 54°C. (C) Viability was assayed immediately before and at different time points after the addition of ethanol to a final concentration of 9% (vol/vol).

the suggestion that σ^B is responsible for the development of multiple stress resistance in nongrowing *B. subtilis* cells.

MATERIALS AND METHODS

Bacterial strains. The *B. subtilis* wild-type strain 168 (*trpC2*), its isogenic *sigB* mutant ML6 (*trpC2 sigB::ΔHindIII-EcoRV::cat*) (27), and BSA115 were grown either in LB or in a synthetic medium described previously (42). In BSA115 (*trpC2 rsbU::kan P_BΔ28::P_{spac} rsbW313 pTET-I SPβ ctc::lacZ*) (44), *rsbW* is inactivated by a frameshift mutation and the downstream half of the *sigB* operon is placed under the control of the isopropyl-β-D-thiogalactopyranoside (IPTG)-inducible promoter *P_{spac}*. Inclusion of IPTG triggers the production of active σ^B and hyperinduction of the σ^B regulon in this strain.

Growth conditions and viability assay. The cultures were inoculated into prewarmed LB or synthetic medium; during exponential growth, the cultures (6×10^7 to 8×10^7 cells per ml) were exposed to the stresses according to the following scheme: for heat shock, the culture was transferred to 54°C; for ethanol stress, ethanol was added to a final concentration of 9% (vol/vol); for salt stress, solid NaCl was added to a final concentration of 10% (wt/vol); for acid shock, 1 N HCl was added to shift the pH of the culture from 7.5 to 4.3; for freezing, aliquots of the culture were rapidly frozen at -20°C, thawed, and plated at the time indicated. Cells were preadapted if indicated by incubation for 30 min either in the presence of 4% NaCl, at 48°C, or at pH 5.2. Thereafter, the stress strength was raised to the level described above. The influence of preloading cells with σ^B -dependent stress proteins was investigated by dividing an exponentially growing culture of BSA115 60 min (LB) or 120 min (synthetic medium) prior to stress and including 1 mM IPTG (final concentration) in one half of the culture. Both parts were exposed to the stresses as described above. The osmoprotectant glycine betaine was added in some experiments to a final concentration of 1 mM.

At the time points indicated, aliquots of the culture were sampled and diluted in 0.9% (wt/vol) NaCl solution, and appropriate dilutions were plated on LB agar plates to determine cell viability. All experiments were performed at least in triplicate, and the results of typical experiments are displayed in the figures.

RESULTS

Role of σ^B in heat and ethanol stress resistance. Transferring *B. subtilis* from 37 to 48°C is commonly applied to induce the expression of the σ^B regulon (10, 45), but the *sigB* mutant displayed no difference in either growth or survival compared with the wild type (data not shown). However, assuming that σ^B -dependent gene expression serves to prepare the cell for more severe stress conditions, then the effect of the lack of σ^B could be more pronounced at harsher conditions. Therefore, the heat shock was repeated by shifting bacteria from 37 to 52°C. Whereas wild-type bacteria continued to grow, albeit at a reduced rate, the *sigB* mutant strain ceased growth and

started to lyse (Fig. 1A). Since this reduction in optical density of the *sigB* mutant was accompanied by a slight decrease in survival only (data not shown), the heat shock was extended to 54°C, which also prevented growth of the wild type (Fig. 1A). At this critical point when even the wild-type strain was severely impaired, viability declined dramatically within 60 min for both strains, but striking differences in survival between the wild type and the *sigB* mutant became visible (Fig. 1B). A mild heat preadaptation period at 48°C prevented the decrease in viability in the wild type but not in the *sigB* mutant. As a result, the differences in survival between the two strains became much more pronounced: after 5 h, a more than 1,000-fold difference in heat stress survival between the wild type and the *sigB* mutant was found. Nevertheless, we observed a protective effect by mild heat stress also in the *sigB* mutant, most likely because of the induction of the σ^B -independent chaperone-encoding genes (Fig. 1B).

Ethanol treatment seems to induce cell damage similar to that caused by heat shock (36), and ethanol-treated cells of a *sigB* mutant displayed a lower growth yield than the corresponding wild-type strain (22). Therefore, results similar to those observed for heat shock were expected for ethanol. The *sigB* mutant showed indeed an approximately 100-fold greater drop in cell survival in response to treatment with 9% ethanol than its wild-type counterpart (Fig. 1C).

Role of σ^B in resistance to salt stress. Comparison of the growth patterns of *sigB* mutant cells and wild-type bacteria after exposure to NaCl in the range from 4 to 16% (final concentration) revealed no difference between the two strains (data not shown). However, the strains displayed clear differences in survival of such harsh conditions. The survival rate of the wild-type strain was approximately 50-fold higher than that of the *sigB* mutant (Fig. 2A). An attempt to protect these cells by a mild (4% NaCl) salt preadaptation for 30 min increased the viability of the wild type 10-fold but had no effect on the *sigB* mutant (Fig. 2A). In contrast to heat shock, the pretreatment did not completely protect the wild-type bacteria.

Glycine betaine is a very effective osmoprotectant in *B. subtilis* (30). We wanted to determine whether glycine betaine can compensate for the missing induction of the σ^B response in the




FIG. 2. Survival of *B. subtilis* strains during salt stress. The wild-type strain 168 (squares) and its isogenic *sigB* mutant ML6 (triangles) were grown in a synthetic medium and exposed to salt stress. Survival was determined as described in the legend to Fig. 1. (A) Open symbols represent cultures exposed to 10% (wt/vol) sodium chloride at time zero; filled symbols represent cultures pretreated with a mild salt stress of 4% NaCl for 30 min before the sodium chloride concentration was raised to 10% (wt/vol) at time zero. (B) The *sigB* mutant ML6 was grown in synthetic medium with (closed triangles) or without (open triangles) 1 mM glycine betaine; 30 min before the sodium chloride concentration was increased to 10% (wt/vol), the bacteria were exposed to a mild salt stress of 4% NaCl.

sigB mutant. Addition of 1 mM glycine betaine to growing *sigB* mutant cells before the imposition of salt stress triggered an almost complete protection against the otherwise lethal salt stress (Fig. 2B). This result indicates that induction of σ^B -dependent general stress genes and accumulation of osmoprotective substances constitute alternative mechanisms for protection against the deleterious effects of growth-inhibiting salt concentrations. Inclusion of 1 mM glycine betaine in the culture medium did not protect heat-damaged cells (data not shown), indicating that heat and salt seem to cause distinctively different types of damage.

Artificial induction of the σ^B response and cross-adaptation produce stress resistance. To separate the effects of the σ^B response from protection triggered by induction of specific stress proteins, we analyzed the effects of cross-adaptation and preloading cells with σ^B -dependent general stress proteins.

A mild heat preadaptation which also induced the σ^B -dependent stress genes was almost as effective as a mild salt pretreatment in protection against severe salt stress at least within 60 min after imposition of the salt stress (Fig. 3B). As expected, this cross-protection occurred in the wild type but not in the *sigB* mutant. Preincubating wild-type cells with 4% NaCl for 30 min prior to a heat treatment at 54°C similarly provided a significant but incomplete protection (Fig. 3A).

In another set of experiments, the level of σ^B -dependent stress proteins was varied without exposing bacteria to stress. In *B. subtilis* BSA115, a *sigB* operon with a null mutation in the primary negative regulator *rsbW* is placed under the control of the IPTG-inducible promoter P_{spac} . Addition of IPTG prior to heat shock triggered the accumulation of large amounts of general stress proteins in the absence of stress and partially prevented the sharp decline in cell survival at 54°C observed in the absence of IPTG (Fig. 3C). In the case of severe salt stress, the addition of IPTG almost completely restored cell viability, indicating that the σ^B -dependent stress response seems to be mainly responsible for the resistance against severe salt stress (Fig. 3D).

σ^B and acid stress resistance. Exponentially growing cells treated at pH 4.3 showed a dramatic drop in cell survival within 60 min, but the *sigB* mutant was about 50- to 100-fold more sensitive than its wild-type counterpart (Fig. 4). Both strains,

however, exhibited an about 10-fold increase in cell viability after a 30-min preexposure to mild acid stress (pH 5.2), indicating that this protective acid-adaptive response does not depend solely on σ^B .

Influence of σ^B on survival of freezing and thawing. The effect on survival of *B. subtilis* of changing the water activity inside the cell during freezing and thawing was tested. Samples of the *B. subtilis* wild-type strain 168 and its isogenic *sigB* mutant strain ML6 were taken 1 h after the cultures had entered the stationary phase as a result of the exhaustion of glucose from a synthetic medium and immediately frozen at -20°C without the addition of a cryoprotective substance. Over a period of 14 days, we thawed aliquots of the samples each day at room temperature and assayed viability by plating on LB agar plates. The results shown in Fig. 5 indicate that the wild-type cells survived the treatment 50- to 100-fold better than the *sigB* mutant. These findings were supported by desiccation experiments: the wild-type strain survived lyophilization at least 10-fold better than the *sigB* strain (data not shown).

Role of σ^B in starvation survival. σ^B does not seem to be essential for starvation survival under glucose or phosphate limitation in a minimal medium. Similar starvation survival kinetics were measured for the wild type and the *sigB* mutant (data not shown). It is well known that glucose-starved cells become resistant to oxidative stress (for a review, see reference 18). This nonspecifically acquired stress resistance depends on σ^B because glucose-starved cells of a *sigB* mutant were much more sensitive to oxidative stress than starved wild-type cells (21). Because glucose starvation activates the σ^B response, we wondered if the resistance to other stresses in glucose-starved cells is also enhanced under these conditions. Surprisingly, glucose-starved cells were as sensitive to heat stress as exponentially growing cells (not shown). In the case of salt stress resistance, the difference between growing and glucose-starved cells was at least 10-fold (Fig. 6). However, glucose-starved wild-type and *sigB* mutant cells differed only two- to threefold in salt stress resistance, because of the development of a σ^B -independent salt resistance in both strains during glucose starvation (Fig. 6).




FIG. 3. Influence of cross-adaptation (A and B) and preloading bacteria with general stress proteins (C and D) on survival of heat shock (A and C) or salt stress (B and D). Bacteria were grown in a synthetic medium, and during exponential growth the cultures were exposed to heat shock or salt stress. Survival was determined as described in the legend to Fig. 1. (A) The wild-type strain 168 was pretreated for 30 min with either a mild heat shock of 48°C (black squares) or a mild salt stress of 4% NaCl (gray squares) prior to the shift to 54°C or directly exposed to 54°C (open squares) at time zero. (B) The wild-type strain 168 (squares) or its isogenic *sigB* counterpart ML6 (triangles) were exposed to 10% (wt/vol) sodium chloride at time zero. Cells were immediately exposed to the severe salt stress (open symbols) or pretreated for 30 min with 4% NaCl (black symbols) or a mild heat shock of 48°C (gray symbols). (C and D) *B. subtilis* BSA115 was grown in synthetic medium. During exponential growth, the culture was divided and IPTG was added to one half to a final concentration of 1 mM; 120 min after the addition of IPTG (time zero), both cultures were exposed to either 54°C (C) or 10% (wt/vol) NaCl (D). Open triangles, no IPTG; closed triangles, 1 mM IPTG added.

DISCUSSION

σ^B was identified as the first bacterial alternative sigma factor almost 20 years ago, but interest in σ^B declined after it was determined that this sigma factor is not involved in sporulation (12, 20, 27, 28). The discovery that σ^B , whose activity itself is tightly regulated by a set of different environmental stimuli, controls a large general stress and starvation regulon activated in nongrowing cells may have been responsible for the revival of attention to σ^B (10, 13, 14, 45). Several laboratories started to investigate not only the role of σ^B in the adaptation to growth-restricting conditions but also the pathogenicity of gram-positive bacteria such as *Staphylococcus aureus*, *Listeria monocytogenes*, and mycobacteria (7, 17, 32, 33, 50).

The discrepancy between a constantly increasing list of newly described σ^B -dependent stress genes (26) and the lack of any altered phenotype of *sigB* mutants (12, 20) called for a careful reinvestigation of the physiological role of σ^B . It was tempting to speculate that this large stress regulon might fulfill a crucial adaptive function under stress, but experimental evidence was lacking. Recently, new data revealed that σ^B is

necessary for the adaptation to oxidative stress (21) as well as to acid or alkaline stress (22). It is interesting that the σ^B regulon is clearly involved in adaptation to oxidative and alkaline stress, but neither induces this σ^B regulon. On the other hand, no evidence has been presented for any role of σ^B in protection against heat and salt stress, which are excellent inducers of the more than 100 genes of the σ^B regulon. The results presented in this report suggest that general stress proteins are required during growth inhibition resulting from severe heat or salt stress. Conditions which reduce but still permit growth do not require σ^B -dependent stress proteins for the development of resistance. Instead, specific stress proteins such as the systems used for the uptake and synthesis of osmoprotective substances sufficiently protect the cells during, for example, less severe osmotic stress (49; for a review, see reference (30)).

Schulz et al. (40) and Li and Wong (34) found that *dnaK* or *groESL* mutants of *B. subtilis* were impaired in thermotolerance. Our data clearly demonstrate that besides chaperones, σ^B -dependent stress proteins are also involved in heat stress





FIG. 4. Influence of σ^B on survival of acid shock. During exponential growth in LB, cultures of the wild-type strain 168 (squares) and the *sigB* mutant ML6 (triangles) were exposed to acid shock. Survival was determined as described in the legend to Fig. 1. Acid shock was imposed either directly at time zero by shifting the pH of the culture medium from 7.5 to 4.3 (open symbols) or after a preadaptation for 30 min with a intermediate acid shock at pH 5.2 (filled symbols).

resistance in *B. subtilis*. The increase in the survival rates at 54°C of *sigB* mutant cells after pretreatment at 48°C (Fig. 1) might be explained by the induction of the DnaK and GroEL and also of the Clp machinery in the *sigB* mutant (11, 31). The much stronger, almost complete protective effect of the same mild heat shock pretreatment of the wild type argues for a major role of general stress proteins in this heat stress resistance (Fig. 1). Such a conclusion gains support from the protection from severe heat shock by cross-adaptation with a mild salt stress or preloading cells of BSA115 with σ^B -dependent stress proteins in the absence of stress (Fig. 3). Both variants yield only partial protection since they failed to induce the heat-specific chaperones, corroborating the essential function of chaperones in this adaptation (34, 40, 46).



in a glucose-limited chemostat in the absence of additional stresses (41). These results indicate that σ^B -dependent stress proteins do not contribute to the starvation survival potential of glucose-starved cells; rather, the main function of σ^B is to protect the starved cell against future stress that may emerge during long-term survival. Glucose-starved cells develop an oxidative stress resistance that depends on σ^B (21). This non-specific and multiple stress resistance of glucose-starved cells also includes salt stress resistance that increased more than 10-fold during starvation. Surprisingly, the salt resistance acquired during glucose starvation is only partially σ^B dependent (Fig. 6). Thermal resistance, however, does not significantly increase after glucose exhaustion compared to exponentially growth despite the fact that the σ^B regulon is fully active. Failure of development of a full heat stress resistance might at least partially be accounted for by the lack of induction of chaperones in glucose-starved cells (11).

Therefore, the main function of σ^B during energy depletion should be to equip glucose-starved cells with nonspecific resistance to oxidative stress, including resistance to freezing and thawing or to UV light damage (not shown here). In addition to this function, σ^B is essential for the interaction with a severe heat, ethanol, acid, or salt challenge, cross-protection between these stimuli included. Oxidative stress, changing osmotic or salt stress conditions triggered by desiccation and rain, and also acid or heat stress may be typical growth-restricting conditions in the upper layers of soil, the natural habitat of *B. subtilis*. The σ^B -dependent stress response could be particularly relevant for cell survival in soil under conditions that do not allow efficient sporulation such as high osmolarity or low cell density (23, 39). Therefore, it is tempting to speculate that *sigB* mutants in soil have a survival disadvantage compared to the wild type, but experimental evidence is still lacking.

The function of the *sigB* regulon in multiple and prospective stress resistance is in good accordance with recent data on the function of σ^B -dependent genes (for a review, see reference 26). These genes form a complex regulon that may comprise more than 100 genes. The identification of their function was one of our main strategies to uncover the function of the regulon. It is an attractive goal for future research to assign the single general stress proteins to the different components of the multiple stress resistance network typical of stressed or starving *B. subtilis* cells.

ACKNOWLEDGMENTS

We thank Anita Harang for excellent technical assistance throughout this investigation. We also thank E. Bremer for stimulating discussion.

This work was supported by grants from the Deutsche Forschungsgemeinschaft to M.H. and U.V. and by a grant from the Fonds der Chemischen Industrie to M.H.

REFERENCES

- Akbar, S., C. M. Kang, T. A. Gaidenko, and C. W. Price. 1997. Modulator protein RsbR regulates environmental signalling in the general stress pathway of *Bacillus subtilis*. *Mol. Microbiol.* **24**:567–578.
- Alper, S., A. Dufour, D. A. Garsin, L. Duncan, and R. Losick. 1996. Role of adenosine nucleotides in the regulation of a stress-response transcription factor in *Bacillus subtilis*. *J. Mol. Biol.* **260**:165–177.
- Alper, S., L. Duncan, and R. Losick. 1994. An adenosine nucleotide switch controlling the activity of a cell type-specific transcription factor in *B. subtilis*. *Cell* **77**:195–205.
- Antelmann, H., J. Bernhardt, R. Schmid, H. Mach, U. Völker, and M. Hecker. 1997. First steps from a two-dimensional protein index towards a response-regulation map for *Bacillus subtilis*. *Electrophoresis* **18**:1451–1463.
- Antelmann, H., S. Engelmann, R. Schmid, and M. Hecker. 1996. General and oxidative stress responses in *Bacillus subtilis*: cloning, expression, and mutation of the alkyl hydroperoxide reductase operon. *J. Bacteriol.* **178**:6571–6578.
- Antelmann, H., S. Engelmann, R. Schmid, A. Sorokin, A. Lapidus, and M. Hecker. 1997. Expression of a stress- and starvation-induced *dps/pexB*-homologous gene is controlled by the alternative sigma factor σ^B in *Bacillus subtilis*. *J. Bacteriol.* **179**:7251–7256.
- Becker, L. A., M. S. Cetin, R. W. Hutkins, and A. K. Benson. 1998. Identification of the gene encoding the alternative sigma factor σ^B from *Listeria monocytogenes* and its role in osmotolerance. *J. Bacteriol.* **180**:4547–4554.
- Benson, A. K., and W. G. Haldenwang. 1993. *Bacillus subtilis* σ^B is regulated by a binding protein (RsbW) that blocks its association with core RNA polymerase. *Proc. Natl. Acad. Sci. USA* **90**:2330–2334.
- Benson, A. K., and W. G. Haldenwang. 1992. Characterization of a regulatory network that controls σ^B expression in *Bacillus subtilis*. *J. Bacteriol.* **174**:749–757.
- Benson, A. K., and W. G. Haldenwang. 1993. The σ^B -dependent promoter of the *Bacillus subtilis* *sigB* operon is induced by heat shock. *J. Bacteriol.* **175**:1929–1935.
- Bernhardt, J., U. Völker, A. Völker, H. Antelmann, R. Schmid, H. Mach, and M. Hecker. 1997. Specific and general stress proteins in *Bacillus subtilis*—a two-dimensional protein electrophoresis study. *Microbiology* **143**:999–1017.
- Binnie, C., M. Lampe, and R. Losick. 1986. Gene encoding the sigma 37 species of RNA polymerase sigma factor from *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **83**:5943–5947.
- Boylan, S. A., A. R. Redfield, M. S. Brody, and C. W. Price. 1993. Stress-induced activation of the σ^B transcription factor of *Bacillus subtilis*. *J. Bacteriol.* **175**:7931–7937.
- Boylan, S. A., A. R. Redfield, and C. W. Price. 1993. Transcription factor σ^B of *Bacillus subtilis* controls a large stationary-phase regulon. *J. Bacteriol.* **175**:3957–3963.
- Boylan, S. A., A. Rutherford, S. M. Thomas, and C. W. Price. 1992. Activation of *Bacillus subtilis* transcription factor σ^B by a regulatory pathway responsive to stationary-phase signals. *J. Bacteriol.* **174**:3695–3706.
- Chan, P. F., S. J. Foster, E. Ingham, and M. O. Clements. 1998. The *Staphylococcus aureus* alternative sigma factor σ^B controls the environmental stress response but not starvation survival or pathogenicity in a mouse abscess model. *J. Bacteriol.* **180**:6082–6089.
- DeMaio, J., Y. Zhang, C. Ko, and W. Bshai. 1997. *Mycobacterium tuberculosis* *sigF* is part of a gene cluster with similarities to the *Bacillus subtilis* *sigF* and *sigB* operons. *Tuberc. Lung Dis.* **78**:3–12.
- Dowds, B. C. A. 1994. The oxidative stress response in *Bacillus subtilis*. *FEMS Microbiol. Lett.* **124**:255–263.
- Dufour, A., and W. G. Haldenwang. 1994. Interactions between a *Bacillus subtilis* anti-sigma factor (RsbW) and its antagonist (RsbV). *J. Bacteriol.* **176**:1813–1820.
- Duncan, M. L., S. S. Kalman, S. M. Thomas, and C. W. Price. 1987. Gene encoding the 37,000-dalton minor sigma factor of *Bacillus subtilis* RNA polymerase: isolation, nucleotide sequence, chromosomal locus, and cryptic function. *J. Bacteriol.* **169**:771–778.
- Engelmann, S., and M. Hecker. 1996. Impaired oxidative stress resistance of *Bacillus subtilis* *sigB* mutants and the role of *katA* and *kate*. *FEMS Microbiol. Lett.* **145**:63–69.
- Gaidenko, T. A., and C. W. Price. 1998. General stress transcription factor- σ^B and sporulation transcription factor- σ^{H} each contribute to survival of *Bacillus subtilis* under extreme growth conditions. *J. Bacteriol.* **180**:3730–3733.
- Grossman, A., and R. Losick. 1988. Extracellular control of spore formation in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **85**:4369–4373.
- Haldenwang, W. G. 1995. The sigma factors of *Bacillus subtilis*. *Microbiol. Rev.* **59**:1–30.
- Hecker, M., W. Schumann, and U. Völker. 1996. Heat-shock and general stress response in *Bacillus subtilis*. *Mol. Microbiol.* **19**:417–428.
- Hecker, M., and U. Völker. 1998. Non-specific, general and multiple stress resistance of growth-restricted *Bacillus subtilis* cells by the expression of the σ^B regulon. *Mol. Microbiol.* **29**:1129–1136.
- Igo, M., M. Lampe, C. Ray, W. Schafer, C. P. Moran, Jr., and R. Losick. 1987. Genetic studies of a secondary RNA polymerase sigma factor in *Bacillus subtilis*. *J. Bacteriol.* **169**:3464–3469.
- Kalman, S., M. L. Duncan, S. M. Thomas, and C. W. Price. 1990. Similar organization of the *sigB* and *spoIIA* operons encoding alternate sigma factors of *Bacillus subtilis* RNA polymerase. *J. Bacteriol.* **172**:5575–5585.
- Kang, C. M., K. Vijay, and C. W. Price. 1998. Serine kinase activity of a *Bacillus subtilis* switch protein is required to transduce environmental stress signals but not to activate its target PP2C phosphatase. *Mol. Microbiol.* **30**:189–196.
- Kempf, B., and E. Bremer. 1998. Uptake and synthesis of compatible solutes as microbial stress responses to high-osmolarity environments. *Arch. Microbiol.* **170**:319–330.
- Krüger, E., T. Msadek, and M. Hecker. 1996. Alternate promoters direct stress-induced transcription of the *Bacillus subtilis* *clpC* operon. *Mol. Microbiol.* **20**:713–723.
- Kullik, I., and P. Giachino. 1997. The alternative sigma factor σ^B in *Staphylococcus aureus*—regulation of the *sigB* operon in response to growth phase and heat shock. *Arch. Microbiol.* **167**:151–159.
- Kullik, I., P. Giachino, and T. Fuchs. 1998. Deletion of the alternative sigma

- factor σ^B in *Staphylococcus aureus* reveals its function as a global regulator of virulence genes. *J. Bacteriol.* **180**:4814–4820.
34. Li, M., and S. Wong. 1992. Cloning and characterization of the *groESL* operon from *Bacillus subtilis*. *J. Bacteriol.* **174**:3981–3992.
 35. Matin, A., E. A. Auger, P. H. Blum, and J. E. Schultz. 1989. Genetic basis of starvation survival in nondifferentiating bacteria. *Annu. Rev. Microbiol.* **43**: 293–316.
 36. Mogk, A., A. Völker, S. Engelmann, M. Hecker, W. Schumann, and U. Völker. 1998. Nonnative proteins signal induction of the *Bacillus subtilis* CIRCE regulon. *J. Bacteriol.* **180**:2895–2900.
 37. Park, J., C. Grant, M. Davies, and I. Dawes. 1998. The cytoplasmic Cu,Zn superoxide dismutase of *Saccharomyces cerevisiae* is required for resistance to freeze-thaw stress—generation of free radicals during freezing and thawing. *J. Biol. Chem.* **273**:22921–22928.
 38. Richter, A., and M. Hecker. 1986. Heat-shock proteins in *Bacillus subtilis*: a two-dimensional gel electrophoresis study. *FEMS Microbiol. Lett.* **36**:69–71.
 39. Ruzal, S. M., C. Lopez, E. Rivas, and C. Sanchez-Rivas. 1998. Osmotic strength blocks sporulation at stage II by impeding activation of early sigma factors in *Bacillus subtilis*. *Curr. Microbiol.* **36**:75–79.
 40. Schulz, A., B. Tzschaschel, and W. Schumann. 1995. Isolation and analysis of mutants of the *dnaK* operon of *Bacillus subtilis*. *Mol. Microbiol.* **15**:421–429.
 41. Schweder, T., A. Kolyschcow, U. Völker, and M. Hecker. Analysis of the expression and function of the σ^B -dependent general stress regulon of *Bacillus subtilis* during slow growth. *Arch. Microbiol.*, in press.
 42. Stülke, J., R. Hanschke, and M. Hecker. 1993. Temporal activation of beta-glucanase synthesis in *Bacillus subtilis* is mediated by the GTP pool. *J. Gen. Microbiol.* **139**:2041–2045.
 43. Truitt, C. L., E. A. Weaver, and W. G. Haldenwang. 1988. Effects on growth and sporulation of inactivation of a *Bacillus subtilis* gene (*ctc*) transcribed in vitro by minor vegetative cell RNA polymerases ($E\text{-}\sigma^{37}$, $E\text{-}\sigma^{32}$). *Mol. Gen. Genet.* **212**:166–171.
 44. Völker, U., A. Dufour, and W. G. Haldenwang. 1995. The *Bacillus subtilis* *rsbU* gene product is necessary for RsbX-dependent regulation of σ^B . *J. Bacteriol.* **177**:114–122.
 45. Völker, U., S. Engelmann, B. Maul, S. Riethdorf, A. Völker, R. Schmid, H. Mach, and M. Hecker. 1994. Analysis of the induction of general stress proteins of *Bacillus subtilis*. *Microbiology* **140**:741–752.
 46. Völker, U., H. Mach, R. Schmid, and M. Hecker. 1992. Stress proteins and cross-protection by heat shock and salt stress in *Bacillus subtilis*. *J. Gen. Microbiol.* **138**:2125–2135.
 47. Völker, U., A. Völker, and W. G. Haldenwang. 1996. Reactivation of the *Bacillus subtilis* anti- σ^B antagonist, RsbV, by stress- or starvation-induced phosphatase activities. *J. Bacteriol.* **178**:5456–5463.
 48. Völker, U., A. Völker, B. Maul, M. Hecker, A. Dufour, and W. G. Haldenwang. 1995. Separate mechanisms activate σ^B of *Bacillus subtilis* in response to environmental and metabolic stresses. *J. Bacteriol.* **177**:3771–3780.
 49. VonBlohn, C., B. Kempf, R. M. Kappes, and E. Bremer. 1997. Osmostress response in *Bacillus subtilis*—characterization of a proline uptake system (OpuE) regulated by high osmolarity and the alternative transcription factor sigma B. *Mol. Microbiol.* **25**:175–187.
 50. Wu, S. W., H. Delencastre, and A. Tomasz. 1996. Sigma-B, a putative operon encoding alternate sigma factor of *Staphylococcus aureus* RNA polymerase—molecular cloning and DNA sequencing. *J. Bacteriol.* **178**:6036–6042.
 51. Yang, X. F., C. M. Kang, M. S. Brody, and C. W. Price. 1996. Opposing pairs of serine protein kinases and phosphatases transmit signals of environmental stress to activate a bacterial transcription factor. *Genes Dev.* **10**:2265–2275.