

NOTES

Characterization of a *Pseudomonas aeruginosa* Fatty Acid Biosynthetic Gene Cluster: Purification of Acyl Carrier Protein (ACP) and Malonyl-Coenzyme A:ACP Transacylase (FabD)

ALECKSANDR J. KUTCHMA,[†] TUNG T. HOANG, AND HERBERT P. SCHWEIZER*

Department of Microbiology, Colorado State University, Fort Collins, Colorado 80523

Received 8 April 1999/Accepted 14 June 1999

A DNA fragment containing the *Pseudomonas aeruginosa* *fabD* (encoding malonyl-coenzyme A [CoA]:acyl carrier protein [ACP] transacylase), *fabG* (encoding β -ketoacyl-ACP reductase), *acpP* (encoding ACP), and *fabF* (encoding β -ketoacyl-ACP synthase II) genes was cloned and sequenced. This *fab* gene cluster is delimited by the *plsX* (encoding a poorly understood enzyme of phospholipid metabolism) and *pabC* (encoding 4-amino-4-deoxychorismate lyase) genes; the *fabF* and *pabC* genes seem to be translationally coupled. The *fabH* gene (encoding β -ketoacyl-ACP synthase III), which in most gram-negative bacteria is located between *plsX* and *fabD*, is absent from this gene cluster. A chromosomal temperature-sensitive *fabD* mutant was obtained by site-directed mutagenesis that resulted in a W258Q change. A chromosomal *fabF* insertion mutant was generated, and the resulting mutant strain contained substantially reduced levels of *cis*-vaccenic acid. Multiple attempts aimed at disruption of the chromosomal *fabG* gene were unsuccessful. We purified FabD as a hexahistidine fusion protein (H₆-FabD) and ACP in its native form via an ACP-intein-chitin binding domain fusion protein, using a novel expression and purification scheme that should be applicable to ACP from other bacteria. Matrix-assisted laser desorption-ionization spectroscopy, native polyacrylamide electrophoresis, and amino-terminal sequencing revealed that (i) most of the purified ACP was properly modified with its 4'-phosphopantetheine functional group, (ii) it was not acylated, and (iii) the amino-terminal methionine was removed. In an *in vitro* system, purified ACP functioned as acyl acceptor and H₆-FabD exhibited malonyl-CoA:ACP transacylase activity.

Fatty acid biosynthesis in eukaryotes is catalyzed by a multienzyme complex encoded by a single gene and known as the type I system. In contrast, eubacteria contain a type II or dissociated Fab (fatty acid biosynthesis) system, where the reactions are carried out by proteins encoded by multiple genes (for a review, see reference 6). Bacterial fatty acid biosynthesis necessitates a three-carbon precursor, malonyl-coenzyme A (CoA), which is derived from acetyl-CoA by the action of acetyl-CoA carboxylase. The malonyl-CoA is transferred to the acyl carrier protein (ACP) by malonyl-CoA:ACP acyltransferase (FabD). Fatty acid elongation involves four reactions: (i) a condensation reaction catalyzed by one of three β -ketoacyl-ACP synthases, FabB, FabF, or FabH; (ii) a reduction involving a NADPH-dependent β -ketoacyl-ACP reductase (FabG); (iii) a dehydration reaction catalyzed by either FabA or FabZ, both of which are β -hydroxyacyl-ACP dehydratases with broad, overlapping chain length specificities (15); and (iv) a second reduction reaction catalyzed by NADH-dependent enoyl-ACP reductase (FabI).

In *Escherichia coli* (28) and other bacteria (9, 25, 38), the genes *acpP*, *fabD*, *fabF*, *fabG*, and *fabH*, encoding ACP, FabD, FabF, FabG, and FabH, respectively, are contained in a *fab*

gene cluster. In *Bacillus subtilis*, *fabH* is not part of the major *fab* gene cluster (25).

Besides the role of ACP in phospholipid (6) and rhamnolipid (3, 27) synthesis, ACPs play central roles in a broad range of other biosynthetic pathways that depend on acyl transfer reactions, including polyketide (35), nonribosomal peptide (1), and depsipeptide biosynthesis (31), as well as in the transacylation of oligosaccharides (8, 11) and proteins (22). More recently, acyl-ACPs derived from the Fab pathway have been proposed to be the acyl donors for synthesis of acylated homoserine lactones (AHLs) (26, 32, 40). The AHLs (or autoinducers) have received considerable attention in recent years since they are required for a regulatory phenomenon termed quorum sensing (10). Characterization of the two known AHL-producing systems of *Pseudomonas aeruginosa*, *rhl* and *las*, has shown that the respective autoinducers, *N*-butyryl homoserine lactone (C₄-HSL) and *N*-(3-oxo)-dodecanoyl homoserine lactone (3-oxo-C₁₂-HSL), are required for expression of a multitude of virulence factors and secondary metabolites (41).

We recently began characterization of the unique aspects of the *P. aeruginosa* Fab system (20), and in this report we describe the characterization of the *P. aeruginosa* *fabD-fabG-acpP-fabF* gene cluster, as well as the purification and characterization of FabD and ACP.

Cloning and characterization of the *acpP*-containing region. When we initiated these experiments, the sequence of only a fragmented *acpP*-containing region was available from the *P. aeruginosa* genome database, and it contained no intact *acpP* gene. This *acpP*-containing gene cluster was cloned into M13,

* Corresponding author. Mailing address: Department of Microbiology, Colorado State University, Fort Collins, CO 80523. Phone: (970) 491-3536. Fax: (970) 491-1815. E-mail: hschweiz@cvmbs.colostate.edu.

[†] Present address: Myriad Genetics, Salt Lake City, UT 84108.

TABLE 1. Strains, plasmids, and primers used in this study

Strain, plasmid, or primer	Relevant properties or sequence ^a	Reference or origin
<i>E. coli</i>		
BL21(DE3)	<i>E. coli</i> B; F ⁻ <i>ompT</i> r _B ⁻ m _B ⁻ (λDE3)	37
SA1503(DE3)	<i>lon-100 his-87 ompT::Km^r relA1 rpsL781 spoT1 thi-1</i> (λDE3)	19a
<i>P. aeruginosa</i>		
PAO1	Prototroph	B. H. Holloway
PAO198	PAO1 with <i>fabF::Gm^r</i>	This study
PAO204	PAO1 with <i>fabD</i> (Ts)	This study
Plasmids		
pEX100T	Ap ^r <i>sacB</i> ⁺ ; gene replacement vector	34
pUCGM	Ap ^r Gm ^r ; source of Gm ^r cassette	33
pWSK29 and pWKS30	Ap ^r ; low-copy-number cloning and T7 expression vectors	44
pCYB1	Ap ^r ; intein-chitin binding domain expression vector	New England Biolabs
pET-15b	Ap ^r ; hexahistidine fusion and expression vector	Novagen
pGEM-T	Ap ^r ; TA-cloning vector	Promega
pNam	Ap ^r ; low-copy-number intein-chitin binding domain expression vector	19a
pT7-7	Ap ^r ; T7 promoter expression vector	39
pPS671	Ap ^r <i>fabF</i> ⁺ <i>pabC</i> ⁺ (ligation of chromosomal 4-kb <i>EcoRI-HindIII</i> fragment between the same sites of pWSK29)	This study
pPS681	Ap ^r <i>fabD</i> ⁺ <i>fabG</i> ⁺ <i>acpP</i> ⁺ <i>fabF</i> ⁺ <i>pabC</i> ⁺ (ligation of 3.4-kb <i>BamHI-EcoRV</i> fragment from pPS840 between the <i>BamHI</i> and <i>XbaI</i> [blunt-ended] sites of pPS671)	This study
pPS831	Ap ^r ; PCR-amplified 147-bp genomic segment from PAO1 cloned into pGEM-T	This study
pPS840	Ap ^r <i>fabD</i> ⁺ <i>fabG</i> ⁺ <i>acpP</i> ⁺ (3.7-kb chromosomal <i>EcoRV-BamHI</i> fragment cloned between the same sites of pWSK29)	This study
pPS966	Ap ^r ; ligation of 1.8-kb <i>NdeI-PstI</i> fragment from pPS981 between the same sites of pT7-7	This study
pPS981	Ap ^r <i>acpP</i> ⁺ ; ligation of <i>NdeI-SapI</i> -digested PCR fragment between the same sites of pCYB1	This study
pPS1096	Ap ^r <i>acpP</i> ⁺ (ligation of 735-bp <i>HindIII-XbaI</i> fragment from pPS966 between the same sites of pNam)	This study
Primers		
ACP1	ATCGAGGAGCG(A/C/T/G)GT(G/C)AAGAAGAT(A/C)AT	
ACP2	GTCGAACTCCTC(G/C)TC(A/C/T/G)AG(G/C)GCCAT	
FabD1	TGGATGCTCTCCACCTgGCGACCCGGGCTGT	
ACP-Nde	<i>NdeI</i> -TGAAAACAACATatgAGCACCATCGAAGAACGCGTT	
ACP-Sap	<i>SapI</i> -ATCCGGCTCTTCCGCAatgCTGGGGAGC	
FabD-Nde	<i>NdeI</i> -GGGACCTATcatATGTCTGCATCCCTCGCATTTCGTC	
FabD-Bam	<i>BamHI</i> -CCCAGGatCCCCAGTTCCAGCGCAATCGCC	

^a Primer sequences are printed 5' to 3'; lowercase letters indicate nonmatching oligonucleotides used to either form the indicated motif as underlined or introduce other nonmotif changes indicated in lowercase letters.

which led to expression of toxic levels of apo-ACP. Previous attempts to clone intact *acpP* from *E. coli* (42), as well as from a number of other bacteria (25, 36), into multicopy plasmids were hampered by the toxic effects of overexpressed apo-ACP (23). We therefore chose to reclone the chromosomal *acpP*-containing region into the low copy-number pWSK vectors (typically exhibiting only five to eight copies per cell [44]). A 147-bp sequence containing a partial *acpP* coding sequence was PCR amplified from *P. aeruginosa* chromosomal DNA by using the two primers ACP1 and ACP2 (Table 1). These partially degenerate primers were modeled after conserved amino acid sequence regions found in the *E. coli*, *Haemophilus influenzae* and *Vibrio harveyi* ACP homologs. PCR was performed on a PTC-100 PCR system thermocycler (MJ Research, Watertown, Mass.), using *Taq* polymerase from Qiagen (Santa Clarita, Calif.) and previously described conditions (19). The PCR fragment was cloned into pGEM-T (Promega, Madison, Wis.) to form pPS831. Nucleotide sequence analysis revealed the presence of a partial *acpP* gene on the PCR fragment. This fragment was used as a probe to generate a partial restriction map of the PAO1 chromosomal *acpP* region, which revealed a

3.4-kb *EcoRV-BamHI* fragment which was cloned into the low-copy-number cloning vector pWKS30 by using previously described strategies (17, 20), yielding pPS840 (Fig. 1A).

The entire nucleotide sequence of both strands of a 3,433-bp chromosomal *EcoRV-BamHI* fragment was determined (relevant segments of the sequence are shown in Fig. 1B). BLAST searches revealed the presence of the 3' end of *plsX*, complete sequences of *fabD*, *fabG*, and *acpP*, and the 5' end of *fabF* (Fig. 1A). To obtain the 3' end of *fabF*, a 1.1-kb *BamHI-EcoRI* fragment from pPS840 was used as a probe to identify a ~4-kb *EcoRI-HindIII* fragment, which was cloned into pWSK29 to generate pPS671 (Fig. 1A).

Sequence analyses of pPS671 and several subclones derived from it established the entire sequence of both strands from the *BamHI* site within *fabF* to the *SalI* site immediately downstream of *fabF* (Fig. 1A). The *fab* gene cluster contains *fabF*, *acpP*, *fabG*, and *fabD* but not *fabH*. Many bacteria, i.e., *B. subtilis* (25), *V. harveyi* (36), *H. influenzae* (9), and *Streptomyces glaucescens* (38), contain similarly organized *fab* gene clusters, but *P. aeruginosa* provides the first example of a gram-negative bacterium where the *fabH* gene is absent from the




FIG. 1. The *P. aeruginosa* *acpP* region. (A) Maps of plasmids containing a *fab* gene cluster and flanking genes. The chromosomal inserts of pPS840 and pPS671 were cloned individually, and fusion of these two clones at the common *Bam*HI site yielded pPS681, containing a continuous sequence of this region. Abbreviations: Ba, *Bam*HI; Ec, *Eco*RI; Ev, *Eco*RV; Hd, *Hind*III; Ps, *Pst*I; Sa, *Sal*I; Xb, *Xba*I; Xh, *Xho*I. The following genes and their products were identified by nucleotide sequencing: *plsX*, encoding a poorly understood protein involved in phospholipid biosynthesis; *fabD*, malonyl-CoA:ACP transacylase; *fabG*, β -ketoacyl-ACP reductase; *acpP*, ACP; *fabF*, β -ketoacyl-ACP synthase II; *pabC*, 4-amino-deoxychorismate lyase; *tmk*, thymidylate kinase. (B) Partial nucleotide sequences of the *P. aeruginosa* *plsX*, *fabD*, *fabG*, *acpP*, *fabF*, and *pabC* genes. Most of the sequences within the structural genes are omitted, as indicated by dashes. The deduced amino acid sequences are given in one-letter code below the nucleotide sequence. Putative ribosome binding sites (RBS) are labeled. Numbers above the sequence mark the first nucleotides of the initiation and last codons, and stop codons are marked with asterisks. The TG-to-CA nucleotide changes in *fabD* that were introduced by site-directed mutagenesis and resulted in a W258Q change and a FabD(Ts) phenotype are indicated above the nucleotide sequence.

acpP-containing cluster; the only other known example is *B. subtilis* (25).

The *fab* cluster is delimited by *pabC* and *plsX* homologs, encoding 4-amino-deoxychorismate lyase and a protein of un-

known function in phospholipid synthesis, respectively. We recently confirmed that *pabC* encodes *P. aeruginosa* aminodeoxychorismate lyase (13), which is required for *p*-aminobenzoic acid (PABA) synthesis, since *pabC* mutants are PABA auxo-

trophs (18). Unlike *E. coli*, the ATG start codon of the *P. aeruginosa* *pabC* gene overlaps the TGA stop codon of *fabF*. Although the two genes are translationally coupled, it is unclear whether both genes are cotranscribed from a single promoter upstream of *fabF*. The *fabF* mutant PAO198 (Table 1), containing a nonpolar gentamicin resistance (Gm^r) cassette in a transcriptional orientation opposite *fabF* and *pabC*, was not a PABA auxotroph, and we have not yet isolated mutants containing polar *fabF* mutations.

After our studies were completed, a contig appeared in the *Pseudomonas* genome project database; albeit not annotated, this entry confirmed our results and the gene order *plsX-fabD-fabG-*acpP-fabF-pabC**, as well as the absence of *fabH* from this cluster. Searches of this database revealed several possible *fabH* homologs, which are located elsewhere in the chromosome, and we are in the process of characterizing the most likely *fabH* homolog.

The deduced amino acid sequences of the *fabD*, *fabG*, *acpP*, and *fabF* genes were analyzed, and, in general, found to be most similar to the respective *E. coli* homologs. The identities were 66% for FabF, 65% for FabG, and 56% for FabD. However, other bacterial Fab proteins also exhibited considerable similarities to the deduced *P. aeruginosa* protein sequences.

The high degree of primary amino acid sequence conservation was especially evident with ACP. *P. aeruginosa* ACP was 90% identical to *E. coli* ACP, 83% identical to *H. influenzae* ACP, and 80% identical to *V. harveyi* ACP. All ACPs consisted of between 76 to 78 amino acids, and they contained (i) the consensus sequence, DSLD, for attachment of the 4'-phosphopantetheine and (ii) a large proportion of acidic amino acids. The calculated isoelectric point of PAO1 ACP (pH 3.8) confirmed the acidic nature of this protein.

Characteristics of *fab* genes and their products. Several lines of evidence indicated that the *fabF*, *fabD*, and *fabG* genes cloned in this work encode β -ketoacyl-ACP synthase II (FabF), malonyl-CoA:ACP transacylase (FabD), and β -ketoacyl-ACP reductase (FabG), respectively.

The evidence for FabF includes the following: (i) there is a high degree of similarity or identity to the well-characterized homologs from *E. coli*; (ii) translation of both *E. coli* and *P. aeruginosa* FabF coding sequences is apparently initiated at a GTG because the first Met codon found in the *P. aeruginosa* FabF protein is not preceded by a suitable Shine-Dalgarno sequence and the first 10 NH₂-terminal amino acids of the *P. aeruginosa* FabF protein (MSRRRVVITG) are 80% identical to those of the *E. coli* FabF protein (MSKRRVVVTG), which has a GTG start (GenBank accession no. AE000210); and (iii) a *P. aeruginosa* *fabF* mutant was constructed by deletion of a 597-bp *NcoI* fragment from within the *fabF* gene and replacing it with a 830-bp blunt-ended Gm^r encoding cassette from plasmid pUCGM (33). The mutated *fabF* sequence was returned to the PAO1 chromosome via the gene replacement vector pEX100T (34). Gas chromatography-mass spectrometric (12) analysis of the fatty acid content in cell extracts of the *fabF* knockout mutant PAO198 (*fabF::Gm^r*) demonstrated eight-fold lower levels of *cis*-vaccenic acid compared to wild-type PAO1 (data not shown). This compares favorably with a 10-fold reduction of *cis*-vaccenic acid levels seen in an *E. coli* *fabF* mutant (12).

The following evidence suggests that the *fabD* gene encodes malonyl-CoA:ACP transacylase (FabD). (i) FabD not only is 56% identical (78% similar) to its counterpart from *E. coli* but also contains the critical catalytic domain, the conserved pentapeptide GHSLG, which is the GX SXG signature motif of serine-dependent acylhydrolases (2). Characterization of an *E. coli* *fabD*(Ts) mutant showed that the temperature-sensitive

(Ts) phenotype resulted from a mutation that led to a W257Q substitution (43). We changed a conserved Trp²⁵⁸ to a Gln residue in the *P. aeruginosa* FabD protein by mutating two nucleotides (Fig. 1B) by site-specific mutagenesis using the Altered Sites mutagenesis system (Promega). Primer FabD1 (Table 1) introduced a double TG-to-CA mismatch that resulted in a W258Q change in the FabD amino acid sequence, which was confirmed by nucleotide sequence analysis. Return of the *fabD*(Ts) allele to the PAO1 chromosome was achieved by using a previously described strategy (20). FabD(Ts) mutants, including PAO204, were obtained at a frequency of 23%, demonstrating the essentiality of FabD in *P. aeruginosa*. (iii) A hexahistidine (H₆)-tagged FabD protein was overexpressed and purified (Fig. 2A). Its observed molecular mass of ~33 kDa matches closely that for FabD (32,442 Da) plus the H₆-containing NH₂-terminal extension (2,181 Da). (iv) The purified FabD protein exhibits malonyl-CoA:ACP transacylase activity (see below).

Besides the high homology to other bacterial FabG proteins and the presence of the amino acids of the NADPH binding signature AX₂GXGX₂AX₆G near the NH₂ terminus, other evidence suggests its essential role in the cycles of fatty acid elongation in *P. aeruginosa*. First, despite repeated attempts, we were unable to isolate a *fabG* knockout mutation, indicating the gene's essential nature. Second, the purified FabG protein is essential for fatty acid elongation in a reconstituted *in vitro* Fab synthesis system (21). A separate NADPH-dependent β -ketoacyl-ACP reductase, RhlG, that is specifically involved in rhamnolipid synthesis has recently been described (3), although no biochemical data supporting its presumed function were presented.

Expression and purification of ACP. The ACP was overexpressed and purified by using the intein-chitin binding domain (CDB) system. An ACP-intein-CBD fusion protein was constructed according to the basic protocol provided by New England Biolabs (Beverly, Mass.). Two PCR primers, ACP-Nde and ACP-Sap, were designed to introduce an *NdeI* site at the ACP initiation codon and a *SapI* site immediately downstream of the last codon of *acpP*. These primers were used to amplify a ~270-bp fragment by using pPS681 (Table 1) DNA as the template and standard PCR conditions (18). The PCR fragment was cloned into pCYB1 DNA. This procedure yielded pPS981. Using pPS981, various *E. coli* host strains, and different induction conditions, we detected no expression of ACP-intein-CBD unless the *lac* promoter was replaced with the more powerful T7 promoter from pT7-7 (39), a step that yielded pPS966. Optimization of the expression conditions by using *E. coli* BL21(DE3)/pPS966 revealed that an overnight induction and growth at room temperature (RT) in Luria-Bertani (LB) medium (Gibco-BRL, Gaithersburg, Md.) led to substantial overproduction of the ~66,000-M_r ACP-intein-CBD fusion protein.

For purification of ACP-intein-CBD from a high-copy-number expression vector, 4-liter LB-ampicillin (100 μ g/ml) cultures of BL21(DE3)/pPS966 were grown at 37°C to log phase (A_{600} of ~1.0), and gene expression was induced by addition of 0.5 mM isopropyl- β -D-thiogalactopyranoside (IPTG). The cells were shaken at RT and harvested after a 9-h induction period. The cells were resuspended in 4 liters of fresh LB medium without IPTG and further incubated for 1.5 h at 37°C before they were harvested. During this recovery period, most of the apo-ACP was converted to holo-ACP by ligation of the 4'-phosphopantetheine. When this period was omitted and the cells were harvested after a 13-h incubation at RT in the presence of IPTG, ~90 to 95% of the ACP preparation was apo-ACP which did not serve as a FabD substrate. The cell




FIG. 2. Purification of ACP and FabD, and assay of FabD activity. (A) The FabD protein was expressed as an H_6 -FabD protein and purified by Ni^{2+} -agarose affinity chromatography. ACP was expressed as an ACP-intein-CBD fusion protein which was adsorbed to a chitin-agarose affinity column, from which the ACP was eluted after self-cleavage of the fusion protein by DTT. Each protein (3 μ g) was analyzed by electrophoresis on a 0.1% SDS–10% polyacrylamide gel, and protein bands were visualized by Coomassie blue staining. The positions of two molecular weight markers, carbonic anhydrase (31 kDa) and trypsin inhibitor (21.5 kDa), are indicated on the left. (B) Acylation of ACP by H_6 -FabD. The reaction mixtures (20 μ l) contained ACP (2 μ g), H_6 -FabD (0.25 μ g), and 0.1 mM malonyl-CoA in the indicated combinations. The products were analyzed by electrophoresis on 20% native polyacrylamide gels followed by staining with Coomassie blue. (C) The acylation reaction and detection were performed as described for panel B except that the ACP was purified from cultures grown under conditions that did not allow for proper modification of apo-ACP with 4'-phosphopantetheine (see text for details). (D) The same experiment as shown in panel B except that the ACP was purified from a low-copy-number expression construct.

pellet was suspended in 50 ml of column buffer (20 mM Tris-HCl [pH 8.0], 500 mM NaCl, 0.1 mM EDTA, 0.1% Triton X-100, 1 mM phenylmethylsulfonyl fluoride) and disrupted by French press treatment (19,000 lb/in²). All subsequent steps were performed at 4°C. Cell debris was removed by ultracentrifugation for 1 h at 260,000 $\times g$. The cell extract was applied to a 10-ml bed volume of chitin beads (New England Biolabs) in a 30-ml column and washed with 15 volumes of column buffer. The column buffer was exchanged with cleavage buffer (20 mM Tris-HCl [pH 8.0], 50 mM NaCl, 0.1 mM EDTA, 30 mM dithiothreitol [DTT]), and cleavage of the fusion protein on the column was achieved by overnight incubation. The cleaved ACP was eluted with 30 ml of cleavage buffer without DTT, and fractions were collected. The ACP content of the fractions was analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (24), and protein concentrations were determined by using the Bradford dye binding assay (Bio-Rad Laboratories) and bovine serum albumin as the standard.

For purification of ACP from *E. coli* cells containing the low-copy-number expression vector pPS1096, 250-ml LB-ampicillin cultures of *E. coli* SA1503(DE3)/pPS1096 were grown at 37°C to log phase (A_{600} of ~ 1.0) and gene expression was induced by addition of 0.5 mM IPTG. The cells were shaken at RT for 12 h, and ACP was purified as described above. The low-copy-number expression vector allowed for a simpler induction scheme and yielded an ACP preparation with a holo-ACP content of $\sim 95\%$. This amount could not be boosted by growing the cells in the presence of pantothenic acid. In con-

trast to expression experiments using the high-copy-number expression construct, the low-copy-number expression vector did not lead to an observable cessation of cell growth.

SDS-PAGE analysis revealed that ACP had been purified to near homogeneity (Fig. 2A). On this gel, ACP migrated at the position of a 22.5-kDa protein, significantly larger than its calculated molecular mass of 8.7 kDa. The anomalous migration of PAO1 ACP on SDS-PAGE is consistent with observations for ACPs from other bacteria (25, 28, 35) and has been attributed to the protein's high charge-to-mass ratio (with ACP being highly acidic; calculated pI 3.8) as well as its low hydrophobic amino acid content, two factors that have a considerable influence on SDS binding (30). When the same ACP preparation was analyzed on a 0.1% SDS–13% polyacrylamide gel containing 5 M urea, we observed a single protein band of ~ 9 kDa (data not shown), a value that closely matches its calculated mass of 8.7 kDa.

Characterization of ACP. Matrix-assisted laser desorption-ionization mass spectrometric analysis (4) (performed at the Colorado State University Macromolecular Resource Facility) of the purified ACP fraction revealed three species with molecular masses of 8,583 Da (minor peak), 8,934 Da (major peak), and 17,844 Da (minor peak) (data not shown). These values correspond to the three ACP species commonly found in ACP preparations, apo-ACP (without 4'-phosphopantetheine), holo-ACP (with 4'-phosphopantetheine), and the ACP dimer. ACP dimerization was an artifact of the precipitation step used in matrix-assisted laser desorption-ionization sample preparation, as no dimers were present in our ACP preparation, as judged by native PAGE (data not shown). The data confirm that the majority of our recombinant ACP (the major peak at 8,934 Da) contains the 4'-phosphopantetheine group required for its activity, presumably attached to Ser³⁶, and that the NH₂-terminal methionine of ACP is posttranslationally removed by an aminopeptidase (16), as has been observed with other ACPs (25, 28, 35). The latter was verified by NH₂-terminal amino acid sequence analysis of purified *P. aeruginosa* ACP (performed at the Peptide Sequencing Facility at the University of Victoria, Victoria, British Columbia, Canada), which revealed the sequence STIEE, corresponding to amino acids 2 to 6 deduced from the nucleotide sequence (Fig. 1B).

Native PAGE is a powerful means for characterization of modified and unmodified ACP (29). It can be used to assess ACP dimer content and the ratio of apo-ACP to holo-ACP-SH. Since acylation alters the Stokes radius of ACP, it can also be used to assess the fraction of acyl-ACP in ACP preparations (29). Using native PAGE, we showed that our ACP preparation (i) contained $\sim 95\%$ holo-ACP and $\sim 5\%$ apo-ACP and (ii) contained no detectable acyl-ACPs (Fig. 2B). We do not know whether nonacylated ACP is due to masking of the 4'-phosphopantetheine prosthetic group in the ACP-intein-CBD fusion protein or due to efficient removal of acyl groups by the DTT used for cleavage of the fusion protein.

The identity of the holo-ACP was confirmed in a malonyl-CoA:ACP transferase assay by incubation of our ACP preparation with malonyl-CoA and purified FabD. For purification of FabD, a H_6 -FabD expression vector was constructed. The *fabD* coding sequence was PCR amplified from PAO1 genomic DNA with primers FabD-Nde, creating a *Nde*I site at the *fabD* ATG initiation codon, and FabD-Bam, which creates a *Bam*HI site immediately downstream of *fabD*. The PCR fragment was cloned into pET-15b (Novagen, Madison, Wis.) to form pPS979, which was then transformed into *E. coli* BL21(DE3). Expression of H_6 -FabD, cell lysis, and purification of the soluble fusion protein on a Ni^{2+} -agarose column (Qiagen) were performed as previously described (14) except that the cells

were grown in LB-ampicillin medium. Purified FabD was used in acylation reaction mixtures (20 μ l) that contained 2 μ g of ACP, 0.25 μ g of FabD, and 0.1 mM malonyl-CoA in 20 mM Tris-HCl (pH 7.2)–100 mM NaCl–10% glycerol–1 mM DTT–2 mM EDTA–25 mM MgSO₄–0.1 mM FeSO₄. The mixtures were incubated at RT for 5 min, and products were analyzed by electrophoresis on 20% native polyacrylamide gels and followed by staining with Coomassie blue R-250 as previously described (29). In this assay, at least 50% of the holo-ACP was converted to malonyl-ACP by FabD (Fig. 2B, lane 1); conversion was not complete since the reaction catalyzed by FabD is reversible. This reaction required FabD and malonyl-CoA since neither FabD alone (lane 2) nor malonyl-CoA alone (lane 1) led to formation of malonyl-ACP after a 5-min incubation period. Longer incubation times and higher malonyl-CoA concentrations resulted in some spontaneous acylation of ACP by malonyl-CoA alone. As shown in Fig. 2C, apo-ACP did not serve as a FabD substrate. Although of the ~10% holo-ACP present about half was converted to malonyl-ACP, the ACP preparation contained ~90% apo-ACP which did not serve as a FabD substrate. Similar results were obtained with ACP purified from a strain carrying a low-copy-number ACP overproducer (Fig. 2D).

Conclusions. The ACP purification procedure described herein has several distinct advantages over other purification methods (7, 8). (i) It is rapid, and yields are comparable to those obtained by other methods (15, 25, 29). We routinely isolate in excess of 5 mg of ACP from 4 liter of induced culture, containing either low- or high-copy-number expression vectors, using a single 5- to 10-ml chitin-agarose affinity column on which all washing steps and the cleavage step are performed. We only know of one other method that yields more ACP when expressed from an overproducer, but it is much more involved with respect to both labor and instrumentation requirements (15). (ii) Our purification procedure yields >95% holo-ACP, whereas other procedures relying on expression constructs yield mostly apo-ACP and little to almost no holo-ACP (7, 8). Although apo-ACP can efficiently be converted into holo-ACP by purified holo-ACP synthase (8), this step contaminates the ACP preparation, and thus additional steps are required to purify the holo-ACP before further use. (iii) ACP purified by our procedure is not acylated and therefore does not require the deacylation steps prior to its use as holo-ACP in the synthesis of defined acyl-ACP substrates, compared to ACP obtained with more traditional purification procedures (5). (iv) Our purification method is inexpensive and can be performed in laboratories that have neither the expertise nor the equipment necessary for traditional protein purification schemes.

We have successfully used the *P. aeruginosa* ACP prepared in this manner for the synthesis of defined acyl-ACP substrates (19); together with purified H₆-FabD and H₆-FabG, it functions in a reconstituted enzyme system leading to synthesis of biologically active 3-oxo-C₁₂-HSL from simple metabolic precursors (21).

Nucleotide sequence accession number. The complete sequence of the 3,811-bp *EcoRV*-*SalI* fragment (Fig. 1A) was deposited in GenBank and assigned accession no. U91631.

This work was supported by Public Health Service grant GM56685 from the National Institutes of Health and by a grant from the CSU College of Veterinary Medicine and Biomedical Sciences. A.J.K. was supported by a graduate teaching assistantship from the Department of Microbiology.

A.J.K. and T.T.H. contributed equally to this work.

REFERENCES

- Baldwin, J. E., J. W. Bird, R. A. Field, N. M. O'Callaghan, C. J. Schofield, and A. C. Willis. 1991. Isolation and partial characterization of ACV synthetase from *Cephalosporium acremonium* and *Streptomyces clavuligerus*. Evidence for the presence of phosphopantothinate in ACV synthase. *J. Antibiot.* **44**:241–248.
- Brenner, S. 1988. The molecular evolution of genes and proteins: a tale of two serines. *Nature* **334**:528–530.
- Campos-Garcia, J., A. D. Caro, R. Najera, R. M. Miller-Maier, R. A. Al-Tahhan, and G. Soberon-Chavez. 1998. The *Pseudomonas aeruginosa* *rhlG* gene encodes an NADPH-dependent β -ketoacyl reductase which is specifically involved in rhamnolipid synthesis. *J. Bacteriol.* **180**:4442–4451.
- Cottrell, J. S., and C. W. Sutton. 1998. The identification of electrophoretically separated proteins by peptide mass fingerprinting, p. 1–16. *In* J. M. Walker (ed.), *Protein protocols on CD-ROM*. Humana Press, Totowa, N.J.
- Cronan, J. E., and A. L. Klages. 1981. Chemical synthesis of acyl thioesters of acyl carrier protein with native structure. *Proc. Natl. Acad. Sci. USA* **78**:5440–5444.
- Cronan, J. E., and C. O. Rock. 1996. Biosynthesis of membrane lipids, p. 612–636. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.
- Crosby, J., D. H. Sherman, M. J. Bibb, W. P. Revill, D. A. Hopwood, and T. J. Simpson. 1995. Polyketide synthase acyl carrier proteins from *Streptomyces*: expression in *Escherichia coli*, purification and partial characterization. *Biochim. Biophys. Acta* **1251**:32–42.
- Eppele, G., K. M. G. M. Van der Drift, J. E. Thomas-Oates, and O. Geiger. 1998. Characterization of a novel acyl carrier protein, RkpF, encoded by an operon involved in capsular polysaccharide biosynthesis in *Sinorhizobium meliloti*. *J. Bacteriol.* **180**:4950–4954.
- Fleischmann, R. D., M. D. Adams, O. White, R. A. Clayton, E. F. Kirkness, A. R. Kerlavage, C. J. Bult, J.-F. Tomb, B. A. Dougherty, J. M. Merrick, K. McKenney, G. Sutton, W. FitzHugh, C. Fields, J. D. Gocayne, J. Scott, R. Shirley, L.-I. Liu, A. Glodek, J. M. Kelley, J. F. Weidman, C. A. Phillips, T. Spriggs, E. Hedblom, M. D. Cotton, T. R. Utterback, M. C. Hanna, D. T. Nguyen, D. M. Saudeck, R. C. Brandon, L. D. Fine, J. L. Fritchman, J. L. Fuhrmann, N. S. M. Geoghagen, C. L. Gnehm, L. A. McDonald, K. V. Small, C. M. Fraser, H. O. Smith, and J. C. Venter. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496–512.
- Fuqua, W. C., S. C. Winans, and E. P. Greenberg. 1996. Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu. Rev. Microbiol.* **50**:727–751.
- Geiger, O., H. P. Spaink, and E. P. Kennedy. 1991. Isolation of the *Rhizobium leguminosarum* NodF nodulation protein: NodF carries a 4'-phosphopantetheine prosthetic group. *J. Bacteriol.* **173**:2872–2878.
- Gelman, E. P., and J. E. Cronan. 1972. Mutant of *Escherichia coli* deficient in the synthesis of *cis*-vaccenic acid. *J. Bacteriol.* **112**:381–387.
- Green, J. M., W. K. Merkel, and B. P. Nichols. 1992. Characterization and sequence of *Escherichia coli* *pabC*, the gene encoding aminodeoxychorismate lyase, a pyridoxal phosphate-containing enzyme. *J. Bacteriol.* **174**:5317–5323.
- Heath, R. J., and C. O. Rock. 1996. Roles of the FabA and FabZ β -hydroxyacyl-acyl carrier protein dehydratase in *Escherichia coli* fatty acid biosynthesis. *J. Biol. Chem.* **271**:27795–27801.
- Hill, R. B., K. R. MacKenzie, J. M. Flanagan, J. E. Cronan, and J. H. Prestegard. 1995. Overexpression, purification and characterization of *Escherichia coli* acyl carrier protein and two mutant proteins. *Protein Expression Purif.* **6**:394–400.
- Hirel, P. H., J. M. Schmitter, P. Dessen, K. Fayat, and T. S. Banque. 1989. Extent of N-terminal methionine excision from *Escherichia coli* proteins is governed by the side-chain length of the penultimate amino acid. *Proc. Natl. Acad. Sci. USA* **86**:8274–8251.
- Hoang, T., S. Williams, and H. P. Schweizer. 1997. Molecular genetic analysis of the region containing the essential *Pseudomonas aeruginosa* *asd* gene encoding aspartate- β -semialdehyde dehydrogenase. *Microbiology* **143**:899–907.
- Hoang, T. T., R. R. Karkhoff-Schweizer, A. J. Kutchma, and H. P. Schweizer. 1998. A broad-host-range Flp-*FRT* recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**:77–86.
- Hoang, T. T., and H. P. Schweizer. 1999. Characterization of *Pseudomonas aeruginosa* enoyl-acyl carrier protein reductase: a target for the antimicrobial triclosan and its role in acylated homoserine lactone synthesis. *J. Bacteriol.* **181**:5489–5497.
- Hoang, T. T., Y. Ma, R. J. Stern, M. R. McNeil, and H. P. Schweizer. Construction and use of low-copy number T7 expression vectors for purification of problem proteins: purification of *Mycobacterium tuberculosis* RmlD and *Pseudomonas aeruginosa* LasI and RhlI, and functional analysis of purified RhlI. *Gene*, in press.
- Hoang, T. T., and H. P. Schweizer. 1997. Fatty acid biosynthesis in *Pseudomonas aeruginosa*: cloning and characterization of the *fabAB* operon encod-

- ing β -hydroxyacyl-acyl carrier protein dehydratase (*fabA*) and β -ketoacyl-acyl carrier protein synthase I (*fabB*). *J. Bacteriol.* **179**:5326–5332.
21. **Hoang, T. T., and H. P. Schweizer.** 1999. Synthesis of the *Pseudomonas aeruginosa* autoinducer *N*-[3-oxo-dodecanoyl]-L-homoserine lactone from metabolic precursors in a reconstituted enzyme system, abstr. B/D309, p. 90. In Abstracts of the 99th General Meeting of the American Society for Microbiology, 1999. American Society for Microbiology, Washington, D.C.
 22. **Issartel, J. P., V. Koronakis, and C. Hughes.** 1991. Activation of *Escherichia coli* prohaemolysin to the mature toxin by acyl carrier protein-dependent fatty acylation. *Nature* **351**:759–761.
 23. **Keating, D. H., M. R. Carey, and J. E. Cronan.** 1995. The unmodified (apo) form of *Escherichia coli* acyl carrier protein is a potent inhibitor of cell growth. *J. Biol. Chem.* **270**:22229–22235.
 24. **Makowski, G. S., and M. L. Ramsby.** 1993. pH modification to enhance the molecular sieving properties of sodium dodecyl sulfate-10% polyacrylamide gel. *Anal. Biochem.* **212**:283–285.
 25. **Morbidoni, H. R., D. De Mendoza, and J. E. Cronan.** 1996. *Bacillus subtilis* acyl carrier protein is encoded in a cluster of lipid biosynthesis genes. *J. Bacteriol.* **178**:4794–4800.
 26. **Moré, M. I., D. Finger, J. L. Stryker, C. Fuqua, A. Eberhard, and S. C. Winans.** 1996. Enzymatic synthesis of a quorum-sensing autoinducer through use of defined substrates. *Science* **272**:1655–1658.
 27. **Ochsner, U. A., A. Fiechter, and J. Reiser.** 1994. Isolation, characterization, and expression in *Escherichia coli* of the *Pseudomonas aeruginosa* *rhlAB* genes encoding a rhamnosyltransferase involved in rhamnolipid biosurfactant synthesis. *J. Biol. Chem.* **269**:19787–19795.
 28. **Rawlings, M., and J. E. Cronan.** 1992. The gene encoding *Escherichia coli* acyl carrier protein lies within a cluster of fatty acid biosynthetic genes. *J. Biol. Chem.* **267**:5751–5754.
 29. **Rock, C. O., and J. E. Cronan.** 1981. Acyl carrier protein from *Escherichia coli*. *Methods Enzymol.* **71**:341–351.
 30. **Rock, C. O., and J. E. Cronan.** 1979. Re-evaluation of the solution structure of acyl carrier protein. *J. Biol. Chem.* **254**:9778–9785.
 31. **Rusnak, F., M. Sakaitani, D. Drucekhammer, J. Reichert, and C. T. Walsh.** 1991. Biosynthesis of the *Escherichia coli* siderophore enterobactin: sequence of the *entF* gene, expression and purification of EntF, and analysis of covalent phosphopantetheine. *Biochemistry* **30**:2916–2927.
 32. **Schaefer, A. L., D. L. Val, B. L. Hanzelka, J. E. Cronan, and E. P. Greenberg.** 1996. Generation of cell-to-cell signals in quorum sensing: acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. *Proc. Natl. Acad. Sci. USA* **93**:9505–9509.
 33. **Schweizer, H. P.** 1993. Small broad-host-range gentamycin resistance cassettes for site-specific insertion and deletion mutagenesis. *BioTechniques* **15**:831–833.
 34. **Schweizer, H. P., and T. Hoang.** 1995. An improved system for gene replacement and *xylE* fusion analysis in *Pseudomonas aeruginosa*. *Gene* **158**:15–22.
 35. **Shen, B., R. G. Summers, H. Gramajo, M. J. Bibb, and C. R. Hutchinson.** 1992. Purification and characterization of the acyl carrier protein of the *Streptomyces glaucescens* tetracenomycin C polyketide synthase. *J. Bacteriol.* **174**:3818–3821.
 36. **Shen, Z., and D. M. Byers.** 1996. Isolation of *Vibrio harveyi* acyl carrier protein and the *fabG*, *acpP*, and *fabF* genes involved in fatty acid biosynthesis. *J. Bacteriol.* **178**:571–573.
 37. **Studier, F. W., A. H. Rosenberg, J. J. Dunn, and J. W. Dubendorff.** 1990. Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.* **185**:60–89.
 38. **Summers, R. G., A. Ali, B. Shen, W. A. Wessel, and C. R. Hutchinson.** 1995. Malonyl-coenzyme A:acyl carrier protein acyltransferase of *Streptomyces glaucescens*: a possible link between fatty acid and polyketide biosynthesis. *Biochemistry* **34**:9389–9402.
 39. **Tabor, S.** 1995. Expression using the T7 RNA polymerase/promoter system, p. 16.2–16.9. In F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl (ed.), *Short protocols in molecular biology*, 3 ed. John Wiley & Sons, New York, N.Y.
 40. **Val, D. L., and J. E. Cronan.** 1998. In vivo evidence that *S*-adenosylmethionine and fatty acid intermediates are the substrates for the LuxI family of autoinducer synthases. *J. Bacteriol.* **180**:2644–2651.
 41. **Van Delden, C., and B. H. Iglewski.** 1998. Cell-to-cell signaling and *Pseudomonas aeruginosa* infections. *Emerg. Infect. Dis.* **4**:551–560.
 42. **Van Den Boom, T., and J. E. Cronan.** 1989. Genetics and regulation of bacterial lipid metabolism. *Annu. Rev. Microbiol.* **43**:317–343.
 43. **Verwoert, I. G. S., E. F. Verhagen, K. H. van der Linden, H. Verbree, J. J. Nijkamp, and A. R. Stuitje.** 1994. Molecular characterization of an *Escherichia coli* mutant with a temperature-sensitive malonyl coenzyme A:acyl carrier protein transacylase. *FEBS Lett.* **348**:311–316.
 44. **Wang, R. F., and S. R. Kushner.** 1991. Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. *Gene* **100**:195–199.