

## Nitrite and Nitrous Oxide Reductase Regulation by Nitrogen Oxides in *Rhodobacter sphaeroides* f. sp. *denitrificans* IL106

MONIQUE SABATY,\* CAROLE SCHWINTNER,† SANDRINE CAHORS, PIERRE RICHAUD,  
AND ANDRE VERMÉGLIO

Commissariat à l'Énergie Atomique/Cadarache DSV, DEVM, Laboratoire de Bioénergétique  
Cellulaire, 13108 St. Paul lez Durance Cedex, France

Received 16 February 1999/Accepted 19 July 1999

**We have cloned the *nap* locus encoding the periplasmic nitrate reductase in *Rhodobacter sphaeroides* f. sp. *denitrificans* IL106. A mutant with this enzyme deleted is unable to grow under denitrifying conditions. Biochemical analysis of this mutant shows that in contrast to the wild-type strain, the level of synthesis of the nitrite and N<sub>2</sub>O reductases is not increased by the addition of nitrate. Growth under denitrifying conditions and induction of N oxide reductase synthesis are both restored by the presence of a plasmid containing the genes encoding the nitrate reductase. This demonstrates that *R. sphaeroides* f. sp. *denitrificans* IL106 does not possess an efficient membrane-bound nitrate reductase and that nitrate is not the direct inducer for the nitrite and N<sub>2</sub>O reductases in this species. In contrast, we show that nitrite induces the synthesis of the nitrate reductase.**

Complete denitrification, i.e., reduction of nitrate into nitrous oxide or dinitrogen, is a bioenergetic process used by several species of bacteria. Four nitrogen oxide (N oxide) reductases (nitrate, nitrite, NO, and N<sub>2</sub>O reductases) are necessary to complete this reaction. The systems that regulate the synthesis of these enzymes are complex and vary from one denitrifier to another. In general, nitrate and N<sub>2</sub>O reductases are regulated independently with special regulators. In contrast, the regulation of nitrite and NO reductases is often linked at both the transcription and enzyme activity levels (45). Anaerobiosis and the presence of N oxides are the two essential factors that control the synthesis of the N oxide reductases (reviewed in reference 45).

Fumarate nitrate reductase and homologues are important elements of the denitrification regulation (45) and have been extensively studied in *Escherichia coli* (38). These *trans*-acting proteins activate, under anaerobic conditions, expression of operons such as *nar*. This operon encodes the membrane-bound nitrate reductase (5), an enzyme generally synthesized under anaerobic conditions. On the other hand, the expression of the periplasmic nitrate reductase, an enzyme first discovered in photosynthetic bacteria (21, 33, 35), is repressed during anaerobic growth for most of the denitrifiers (1, 42). This suggests a putative role for this enzyme in adaptation during a shift from aerobiosis to anaerobiosis (2, 36).

Anaerobic shift is sometimes not sufficient for the induction of denitrification enzymes (12, 17, 41). For example, the presence of N oxide is required, in addition to anaerobiosis, to induce the synthesis of the denitrification reductases in *Pseudomonas stutzeri* (17). More generally, nitrate acts as a good inducer for all N oxide reductases (17, 22), while nitrite, nitric oxide, and nitrous oxide at least induce their corresponding reductases (17, 18).

Regulations by nitrate and nitrite have been extensively

studied in *E. coli*. Two two-component regulatory systems, NarXL and NarQP, have been shown to regulate the membrane-bound and periplasmic nitrate reductases (9, 39). NarX and NarQ are two sensors that can phosphorylate the two regulators NarL and NarP (25). In *P. stutzeri*, *Paracoccus denitrificans*, and *Rhodobacter sphaeroides* 2.4.3, coregulation of the nitrite and nitric oxide reductases has been demonstrated by the observation that mutation in the nitrite reductase gene affects *norCB* transcription (41). Fumarate nitrate reductase-like factors DNR, NNR, NnrR, and FnrD, belonging to the FixK group (45) and generally flanking the *nor* region, modulate both *nirS* and *norCB* genes (1, 44, 45). For *R. sphaeroides* 2.4.3, Kwiatkowski et al. (19) showed that NnrR, which is also present in *R. sphaeroides* 2.4.1 (19), regulates *nirK* and *norCB* in response to the presence of NO (18, 19, 40).

The presence of N<sub>2</sub>O can also induce the synthesis of some reductases: nitrate and N<sub>2</sub>O reductases in *P. stutzeri* (17) and N<sub>2</sub>O reductase in *Rhodobacter capsulatus* MT1131 (28). The membrane-bound component, NosR, necessary for the expression of the nitrous oxide reductase in this strain, may be implicated in this regulation (8).

*R. sphaeroides* f. sp. *denitrificans* IL106 is one of the few purple, nonsulfur denitrifying photosynthetic bacteria able to perform a complete denitrification process (16, 23, 29, 32, 35). A property of this bacterium is that the periplasmic nitrate reductase is synthesized under both aerobic and anaerobic conditions (20, 30). In contrast to the consensus reached for the other denitrifying species, the presence of a membrane-bound nitrate reductase is controversial for *R. sphaeroides* f. sp. *denitrificans* IL106.

In this work, we created a mutant deficient in the periplasmic nitrate reductase for *R. sphaeroides* f. sp. *denitrificans* IL106. The aim was to determine the role of this enzyme and to study the effect of nitrate on N oxide reductase synthesis.

### MATERIALS AND METHODS

**Bacterial strains, plasmids, and growth conditions.** *R. sphaeroides* strains were grown at 30°C in Sistro minimal medium supplemented with succinate as the carbon source (7) under anaerobic conditions, in the absence or presence (phototrophic conditions) of light (75 mol of photons m<sup>-2</sup> s<sup>-1</sup>), or under aerobic conditions (100 ml of culture in 250-ml conical flasks, 275 rpm). Where indicated, the medium was supplemented with 20 mM KNO<sub>3</sub> or NaNO<sub>2</sub> or sparged with N<sub>2</sub>O (to saturation). *E. coli* strains were grown at 37°C in Luria-Bertani medium.

\* Corresponding author. Mailing address: CEA/Cadarache DSV, DEVM, Laboratoire de Bioénergétique Cellulaire, Bât. 156, 13108 St. Paul lez Durance Cedex, France. Phone: 33 4 42 25 35 70. Fax: 33 4 42 25 47 01. E-mail: msabaty@cea.fr.

† Present address: Laboratoire de Génétique Moléculaire de la Recombinaison, Service Recherche, Institut Curie, 26 rue d'Ulm, 75231 Paris Cedex, France.

When appropriate, tetracycline, spectinomycin, streptomycin, and kanamycin were added at concentrations of 1, 50, 50, and 25  $\mu\text{g/ml}$ , respectively, for *R. sphaeroides* and at concentrations of 20, 50, 50, and 25  $\mu\text{g/ml}$  for *E. coli*.

**Preparation of cell extracts for electrophoresis.** Preparation of cell extracts, nondenaturing gel electrophoresis, and activity staining were performed as previously described (31).

**Assays for nitrate reductase activity.** Cells were grown 48 h photosynthetically or under aerobic conditions, in the presence of N oxide, washed with 50 mM Tris-HCl, pH 8.2, and resuspended in the same buffer. They were broken up with a French press and centrifuged for 1 h 30 min at  $200,000 \times g$ . The nitrate reductase activity of the supernatant was measured as follows: the reaction mixture contained 0.85 ml of 50 mM Tris-HCl (pH 8.2), 2 mM methyl viologen, 20 mM  $\text{KNO}_3$ , 100  $\mu\text{l}$  of soluble fraction, and 50  $\mu\text{l}$  of 10 mg of Na dithionite per ml of 200 mM  $\text{NaHCO}_3$ . After 5 or 10 min at  $30^\circ\text{C}$ , the reaction was stopped by vigorous agitation until complete oxidation of methyl viologen. The nitrite formed was assayed by the diazo-coupling method (24).

**Mass spectrometry measurements.** Mass spectrometry measurements were performed as previously described (30).

**DNA manipulation and sequence analyses.** Isolation of plasmid DNA and restriction endonucleases and other enzymatic treatments of DNA were carried out according to standard protocols or manufacturers' instructions. Sequence determination was performed by Genome Express S.A. (Grenoble). Sequence analyses were performed with the BISANCE computer program (Infobiogen).

**Cosmid bank construction.** The *cos* vector SuperCos 1 and packaging extracts were obtained from Stratagene (La Jolla, Calif.). *R. sphaeroides* f. sp. *denitrificans* genomic DNA was partially digested with *Bam*HI. The digested DNA was sized to yield 40-kb fragments, dephosphorylated, and then ligated into the *Bam*HI site of SuperCos 1, previously linearized with *Xba*I. Packaging of the cosmids into phage heads and their subsequent infection in the *E. coli* strain XL1-Blue MR were performed as described by the manufacturer (SuperCos 1 and Gigapack II XL kits; Stratagene).

**Probes.** From the sequence of two nitrate reductase peptides (31) and taking into account the codon bias of known *R. sphaeroides* genes, two degenerate primers, GA(C/T)TGGA(G/T)GA(G/A)GC(G/C)TT(C/T)GA(C/T)GT and TC(G/A)AACCA(C/G)GG(C/G)AC(G/A)AA(C/G)AC(C/G)AC, were designed and used for PCR with *Taq* polymerase. The 1.9-kb product obtained was sequenced. From this sequence, a 104-bp oligonucleotide was synthesized and used as a probe for *napA* (Nitra104; nucleotides 2062 to 2165 from *napA*). Another oligonucleotide containing the first 52 nucleotides from *napA* was also synthesized and used as a probe (Nitra52; OligoExpress, Paris, France).

**Southern DNA analysis.** DNA was transferred to nylon Hybond N+ membranes (Amersham) with a TE 80 Transvac vacuum blotter (Hofer Scientific Instruments). The probes were labelled with digoxigenin (DIG)-dUTP (DIG high-prime or DIG oligonucleotide 3'-end tailing kits from Boehringer). Hybridizations and detection of hybridizing sequences by chemiluminescence with CDP-Star were performed according to the manufacturer's protocols (Boehringer).

**Nucleotide sequence accession numbers.** The nucleotide sequences of the *nap*, *nos*, and *nor* regions have been submitted to the GenBank and EMBL databases and were given accession no. AF069545, AF125260, and AF126490, respectively.

## RESULTS

**Cloning and sequencing of the periplasmic nitrate reductase gene of *R. sphaeroides* f. sp. *denitrificans* IL106.** In *R. sphaeroides* f. sp. *denitrificans* IL106, both a membrane-bound and a periplasmic nitrate reductase have been described and purified (6, 33). The evidence for two nitrate reductase activities is, however, controversial. Sawada and Satoh found 93% of nitrate reductase activity in the soluble fraction (33), whereas Byrne and Nicholas, using the same strain, detected 97% of the activity in the membrane fraction for cultures grown in the same conditions (6). Like Sawada and Satoh, we always observed a nitrate reductase activity in the periplasmic fraction of *R. sphaeroides* f. sp. *denitrificans* (31). On the other hand, we never detected any nitrate reductase activity in the membrane fraction that was not imputable to a contamination by the soluble fraction trapped in the chromatophores, even when we used the same procedure as Byrne and Nicholas. To determine the importance of the periplasmic enzyme in the denitrification pathway, we cloned, sequenced, and disrupted the *nap* locus.

*NapA* peptide sequences (31) provided the basis for the construction of one pair of degenerated oligonucleotides. From the sequence of the PCR product obtained, a 104-bp oligonucleotide was synthesized and used as a probe for *napA*.




FIG. 1. Growth curves of *R. sphaeroides* f. sp. *denitrificans* wild type (□), MS523 mutant complemented with plasmid pRK415 (●), and MS523 mutant complemented with plasmid pMS538 (▲) under denitrifying conditions in the presence of 50 mM nitrate. OD<sub>660nm</sub>, optical density at 660 nm.

A cosmid library of genomic DNA (SuperCos 1 and Gigapack II XL kits; Stratagene) was screened. A positive signal was obtained with the cosmid pCOSIXE11, which was subcloned. A 4.6-kb *Eco*RI fragment was cloned into pUC18 (plasmid pCS1). Comparison of the sequence obtained with the sequence of *R. sphaeroides* 2.4.1 *NapA* (26) showed that the 5'-terminal part of *napA* was missing on the 4.6-kb *Eco*RI fragment. A 2.8-kb *Kpn*I-*Eco*RI fragment was cloned into pBluescript (plasmid pMS578). The sequence of the *nap* locus of *R. sphaeroides* f. sp. *denitrificans* was obtained by sequencing the inserts of plasmids pCS1 and pMS578 (accession no. AF069545). This locus contains seven open reading frames, *napKEFDABC*, transcribed in the same direction. The entire *nap* operon has been cloned and sequenced in several bacteria (3, 11, 27). As expected, the closest similarities were found with the *R. sphaeroides* 2.4.1 locus (27), with 93, 98, 91, 89, 98, 95, and 98% of the amino acids identical for *NapK*, -E, -F, -D, -A, -B, and -C, respectively (data not shown).

**Disruption of *napA*: effects on growth.** A 4.6-kb *Eco*RI fragment from cosmid pCOSIXE11, containing the last 798 nucleotides of *napA* and the entire *napB* and *napC* genes, was cloned into pSUP202Km (37) to yield pMS503 (*NapC* is a tetraheme cytochrome acting as an electron donor to the nitrate reductase) (19). An omega cartridge encoding resistance to spectinomycin and streptomycin was then cloned into the *Sac*I site of *napA*. The resulting plasmid, pMS507, unable to replicate into *R. sphaeroides*, was moved from *E. coli* to *R. sphaeroides* f. sp. *denitrificans* by conjugation (10). The double crossover event was confirmed by Southern hybridization analysis. The resulting mutant, MS523, grew under both aerobic and phototrophic conditions but not under dark anaerobic conditions with nitrate as the electron acceptor (Fig. 1). When the plasmid pMS538 containing *napABC* (4.6-kb *Sma*I fragment cloned into pRK415 [14]) was introduced into the MS523 mutant, growth with nitrate was restored and the observed growth rate was even faster than it was for the wild type (Fig. 1). As expected, the mutant displays no nitrate reductase activity (Fig. 2), but the synthesis of the nitrate reductase was restored when the plasmid pMS538 was present in the MS523 strain (Fig. 3). The level of synthesis was high even in the absence of nitrate (compare Fig. 2, lane 1, and Fig. 3, lane 3).




FIG. 2. Nondenaturing electrophoresis of periplasmic extracts (50  $\mu$ g of protein) of *R. sphaeroides* f. sp. *denitrificans* wild type (WT) and MS523 mutant grown under phototrophic conditions in the presence (+) or absence (-) of 20 mM nitrate. Gels were stained for nitrate (A), nitrite (B), and  $N_2O$  (C) reductase activities with dithionite-reduced methyl viologen as the electron donor or with Coomassie R250 for protein detection (D).

This may be either because multicopies (four to six) of the *napAB* genes are present with pMS538, compared with the wild-type strain without plasmid, or because some regulatory sequences upstream of *napA* are missing in the plasmid construct (the insert contains only 173 bp upstream of the *napA* start codon). We deduced from this series of experiments that the periplasmic nitrate reductase of *R. sphaeroides* f. sp. *denitrificans* is necessary for growth under denitrifying conditions.

**Effects of *napA* disruption on enzyme synthesis.** To further characterize this MS523 strain, nitrite reductase and  $N_2O$  reductase activities of periplasmic extracts were analyzed, in addition to nitrate reductase activity, by nondenaturing polyacrylamide gel electrophoresis. In the wild type, the presence of nitrate in the medium strongly induced the synthesis of nitrate, nitrite, and  $N_2O$  reductases (Fig. 2). An unexpected result of the disruption of the *napA* gene was that the synthesis of the nitrite and  $N_2O$  reductases was no longer induced by the presence of nitrate in the medium (Fig. 2, lanes 8 and 12). These inductions were restored when the plasmid pMS538 was present in the MS523 strain (Fig. 3). These results show that nitrate itself is unable to induce the synthesis of nitrite and  $N_2O$  reductases under anaerobic conditions, contrary to previous claims (22, 31). The induction of the synthesis of the nitrite and  $N_2O$  reductases is observed only when both nitrate

and nitrate reductase are present. This suggests that the real inducer is one of the products of nitrate reduction, i.e., nitrite, NO, or  $N_2O$ . We investigated the effects of the presence of nitrite and  $N_2O$  on reductase synthesis to test this hypothesis. Cells were grown under photosynthetic conditions for 18 h, and different concentrations of nitrite or  $N_2O$  were then added. The cells were harvested 5 h after the addition of nitrite or  $N_2O$ , and cell extracts were prepared. The addition of nitrite or  $N_2O$  to the medium induced the synthesis of nitrate, nitrite, and  $N_2O$  reductases. The dependence on nitrite concentration of the nitrate and nitrite reductase activities (maximal at 0.1 mM) was different from that observed for the  $N_2O$  reductase synthesis, which peaked at 1 mM (Fig. 4). This experiment, however, does not enable us to differentiate between a direct effect of nitrite and an effect due to the NO or  $N_2O$  produced during nitrite reduction. Analysis of the induction of the synthesis of the different reductases in the presence of N oxides for a mutant of *R. sphaeroides* f. sp. *denitrificans* with nitrite reductase deleted will be necessary to clarify this point.

A partial answer can, however, be obtained by studying the induction of the nitrate reductase for *R. sphaeroides* 2.4.1 cells grown in the presence of nitrite. This strain possesses no nitrite reductase activity (15), and the induction effect cannot be attributed to the NO produced during the enzymatic reduction of nitrite. To avoid a chemical reduction of nitrite into NO, as suggested by Tosques et al. (41), we performed the experiment with aerobic cultures because NO reacts immediately with oxygen and thus cannot accumulate. As expected, in these cultures the  $^{15}NO$  concentration, measured with a mass spectrometer, was undetectable (lower than 0.2  $\mu$ M) for cells grown for 48 h in the presence of  $^{15}NO_2^-$  or  $^{15}NO_3^-$ . Under such conditions, the presence of 1 mM nitrite doubled the level of nitrate reductase, from 21 to 41.5 nmol of nitrite formed per min per mg of protein. It is deduced that nitrite is a good inducer of the nitrate reductase synthesis.

## DISCUSSION

The inability of *R. sphaeroides* f. sp. *denitrificans* to grow on nitrate in the absence of the periplasmic enzyme, plus the fact that we could detect no nitrate reductase activity in the membrane fraction, leads us to conclude that the membrane-bound nitrate reductase is absent in this strain. This conclusion is




FIG. 3. Nondenaturing electrophoresis of periplasmic extracts (50  $\mu$ g of protein) of the MS523 mutant containing pRK415 or pMS538 *in trans*. Cells were grown under phototrophic conditions in the presence (+) or absence (-) of 20 mM nitrate. Gels were stained for nitrate (A), nitrite (B), and  $N_2O$  (C) reductase activities with dithionite-reduced methyl viologen as the electron donor.




FIG. 4. Nondenaturing electrophoresis of periplasmic extracts (50  $\mu$ g of protein) of *R. sphaeroides* f. sp. *denitrificans* grown under phototrophic conditions. Added to 18-h grown cultures were 0, 0.1, or 1 mM  $\text{NaNO}_2$  or 20 mM  $\text{KNO}_3$  or  $\text{N}_2\text{O}$ . Cell extracts were prepared 5 h after addition of the N oxides. Gels were stained for nitrate (A), nitrite (B), and  $\text{N}_2\text{O}$  (C) reductase activities with dithionite-reduced methyl viologen as the electron donor.

further substantiated by the absence of hybridization between genomic DNA of *R. sphaeroides* f. sp. *denitrificans* and the *narG* probe from *E. coli* (data not shown).

This shows that the periplasmic nitrate reductase is essential for this strain to grow on nitrate. What is this essential role? Generation of a proton motive force (PMF) or simply reduction of nitrate into nitrite, with the following reductions of nitrite into NO, NO into  $\text{N}_2\text{O}$ , and  $\text{N}_2\text{O}$  into  $\text{N}_2$  being the major components of the PMF? Experiments with the related strain *R. sphaeroides* 2.4.1, which possesses the periplasmic reductase but not the nitrite reductase, have shown that the reduction of nitrate by the periplasmic enzyme generates a PMF most probably formed by the coupling of the NADH-dehydrogenase to the periplasmic reduction via the quinone pool (4). This PMF is, however, insufficient to allow growth of this bacterium with nitrate as the sole electron acceptor (15). We obtained the same result even with an *R. sphaeroides* 2.4.1 strain containing multicopies of the reductase gene of *R. sphaeroides* f. sp. *denitrificans* (on plasmid pMS538) (data not shown). Similar behavior has been observed by McEwan et al. (21) for *R. capsulatus* N22DNAR<sup>+</sup>. We conclude that the main role of the periplasmic nitrate reductase is only to reduce nitrate into nitrite. The reduction of nitrite into  $\text{N}_2$  will then produce a PMF allowing bacterial growth.

Although the regulation of the synthesis of enzymes involved in denitrification appears to be quite different from one denitrifier to another, this synthesis is always increased by the addition of nitrate. In general, nitrate is assumed to be the direct inducer. In this study, however, we showed that in *R. sphaeroides* f. sp. *denitrificans*, nitrate is not the effector molecule for the nitrite and  $\text{N}_2\text{O}$  reductase induction. The presence of nitrate no longer increases the level of synthesis of these enzymes in a mutant deficient in nitrate reductase activity (Fig. 2). This suggests that the real inducer for these enzymes is a product of nitrate reduction, i.e.,  $\text{N}_2\text{O}$ , NO, or nitrite.

In support of this hypothesis, we obtained experimental evidence that these compounds act as inducers of denitrifying enzymes.  $\text{N}_2\text{O}$  is able to induce the synthesis of nitrite and  $\text{N}_2\text{O}$  reductases under photosynthetic growth conditions to a small extent (Fig. 4). How  $\text{N}_2\text{O}$  is sensed in the cell remains unknown. It has been suggested for other denitrifiers that NosR might be involved (8). *nosR* is present upstream of *nosZ* (encoding the  $\text{N}_2\text{O}$  reductase) in *P. stutzeri* and *Rhizobium meliloti* (13, 43). We cloned and sequenced the *nos* locus (*nosZDFYL*)

in *R. sphaeroides* f. sp. *denitrificans* (34), but the sequence upstream of *nosZ* is missing (GenBank accession no. AF125260). The identical organization of this locus and the close similarity between the deduced protein sequences for *P. stutzeri* or *R. meliloti* and *R. sphaeroides* suggest that *nosR* may be present in this latter species also. This still has to be verified.

The presence of nitrite strongly increases the synthesis of nitrate, nitrite, and  $\text{N}_2\text{O}$  reductases (Fig. 4). However, we never obtained the same level of induction with nitrite as with nitrate, possibly owing to the accumulation of toxic concentrations of NO. We observed that adding nitrite in the millimolar range to growing cultures induced the production of NO. Such NO production was not induced by the addition of nitrate (data not shown). This observation can be readily explained given that the reduction of nitrite into NO is fast, compared with the reduction of NO into  $\text{N}_2\text{O}$  and the reduction of nitrate into nitrite. In other words, to have a good induction of the synthesis of the reductases, it is necessary to have an adequate nitrite concentration but also a concentration of NO reductase relative to the concentration of nitrite reductase such that there is no toxic accumulation of NO. These conditions are reached when nitrate is added to the inoculum.

In *R. sphaeroides* 2.4.3, the transcription of *nirK* (encoding nitrite reductase) was reported to be increased in the presence of nitrite (41). However, Tosques et al. propose that the effector molecule is not nitrite but NO produced enzymatically or chemically from nitrite (41). They showed that NO was able to activate the transcription of the genes encoding nitrite and NO reductases (18). The NO-sensitive regulator is NnrR (18, 19, 40). We sequenced part of the *nor* cluster in *R. sphaeroides* f. sp. *denitrificans* (GenBank accession no. AF126490). It presents high homology with *R. sphaeroides* 2.4.3 (data not shown). An *nnrR* homolog was found upstream of *norC*. This suggests that NO also may be an inducer of nitrite and NO reductases in *R. sphaeroides* f. sp. *denitrificans*. However, experiments with *R. sphaeroides* 2.4.1 show that in *Rhodobacter* species, nitrite is probably an effector molecule for nitrate reductase induction.

Is the inability of nitrate to induce the synthesis of nitrite and  $\text{N}_2\text{O}$  reductases a general feature of denitrifiers? In most of the studies concerning other denitrifiers, the enhancement of nitrite and  $\text{N}_2\text{O}$  reductase synthesis by the addition of nitrate has been observed with wild-type strains possessing a nitrate reductase activity. It is therefore possible that in these strains, like in *R. sphaeroides* f. sp. *denitrificans*, the real effector molecule is not nitrate but a product of nitrate reduction, i.e., nitrite, nitric oxide, or nitrous oxide. To obtain a definite answer, the experiments we have conducted with the MS523 mutant of *R. sphaeroides* f. sp. *denitrificans* will have to be performed with similar mutants of other denitrifying species.

#### REFERENCES

1. Arai, H., T. Kodama, and Y. Igarashi. 1997. Cascade regulation of the two CRP/FNR-related transcriptional regulators (ANR and DNR) and the denitrification enzymes in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **25**:1141–1148.
2. Bell, L. C., D. J. Richardson, and S. J. Ferguson. 1990. Periplasmic and membrane-bound nitrate reductases in *Thiosphaera pantotropha*: the periplasmic enzyme catalyses the first step in aerobic denitrification. *FEBS Lett.* **265**:85–87.
3. Berks, B. C., D. J. Richardson, A. Reilly, A. C. Willis, and S. J. Ferguson. 1995. The *napEADBC* gene cluster encoding the periplasmic nitrate reductase system of *Thiosphaera pantotropha*. *Biochem. J.* **309**:983–992.
4. Berks, B. C., S. J. Ferguson, J. W. B. Moir, and D. J. Richardson. 1995. Enzymes and associated electron transport systems that catalyse the respiratory reduction of nitrogen oxides and oxyanions. *Biochim. Biophys. Acta* **1232**:97–173.
5. Blasco, F., G. Giordano, M. Chippaux, and V. Bonnefoy. 1989. Nitrate reductase of *Escherichia coli*: completion of the nucleotide sequence of the *nar* operon and reassessment of the role of the  $\alpha$  and  $\beta$  subunits in iron

- binding and electron transfer. *Mol. Gen. Genet.* **218**:249–256.
6. **Byrne, M. D., and D. J. D. Nicholas.** 1987. A membrane dissimilatory nitrate reductase from *Rhodobacter sphaeroides* f. sp. *denitrificans*. *Biochim. Biophys. Acta* **915**:120–124.
  7. **Cohen-Bazire, G., W. R. Sistrom, and R. Y. Stanier.** 1956. Kinetic studies of pigment synthesis by non-sulfur purple bacteria. *J. Cell. Comp. Physiol.* **49**:25–68.
  8. **Cuypers, H., A. Viebrock-Sambale, and W. G. Zumft.** 1992. NosR, a membrane-bound regulatory component necessary for expression of nitrous oxide reductase in denitrifying *Pseudomonas stutzeri*. *J. Bacteriol.* **174**:5332–5339.
  9. **Darwin, A. J., and V. Stewart.** 1995. Nitrate and nitrite regulation of the Fnr-dependent *aeg-46.5* promoter of *Escherichia coli* K-12 is mediated by competition between homologous response regulators (NarL and NarP) for a common DNA-binding site. *J. Mol. Biol.* **251**:15–29.
  10. **Davis, J., T. J. Donohue, and S. Kaplan.** 1988. Construction, characterization, and complementation of a *puf* mutant of *Rhodobacter sphaeroides*. *J. Bacteriol.* **170**:320–329.
  11. **Fleischmann, R. D., M. D. Adams, O. White, R. A. Clayton, et al.** 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496–512.
  12. **Härtig, E., and W. G. Zumft.** 1999. Kinetics of *nirS* expression (cytochrome *cd<sub>1</sub>* nitrite reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to denitrification: evidence for a denitrification-specific nitrate- and nitrite-responsive regulatory system. *J. Bacteriol.* **181**:161–166.
  13. **Holloway, P., W. McCormick, R. J. Watson, and Y. K. Chan.** 1996. Identification and analysis of the dissimilatory nitrous oxide reduction genes *nosRZDFY* of *Rhizobium meliloti*. *J. Bacteriol.* **178**:1505–1514.
  14. **Keen, N. T., D. Tamaki, D. Kobayashi, and D. Trollinger.** 1988. Improved broad-host-range plasmids for DNA cloning in gram-negative bacteria. *Gene* **70**:191–197.
  15. **Kerber, N. L., and J. Cardenas.** 1982. Nitrate reductase from *Rhodospseudomonas sphaeroides*. *J. Bacteriol.* **150**:1091–1097.
  16. **Klemme, J. H., I. Chyla, and M. Preuss.** 1980. Dissimilatory nitrate reduction by strains of the facultative phototrophic bacterium *Rhodospseudomonas palustris*. *FEMS Microbiol. Lett.* **9**:137–140.
  17. **Körner, J. M., and W. G. Zumft.** 1989. Expression of denitrification enzymes in response to the dissolved oxygen level and respiratory substrate in continuous culture of *Pseudomonas stutzeri*. *Appl. Environ. Microbiol.* **55**:1670–1676.
  18. **Kwiatkowski, A. V., and J. P. Shapleigh.** 1996. Requirement of nitric oxide for induction of genes whose products are involved in nitric oxide metabolism in *Rhodobacter sphaeroides* 2.4.3. *J. Biol. Chem.* **271**:24382–24388.
  19. **Kwiatkowski, A. V., W. P. Laratta, A. Toffanin, and J. P. Shapleigh.** 1997. Analysis of the role of the *nnrR* gene product in the response of *Rhodobacter sphaeroides* 2.4.1 to exogenous nitric oxide. *J. Bacteriol.* **179**:5618–5620.
  20. **Martinez-Luque, M., M. Dobao, and F. Castillo.** 1991. Characterization of the assimilatory and dissimilatory nitrate-reducing systems in *Rhodobacter*: a comparative study. *FEMS Microbiol. Lett.* **83**:329–334.
  21. **McEwan, A. G., C. L. George, S. J. Fergusson, and J. B. Jackson.** 1982. A nitrate reductase activity in *Rhodospseudomonas capsulata* linked to electron transfer and generation of a membrane potential. *FEBS Lett.* **150**:277–280.
  22. **Michalski, W. P., and D. J. D. Nicholas.** 1984. The adaptation of *Rhodospseudomonas sphaeroides* forma sp. *denitrificans* for growth under denitrifying conditions. *J. Gen. Microbiol.* **130**:155–165.
  23. **Michalski, W. P., and D. J. D. Nicholas.** 1988. Identification of two denitrifying strains of *Rhodobacter sphaeroides*. *FEMS Microbiol. Lett.* **52**:239–244.
  24. **Nicholas, D. J. D., and A. Nason.** 1957. Determination of nitrate and nitrite. *Methods Enzymol.* **3**:981–984.
  25. **Rabin, R. S., and V. Stewart.** 1993. Dual response regulators (NarL and NarP) interact with dual sensors (NarX and NarQ) to control nitrate and nitrite-regulated gene expression in *Escherichia coli* K12. *J. Bacteriol.* **175**:3259–3268.
  26. **Reyes, F., M. D. Roldán, W. Klipp, F. Castillo, and C. Moreno-Vivián.** 1996. Isolation of periplasmic nitrate reductase genes from *Rhodobacter sphaeroides* DSM 158: functional and structural differences among prokaryotic nitrate reductases. *Mol. Microbiol.* **19**:1307–1318.
  27. **Reyes, F., M. Gavira, F. Castillo, and C. Moreno-Vivián.** 1998. Periplasmic nitrate-reducing system of the phototrophic bacterium *Rhodobacter sphaeroides* DSM 158: transcriptional and mutational analysis of the *napKEFD-ABC* gene cluster. *Biochem. J.* **331**:897–904.
  28. **Richardson, D. J., L. C. Bell, A. G. McEwan, J. B. Jackson, and S. J. Ferguson.** 1991. Cytochrome *c<sub>2</sub>* is essential for electron transfer to nitrous oxide reductase from physiological substrates in *Rhodobacter capsulatus* and can act as electron donor to the reductase in vitro. *Eur. J. Biochem.* **199**:677–683.
  29. **Richardson, D. J., L. C. Bell, J. W. B. Moir, and S. J. Ferguson.** 1994. A denitrifying strain of *Rhodobacter capsulatus*. *FEMS Microbiol. Lett.* **120**:323–328.
  30. **Sabaty, M., P. Gans, and A. Verméglio.** 1993. Inhibition of nitrate reduction by light and oxygen in *Rhodobacter sphaeroides* f. sp. *denitrificans*. *Arch. Microbiol.* **159**:153–159.
  31. **Sabaty, M., J. Gagnon, and A. Verméglio.** 1994. Induction by nitrate of cytoplasmic and periplasmic proteins in the photodenitrifier *Rhodobacter sphaeroides* forma sp. *denitrificans* under anaerobic or aerobic condition. *Arch. Microbiol.* **162**:335–343.
  32. **Satoh, T., Y. Hoshino, and H. Kitamura.** 1976. *Rhodospseudomonas sphaeroides* forma sp. *denitrificans*, a denitrifying strain as a subspecies of *Rhodospseudomonas sphaeroides*. *Arch. Microbiol.* **108**:265–269.
  33. **Sawada, E., and T. Satoh.** 1980. Periplasmic location of dissimilatory nitrate and nitrite reductase in a denitrifying phototrophic bacterium. *Plant Cell Physiol.* **21**:205–210.
  34. **Schwintner, C.** 1997. Ph.D. thesis. Etude moléculaire de la dénitrification chez *Rhodobacter sphaeroides* f. sp. *denitrificans*, rôle de deux protéines inconnues, Université d'Aix-Marseille II, Marseille, France.
  35. **Shioi, Y., M. Doi, H. Arata, and K. Takamiya.** 1988. A denitrifying activity in an aerobic photosynthetic bacterium, *Erythrobacter* sp. strain OCh 114. *Plant Cell Physiol.* **29**:861–865.
  36. **Siddiqui, R. A., U. Warnecke-Eberz, A. Hengsberger, B. Schneider, and B. Friedrich.** 1993. Structure and function of a periplasmic nitrate reductase in *Alcaligenes eutrophus* H16. *J. Bacteriol.* **175**:5867–5876.
  37. **Simon, R., U. Priefer, and A. Puhler.** 1983. A broad-host-range mobilization system for *in vivo* genetic engineering; transposon mutagenesis in gram-negative bacteria. *Bio/Technology* **1**:37–45.
  38. **Spiro, S., and J. R. Guest.** 1990. FNR and its role in oxygen-regulated gene expression in *Escherichia coli*. *FEMS Microbiol. Rev.* **75**:399–428.
  39. **Stewart, V.** 1983. Nitrate regulation of anaerobic respiratory gene expression in *Escherichia coli*. *Mol. Microbiol.* **9**:425–434.
  40. **Tosques, I. E., J. Shi, and J. P. Shapleigh.** 1996. Cloning and characterization of *nnrR*, whose product is required for the expression of proteins involved in nitric oxide metabolism in *Rhodobacter sphaeroides* 2.4.3. *J. Bacteriol.* **178**:4958–4964.
  41. **Tosques, I. E., A. V. Kwiatkowski, J. Shi, and J. P. Shapleigh.** 1997. Characterization and regulation of the gene encoding nitrite reductase in *Rhodobacter sphaeroides* 2.4.3. *J. Bacteriol.* **179**:1090–1095.
  42. **Warnecke-Eberz, U., and B. Friedrich.** 1993. Three nitrate reductase activities in *Alcaligenes eutrophus*. *Arch. Microbiol.* **159**:405–409.
  43. **Zumft, W. G., A. Viebrock-Sambale, and C. Braun.** 1990. Nitrous oxide reductase from denitrifying *Pseudomonas stutzeri*: genes for copper-processing and properties of the deduced products, including a new member of the family of ATP/GTP-binding proteins. *Eur. J. Biochem.* **192**:591–599.
  44. **Zumft, W. G., C. Braun, and H. Cuypers.** 1994. Nitric oxide reductase from *Pseudomonas stutzeri*: primary structure and gene organization of a novel bacterial cytochrome *bc* complex. *Eur. J. Biochem.* **219**:481–490.
  45. **Zumft, W. G.** 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* **61**:533–616.