

Sporadic Distribution of tRNA^{Arg}_{CCU} Introns among α -Purple Bacteria: Evidence for Horizontal Transmission and Transposition of a Group I Intron

BRUNO PAQUIN,[†] ANNETTE HEINFLING,[‡] AND DAVID A. SHUB*

Department of Biological Sciences and Center for Molecular Genetics,
University at Albany—SUNY, Albany, New York 12222

Received 8 June 1998/Accepted 17 November 1998

A group I intron interrupts the tRNA^{Arg}_{CCU} gene of the α -purple bacterium *Agrobacterium tumefaciens* (B. Reinhold-Hurek and D. A. Shub, *Nature* [London] 357:173–176, 1992). In this study, we assess the distribution of the corresponding intron among 12 additional species of α -purple bacteria. Of 10 newly identified tRNA^{Arg}_{CCU} genes, we found only two that contained an intron homologous to that of the *Agrobacterium* tRNA^{Arg}_{CCU} intron. This restricted and scattered distribution of the tRNA^{Arg}_{CCU} intron among α -purple bacteria is consistent with a recent origin and horizontal transmission. Primary and secondary structural similarities between tRNA^{Leu}_{UAA} introns found in strains of the cyanobacterium *Microcystis aeruginosa* (K. Rudi and K. S. Jacobsen, *FEMS Microbiol. Lett.* 156:293–298, 1997) and α -purple tRNA^{Arg}_{CCU} introns suggest that these introns share a more recent common ancestor than either does with other known cyanobacterial tRNA^{Leu}_{UAA} introns.

Group I introns are present in cellular and viral genes in eukaryotes and eubacteria (6, 15, 16). Although these introns interrupt a number of different protein-coding and RNA genes of eukaryotes and bacteriophages, the only insertion site observed so far in eubacteria is the sequence of the anticodon loop of tRNA genes. Eubacterial group I introns were first identified in the tRNA^{Leu}_{UAA} genes of five cyanobacterial species (13, 22), prompting speculation that these introns would be widely distributed among cyanobacteria. In conjunction with the observation that most plastid genomes also possessed a homologous intron, an ancient origin predating the endosymbiotic event that gave rise to plastids was proposed for the tRNA^{Leu}_{UAA} intron (13, 22). The phylogenetic distribution of tRNA^{Leu}_{UAA} introns among cyanobacteria was later determined, and it was shown that the distribution was not universal but, nevertheless, was consistent with an ancient origin (17). This conclusion was challenged by the discovery of tRNA^{Leu}_{UAA} introns in some strains of the cyanobacterium *Microcystis aeruginosa* which, however, seem to have originated independently of the previously characterized cyanobacterial tRNA^{Leu}_{UAA} introns through horizontal transfer (19). Conversely, tRNA fMet group I introns are sporadically distributed among cyanobacteria (17), a finding that corroborated the initial suggestion of a relatively recent origin of these introns during cyanobacterial evolution (3).

It has previously been demonstrated that two other eubacterial tRNA genes are interrupted by group I introns, the tRNA^{Ile}_{CAU} gene of *Azoarcus* sp. BH72 (a β -purple bacterium) and the tRNA^{Arg}_{CCU} gene of *Agrobacterium tumefaciens* A136 (an α -purple bacterium) (18). Initial data suggested that similar introns are likely to be widespread among proteobacteria,

as the *Azoarcus* and *Agrobacterium* intron probes cross-hybridized to genomic DNA from a number of purple bacteria (11, 18). Because determining their phylogenetic distribution is a valuable tool to assess the evolutionary history of group I introns (2, 17), we decided to survey the distribution of the tRNA^{Arg}_{CCU} intron among α -purple bacteria.

A tRNA^{Arg}_{CCU} intron in *Azospirillum halopraeferens*. It has been shown that a restriction fragment from the genomic DNA of the α -purple bacterium *A. halopraeferens* Au5 hybridized to an *Azoarcus* intron probe (18). The signal was also observed when the Southern hybridization was repeated with an *Agrobacterium* tRNA^{Arg}_{CCU} intron probe (11). The hybridizing region was cloned in pBSM13 as a 4.3-kb *Pst*I-*Sst*I restriction fragment (pAGAU1.1) and subsequently subcloned as a 400-bp *Ava*II fragment (pAGAU1.2). The latter insert was completely sequenced on both strands, revealing a tRNA^{Arg}_{CCU} gene interrupted by a potential group I intron inserted after the U of the CCU anticodon (the same position as the *Agrobacterium* intron). The 217-bp *Azospirillum* intron sequence folds into a bona fide group I secondary structure (Fig. 1) and shares 69% identity with the *Agrobacterium* intron (Table 1). In addition, like the *Agrobacterium* intron, it self-splices in vitro (11, 18).

PCR amplification of tRNA^{Arg}_{CCU} genes from various α -purple bacteria. Based upon the exonic tRNA sequences of *Agrobacterium* and *Azospirillum*, we designed degenerate primers for amplifying tRNA^{Arg}_{CCU} genes from various α -purple bacteria. We performed PCRs (17) on either the extracted DNAs or cell pellets from 11 species, with the primers ARG-5' (5'-GTCC[G/A/T]CGATGCTCA[G/A][C/T]A[A/G]GATA-3') and ARG-3' (5'-TGGTGTCC[G/A/C]C[G/T][G/A][G/C][G/A/T]GGA[A/T]TCGAACC-3'). These primers were expected to amplify a sequence that includes the entire anticodon stem and loop of the tRNA. Although PCR products of ~75 bp, the expected size of an uninterrupted tRNA gene, were amplified from all the samples, a PCR product of approximately 300 bp, the expected size of an intron-containing gene was also amplified from the DNA of *Anaplasma marginale* (Fig. 2). Cloning and sequencing of three independent clones confirmed the identity of this product as a tRNA^{Arg}_{CCU} gene interrupted by a group I intron inserted after the U of the CCU anticodon. The *Anaplasma* intron is similar both in

* Corresponding author. Mailing address: Department of Biological Sciences and Center for Molecular Genetics, University at Albany—SUNY, Albany, NY 12222. Phone: (518) 442-4324. Fax: (518) 442-4767. E-mail: shub@cnsunix.albany.edu.

[†] Present address: Département de Biochimie, Université de Montréal, Montréal, QC, H3C 3J7, Canada.

[‡] Present address: FG Microbial Ecology, Technical University Berlin, D-10587 Berlin, Germany.

TABLE 2. PCR results and distribution of the tRNA^{Arg}_{CCU} intron

Species	tRNA ^{Arg} genes amplified ^a					Total	Intron in tRNA ^{Arg} _{CCU}
	CCU	UCU	ACG	CCG			
<i>Agrobacterium tumefaciens</i> A136 ^b							+ ^d
<i>Anaplasma marginale</i>	3	0	0	3	6		+
<i>Azospirillum halopraeferens</i> Au5 ^b							+
<i>Bradyrhizobium japonicum</i>	1	2	4	0	7		-
<i>Ehrlichia risticii</i>	1	2	1	0	4		-
<i>Erythrobacter longus</i> OCh 119	0 (1)	0 (1)	6 (1)	4	13		-
<i>Methylobacterium extorquens</i>	1	2	7	0	10		-
<i>Paracoccus denitrificans</i>	1	1	0	2	4		-
<i>Rhizobium etli</i> CE3	0	3 (4)	0	7 (2)	17 ^c		?
<i>Rhodobacter sphaeroides</i> ATH 2.4.1	0 (2)	10	0	0	12		-
<i>Rhodocista centenaria</i>	1	0	0	0	1		-
<i>Rhodospirillum rubrum</i> S.1	2	0	0	0	2		-
<i>Rickettsia prowazekii</i>	0	3 (7)	7	0	18 ^c		- ^e

^a The identities of the amplified genes were determined by the anticodon sequence of their encoded tRNAs. PCR was performed with the pair of primers ARG-5' and ARG-3' or ARG-5'EXT and ARG-3' (numbers in parentheses). After these data were obtained we discovered that the nucleotides complementary to positions 8 and 9 of the *Agrobacterium* tRNA of both 5' PCR primers were transposed, resulting in potential mismatches. Both primers were resynthesized to be complementary at those positions, and PCR revealed no new products indicating the presence of an intron. Sequencing of 12 clones from these PCRs of *R. etli* DNA revealed the same anticodon species that had been determined previously. Two clones were obtained from PCR of *R. prowazekii* DNA; neither had an anticodon for Arg.

^b These species were not screened by PCR. The data were deduced from genomic clones.

^c A tRNA^{Lys}_{UUU} gene (*R. etli*) and a tRNA^{Asn}_{GCU} gene (*R. prowazekii*) were also amplified by PCR with the ARG-5'EXT and ARG-3' primers.

^d Data from reference 18.

^e The tRNA^{Arg}_{CCU} gene is lacking from the *R. prowazekii* genome (1).

nome of *R. prowazekii* has been completely sequenced and, as in some other bacterial genomes (e.g., *Mycoplasma genitalium* [10] and *Haemophilus influenzae* [9]), the tRNA^{Arg}_{CCU} gene is lacking (1). Assuming a G-U wobble, tRNA^{Arg}_{UCU} can substitute for tRNA^{Arg}_{CCU} in decoding AGG triplets. Therefore, it is possible that the tRNA^{Arg}_{CCU} gene is also missing from the genome of *R. etli*.

Distribution and evolution of the tRNA^{Arg}_{CCU} intron among α -purple bacteria. The 13 species we have selected for this study represent only a sampling of the diversity of α -purple bacteria. Nevertheless, the distribution of the tRNA^{Arg}_{CCU} intron is informative because of the phylogenetic relationship among the species we surveyed. We believe the distribution of tRNA^{Arg}_{CCU} introns is best explained by a recent origin and horizontal transmission, rather than by an ancient origin and differential loss in various α -purple lineages. Our reasoning is as follows. First, the three species possessing a tRNA^{Arg}_{CCU} intron, *A. tumefaciens*, *A. halopraeferens*, and *A. marginale*, are widely divergent α -purple bacteria (Fig. 4; see also the phylogenetic tree compiled by the Ribosomal Database Project [RDP] in reference 14). Second, *A. marginale* and *Ehrlichia risticii* branch within a monophyletic clade of closely related species, but only the *Anaplasma* tRNA^{Arg}_{CCU} gene contains an intron. The same is also true for *A. halopraeferens* (intron present) and *Rhodocista centenaria* (also known as *Rhodospirillum centenum*) (no intron).

Group I intron mobility is mediated by intron-encoded homing endonucleases or, alternatively, occurs via reverse splicing (for a review of group I intron mobility, see reference 21). Unlike reverse splicing, in which only the intron sequence is transferred, endonuclease-dependent mobility is accompanied by coconversion of flanking sequences. Interestingly, pairwise comparisons of the amplified regions of the tRNA genes shown in Fig. 3 reveal a striking similarity among the three intron-containing species (1, 2, or 3 differences, respectively, in 30 bp). On the other hand, with the exception of two identical pairs (*E. risticii*-*Bradyrhizobium japonicum* and *Methylobacterium extorquens*-*R. centenaria*) all of the other pairwise comparisons display 6 to 17 differences with a mean of 11 differ-

ences (note that, as expected, the 7-bp sequence defining the anticodon loop is virtually identical among all 11 species). Considering the fact that the three intron-containing species are not particularly closely related (Fig. 4), this observation is reminiscent of a recent endonuclease-dependent invasion. Although none of the introns described in this study contains an open reading frame (ORF), it would not be surprising if an ORF-containing intron is present in some other, unsurveyed species. For example, in a recent survey of cyanobacteria, tRNA-fMet genes of seven distantly related species were interrupted by highly similar introns. However, only one of these introns contained an endonuclease-encoding ORF (4, 17).

Recently, Rudi and Jacobsen showed that group I introns in tRNA^{Lcu}_{UAA} genes were sporadically distributed in strains of the cyanobacterium *M. aeruginosa*; six introns were found in 16 strains (19). Three of these introns were sequenced and shown to be almost identical (>99.5% identity) but only <61.5% identical to the previously described cyanobacterial tRNA^{Lcu}_{UAA} introns (19), whereas the latter are highly similar among themselves (Table 1; see also reference 17). Therefore, it was sug-




FIG. 2. PCR amplification of tRNA^{Arg} genes. The migration in a 1% agarose gel of the PCR amplification products from DNA or cell pellets of various α -purple bacteria is shown. Am, *A. marginale*; Bj, *B. japonicum*; Er, *E. risticii*; El, *Erythrobacter longus*; Me, *M. extorquens*; Pd, *Paracoccus denitrificans*; Re, *R. etli*; Rs, *Rhodobacter sphaeroides*; Rc, *R. centenaria*; Rr, *Rhodospirillum rubrum*; Rp, *R. prowazekii*; -, no DNA (negative control); M, 100-bp DNA marker (Gibco/BRL) (sizes are indicated on the right side of the gel). Some of the samples shown here are the results of a reamplification (Am, Bj, Me, and Pd). *R. etli* and *R. prowazekii* are the two species from which the tRNA^{Arg}_{CCU} genes could not be detected.




FIG. 3. Alignment of tRNA^{Arg}_{CCU} gene sequences. Base-pairing in the acceptor stem (A), anticodon stem (Ac), D stem (D), and T stem (T), the intron insertion site (arrow), the anticodon sequence (underline), and the position used to restrict amplification to tRNA^{Arg}_{CCU} genes (double underline) are indicated. N, any of the four possible nucleotides. I+ and I-, intron plus and minus, respectively. At, *A. tumefaciens*; Ah, *A. halopraeferens*; Am, *A. marginale*; Bj, *B. japonicum*; Er, *E. risticii*; El, *E. longus*; Me, *M. extorquens*; Pd, *P. denitrificans*; Rs, *R. sphaeroides*; Rc, *R. centenaria*; Rr, *R. rubrum*. Except for At and Ah, the tRNA^{Arg}_{CCU} genes were amplified by PCR. Sequences of the introns (shown in Fig. 1) and primers are omitted. The shorter sequences for *E. longus* and *R. sphaeroides* denote the use of the longer primer, ARG-5'EXT.

gested that the tRNA^{Leu}_{UAA} introns in cyanobacteria are polyphyletic and that the *Microcystis* introns originated independently through horizontal transfer (19). The *Microcystis* tRNA^{Leu}_{UAA} introns most likely share a more recent common ancestor with the α -purple tRNA^{Arg}_{CCU} introns than either does with the other cyanobacterial tRNA^{Leu}_{UAA} introns. This is supported by comparisons of primary sequences (Table 1; see also reference 19) and secondary structures (Fig. 1), as well as phylogenetic analyses (19 and data not shown). However, the origin of the tRNA^{Leu}_{UAA} introns, excluding the *Microcystis* introns, appears to be monophyletic (17). Taken together, these results suggest that the *Microcystis* introns originated from a tRNA^{Arg}_{CCU}-like intron by horizontal transfer. This represents an interesting case of intron transposition and horizontal transmission between widely divergent species (cyanobacteria and α -purple bacteria).

Nucleotide sequence accession numbers. The intron sequences reported here have been deposited in GenBank under

accession numbers AF081791 (*A. marginale*) and AF081792 (*A. halopraeferens*).

We thank David Edgell and an anonymous reviewer for helpful comments and critical readings of the manuscript, David Mulbauer and Markus Landthaler for assistance in cloning and sequencing some of the clones, and Siv Andersson for sharing unpublished data.

This work was supported by NIH grant GM37746. B.P. was a postdoctoral fellow of NSERC (Canada).

REFERENCES

- Andersson, S. G. E., A. Zomorodipour, J. O. Andersson, T. Sicheritz-Pontén, U. C. M. Alsmark, R. M. Podowski, A. K. Näslund, A.-S. Eriksson, H. H. Winkler, and C. G. Kurland. 1998. The genome sequence of *Rickettsia prowazekii* and the origin of mitochondria. *Nature (London)* **396**:133–140.
- Bhattacharya, D., B. Surek, M. Rüsing, S. Damberger, and M. Melkonian. 1994. Group I introns are inherited through common ancestry in the nuclear-encoded rRNA of Zygnematales (Charophyceae). *Proc. Natl. Acad. Sci. USA* **91**:9916–9920.
- Biniszkievicz, D., E. Cesnaviciene, and D. A. Shub. 1994. Self-splicing group I intron in cyanobacterial initiator methionine tRNA: evidence for lateral transfer of introns in bacteria. *EMBO J.* **13**:4629–4635.
- Bonocora, R. P., and D. A. Shub. Unpublished results.
- Cech, T. R., S. H. Damberger, and R. R. Gutell. 1994. Representation of the secondary structure of group I introns. *Nat. Struct. Biol.* **1**:273–280.
- Damberger, S. H., and R. R. Gutell. 1994. A comparative database of group I intron structures. *Nucleic Acids Res.* **22**:3508–3510.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**:783–791.
- Felsenstein, J. 1993. PHYLIP (Phylogeny Inference Package) version 3.5c. Department of Genetics, University of Washington, Seattle.
- Fleischmann, R. D., M. D. Adams, O. White, R. A. Clayton, E. F. Kirkness, A. R. Kerlavage, C. J. Bult, J.-F. Tomb, B. A. Dougherty, J. M. Merrick, K. McKenney, G. Sutton, W. FitzHugh, C. Fields, J. D. Gocayne, J. Scott, R. Shirely, L.-I. Liu, A. Glodek, J. M. Kelley, J. F. Weidman, C. A. Phillips, T. Spriggs, E. Hedblom, M. D. Cotton, T. R. Utterback, M. C. Hanna, D. T. Nguyen, D. M. Saudek, R. C. Brandon, L. D. Fine, J. L. Fritchman, J. L. Fuhrmann, N. S. M. Geoghagen, C. L. Gnehm, L. A. McDonald, K. V. Small, C. M. Fraser, and J. C. Venter. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**:496–512.
- Fraser, C. M., J. D. Gocayne, O. White, M. D. Adams, R. A. Clayton, R. D. Fleischmann, C. J. Bult, A. R. Kerlavage, G. Sutton, J. M. Kelley, J. L. Fritchman, J. F. Weidman, K. V. Small, M. Sandusky, J. Fuhrmann, D. Nguyen, T. R. Utterback, D. M. Saudek, C. A. Phillips, J. M. Merrick, J.-F. Tomb, B. A. Dougherty, K. F. Bott, P.-C. Hu, T. S. Lucier, S. N. Peterson, H. O. Smith, C. A. Hutchinson III, and J. C. Venter. 1995. The minimal gene complement of *Mycoplasma genitalium*. *Science* **270**:397–403.
- Heinfling, A., and D. A. Shub. Unpublished results.
- Kimura, M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **16**:111–120.
- Kuhse, M. G., R. Strickland, and J. D. Palmer. 1990. An ancient group I intron shared by eubacteria and chloroplasts. *Science* **250**:1570–1573.
- Maidak, B. L., G. J. Olsen, N. Larsen, R. Overbeek, M. J. McCaughey, and C. R. Woese. 1997. The RDP (Ribosomal Database Project). *Nucleic Acids Res.* **25**:109–111.




FIG. 4. rRNA phylogenetic tree. Prealigned small subunit rRNA sequences were extracted from the RDP web site (14) for the 11 species whose tRNA^{Arg}_{CCU} genes were detected. The tree was inferred with DNADIST (Kimura two-parameter model [12], with a transition/transversion ratio of 2) and NEIGHBOR (20) as implemented in PHYLIP (8). Bootstrap (7) values were deduced from 100 replicates (only those values higher than 50 are shown). Branches conserved between this tree and that published on the RDP web site (14) are indicated (*), and species harboring a tRNA^{Arg}_{CCU} intron are boxed. Note the scattered distribution of the three tRNA^{Arg}_{CCU} intron-containing species; a monophyletic clade grouping these three species is supported by a bootstrap value of <1%. The topology shown has been suggested by the branching position of an outgroup (γ -purple bacterium *Ectothiorhodospira shaposhnikovii*).

15. **Michel, F., and E. Westhoff.** 1990. Modelling of the three-dimensional architecture of group I catalytic introns based on comparative sequence analysis. *J. Mol. Biol.* **216**:585–610.
16. **Nishida, K., S. Suzuki, Y. Kimura, N. Nomura, M. Fujie, and T. Yamada.** 1998. Group I introns found in *Chlorella* viruses: biological implications. *Virology* **242**:319–326.
17. **Paquin, B., S. D. Kathe, S. A. Nierzwicki-Bauer, and D. A. Shub.** 1997. Origin and evolution of group I introns in cyanobacterial tRNA genes. *J. Bacteriol.* **179**:6798–6806.
18. **Reinhold-Hurek, B., and D. A. Shub.** 1992. Self-splicing introns in tRNA genes of widely divergent bacteria. *Nature (London)* **357**:173–176.
19. **Rudi, K., and K. S. Jacobsen.** 1997. Cyanobacterial tRNA^{Leu}_{UAA} group I introns have polyphyletic origin. *FEMS Microbiol. Lett.* **156**:293–298.
20. **Saitou, N., and M. Nei.** 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**:406–425.
21. **Saldanha, R., G. Mohr, M. Belfort, and A. M. Lambowitz.** 1993. Group I and group II introns. *FASEB J.* **7**:15–24.
22. **Xu, M.-Q., S. D. Kathe, H. Goodrich-Blair, S. A. Nierzwicki-Bauer, and D. A. Shub.** 1990. Bacterial origin of a chloroplast intron: conserved self-splicing group I introns in cyanobacteria. *Science* **250**:1566–1570.