

Gene Duplication and Multiplicity of Collagenases in *Clostridium histolyticum*

OSAMU MATSUSHITA,¹ CHANG-MIN JUNG,¹ SEIICHI KATAYAMA,¹
JUNZABURO MINAMI,¹ YUKIE TAKAHASHI,²
AND AKINOBU OKABE^{1*}

*Department of Microbiology, Faculty of Medicine, Kagawa Medical University, Kagawa 761-0793,¹
and Faculty of Pharmacy and Pharmaceutical Sciences, Fukuyama University,
Fukuyama, Hiroshima 729-0292,² Japan*

Received 30 June 1998/Accepted 16 November 1998

Clostridium histolyticum collagenase contains a number of different active components. Previously we have shown that *colH* encodes a 116-kDa collagenase (ColH) and a 98-kDa gelatinase. We purified a different 116-kDa collagenase (ColG) from the culture supernatant and sequenced its gene (*colG*). We also identified four other gelatinases (105, 82, 78, and 67 kDa) and determined their N-terminal amino acid sequences, all of which coincided with that of either ColG or ColH. Hybridization experiments showed that each gene is present in a single copy and each gene is transcribed into a single mRNA. These results suggest that all the gelatinases are produced from the respective full-length collagenase by the proteolytic removal of C-terminal fragments. The substrate specificities of the enzymes suggest that *colG* and *colH* encode class I and class II enzymes, respectively. Analysis of their DNA locations by pulsed-field gel electrophoresis and nucleotide sequencing of their surrounding regions revealed that the two genes are located in different sites on the chromosome. *C. histolyticum colG* is more similar to *C. perfringens colA* than to *colH* in terms of domain structure. Both *colG* and *colA* have a homologous gene, *mscL*, at their 3' ends. These results suggest that gene duplication and segment duplication have occurred in an ancestor cell common to *C. histolyticum* and *C. perfringens* and that further divergence of the parent gene produced *colG* and *colA*.

Clostridium histolyticum, a pathogenic clostridium causing myonecrosis, produces collagenase, a zinc metalloproteinase that degrades various types of collagen and gelatin. This enzyme has a broad substrate specificity and potent collagenolytic activity compared to vertebrate collagenases (38). Furthermore, *C. histolyticum* can grow well in simple media, producing fairly large amounts of enzyme in the culture medium. Because of these characteristics, this enzyme is widely used for biochemical and physiological studies, e.g., collagen depletion from vertebrate tissues (40, 46), cleavage of collagen linkers (3, 15), and therapeutic purposes (14, 41).

The biochemical and physicochemical properties of this collagenase have been extensively studied (34). Several lines of evidence show that it has multiple forms (7, 23, 26, 27, 33, 38, 48), with molecular masses ranging from 68 to 130 kDa and isoelectric points between pH 5.35 and 6.20 (34). On the basis of the ratio of the activities toward collagen and synthetic peptide substrates, the forms are divided into two classes (6). Amino acid analysis (6), peptide mapping (7), and circular dichroism spectroscopy (7) have revealed that there is extensive similarity between the enzymes within the same class and that there are distinct differences between the two classes in both their primary and secondary structures. This led to the prediction that the two classes are encoded by different genes and that one class evolved from the other by gene duplication followed by divergent evolution (7, 34). For the origin of the multiple enzymes in each class, the following three possibilities were postulated: they are encoded by different genes having extensive similarity, they result from different transcripts from

a single gene, or they are produced by proteolytic cleavage of a single precursor protein encoded by a single gene (34).

In a previous study, we have cloned and sequenced the *colH* gene, encoding a 116-kDa collagenase (ColH), and obtained evidence that a 98-kDa gelatinase is derived from the *colH* gene (49). Gelatin zymography showed the presence of two additional enzymes (78 and 67 kDa), which are highly gelatinolytic compared to the two ColH enzymes, in our ColH preparation (49). This observation led us to suspect that the two smaller gelatinases are encoded by a gene(s) other than *colH*. However, only one gene was detected when *C. histolyticum* DNA was analyzed by Southern hybridization with a *colH* probe. Furthermore, an 80-kDa recombinant ColH (rColH) protein lacking a C-terminal peptide retained activity toward water-soluble substrates (29). Thus, these gelatinases could be truncated forms of ColH. One direct approach to solving these questions is to identify the enzymes by their primary sequence and to clone and characterize their genes. This study was designed to show the relationships between all the gelatinolytic and collagenolytic enzymes present in the culture supernatant of *C. histolyticum*. We cloned a novel collagenase gene, determined its nucleotide sequence, and investigated locations of the collagenase genes. Based on these observations, we discuss a likely explanation for the multiplicity of *C. histolyticum* collagenases and discuss their molecular evolution and the structure-function relationship of these multidomain enzymes (29, 36).

MATERIALS AND METHODS

Bacterial strains, plasmids, and bacteriophage. *C. histolyticum* JCM 1403 (ATCC 19401) was obtained from the Institute of Physical and Chemical Research (Saitama, Japan) and used throughout this study. pT7Blue T vector and the host strain, *Escherichia coli* NovaBlue (Novagen, Madison, Wis.), were used for cloning DNA fragments obtained by PCR. The plasmid vector pUC19 (32)

* Corresponding author. Mailing address: Department of Microbiology, Faculty of Medicine, Kagawa Medical University, 1750-1 Ikenobe, Miki-cho, Kagawa 761-0793, Japan. Phone: 81 (87) 891-2129. Fax: 81 (87) 898-7109. E-mail: microbio@kms.ac.jp.

and the host strain *E. coli* DH5 α (4) were used for all other recombinant DNA experiments.

Media and culture conditions. *C. histolyticum* was precultured in cooked meat medium (Nissui, Tokyo, Japan) at 37°C for 12 h and grown in Warren and Gray's medium (44) at 37°C for the preparation of enzymes and total RNA. Transformed *E. coli* cells were selected on Luria-Bertani plates containing 20 g of LB broth base (Gibco, Paisley, United Kingdom) and 15 g of agar per liter, supplemented with 100 μ g of ampicillin per ml, 0.2 mM isopropyl- β -D-thiogalactopyranoside (IPTG; Wako Pure Chemical Industries, Ltd., Osaka, Japan), and 0.004% 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-Gal; Wako).

Assay for collagenase. Collagenase activity was determined with azocoll as described previously (31), 4-phenylazobenzoyloxycarbonyl-Pro-Leu-Gly-Pro-D-Arg (PZ-peptide; Sigma Chemical Co., St. Louis, Mo.) as described by Wünsch and Heidrich (47), or insoluble collagen as described previously (19). Protein concentrations were determined by using the bicinchoninic acid protein assay reagent (Pierce, Rockford, Ill.), with bovine serum albumin as a standard. All assays were carried out in triplicate.

Zymography. Collagen zymography was performed by the method of Birkedal-Hansen and Taylor (5), using acid soluble type I collagen (Sigma) as described previously (31). Zymography with gelatin (Wako) was carried out by the method of Wilson et al. (45). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) with a 7.5% polyacrylamide gel and staining with Coomassie brilliant blue R were performed as described by Laemmli (24).

Purification of the 78-kDa gelatinase and 116-kDa collagenase. *C. histolyticum* was grown in 10 ml of cooked meat medium at 37°C for 12 h. The preculture was diluted 100-fold with 500 ml of Warren and Gray's medium and incubated at 37°C for 16 h. Cells were removed by centrifugation at 12,000 \times g for 10 min at 4°C. Ammonium sulfate was added to 80% saturation, and the precipitate was collected by centrifugation at 12,000 \times g for 30 min at 4°C. The pellet was dissolved in 3.5 ml of 50 mM Tris-HCl buffer (pH 7.5), and then it (approximately 5 ml) was applied to a Sephacryl S-200 column (2.2 by 70 cm). Proteins were eluted with the same buffer at a flow rate of 20 ml/h. Enzyme activity was monitored by determination of azocoll-hydrolyzing activity. The fractions which were eluted in the first active peak were collected.

For purification of the 78-kDa gelatinase, a one-third volume (approximately 6.7 ml) of the pooled fraction was applied to an ion-exchange fast protein liquid chromatography column (MonoQ; bed volume, 1 ml; Pharmacia LKB Biotechnology, Uppsala, Sweden) which had been pre-equilibrated with 50 mM Tris-HCl (pH 7.5). Proteins were eluted with a 20-ml linear gradient of 0 to 0.5 M NaCl in 50 mM Tris-HCl (pH 7.5). The 78-kDa enzyme was eluted in a single peak at 192 mM NaCl. This step was repeated three times, and the active fractions were combined.

The 116-kDa ColG collagenase was purified in the same manner by ammonium sulfate precipitation, gel filtration, and ion-exchange chromatography. A one-fourth volume (5 ml) of the pooled collagenase fraction from the gel filtration column was applied to the ion-exchange fast protein liquid chromatography column. The collagenase was eluted in a single peak at 214 mM NaCl. This step was repeated four times, and the active fractions were combined.

Determination of the N-terminal amino acid sequence. Enzyme fractions were applied to an SDS-7.5% polyacrylamide gel. Electrophoresis was carried out at 100 V for 3.5 h so that the band moved to the middle of the gel. Proteins were electrophoretically transferred to a sheet of polyvinylidene difluoride membrane (Trans-Blot transfer medium; Bio-Rad Laboratories, Hercules, Calif.) and visualized by staining with Coomassie brilliant blue R as described by the supplier. The area corresponding to the band was cut out and subjected to N-terminal amino acid sequence analysis on a protein sequencer (model 492; Perkin-Elmer, Foster City, Calif.).

Amino acid sequence similarity search. A search for similar protein sequences was carried out by using the Blastp World Wide Web server (2) in the DNA Data Bank of Japan at the Center for Information Biology, National Institute of Genetics (Mishima, Japan). The N-terminal amino acid sequence determined for the 78-kDa ColG gelatinase was used as the query sequence, and this was searched against the nonredundant protein database, including SWISS-PROT, PIR, GenPept, and GenPept updates of the databases. Default parameters of the program were used for this search. A similar search using the amino acid sequences deduced from *colG* and the open reading frames (ORFs) found around the two collagenase genes was carried out in the same way.

DNA manipulations. Restriction endonucleases were purchased from Takara Shuzo Co. (Kyoto, Japan), Toyobo (Osaka, Japan), and New England Biolabs (Beverly, Mass.). The DNA ligation kit was a product of Takara Shuzo. All recombinant DNA procedures were carried out as described by Maniatis et al. (28).

Cloning of a portion of *colG*. A pair of degenerate oligonucleotide primers were designed to amplify a portion of *colG* by PCR. Their sequences, 5'-GA(A/G)AA(A/G)TA(C/T)GA(C/T)TT(C/T)GA(A/G)TA-3' and 5'-TG(A/G)TTC CA(C/T)TT(A/G/T)AT(A/G)TT(C/T)TT-3', correspond to the N-terminal amino acid sequence (Glu7 to Gln33) of the ColG enzymes. PCR was performed in a 100- μ l mixture containing a 1 μ M concentration of each primer, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 0.01% gelatin, 2.5 mM MgCl₂, a 0.2 mM concentration of each deoxynucleotide triphosphate, and 2.5 U of *Taq* DNA polymerase (Takara) by using a thermal cycler (model TC1; Perkin-Elmer). *C. histolyticum* DNA was prepared as described previously (49) and used as the

template. After a denaturation step at 94°C for 5 min, PCR (30 cycles) was carried out as follows: 94°C for 30 s, 45°C for 30 s, and 72°C for 30 s. The PCR product was purified from an acrylamide gel and cloned into pTTBlue T vector (Novagen). The resulting plasmid was designated pCHG1.

Construction of partial genomic libraries and their screening by PCR. The insert DNA in pCHG1 was amplified by PCR and isolated by PAGE. The fragment was labelled with digoxigenin-11-dUTP (Boehringer Mannheim Biochemicals, Indianapolis, Ind.) by PCR as described by Lanzillo (25) and used as a probe. Southern hybridization was carried out at 60°C. DNA fragments around the positive signal were recovered from an agarose gel and ligated into the *Hind*III site of pUC19. *E. coli* DH5 α was transformed with the ligation mixture, and colonies were subjected to PCR screening as follows. Fifteen colonies were suspended in one tube containing 20 μ l of distilled water, and 1 μ l of the mixed suspension was added to 19 μ l of the PCR mixture described above. After PCR under the above-described conditions, the products were examined by PAGE. To isolate a positive clone, the 15 clones in a positive group were examined separately by the same PCR screen.

The downstream *colG* fragments were obtained in the same way. A partial *Dra*I library was screened by PCR with a pair of synthetic oligonucleotides, 5'-TGCCTTGGTATGGAAAAATTGA-3' and 5'-TTGGCAGATAATGTTTT TCAGC-3', to clone a middle portion of *colG* (pCHG3). Another pair of synthetic oligonucleotides, 5'-GGATATTTGGCTAAGATAA-3' and 5'-GTGTT TGTAAGAGAAGCAGC-3', were used to screen a partial *Eco*R1 library to clone a 3'-terminal portion of *colG* (pCHG5).

Nucleotide sequencing of *colG*. The nucleotide sequence was determined by the dideoxy chain termination method (39), using an automated nucleotide sequencer (model ABI PRISM 377; Perkin-Elmer). Plasmid template DNA was prepared with the Wizard plus miniprep DNA purification system (Promega, Madison, Wis.). A Thermo Sequenase fluorescently labelled primer cycle sequencing kit with 7-deaza-dGTP (Amersham International plc, Little Chalfont, Buckinghamshire, England) and M13 dye primers (Perkin-Elmer) was used for sequencing. The ABI PRISM dye terminator cycle sequencing ready reaction kit with AmpliTaq DNA polymerase, FS (Perkin-Elmer), and various synthetic primers were also used to determine ambiguous nucleotides and to fill in sequence gaps.

Southern hybridization. Nucleotide fragments corresponding to the N-terminal amino acid sequences of the two collagenases (Ile1 to Tyr40 for the ColG enzymes and Val1 to Tyr39 for the ColH enzymes [49]) were prepared by PCR with two pairs of primers (5'-ATAGCGAATACTAATTCTGA-3' plus 5'-ATA ATTAATAAACCATTA-3' and 5'-GTACAAAATGAAAGTAAGAG-3' plus 5'-ATACTGAAAAGGCTCTGTA-3') and cloned into pTTBlue T vector (Novagen). After their nucleotide sequences were confirmed, they were amplified with the same primers, isolated by PAGE, and labelled with digoxigenin by PCR as described above. *C. histolyticum* DNA (0.5 μ g) was digested with the appropriate restriction enzymes, applied to a 0.8% agarose minigel, and electrophoresed at 100 V for 1 h. Treatment of the gel and transfer of DNA onto a sheet of nylon membrane (Hybond-N; Amersham) were performed as described previously (31). After hybridization at 50°C, the hybridized probes were detected with anti-digoxigenin-alkaline phosphatase Fab fragments (Boehringer) and a chemiluminescent dye (Lumi-Phos 530; Lumigen, Detroit, Mich.).

Northern hybridization. *C. histolyticum* cells were grown in Warren and Gray's medium at 37°C until the culture reached an optical density at 600 nm of 1.8 or 2.7. Total RNA was prepared by the SDS-phenol method (1), and the sample (3 μ g) was separated on a denaturing agarose gel (1%). Hybridization was carried out at 50°C with the same probes as described above.

Pulsed-field gel electrophoresis and Southern hybridization. Genomic DNA of *C. histolyticum* was prepared and digested by the method of Canard and Cole (9). Endonuclease I-CeuI was purchased from New England Biolabs. After the plugs were digested for 6 h at the appropriate temperature, the fragments were separated by contour-clamped homogeneous electric field electrophoresis with a CHEF-DR III apparatus (Bio-Rad). Electrophoresis was carried out with a 1% agarose gel at 5 V/cm with a ramping pulse from 5 to 100 s for 22 h. Gels were calibrated with *Saccharomyces cerevisiae* chromosomes (size range, 225 to 1,900 kb) and a mixture of λ concatemers and *Hind*III fragments (New England Biolabs). To determine the sizes of the larger fragments (above 1 Mb), electrophoresis was carried out under the following conditions: agarose gel, 0.8%; electric field, 3.7 V/cm; ramping pulse, 120 to 500 s; duration, 36 h; and size marker, *Hansenula wingei* chromosomes (size range, 1.05 to 3.13 Mb; Bio-Rad).

Ribosomal DNA probes were used to show the general arrangement of the *C. histolyticum* chromosome. Three *C. histolyticum* *rIB* fragments (a 1.6-kb *Hind*III-I-CeuI fragment containing the upstream *rIB* region, a 1.1-kb I-CeuI-*Hind*III fragment containing the downstream *rIB* region, and a 2.7-kb *Hind*III fragment containing both regions) were isolated by agarose gel electrophoresis, labelled with digoxigenin, and used for hybridization. The *rns* and *rfl* probes were prepared from the clones isolated from *Borrelia burgdorferi* (12). The *colG* and *colH* probes were also used for hybridization.

Nucleotide sequencing of regions adjacent to *colG* and *colH*. The nucleotide sequences of the regions adjacent to *colG* were determined as described above. The following plasmids, derived from a phage clone, *col18* (49), were used for nucleotide sequencing of the regions adjacent to *colH*: pCHC16 carrying a 2.5-kb *Sau*3AI-*Sac*I fragment, pCHC111 carrying a 1.4-kb *Sac*I-*Xba*I fragment, pCHC113 carrying a 1.6-kb *Xba*I fragment, pCHC1145 carrying a 0.5-kb *Xba*I-




FIG. 1. Analysis of purified ColG enzymes by SDS-PAGE and zymography. (A) SDS-PAGE. Lanes: 1, molecular mass markers (sizes are indicated on the left); 2, 78-kDa ColG gelatinase (2 μ g); 3, 116-kDa ColG collagenase (2 μ g). (B) Collagen zymography. Five micrograms of a 116-kDa ColG collagenase sample was electrophoresed on an SDS-polyacrylamide gel, followed by zymography with collagen fibrils (type I). (C) Gelatin zymography. Ten nanograms of the same sample as in panel B was electrophoresed on a gelatin-impregnated SDS-polyacrylamide gel. The gel was stained with Coomassie brilliant blue R to detect gelatinolytic activity as an unstained area.

*Hind*III fragment, pCHC1146 carrying a 2.0-kb *Hind*III fragment, pCHC1147 carrying a 2.3-kb *Hind*III fragment, and pCHC116 carrying a 2.9-kb *Pst*I-*Sau*3AI fragment (49). Sequencing templates were prepared by nested deletion of these plasmids, and their nucleotide sequences were determined with an ABI PRISM BigDye terminator cycle sequencing ready reaction kit with AmpliTaq DNA polymerase, FS (Perkin-Elmer).

Statistical analyses of the deduced amino acid sequence. The presence and location of signal peptide cleavage sites in possible prepropeptides of ColA and ColG were predicted by using the SignalP server (35) at the Center of Biological Sequence Analysis, Department of Biotechnology, The Technical University of Denmark (Lyngby, Denmark). Sequence alignment was carried out with the Blast server or the Clustal W program (43). The statistical significance of the sequence similarity was examined by a Monte Carlo test by using the Lipman-Pearson alignment algorithm with the Dayhoff similarity scoring (37). The best alignment of two original sequences was compared with a control obtained from the best alignments between 1,000 different randomized sequences by using the rfd2 program at the National Institute of Genetics. Default parameters were used except for the k_{rup} value, which was set at 1.

Nucleotide sequence accession numbers. The DNA sequences of *C. histolyticum* *colG* and *colH* are available from the GenBank/EMBL/DBJ databases (accession no. D87215 and AB014075, respectively). The sequences of a part of *C. histolyticum* 23S rRNA, i.e., *r1A*, *r1B*, and *r1C*, are available from the databases (accession no. AB013089, AB013090, and AB013091, respectively).

RESULTS

N-terminal sequence of a 78-kDa gelatinase. ColH collagenase (49) contained two minor gelatinases besides the 116-kDa ColH collagenase and the 98-kDa ColH gelatinase. One of these gelatinases (78 kDa) was purified to homogeneity (Fig. 1A), and its N-terminal amino acid sequence was determined. The sequence, Ile-Ala-Asn-Thr-Asn-Ser-Glu-Lys-Tyr-Asp-Phe-Glu-Tyr-Leu-Asn-Gly-Leu-Ser-Tyr-Thr-Glu-Leu-Thr-Asn-Leu-Ile-Lys-Asn-Ile-Lys-Trp-Asn-Gln-Ile-Asn-Gly-Leu-Phe-Asn-Tyr, was different from that of the ColH enzymes. A Blastp similarity search showed that it is similar (Poisson possibility value, 0.00018) to the N-terminal sequence of the *Clostridium perfringens* collagenase (ColA) (31). This enzyme was designated the 78-kDa ColG gelatinase.

Purification of the *C. histolyticum* collagenase. The ColG collagenase was purified by ammonium sulfate precipitation, gel filtration, and ion-exchange chromatography. The azocoll-hydrolyzing activity was eluted in two peaks near the void volume of the size exclusion column. Fractions in the first peak were combined, and the proteins were separated by ion-exchange chromatography. The activity which was eluted at 214

mM NaCl differed from that of ColH, which was eluted at 100 mM NaCl. The enzyme was purified to near homogeneity (Fig. 1A), with a recovery of 0.53 mg of protein from a 500-ml culture. Its apparent molecular mass was estimated to be 116 kDa by SDS-PAGE, and its hydrolytic activity against insoluble collagen fibrils was shown by collagen zymography (Fig. 1B). Gelatin zymography also showed that it possesses gelatinolytic activity (Fig. 1C). The specific activities of the enzyme against insoluble collagen and PZ-peptide were (means \pm standard deviations) 826 ± 42 and 26.0 ± 2.3 U/mg of protein, respectively. The sequence of its N-terminal 40 amino acid residues was determined to be identical to that of the 78-kDa gelatinase. This enzyme was designated the 116-kDa ColG collagenase.

N-terminal sequencing of various gelatinases. To find other enzymes, gelatin zymography was carried out with the enzyme fractions obtained as described previously (49). There were three bands exhibiting gelatinolytic activity besides the ColG and ColH enzymes. These proteins were separated by SDS-PAGE and transferred onto a polyvinylidene difluoride membrane, and their N-terminal amino acid sequences were determined. Two enzymes (82 and 67 kDa) had the same sequence as ColG, while one (105 kDa) had the same sequence as ColH.

Cloning of the *colG* gene. To amplify a portion of the *colG* gene, a pair of degenerate PCR primers were designed based on the N-terminal amino acid sequence of *colG*. They yielded a single amplification product of the expected size (approximately 80 bp). The fragment was cloned into a T vector, and the nucleotide sequence was determined by using 12 independent clones. The sequences of the 80-bp insert coincided with the N-terminal amino acid sequence of the ColG enzymes except for some variation due to the degeneracy of the primers. One of the plasmids was designated pCHG1. The insert fragment was labelled with digoxigenin and used as a probe for Southern hybridization to detect a single 2.4-kb *Hind*III fragment (data not shown). A partial genomic library was constructed by using 2.2- to 2.7-kb *Hind*III fragments, and 600 recombinant colonies were subjected to screening by PCR with the same degenerate primers. Five clones showed a positive signal, and all but one contained the same 2.4-kb *Hind*III insert. One of these plasmids was chosen and designated pCHG2. Nucleotide sequencing of this plasmid showed that it carries the 5' portion of *colG* (nucleotide positions 1 to 2376 [Fig. 2]).

A 1.6-kb *Dra*I fragment was detected by another round of Southern hybridization with a probe made from a 0.49-kb *Eco*RI-*Hind*III fragment (nucleotide positions 1882 to 2371) (data not shown). A partial genomic library carrying 1.2- to 2.0-kb *Dra*I fragments was constructed and screened by PCR. One positive clone (pCHG3) was obtained from 250 clones; it carries a 1.6-kb *Dra*I fragment (nucleotide positions 1625 to 3203). A fragment of *colG* further downstream was cloned in the same way by using a partial library carrying 3.0- to 4.0-kb *Eco*RI fragments. Four positive clones were obtained from 250 clones by PCR screening, all having the same 3.2-kb *Eco*RI fragment (nucleotide positions 2676 to 5914), designated pCHG5.

Nucleotide sequence of the *colG* gene. The nucleotide sequences of the cloned inserts in pCHG2, pCHG3, and pCHG5 were determined by using various subclones constructed from them. The sequences were aligned by their overlaps to form a single contig (Fig. 2). The sequence which coincides with the N-terminal amino acid sequence of the ColG enzymes was found within an ORF starting at nucleotide position 1002 and ending at 4358. This ORF encodes a protein with a possible prepropeptide of 110 amino acid residues which when removed would give a mature protein of 113,897 Da. This is in




FIG. 2. Nucleotide sequence of the *colG* gene and deduced amino acid sequence. The amino acid sequence determined for the N terminus of the 116-kDa ColG collagenase and the 78-kDa gelatinase is indicated by a thick underline. The potential ribosome binding site is indicated by a thin underline. The putative transcriptional terminator sequence is indicated by arrows. Adjacent regions are also shown. Restriction sites are indicated as follows: Dra, *DraI*; RI, *EcoRI*; and Hd, *HindIII*.

good agreement with the molecular mass determined by SDS-PAGE (116 kDa) for the purified ColG collagenase. There exist three possible ATG initiation codons beginning at nucleotide positions 1002, 1029, and 1110. Since only the first ATG codon is preceded by a possible ribosome binding sequence (AGGGGG) with a 7-bp gap, this could be the translational initiation site. A stem-loop sequence (nucleotide positions 4370 to 4405) with a short run of T's is present downstream of the termination codon; this could be a transcription terminator.

Deduced amino acid sequence of ColG. A database search with the Blastp server revealed that the deduced amino acid sequence of mature ColG aligns well with those of *C. histolyticum* ColH and *C. perfringens* ColA. The latter showed higher similarity (Poisson *P* value, 8.0e-249) than the former (Poisson

P value, 1.8e-215). When the possible prepropeptide of ColG (110 amino acid residues) was included in the query sequence, the *P* value became 1.2e-255 due to the significant alignment of the relevant sequences (Fig. 3A).

The signal peptides of ColA and ColG were predicted by the SignalP server to be a peptide extending from the N terminus to Ala39 (maximum C score, 0.734 at Ala40) and a peptide extending from the N terminus to Ala45 (maximum C score, 0.371 at Lys46), respectively. Although the score of the latter was low compared to that of the former, these predicted sites aligned well (Fig. 3A). This reinforces the proposed translation initiation site and the presence of a long prepropeptide in the ColG precursor. In a previous study we suggested that the PLGP sequence located near the C terminus of the ColA prepropeptide would play a role in postsignal cleavage matu-

AGGGACTTCTACACTTATGAGAGAACCCTCAACAAAGTATATTTAGTCTTGAAGAATGTGTTAGACATGAATATACTCACTATTTACAGCGAGAT 2600
 GTFYTYERTPQQSIFSLLEELFRHEYTHYLQARY
 CTTGTAGATGGTTATGGGGCAAGTCCATTTTGAIAAAATAGATTAACCTGGTTGATGAAGGTACAGCTGAATTCITTTGCAGGATCTACCCGT 2700
 LVDGLWGGQGPFFEKRNRLTWFDEGTAEFFAGSTR
 CATCTGGTGTITACCAAGAAATCAATATAGGATATTGGCTAAGGAATAAGTAGATCATAGATACTCATTAAAGAAGACTCTTAATTCAGGGATGA 2800
 SGVLPKRKSLILGYLAKDKVDHRYSLKKTLSNGYD
 TGACAGTGAITGGAGTTCATAATTATGGATTGTCAGTTGCACATTACCTATATGAAIAAGATAGCCTACATTTATTAAGATGAATAAGCTATATTG 2900
 DSDWMFYNFYAVAHYLYEKDMPFTFIKMGKAIL
 AATACAGATGTAAATCTTATGAAATATAAAAAATTAAGTATGATGCAAAATAAATAACAGAAATCAAAACCATATCAAGAGTTAGCAGATA 3000
 NTDVKSYDEIILKLSDDANKNTEYQNHIOELADK
 AATACAGAGGAGCCGATACCTCTAGTATCAGATGATTAATAAGATCATGGATATAAGAAAGCATCTGAAGTATATCTGAAATTCAAAAGCTGC 3100
 YQGA GIPLVSDDYLDKDHGYKKA SEVYSEISKAA
 TTCTCTTACAACTAGTGTAAACAGCAAAAATCTCAATATTTAACACATTCCTTTAAGAGAACTTATACAGGTGAAACTCTAAAGGTGAATTT 3200
 SLTNTS VTA EKSDYFNTFTLRNTYTG ETSKGEF
 AAGATTGGGATGAAATGAGTAAAAATAGATGGAATTTGGAGTCCCTTGCTAAAAATCTGGAGTGGATACAAAACTTAAACAGCATACTTTACG 3300
 KDWD E M S K K L D G T L E S L A K N S W S G Y K T L T A Y F T N
 ATTATAGAGITACAAGCGATAATAAAGTCAATATGATGATGTTTCCATGGGTTTTAAGCAGATAATGGGATATTAGTAAACAATAAGGCTCCAATAGC 3400
 YRVDKSDNKEVQYDVFVHGVLTDNADISNNKAPIA
 AAAGTAACTGGACCAAGCTGGTGTGAGGAAGAAATTAATGAAATTTAGTGGAAAGATAGTAAAGATGAAGATGGTAAATAGTATCATATGATGG 3500
 KVTG P ST G AV G R N I E F S G K D S K D E D G K I V S Y D W
 GATTTGGCAGTGGTCAACTAGTAGAGGCAAAAATCAGTACATGCTTCAAAAAAGCAGGAACATATATGTTACATTAAGTAACTGACGATAAGG 3600
 DFGD G A T S R G K N S V H A Y K K A G T Y N V T L K V T D D K G
 GTGCAACAGCTACAGAAAGCTTACTATAGAAATAAGAAACGAAGATACAAACACCTATACTAAAGAAATGGAACCTAATGATGATAAAGAGGC 3700
 ATAT E I K N E D T T T P I T K E M E P N D I K E A
 TAATGGTCCATAGTGAAGGTGTTACTGTAAAAGGTGATTAATGGTCTGATGATGCTGATACCTTCTATTTGATGTAAAAGAAGTGGTATGTT 3800
 N G P I V E G V T V K G D L N G S D D A D T F Y F D V K E D G D V
 ACAATGAACTTCCCTATTTCAGGGTCATCTAATTTTCACATGGTGTAGTTATAAAGAGGGAGACGATCAAAACCATATTGCAAGTGGTATAGATAAGAATA 3900
 T I E L P Y S G S S N F T W L V Y K E G D D Q N H I A S M N K A I L N
 ACTCAAAAGTGGAACTTTAAATCTACAAAGGAAGACATTATGTGTTATATATAAAACAGATTCTGCTTCAATATATCTCTTCTTAAACATAAA 4000
 S K V G T F K S T K G R H Y V F I Y K H D S A S N I S Y S L N I K
 AGGATTAGGTAAACGAGAAATTAAGGAAAGAAATAATGATCTTCTGATAAAGCTACAGTTATACCAATTTCAATACCCTATGCAAGGTTCACTT 4100
 G L G N E K L K E K E N D S D K A T V I P N F N T M Q K A I L N
 TTAGTGAATCAAGAGATTATTTCTTTGAGGTTAAGGAAGAGGCGAAGTTAATAGAACTAGATAAAAGGATGAATTTGGTGAACATGGA 4200
 L G D D S R D Y S F E V K E E G E V N I E L D K K D E F G V T W T
 CACTACATCCAGAGTCAAATTAATGACAGAATAACTACGGACAAGTTGATGGTAAATAGGTATCTAATAAAGTAAATTAAGACCAAGGAAATATT 4300
 L H P E S N I N D R I T Y G Q V D G N K V S N K V K L R P G K Y Y
 TCTACTGTTTATAAATACCTCAGGATCAGGAAACTATGAGTTAAGGGTAAATAAATTTATCTTATAAAGAGTGTGCCAATACATGGCACACTC 4400
 L L V Y K Y S G S G N Y E L R V N K *
 TTTTATTATTTTTTCTTTAAAAGATCTCTGATTTCCAAGTAACTCTTCTCTCTGAAATTTCAAGGAACTTAACTTCTTCACTGCTTCTCT 4500
 * K N K E K L L D R I E G L L E E E R S I E P I K A E E V A E E MscL
 TTCTTTTAAATCTGTTTATAGTCTTATTAATAGGAATTTGAAIAAGAAATTTAAGAAAGTCAATATATTTGATTAATTTGACCATAATTAAG 4600
 T I L P F R N I L R I F L F I S F S I L F D L I N Q I F S R G N C D S
 TCAAAGGTTTCTGAAATTAATCCATGAAGTGAAGTTTTCGCTAGTAAAATAATCCACCTAAGATAAGTCTAGATAAGGCATTATAACATCATT 4700
 T L P K E S N L G H L T L K A S T F N I G G L I L G L I P M I V D N
 TACTAAGATGTTACAATCTTTCCGAAAGCCACCCTATGATAACACCTACAGCAAGATCGACTACATTAACCTTTCATGGCAAATTCCTTAAAATCTTT 4800
 V L S T V I K G F A G G I I V G V A L D V V N G K M A F E K F D K
 CACATAAAAATCTCCCTAAAGATTTAATATTAATATTAAATAACAAGTATAATCTTATTTAAATTAACATTAATATACTAAATATAATATGA 4900
 W M
 ATTATAAAAGTTTTACATTTATAGATAAATAAATATGGTATTAATATTATCAGGTTGATTGTTCTTTGATGTTCTTAAATTTCAAAAAATATGA 5000
 ATAATATAAGAGATAGTATCGTTTGAATATCTATTAAAAAAAATTAAGTGTGTTAAGAGGTTTAAATGAAGATATGTTATATCTGA 5100
 M K I M F I S D Orf2d6
 TATTCATGGTCTTTATATTTTTTAAATAAAGCATTAGAAGATTTGAAGAGGAAAAAGCAGATTATATAGGATTTAAGGAGATGATATATCATCGA 5200
 I H G S L Y F L N K A L E R F E E E K A D Y I G I L G D V L Y H G
 CCTAGAAATGATTTACCAAAAAGATATAAATCAAAAGGATGTTGCAAAAATCTAAATAGGTATAAAAATAAATAAGCCTAAGGGGAAATTTGTGATA 5300
 P R N D L P K E Y N P K D V A K I L N R Y K N K I I A V R G N C D S
 GTGAAGTAGATAAATGCTTATAGACTATCAATGCTTATGATGATTATAGTAAATTTTTTAAATAACAAGAAAAATTTTAACTCAGGCACATTTT 5400
 E V D Q M L I D Y P M L S D Y S I I F F N N K K I F L T H G H I F
 TAATAAGATTAATGCCTATTTAATATAGGAGATATGATAAGTGGTCATACTCATATCCAAGTATAGAACATATAGACGGGGTAACTTTTAT 5500
 N K D N M P H F N I G D I M I S G H T H I P S I E H I D G V T F I
 AATCCTGGTCTATCTATACCTAAAGGTGGAAGTGAATAATCTTATGATTTTAAATGAGGATGGAATCTCAATTAAAAATTTAAATGGAAGGTTA 5600
 N P G S I S I P K G G S E N S Y G I L N E D G F S I K N L N G K V I
 TATTAACITTTAAATATATAAGACTAAAGGAGGAATAAAGATGAATACATAAGAAATGGTTTTAAATAGTTTAAAGAGCGAGGGAAACCGCTAAAGGC 5700
 L T L N I * M N T I E M V L N S L K E A G E P L K A Orf3d6
 TGGAGAGATAGCAGAAAGACTGGTATTGACAAAAAGAGTGGATAAAGCTATAAAAAAATAAAGCCGAGAAAGATAACTCTCTCAAGAGGTGT 5800
 G E I A E K T G I D K E V D K A I K K L K A E E K I T S P C K R V
 TATTATACTTTGCATAAATTTATCTCACATGATAAATTAATAATATAAAAAATAAATAAGTAAATGTTTGGTCTTATTTGTAGATTAATAGGG 5900
 Y Y T I A *
 AAGTGGTCAATTC 5914

FIG. 2—Continued.

Downloaded from <http://jfb.asm.org/> on November 26, 2020 by guest

TABLE 1. Similarities among segments of collagenases^a

Segment	Degree of sequence similarity with:											
	ColH1	ColA1	Val1	ColH2a	ColH2b	ColA2	Val2	ColG3b	ColH3	ColA3a	ColA3b	Val3
ColG1	+163.1	+160.1	+18.3									
ColH1		+150.8	+9.4									
ColA1			+10.0									
ColG2				+23.5	+24.6	+23.5	+21.2					
ColH2a					+23.0	+17.1	+21.7					
ColH2b						+20.7	+18.2					
ColA2							+21.0					
ColG3a								+17.0	+24.1	+23.9	+14.9	+0.37
ColG3b									+20.4	+24.5	+26.4	-1.37
ColH3										+22.8	+18.0	+1.67
ColA2a											+29.9	+0.44
ColA2b												+0.89

^a The degree of sequence similarity is expressed as the standard deviation of the alignment score obtained for an original pair of sequences against the scores obtained for 1,000 randomized pairs. Segment designations are composed of the enzyme designation plus the segment number; for example, ColH1 is segment 1 of ColH. Val1, Val2, and Val3 are segments 1, 2, and 3 of the *V. alginolyticus* collagenase, respectively (31, 42, 49).

enzymes showed significant similarity to *Vibrio alginolyticus* collagenase (42) in segments 1 and 2, but not in segment 3 (alignment data not shown). A consensus motif for metalloproteinases, HEXXH, was conserved in segment 1. We proposed that the sequence EEXXXE, which is located at the C-terminal side of the HEXXH sequence with a gap of 26 amino acid residues (20), participates in forming the catalytic center of ColH. This second motif is also conserved in ColG, although the first glutamate residue is replaced with an aspartate residue. Each of the segment 3s of ColG produced in *E. coli* bound to insoluble collagen (unpublished data), suggesting that the region forms a collagen binding domain in ColG like that of ColH (29).

Southern hybridization. *C. histolyticum* DNA was digested with *Hind*III, *Xba*I, *Eco*RI, *Pst*I, *Sau*3AI, or *Ssp*I and separated on a 0.8% agarose gel. *colG*- and *colH*-specific probes were prepared by PCR. Only one band was detected for a given digestion in each hybridization profile (Fig. 4). The sizes of the positive bands were in good agreement with the size of the fragment predicted from the nucleotide sequence (Fig. 4A). The differences between the two hybridization profiles were clearly observed for the *Eco*RI, *Pst*I, and *Sau*3AI digests.

Northern hybridization with the *colG* and *colH* gene probes. Total RNA from *C. histolyticum* was prepared in late logarithmic growth phase, when production of the collagenases continued at a high level. The RNAs were separated on a denaturing agarose gel and subjected to Northern hybridization with the *colG* and *colH* probes. Only one band was detected in each hybridization experiment (Fig. 5), and the sizes of the *colG* and *colH* transcripts were estimated to be 3.7 and 3.25 kb, respectively.

Loci of the collagenase genes. Genomic DNA of *C. histolyticum* was analyzed by pulsed-field gel electrophoresis (Fig. 6A). A 45-kb band was observed in the undigested sample. Endonuclease *I-Ceu*I cleaved the chromosome DNA into 10 fragments (fragment A, 2,030 kb; B, 430 kb; C, 280 kb; D, 100 kb; E, 42 kb; F, 15 kb; G, 11 kb; H, 9.8 kb; I, 8.3 kb, and J, 6.2 kb). The size of fragment A was determined by electrophoresis as described in Materials and Methods. *I-Ceu*I fragment E (42 kb) was broader than the other bands in the digest, indicating that it overlapped the 45-kb band in the undigested DNA. The *C. histolyticum rrl* probe hybridized to all the *I-Ceu*I fragments (data not shown). The upstream and downstream *rrl* probes

hybridized to all the *I-Ceu*I fragments except A and C, respectively (Fig. 6B and C). Similar results were obtained with the *rrs* and *rf* probes (data not shown). The *rrl* probes did not hybridize to the 45-kb band in the undigested sample. Each of the collagenase gene probes hybridized to a single fragment in a given digest (Fig. 6D and E). Although the 2.03-Mb *I-Ceu*I fragment, the 2.23-Mb *Mlu*I fragment, and the 760-kb *Sac*II fragment showed positive signals with both probes, different *Nru*I and *Sma*I fragments were detected by the two probes.

Nucleotide sequences of the adjacent regions of the two collagenase genes. In order to see if the two collagenase genes were adjacent, the nucleotide sequences adjacent to *colG* and *colH* were determined (5.9 kb for *colG* and 14.0 kb for *colH* [Fig. 7]). The two fragments did not form a single contig. Around *colG*, one ORF (*orf1uG*) and three ORFs (*mscL*, *orf2dG*, and *orf3dG*) were found in the upstream (~1.0 kb) and downstream (~1.6 kb) regions, respectively. Around *colH*, nine




FIG. 4. Southern hybridization with *colG*-specific (A) and *colH*-specific (B) probes (corresponding to the N termini). Lanes: 1 and 8, size markers (sizes are indicated on the left [in kilobase pairs] [kb]); 2 and 9, *Hind*III digests; 3 and 10, *Xba*I digests; 4 and 11, *Eco*RI digests; 5 and 12, *Pst*I digests; 6 and 13, *Sau*3AI digests; 7 and 14, *Ssp*I digests.




FIG. 5. Northern hybridization with *colG*-specific (A) and *colH*-specific (B) probes (corresponding to the N termini). Lanes: 1 and 6, RNA markers (sizes are indicated on the left and right); 2 and 4, total RNA isolated at an optical density at 600 nm of 1.8 (3 μg); 3 and 5, total RNA isolated at an optical density at 600 nm of 2.7 (3 μg).

ORFs (*orf9uH*, *hprT*, *hflX*, *orf6uH*, *orf5uH*, *orf4uH*, *orf3uH*, *orf2uH*, and *orf1uH*) and four ORFs (*dcd*, *orf2dH*, *ruvA*, and *ruvB*) were found in the upstream (~8.7 kb) and downstream (~2.3 kb) regions, respectively. Results of Blastp similarity searches with the amino acid sequences deduced from these ORFs as the query sequences are shown in Table 2.

DISCUSSION

The similarity between the N-terminal sequences of the 78-kDa gelatinase and the *C. perfringens* ColA collagenase (31) suggested the presence of an additional collagenase besides ColH (49) in *C. histolyticum*. We purified the 116-kDa collagenase (ColG) from the culture supernatant. The molecular mass of ColG is close to that of ColH. This explains why we detected only a single collagenolytic band in the culture supernatant by collagen zymography. The present study showed that

C. histolyticum possesses the two distinct collagenase genes. We determined the N-terminal sequences of five enzymes in this study in addition to the two ColH enzymes already determined (49). All the N-terminal sequences of the seven enzymes correspond to either *colG* or *colH*. The results of Southern hybridization showed that both *colG* and *colH* are single-copy genes. As shown by northern hybridization, each gene is transcribed into a single message, eliminating the possibility that the smaller enzymes are produced by premature transcription termination. The size of each transcript indicated that each gene is monocistronic. Thus, it is likely that the smaller enzymes are produced from their respective precursor enzymes by proteolytic cleavage at the C terminus, leading to the multiplicity in each class.

Bond and Van Wart isolated six enzymes from a commercial collagenase preparation and divided them into two classes based on substrate specificity and amino acid analysis. Class I enzymes (α , β , and γ) are less active against various peptide substrates, including the PZ-peptide, than class II enzymes (δ , ϵ , and ζ) (6). In our study the PZ-peptidase activity of ColG was 2.96 ± 0.26 mU/pmol of protein, much lower than that of rColH (85.9 ± 3.6 mU/pmol of protein). On the other hand, their activities against insoluble collagen are similar; the activities of ColG and ColH are 94.1 ± 4.8 and 164 ± 6 mU/pmol of protein, respectively (19). The 78-kDa ColG gelatinase also showed low hydrolytic activity against the PZ-peptide (1.32 ± 0.10 mU/pmol of protein). Previously, we showed that two C-terminally truncated forms of rColH (rColH'; 87 and 80 kDa) possess high PZ-peptidase activities (62.9 ± 4.8 and 123 ± 6 mU/pmol of protein, respectively), similar to that of full-length ColH (29). The activity on the peptide substrates regardless of C-terminal truncation can be explained by the presence of the catalytic domains at the N termini of ColG and ColH, which determine their specificities on the water-soluble substrates. The amino acid compositions predicted from *colG* and *colH* were compared with the amino acid analysis data for the largest enzymes in each class, β (class I, 115 kDa) and ζ




FIG. 6. (A) Plugs were subjected to endonuclease digestion, and the DNA fragments were separated by pulsed-field gel electrophoresis. Lanes: 1, size markers consisting of a mixture of λ concatemers and *HindIII* fragments (size range, 6.6 to 145.5 kb) (sizes are indicated on the left); 2 and 10, size markers consisting of *Saccharomyces cerevisiae* chromosomes (size range, 225 to 1,900 kb); 3, undigested DNA; 4, I-CeuI digest; 5, *ApaI* digest; 6, *MluI* digest; 7, *NruI* digest; 8, *SacII* digest; 9, *SmaI* digest; 11, size markers consisting of λ concatemers. (B through E) Southern hybridization with an upstream *rrlB* probe (B), a downstream *rrlB* probe (C), a *colG* probe (D), and a *colH* probe (E).




FIG. 7. Genetic organization of the DNA fragments carrying *colG* and *colH*. The amino acid sequence deduced from the ORF was significantly similar to that of the protein shown below the respective arrow. Some restriction sites are also shown. H, HindIII; E, EcoRI.

(class II, 125 kDa) (6). The combinations of ColG plus β and ColH plus ζ showed higher correlation values ($r^2 = 0.978$ and 0.963) than the alternative combination ($r^2 = 0.937$ and 0.922). Based on these observations, it is highly possible that β (class I) and ζ (class II) enzymes are encoded by *colG* and *colH*, respectively. We can assume that the smaller species of enzymes described by Bond and Van Wart (6) are also the C-terminal truncates of the respective full-length enzyme: α (68 kDa) and γ (79 kDa) from β and δ (100 kDa) and ϵ (110 kDa) from ζ .

We analyzed the loci of the two collagenase genes by pulsed-field gel electrophoresis. An intron-encoded endonuclease, I-CeuI, specifically cleaves DNA at a 26-bp recognition site present in the 23S rRNA genes of many species. We cloned three *rrl* fragments (*rrlA*, *rrlB*, and *rrlC*) from *C. histolyticum* to confirm the presence of an I-CeuI site in each gene (unpublished data). The number and sizes of the I-CeuI fragments of the *C. histolyticum* chromosome were similar to those from

C. perfringens (fragment A, 2,280 kb; B, 400 kb; C, 250 kb; D, 250 kb; E, 150 kb; F, 95 kb; G, 60 kb; H, 25 kb; I, 9.5 kb, and J, 9.5 kb) (22). The total genome size of *C. histolyticum* is estimated to be 2.9 Mb from their sizes, which coincides well with the value calculated from the sizes of the *MluI* fragments. Hybridization with the *rrl*, *rms*, and *rnf* probes indicated the presence of 10 copies of the *rrm* operon on the *C. histolyticum* chromosome, which is similar to findings for many gram-positive species (11). The results suggest that fragment C (280 kb) has two copies of the 5' region of the *rrm* operon outbound at both ends and that fragment A (2,030 kb) has two copies of the 3' region of the *rrm* operons inbound at both ends. All the other fragments have a 3' region of the *rrm* operon at one end and a 5' region at the other end. The number and orientation of *rrm* operons showed that the organization of the *C. histolyticum* chromosome is similar to that of the *C. perfringens* chromosome (13).

The presence of a plasmid (approximately 45 kb) was re-

TABLE 2. Proteins deduced from the ORFs around the two collagenase genes

ORF	Size (aa ^a) of deduced protein	Blast, P value	Accession no.	Description of protein similar to the deduced product
<i>orf1uG</i>	164			None (no sequences with significant similarity)
<i>mscL</i>	133	8.9e-52	E70065	Large conductance-mechanosensitive channel from <i>B. subtilis</i>
<i>orf2dG</i>	180	6.4e-61	B65002	Hypothetical protein b2300 from <i>E. coli</i>
<i>orf3dG</i>	58	9.5e-15	G69538	Conserved hypothetical protein from <i>Archaeoglobus fulgidus</i>
<i>orf9uH</i>	215	1.5e-17	E70015	Hypothetical protein YunB from <i>B. subtilis</i>
<i>hprT</i>	175	2.4e-53	S30100	Hypoxanthine phosphoribosyltransferase from <i>Lactococcus lactis</i>
<i>hflX</i>	596	3.5e-105	S75472	GTP binding protein from a <i>Synechocystis</i> sp.
<i>orf6uH</i>	298			None (no sequences with significant similarity)
<i>orf5uH</i>	466	2.8e-44	I40492	Hypothetical protein 8 (<i>srfA</i> operon) from <i>B. subtilis</i>
<i>orf4uH</i>	224	5.8e-6	S75919	Hypothetical protein from a <i>Synechocystis</i> sp.
<i>orf3uH</i>	216	4.7e-54	A70049	Conserved hypothetical protein YvyE from <i>B. subtilis</i>
<i>orf2uH</i>	246	2.1e-90	B69972	Spore coat protein homolog YrbC from <i>B. subtilis</i>
<i>orf1uH</i>	159	5.7e-63	E64249	Pilin repressor PilB homolog from <i>Mycoplasma genitalium</i>
<i>dcd</i>	172	3.4e-15	F64353	dCTP deaminase from <i>Methanococcus jannaschii</i>
<i>orf2dH</i>	204			None (no sequences with significant similarity)
<i>ruvA</i>	197	5.7e-44	E69702	Holliday junction DNA helicase RuvA from <i>B. subtilis</i>
<i>ruvB</i>	>54	(5.3e-11)	C64652	Holliday junction DNA helicase from <i>Helicobacter pylori</i>

^a aa, amino acids.




FIG. 8. Schematic representation of a plausible evolutionary pathway of the bacterial collagenase genes. Seg., segment.

vealed by the comparison of the electrophoretic patterns of uncut and I-CeuI-cut DNA and the *rfl* hybridization experiment. In *C. perfringens* the collagenase (κ toxin) gene (*colA*) is chromosomal, while some of the other toxin genes (ϵ , ι , β , and λ) are plasmid borne (21). The enterotoxin gene (*cpe*) is either chromosomal or plasmid borne depending on the strain (10). However, Southern hybridization with the *colG* and *colH* probes showed that neither gene is plasmid borne. The locus of the two genes on the *C. histolyticum* chromosome differs from that of *colA* on the *C. perfringens* chromosome. The former is located on I-CeuI fragment A (2,030 kb), while the latter is on I-CeuI fragment E (150 kb) (22). The distance between *colG* and *colH* genes in *C. histolyticum* is less than 760 kb, as shown by the *Sac*II digest, but they are expected to be well separated, since there is at least one *Nru*I site and one *Sma*I site between them. The nucleotide sequences of the regions adjacent to the two collagenase genes revealed that the genes are not tandemly located and that they are separated from each other by at least 3.3 kb.

In pathogenic clostridia some of the toxin genes are located on lysogenized phages or transposable elements (8, 16, 17). However, we could not find any such genes around the collagenase genes by a Blastn similarity search. Downstream of *colH* are two ORFs which encode peptides homologous to a Holliday junction helicase, an enzyme essential for general recombination. The evolution of the clostridial collagenase genes can be explained by gene duplication and divergence as follows: (i) in an ancestral cell common to *C. histolyticum* and *C. perfringens*, an ancestral collagenase gene carrying a single copy of each segment (segments 1, 2, and 3) was duplicated; (ii) segments 3 and 2 were duplicated in each descendant to diverge into the subsequent ancestral genes encoding class I (segments 1, 2, 3a, and 3b) and class II (segments 1, 2a, 2b, and 3) enzymes, respectively; and (iii) species divergence occurred by an unknown mechanism and mutations were accumulated in

each copy. *colG* and *colA* are considered to have derived from the former ancestor gene by divergence. This speculation is supported by the following observations. First, the segmental structure of *C. histolyticum* ColG is the same as that of *C. perfringens* ColA, and their sequences align with a significantly high similarity (Blastp *P* value, $1.2e-255$). Second, both genes have a region encoding a long prepropeptide (110 amino acid residues for ColG and 86 residues for ColA), unlike *colH*. Finally, these genes are accompanied by the homologous *mscL* gene, encoding the mechanosensitive channel homolog (Blastp *P* value for similarity to *C. perfringens* MscL [29], $7.7e-44$). We are now searching for the counterpart of *colH* in *C. perfringens*, tentatively named *colB*, although it might have been lost during evolution. Alternatively, the class divergence could have occurred after the species divergence. If this is the case, ColG should be more closely related to ColH than to ColA. In order to evaluate this hypothesis, the similarity between their segment 1s was examined with the Blastp server, so the effects of their different C-terminal structures could be ignored. Since ColG1 showed higher similarity to ColA1 (*P* value, $1.4e-182$) than to ColH1 (*P* value, $3.6e-175$), this hypothesis seems less likely. The International Polycystic Kidney Disease Consortium suggested that a segment 2-encoding fragment has been horizontally transferred from eukaryotic cells to prokaryotic cells (18). If this is the case, the event should have occurred before these genera diverged. Although the *V. alginolyticus* collagenase (42) has segments 1 and 2 that are significantly similar to those of the clostridial collagenases, its C-terminal segment 3 shows no significant similarity to that of the clostridial enzymes. This suggests that the addition of segment 2 had taken place before these genera diverged and that a different C-terminal segment was added to each. On the basis of this discussion, we propose a hypothesis for the evolution of the bacterial collagenases, as depicted schematically (Fig. 8). More system-

atic investigation of bacterial collagenase genes is necessary to draw a conclusive evolutionary picture.

ACKNOWLEDGMENTS

We thank David B. Wilson (Section of Biochemistry, Molecular and Cell Biology, Cornell University, Ithaca, N.Y.) for invaluable discussions and assistance in preparing the manuscript.

This work was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture in Japan.

REFERENCES

- Aiba, H., S. Adhya, and B. de Crombrughe. 1981. Evidence for two functional *gal* promoters in intact *Escherichia coli* cells. *J. Biol. Chem.* **256**:11905–11910.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Becker, A., F. Theuring, M. Gottwald, K. Kauser, W.-D. Schleuning, and P. Donner. 1994. Purification of human big endothelin 1 derived through cleavage with collagenase and dipeptidylpeptidase IV from a fusion protein expressed in *Escherichia coli*. *Protein Expr. Purif.* **5**:50–56.
- Bethesda Research Laboratories. 1986. BRL pUC host: *E. coli* DH5 α TM competent cells. *Focus* **8**:9.
- Birkedal-Hansen, H., and R. E. Taylor. 1982. Detergent-activation of latent collagenase and resolution of its component molecules. *Biochem. Biophys. Res. Commun.* **107**:1173–1178.
- Bond, M. D., and H. E. Van Wart. 1984. Characterization of the individual collagenases from *Clostridium histolyticum*. *Biochemistry* **23**:3085–3091.
- Bond, M. D., and H. E. Van Wart. 1984. Relationship between the individual collagenases of *Clostridium histolyticum*: evidence for evolution by gene duplication. *Biochemistry* **23**:3092–3099.
- Brynstad, S., B. Synstad, and P. E. Granum. 1997. The *Clostridium perfringens* enterotoxin gene is on a transposable element in type A human food poisoning strains. *Microbiology* **143**:2109–2115.
- Canard, B., and S. T. Cole. 1989. Genome organization of the anaerobic pathogen *Clostridium perfringens*. *Proc. Natl. Acad. Sci. USA* **86**:6676–6680.
- Cornillot, E., B. Saint-Joanis, G. Daube, S. Katayama, P. E. Granum, B. Canard, and S. T. Cole. 1995. The enterotoxin gene (*cpe*) of *Clostridium perfringens* can be chromosomal or plasmid-borne. *Mol. Microbiol.* **15**:639–647.
- de Bruijn, F. J., J. R. Lupski, and G. M. Weinstock. 1998. Bacterial genomes: physical structure and analysis. Chapman & Hall, New York, N.Y.
- Fukunaga, M., Y. Yanagihara, and M. Sohnaka. 1992. The 23S/5S ribosomal RNA genes (*rrl/rrf*) are separate from the 16S ribosomal RNA gene (*rrs*) in *Borrelia burgdorferi*, the aetiological agent of Lyme disease. *J. Gen. Microbiol.* **138**:871–877.
- Garnier, T., B. Canard, and S. T. Cole. 1991. Cloning, mapping, and molecular characterization of the rRNA operons of *Clostridium perfringens*. *J. Bacteriol.* **173**:5431–5438.
- Gelbard, M. K., K. James, P. Riach, and F. Dorey. 1993. Collagenase versus placebo in the treatment of Peyronie's disease: a double-blind study. *J. Urol.* **149**:56–58.
- Hanada, K., I. Yamato, and Y. Anraku. 1988. Purification and reconstitution of *Escherichia coli* proline carrier using a site specifically cleavable fusion protein. *J. Biol. Chem.* **263**:7181–7185.
- Inoue, K., and H. Iida. 1970. Conversion of toxigenicity in *Clostridium botulinum* type C. *Jpn. J. Microbiol.* **14**:87–89.
- Inoue, K., and H. Iida. 1971. Phage-conversion of toxigenicity in *Clostridium botulinum* types C and D. *Jpn. J. Med. Sci. Biol.* **24**:53–56.
- The International Polycystic Kidney Disease Consortium. 1995. Polycystic kidney disease: the complete structure of the *PKD1* gene and its protein. *Cell* **81**:289–298.
- Jung, C.-M., O. Matsushita, S. Katayama, J. Minami, I. Ohhira, and A. Okabe. 1996. Expression of the *colH* gene encoding *Clostridium histolyticum* collagenase in *Bacillus subtilis* and its application to enzyme purification. *Microbiol. Immunol.* **40**:923–929.
- Jung, C.-M., O. Matsushita, S. Katayama, J. Minami, J. Sakurai, and A. Okabe. Analysis of the catalytic center in the *Clostridium histolyticum* ColH collagenase. Submitted for publication.
- Katayama, S., B. Dupuy, G. Daube, B. China, and S. T. Cole. 1996. Genome mapping of *Clostridium perfringens* strains with I-CeuI shows many virulence genes to be plasmid-borne. *Mol. Gen. Genet.* **251**:720–726.
- Katayama, S.-I., B. Dupuy, T. Garnier, and S. T. Cole. 1995. Rapid expansion of the physical and genetic map of the chromosome of *Clostridium perfringens* CPN50. *J. Bacteriol.* **177**:5680–5685.
- Kono, T. 1968. Purification and partial characterization of collagenolytic enzymes from *Clostridium histolyticum*. *Biochemistry* **7**:1106–1114.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680–685.
- Lanzillo, J. J. 1991. Chemiluminescent nucleic acid detection with digoxigenin-labeled probes: a model system with probes for angiotensin converting enzyme which detect less than one attomole of target DNA. *Anal. Biochem.* **194**:45–53.
- Lwebuga-Mukasa, J. S., E. Harper, and P. Taylor. 1976. Collagenase enzymes from *Clostridium*: characterization of individual enzymes. *Biochemistry* **15**:4736–4741.
- Mandl, I., S. Keller, and J. Manahan. 1964. Multiplicity of *Clostridium histolyticum* collagenase. *Biochemistry* **3**:1737–1741.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Matsushita, O., C.-M. Jung, J. Minami, S. Katayama, N. Nishi, and A. Okabe. 1998. A study of the collagen-binding domain of a 116-kDa *Clostridium histolyticum* collagenase. *J. Biol. Chem.* **273**:3643–3648.
- Matsushita, O., C. M. Jung, and A. Okabe. 1995. Identification of the gene encoding a mechanosensitive channel MscL homologue in *Clostridium perfringens*. *Gene* **165**:147–148.
- Matsushita, O., K. Yoshihara, S.-I. Katayama, J. Minami, and A. Okabe. 1994. Purification and characterization of a *Clostridium perfringens* 120-kilodalton collagenase and nucleotide sequence of the corresponding gene. *J. Bacteriol.* **176**:149–156.
- Messing, J. 1983. New M13 vectors for cloning. *Methods Enzymol.* **101**:20–78.
- Miyoshi, M., and J. Rosenbloom. 1974. General proteolytic activity of highly purified preparations of clostridial collagenase. *Connect. Tissue Res.* **2**:77–84.
- Mookhtiar, K. A., and H. E. Van Wart. 1992. *Clostridium histolyticum* collagenases: a new look at some old enzymes. *Matrix Suppl.* **1**:116–126.
- Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne. 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng.* **10**:1–6.
- Nishi, N., O. Matsushita, K. Yuube, H. Miyataka, A. Okabe, and F. Wada. 1998. Collagen-binding growth factors: production and characterization of functional fusion proteins having a collagen-binding domain. *Proc. Natl. Acad. Sci. USA* **95**:7018–7023.
- Pearson, W. R., and D. J. Lipman. 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**:2444–2448.
- Peterkofsky, B. 1982. Bacterial collagenase. *Methods Enzymol.* **82**:453–471.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
- Seglen, P. O. 1976. Preparation of isolated rat liver cells. *Methods Cell Biol.* **13**:29–83.
- Starkweather, K. D., S. Lattuga, L. C. Hurst, M. A. Badalamente, F. Guilak, S. P. Sampson, A. Dowd, and D. Wisch. 1996. Collagenase in the treatment of Dupuytren's disease: an *in vitro* study. *J. Hand Surg.* **21**:490–495.
- Takeuchi, H., Y. Shibano, K. Morihara, J. Fukushima, S. Inami, B. Keil, A.-M. Gilles, S. Kawamoto, and K. Okuda. 1992. Structural gene and complete amino acid sequence of *Vibrio alginolyticus* collagenase. *Biochem. J.* **281**:703–708.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
- Warren, G. H., and J. Gray. 1961. A simplified culture medium for the production of collagenase. *Nature (London)* **192**:755–756.
- Wilson, M. J., M. Strasser, M. M. Vogel, and A. A. Sinha. 1991. Calcium-dependent and calcium-independent gelatinolytic proteinase activities of the rat ventral prostate and its secretion: characterization and effect of castration and testosterone treatment. *Biol. Reprod.* **44**:776–785.
- Wolters, G. H. J., G. H. Vos-Scheperkeuter, H.-C. Lin, and R. Van Schilf-gaarde. 1995. Different roles of class I and class II *Clostridium histolyticum* collagenase in rat pancreatic islet isolation. *Diabetes* **44**:227–233.
- Wünsch, E., and H.-G. Heidrich. 1963. Zur quantitativen Bestimmung der Kollagenase. *Hoppe-Seyler's Z. Physiol. Chem.* **333**:149–151.
- Yoshida, E., and H. Noda. 1965. Isolation and characterization of collagenases I and II from *Clostridium histolyticum*. *Biochim. Biophys. Acta* **105**:562–574.
- Yoshihara, K., O. Matsushita, J. Minami, and A. Okabe. 1994. Cloning and nucleotide sequence analysis of the *colH* gene from *Clostridium histolyticum* encoding a collagenase and a gelatinase. *J. Bacteriol.* **176**:6489–6496.