

Cloning and Characterization of the *Flavobacterium johnsoniae* Gliding-Motility Genes *gldB* and *gldC*

DAVID W. HUNNICUTT AND MARK J. MCBRIDE*

Department of Biological Sciences, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin 53201

Received 25 June 1999/Accepted 17 November 1999

The mechanism of bacterial gliding motility (active movement over surfaces without the aid of flagella) is not known. A large number of mutants of the gliding bacterium *Flavobacterium johnsoniae* (*Cytophaga johnsonae*) with defects in gliding motility have been previously isolated, and genetic techniques to analyze these mutants have recently been developed. We complemented a nongliding mutant of *F. johnsoniae* (UW102-99) with a library of wild-type DNA by using the shuttle cosmid pCP26. The complementing plasmid (pCP200) contained an insert of 26 kb and restored gliding motility to 4 of 50 independently isolated nongliding mutants. A 1.9-kb fragment which encompassed two genes, *gldB* and *gldC*, complemented all four mutants. An insertion mutation in *gldB* was polar on *gldC*, suggesting that the two genes form an operon. Disruption of the chromosomal copy of *gldB* in wild-type *F. johnsoniae* UW101 eliminated gliding motility. Introduction of the *gldBC* operon, or *gldB* alone, restored motility. *gldB* appears to be essential for *F. johnsoniae* gliding motility. It codes for a membrane protein that does not exhibit strong sequence similarity to other proteins in the databases. *gldC* is not absolutely required for gliding motility, but cells that do not produce GldC form colonies that spread less well than those of the wild type. GldC is a soluble protein and has weak sequence similarity to the fungal lectin AOL.

The mechanism of bacterial gliding motility has been an unsolved mystery for over 100 years (45). Gliding motility is defined as smooth translocation of cells over surfaces, generally following the long axes of the cells. Gliding bacteria lack flagella. They produce distinctive colonies with multicellular flares at the spreading edges. A variety of mechanisms have been proposed to explain bacterial gliding motility, but no single model has emerged that explains all of the observations made on the many different gliding bacteria that have been studied (7, 19, 29, 42, 51). Bacteria capable of gliding motility are found in many of the branches of the eubacterial phylogenetic tree, and it is possible that bacteria from different branches use different mechanisms to glide over surfaces.

Flavobacterium johnsoniae (formerly *Cytophaga johnsonae* [4]) is a common soil and aquatic bacterium that exhibits rapid gliding motility (29). The cells glide at rates of up to 600 $\mu\text{m}/\text{min}$ over glass surfaces and up to 60 $\mu\text{m}/\text{min}$ over agar surfaces (29). *F. johnsoniae* is a member of the *Cytophaga-Flavobacterium-Bacteroides* group of bacteria. This large and diverse assemblage of gram-negative organisms contains numerous bacteria that exhibit gliding motility. *F. johnsoniae* has become an attractive model organism for studies of bacterial gliding because of its rapid motility and the ease with which it can be cultivated. Pate and colleagues isolated a large number of nongliding mutants of *F. johnsoniae* (8, 50). Unlike wild-type *F. johnsoniae*, these mutants form nonspreading colonies under all conditions of cultivation tested. They also lack the single-cell gliding movements that are observed with wild-type cells. Techniques to genetically manipulate *F. johnsoniae* were recently developed and used to identify one gene, *gldA*, that is required for gliding motility (2, 26, 27). GldA exhibits sequence similarity to ATP-binding cassette transport proteins, but its exact role in gliding motility has not yet been determined. Here we report the identification and characterization

of two additional genes, *gldB* and *gldC*, that are involved in *F. johnsoniae* gliding motility. *gldB* is required for gliding motility. *gldC* is not required for gliding motility, but cells that do not produce GldC form colonies that spread less well than the wild type.

MATERIALS AND METHODS

Bacterial and bacteriophage strains, plasmids, and growth conditions. *F. johnsoniae* UW101 (ATCC 17061) was the wild-type strain used in these studies, and all mutants were derived from this strain. The 50 nongliding mutants of *F. johnsoniae* (obtained from J. Pate) were previously described (8, 50). The strain designations for each of these mutants carry the prefix 'UW102-'. The strain designations are UW102-9, -15, -21, -25, -33, -34, -39, -40, -41, -42, -48, -52, -53, -55, -56, -57, -58, -61, -64, -66, -68, -69, -75, -77, -78, -80, -81, -85, -86, -90, -92, -94, -95, -96, -97, -98, -99, -100, -101, -103, -107, -108, -140, -141, -146, -154, -300, -301, -302, and -348. The bacteriophage active against *F. johnsoniae* that were used in this study ($\phi\text{Cj}1$, $\phi\text{Cj}7$, $\phi\text{Cj}13$, $\phi\text{Cj}23$, $\phi\text{Cj}28$, $\phi\text{Cj}29$, $\phi\text{Cj}42$, $\phi\text{Cj}48$, and $\phi\text{Cj}54$) have been previously described (8, 32, 50). The *Escherichia coli* strains used were DH5 α MCR (GibcoBRL Life Technologies), HB101 (5), LMG194 (17), S17-1 (40), and BW19851 (28), an S17-1 *pir* strain. *E. coli* strains were grown in Luria-Bertani (LB) medium at 37°C and *F. johnsoniae* strains were grown in Casitone-yeast extract (CYE) medium at 30°C, as previously described (27). To observe colony spreading, *F. johnsoniae* was grown on PY2 medium (2) at 25°C. Antibiotics were used at the following concentrations when needed: ampicillin, 100 $\mu\text{g}/\text{ml}$; chloramphenicol, 30 $\mu\text{g}/\text{ml}$; erythromycin, 100 $\mu\text{g}/\text{ml}$; tetracycline, 15 $\mu\text{g}/\text{ml}$; kanamycin, 30 $\mu\text{g}/\text{ml}$; streptomycin, 30 $\mu\text{g}/\text{ml}$; and trimethoprim, 200 $\mu\text{g}/\text{ml}$. The plasmids used in this study are listed in Table 1.

Construction of the *F. johnsoniae*-*E. coli* shuttle cosmids pCP22 and pCP26. To construct pCP22 (Fig. 1), pCP19 was digested with *Bam*HI and *Eco*RI and the 2.8-kb fragment containing the origin of replication that functions in *F. johnsoniae* was ligated into cosmid pNJR6, which had been cut with the same enzymes. The 2-kb *Sac*I-*Pst*I fragment containing the *tetA* and *tetR* genes (originally from RK2) was excised from pTGL130 and made blunt by treatment with DNA polymerase Klenow fragment. This fragment was cloned into pCP22, which had been cut with *Bam*HI and treated with DNA polymerase Klenow fragment, to generate pCP26 (Fig. 1).

Construction of an *F. johnsoniae* genomic library. *F. johnsoniae* DNA was partially digested with *Sau*3AI and separated by agarose gel electrophoresis. Fragments between 15 and 30 kb were cut from the gel and purified using the Gene Clean spin kit (Bio 101 Inc.). The purified fragments were ligated with the shuttle cosmid pCP26 that had been digested with *Bgl*II and treated with alkaline phosphatase. The DNA was packaged into lambda phage particles (MaxPlax; Epicentre Technologies, Madison, Wis.) and introduced into *E. coli* DH5 α MCR. Cosmid DNA from approximately 10,000 colonies was transferred into the *F. johnsoniae* nongliding mutant UW102-99 by triparental conjugation. Approximately 5×10^9 cells each of recombinant *E. coli* DH5 α MCR, *F. johnsoniae* UW102-99, and *E. coli* HB101 (carrying the helper plasmid R702) were mixed

* Corresponding author. Mailing address: Department of Biological Sciences, University of Wisconsin—Milwaukee, P.O. Box 413, Milwaukee, WI 53201. Phone: (414) 229-5844. Fax: (414) 229-3926. E-mail: mcbride@uwm.edu.

TABLE 1. Plasmids used in this study^a

Plasmid	Description	Source or reference
pBC SK(+)	ColE1 ori; Cm ^r	Stratagene
pCR-Script Amp SK(+)	ColE1 ori; Ap ^r	Stratagene
pMAL-c2	ColE1 ori; Ap ^r ; <i>malE</i> fusion expression vector	New England BioLabs
pBAD/His-C	ColE1 ori; Ap ^r ; His-tag expression vector	Invitrogen
pNJR6	RSF1010 ori; Kn ^r (Em ^r)	44
R702	IncP; Km ^r Sm ^r Tc ^r ; helper plasmid for triparental conjugation	18
pLYL03	ColE1 ori; Ap ^r (Em ^r); <i>Bacteroides-Flavobacterium</i> suicide vector used to make chromosomal insertions	25
pTGL130	ColE1 ori; Ap ^r Tc ^r	11
R751::Tn4351Ω4	IncP; Tp ^r Tc ^r (Em ^r); Vector used for Tn4351 mutagenesis	39
pEP4351	<i>pir</i> -requiring R6K oriV; RP4 oriT; Cm ^r Tc ^r (Em ^r); vector used for Tn4351 mutagenesis	9
pCP11	ColE1 ori; (pCP1 ori); Ap ^r (Em ^r); <i>E. coli-F. johnsoniae</i> shuttle plasmid	27
pCP19	(pCP1 ori); (Em ^r); <i>F. johnsoniae</i> plasmid	2
pCP22	RSF1010 ori; (pCP1 ori); Kn ^r (Em ^r); <i>E. coli-F. johnsoniae</i> shuttle cosmid	This study
pCP23	ColE1 ori; (pCP1 ori); Ap ^r (Tc ^r); <i>E. coli-F. johnsoniae</i> shuttle plasmid	2
pCP26	RSF1010 ori; (pCP1 ori); Kn ^r Tc ^r (Em ^r); <i>E. coli-F. johnsoniae</i> shuttle cosmid	This study
pCP200	RSF1010 ori; (pCP1 ori); Tc ^r (Em ^r); cosmid clone complementing <i>F. johnsoniae</i> UW102-99	This study
pDH65	Nucleotides 145–754 of <i>gldB</i> in pBC SK(+); Cm ^r	This study
pDH222	1.9-kb <i>XbaI-ClaI</i> fragment of pCP200 in pBC SK(+); Cm ^r	This study
pDH223	1.9-kb <i>XbaI-SalI</i> fragment of pDH222 in pCP11; Ap ^r (Em ^r)	This study
pDH226	850-bp <i>PstI-PvuII</i> fragment of pDH65 cloned into <i>PstI-SmaI</i> -cut pBC SK(+); Cm ^r	This study
pDH227	854-bp <i>PstI-BamHI</i> fragment of pDH226 cloned into pLYL03; Ap ^r (Em ^r); plasmid for <i>gldB</i> gene disruption	This study
pDH233	1.9-kb <i>XbaI-KpnI</i> fragment of pDH222 in pCP23; Ap ^r (Tc ^r)	This study
pDH238	497-bp fragment containing 3' end of <i>gldB</i> in <i>SrfI</i> site of pCR-Script SK(+); Ap ^r	This study
pDH240	368-bp fragment containing <i>gldC</i> in <i>SrfI</i> site of pCR-Script SK(+); Ap ^r	This study
pDH241	335-bp <i>PstI-HindIII</i> fragment of pDH240 in pBAD/His-C; Ap ^r ; plasmid for <i>gldC</i> overexpression	This study
pDH242	1-kb fragment containing <i>gldB</i> in <i>SrfI</i> site of pCR-Script SK(+); Ap ^r	This study
pDH243	1-kb fragment containing <i>gldB</i> in pCP11; Ap ^r (Em ^r)	This study
pDH245	1-kb fragment containing <i>gldB</i> in pBC SK(+); Cm ^r	This study
pDH246	1-kb fragment containing <i>gldB</i> in pCP23; Ap ^r (Tc ^r)	This study
pDH250	479-bp <i>PstI-HindIII</i> fragment of pDH238 in pMAL-c2; Ap ^r ; plasmid for <i>gldB</i> overexpression	This study

^a Antibiotic resistance phenotypes: ampicillin, Ap^r; chloramphenicol, Cm^r; erythromycin, Em^r; kanamycin, Kn^r; streptomycin, Sm^r; tetracycline, Tc^r; trimethoprim, Tp^r. Unless indicated otherwise, antibiotic resistance genes and origins of replication are those expressed by *E. coli*. Antibiotic resistance phenotypes and other features listed in parentheses are those expressed by *F. johnsoniae* but not by *E. coli*.

and spotted on CYE medium containing 20 g of agar per liter and 5 mM CaCl₂. (The addition of CaCl₂ to the medium was found to increase conjugative DNA transfer efficiency.) Following overnight incubation at 30°C, cells were scraped off the plates, diluted in TC buffer (27), and plated on PY2 medium containing

100 µg of erythromycin per ml. Colonies were examined after incubation for 2 to 3 days at 25°C.

Subcloning of pCP200. The 26-kb insert from cosmid pCP200, which complemented the nongliding mutant UW102-99, was subcloned using standard proce-




FIG. 1. Maps of the *F. johnsoniae*-*E. coli* shuttle cosmids pCP22 and pCP26.

dures (37). A 2.1-kb *Bgl*II fragment of pCP200 was inserted into the *Bam*HI site of pBC SK(+) to generate pCP216. Nested deletions of this fragment for sequencing were obtained by digesting pCP216 with *Xba*I and *Sac*I followed by partial exonuclease III digestion, S1 nuclease digestion, and ligation.

A 1.9-kb fragment of DNA spanning the *gldBC* operon was obtained by digestion of pCP200 with *Xba*I and partial digestion with *Clal*. This fragment was inserted into pBC SK(+) to generate pDH222, excised as an *Xba*I-*Sal*I fragment, and ligated into pCP11 to generate pDH223. The 1.9-kb *gldBC* region of pDH222 was also inserted as a *Xba*I-*Kpn*I fragment into pCP23 to form pDH233.

PCR amplification was used to obtain a clone containing just *gldB*. The *gldB* gene was amplified using primers 5'-CAGGGATGTATATTTGCAG-3' and 5'-TATTGTATAAGCTTTTATTCTTAGGTTT-3' (*Hind*III site [underlined] added) and cloned into the *Srf*I site of pCR-Script Amp SK(+) to produce pDH242. The *Not*I-*Sal*I fragment of pDH242, which contained *gldB*, was cloned into pCP11 to generate pDH243. To generate convenient restriction sites for subsequent cloning into pCP23, the *Bam*HI-*Sal*I fragment of pDH243 was cloned into *Bam*HI-*Sal*I digested pBC SK(+) to produce pDH245. pDH245 was digested with *Kpn*I and *Xba*I, and the *gldB*-containing fragment was cloned into pCP23, which had been digested with the same enzymes, to generate pDH246. pDH243 and pDH246 contain the entire *gldB* coding region and 89 bp of upstream DNA. Plasmid DNA was transferred to nongliding mutants by conjugation or electroporation as previously described (27), and the transformants were plated on PY2 medium containing the appropriate antibiotic and observed for colony spreading.

Nucleic acid sequencing. Nucleic acid sequencing was performed by the dideoxynucleotide procedure with an automated sequencing system (Applied Biosystems). Sequences were analyzed with the MacVector and AssemblyLign software (Oxford Molecular Group, Campbell, Calif.), and comparisons to database sequences were made using the BLAST (3) and FASTA (33) algorithms. Pairwise sequence alignments were performed with the LALIGN program (20).

Disruption of *gldB*. The suicide vector pLYL03 was used for insertional mutagenesis to disrupt the chromosomal copy of *gldB* in wild-type *F. johnsoniae*. pDH65, a plasmid generated during exonuclease deletion subcloning (see above), contained nucleotides 145 to 754 of the *gldB* coding region in pBC SK(+). The *gldB* fragment of pDH65 was isolated following *Pst*I-*Pvu*II digestion. To generate convenient restriction sites for subsequent cloning into pLYL03, the 850-bp *Pst*I-*Pvu*II fragment containing 609 bp of *gldB* and 241 bp of pBC SK(+) was cloned into *Pst*I-*Sma*I-digested pBC SK(+) to generate pDH226. pDH226 was digested with *Pst*I and *Bam*HI, and the 853-bp fragment was ligated into the *Flavobacterium-Bacteroides* suicide vector pLYL03 to produce pDH227. pDH227 was introduced into *F. johnsoniae* UW-101 by conjugation from *E. coli* S17-1. Insertion of pDH227 into the chromosomal copy of *gldB* was confirmed by Southern blot analysis of the resulting Em^r colonies.

Tn4351 mutagenesis. Tn4351 was introduced into wild-type *F. johnsoniae* by conjugation from *E. coli* HB101 as described previously (27), except that 5 mM CaCl₂ was added to the CYE-2% agar conjugation medium. In some experiments, pEP4351 was used instead of R751::Tn4351Q4 as the vector for Tn4351 mutagenesis, since this resulted in an increased efficiency of transposition. pEP4351 was transferred to *F. johnsoniae* from *E. coli* BW19851. Transconjugants were plated on PY2 medium containing erythromycin and incubated for 2 to 3 days at 25°C.

Expression and purification of recombinant GldB and GldC. A fragment coding for the C-terminal 155 amino acids of GldB was amplified by PCR with primers 5'-AATTATTACTGCAGAAATTTGAGGAAAGA-3' (*Pst*I site [underlined] added) and 5'-TTATTGTATAAGCTTTTATTCTTAGGTTT-3' (*Hind*III site [underlined] added) and cloned into the *Srf*I site of pCR-Script Amp SK(+) to produce pDH238. The *gldB* fragment was isolated from pDH238 as a *Pst*I-*Hind*III fragment and was ligated into the expression vector pMAL-c2 cut with the same enzymes to generate pDH250. Fusion protein production was induced in *E. coli* DH5αMCR by the addition of 0.3 mM isopropyl-β-D-thiogalactopyranoside (IPTG).

The *gldC* gene was amplified by PCR with primers 5'-TAAACCTAAGAAC TGCACTCAAATACAATA-3' (*Pst*I site [underlined] added) and 5'-AAAAT ATTGAAGCTTCTATTATTAGTCAG-3' (*Hind*III site [underlined] added) and cloned into the *Srf*I site of pCR-Script Amp SK(+) to generate pDH240. *gldC* was isolated from pDH240 as a *Pst*I-*Hind*III fragment and was ligated into the expression vector pBAD/His-C to produce pDH241. Fusion protein production was induced in *E. coli* LMG194 with 0.004% arabinose.

The GldB fusion protein was partially purified by preparative sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The protein was visualized by CuCl₂ staining (24), the band was excised, and the protein was electroeluted for 16 h at 30 V into 0.1 M Tris-0.77 M glycine-0.4% SDS using an ISCO Little Blue Tank electroelutor. The recovered protein was dialyzed against 100 mM morpholinopropanesulfonic acid (MOPS) (pH 7.5) and used as the antigen. Recombinant GldC protein was purified by Ni²⁺ affinity chromatography (Probond; Invitrogen) with a stepwise elution of 30 mM, 300 mM, and 3 M imidazole (pH 6.0). The recombinant GldC eluted in the final fraction.

Antibody production. The recombinant GldB and GldC proteins (100 μg each) were mixed with an equal volume of Freund's complete adjuvant and injected into female New Zealand White rabbits. The animals were boosted with 20 to 50 μg of protein in Freund's incomplete adjuvant every 4 weeks until test bleeds showed suitable reactivity with control proteins in Western blot assays. Affinity

columns for purification of antisera were made by coupling 2 mg of recombinant GldB or GldC to 1 ml of Affi-Gel 15 (Bio-Rad). A 30-ml volume of heat-treated antiserum (56°C for 30 min, to inactivate complement) was passed over the appropriate affinity column. The column was washed with 10 ml of 100 mM MOPS (pH 7.5) followed by 4 ml of 0.5 M NaCl. Antibodies were eluted with 4 ml of 0.1 M glycine (pH 2.5) and immediately neutralized with 5 ml of 1 M Tris (pH 7.5).

Western blot analysis. Cells were grown to late log phase, pelleted, and lysed by incubation for 3 min at 100°C in SDS-PAGE sample buffer. Approximately 50 μg (GldC) or 200 μg (GldB) of total protein was loaded per lane. Protein was measured using the bicinchoninic acid reagent (Pierce Chemical Co., Rockford, Ill.). GldB preparations were separated by SDS-PAGE (12% acrylamide running gel, 4% acrylamide stacking gel) essentially as described by Laemmli (22), and GldC preparations were separated by Tricine-SDS-PAGE (10% acrylamide running gel, 4% acrylamide stacking gel) using the method of Schagger and von Jagow (38). Following electrophoresis, the proteins were transferred to polyvinylidene fluoride (PVDF) membranes (MSI) using a Bio-Rad TransBlot electrophoretic transfer cell as specified by the manufacturer. The blots were blocked with 5% skim milk in TBS (20 mM Tris hydrochloride [pH 7.5], 0.15 M NaCl) for 2 h at 25°C and then incubated for 4 h with antiserum diluted 1:1,000 in TBST (TBS, 0.5% Tween 20). The blots were washed three times with TBST and then incubated for 2 h with goat anti-rabbit immunoglobulin G-horse radish peroxidase conjugate (Boehringer Mannheim) diluted 1:10,000 in TBST. Antigens were detected using the enhanced chemiluminescence kit (Boehringer Mannheim) and Kodak X-Omat AR X-ray film.

Fractionation of cells. A 50-ml culture was grown to late exponential phase, and cells were pelleted by centrifugation at 7,000 × g for 10 min. The cells were suspended in 20 mM sodium phosphate (pH 7.5) containing 10 mM EDTA and 1 mM *o*-phenanthroline and lysed by two passages through a French pressure cell. Unbroken cells and debris were removed by centrifugation for 10 min at 2,500 × g. The cell extract was centrifuged for 60 min at 223,160 × g or for 30 min at 352,900 × g. The supernatant was retained as the soluble fraction, and the pellet was retained as the membrane fraction. The membranes were fractionated further essentially as described previously (12) by partial solubilization in 5 g of Sarkosyl per liter for 30 min at 25°C, followed by centrifugation for 60 min at 223,160 × g. All steps of the fractionation were carried out at 0 to 4°C unless indicated otherwise.

Cell fractions were analyzed by Western blotting to determine the localization of GldB and GldC. Total-protein profiles of cell fractions were examined by SDS-PAGE followed by staining with Coomassie blue (37). Cytochrome *c* (cytoplasmic membrane marker) was detected by SDS-PAGE followed by staining with 3,3',5,5'-tetramethyl benzidine dihydrochloride for heme-associated peroxidase activity as described by Thomas (46), except that 5% β-mercaptoethanol was included in the loading buffer. Lipopolysaccharide (LPS) (outer membrane marker) was detected by digestion with protease K, separation by Tricine-SDS-PAGE, and silver staining as described previously (43).

Microscopic observations of cell movement and bead movement. Wild-type and mutant cells of *F. johnsoniae* were examined for movement over glass and agar surfaces by phase-contrast microscopy at 25°C using two different assays. Cells were grown to late exponential phase (about 3 × 10⁹ cells/ml) in CYE broth. To observe movement over glass, a drop of the cell culture was placed on a microscope slide, covered with an oxygen-permeable membrane (Yellow Springs Instrument Co., Inc., Yellow Springs, Ohio), which functioned as a coverslip, and examined by phase-contrast microscopy. To observe movement of cells on agar, 5 μl of the cell culture was spotted onto a layer of PY2 agar on a microscope slide. The spot was allowed to partially dry (less than 5 min), after which it was covered with an oxygen-permeable membrane. After 1 h of incubation at 25°C in a moist chamber, the cells were examined by time-lapse videomicroscopy at 25°C. The videotape was played back at 60 times the recording speed to allow detection of slow or intermittent movements. We examined the ability of cells to bind to and propel 0.5-μm polystyrene latex spheres (Seradyn, Indianapolis, Ind.) as previously described (2).

Measurements of phage sensitivity. Sensitivity to *F. johnsoniae* phage was determined essentially as previously described by spotting 10 μl of phage lysates (2 × 10⁷ phage/ml) onto lawns of cells in CYE overlay agar (50). The plates were incubated for 24 h at 25°C to observe lysis.

Genetic nomenclature. Genes involved in gliding motility were given the name *gld* followed by a letter. Open reading frames (ORFs) of unknown function that did not exhibit strong similarity to previously described genes were given the provisional name *fjo* (for "*Flavobacterium johnsoniae* open reading frame") followed by a number.

Nucleotide sequence accession number. The sequence reported in this paper has been deposited in the GenBank database (accession no. AF158372).

RESULTS AND DISCUSSION

Construction of the *F. johnsoniae*-*E. coli* shuttle cosmids pCP22 and pCP26. The *F. johnsoniae*-*E. coli* shuttle cosmid pCP17 was recently developed as a vector to allow the construction of libraries of *F. johnsoniae* DNA (27). Libraries




FIG. 2. Map of the *gldBC* region. The sites of the Tn4351 insertions in CJ562, CJ567, CJ569, CJ578, and CJ683 are indicated by triangles. Regions of *F. johnsoniae* DNA contained in pDH223, pDH233, and pDH246 are shown beneath the map.

produced in pCP17 can be propagated in *E. coli* and can be transferred into *F. johnsoniae* by conjugation. pCP17 was used to identify the *gldA* gene of *F. johnsoniae*, which restored motility to several nongliding mutants (2). pCP17 has one major drawback as a tool for cloning *F. johnsoniae* genes; it has a very high copy number in *E. coli* (greater than 100 copies per cell). This makes it difficult to use pCP17 to create random genomic libraries, since many genomic fragments of *F. johnsoniae* DNA appear to be toxic to *E. coli* when present in many copies (unpublished observations). For this reason, we constructed two new shuttle cosmids, pCP22 and pCP26 (Fig. 1), as described in Materials and Methods. These vectors carry the RSF1010 origin of replication, which yields copy numbers of approximately 10 in *E. coli* (16). Both vectors rely on the pCP1 (27) origin to replicate in *F. johnsoniae* (copy number, approximately 10/cell).

Complementation of nongliding mutants. To identify and clone genes involved in gliding motility, we constructed a library of wild-type *F. johnsoniae* DNA in pCP26 as described in Materials and Methods. The cosmid library was transferred to the nongliding mutant UW102-99, and we identified one colony that exhibited spreading. The cells of this colony carried the cosmid clone pCP200, which had a 26-kb insert of *F. johnsoniae* DNA. We introduced pCP200 into UW102-99 by electroporation and verified that pCP200 complemented the gliding motility defects of this mutant. All of the Em^r colonies carrying pCP200 exhibited spreading, and individual cells exhibited rapid movement over glass or agar surfaces when examined by phase-contrast microscopy. pCP200 was introduced into each of 50 independently isolated nongliding mutants (listed in Materials and Methods). pCP200 complemented UW102-90, UW102-99, UW102-103, and UW102-154 but did not complement the other 46 mutants. UW102-99 was originally isolated by Chang et al. (8) as a nongliding mutant following nitrosoguanidine mutagenesis. The other three mutants were isolated as spontaneous mutants by selecting for resistance to phage (8).

Subcloning of pCP200 and analysis of the *gldBC* sequence. The insert DNA in pCP200 was subcloned to determine which region was required for complementation. A 3.4-kb *Xba*I-*Sal*I fragment of pCP200 was found to be sufficient for complementation of UW102-90, UW102-99, UW102-103, and UW102-154. This region was sequenced, and five ORFs (*fjo5*, *gldB*, *gldC*, *fjo6*, and *fjo7*) were identified (Fig. 2). Further subcloning revealed that the 1.9-kb *Xba*I-*Cl*aI fragment carried by pDH223, which spanned only the *gldB* and *gldC* genes, was sufficient for complementation. pDH223 restored full motility to UW102-90, UW102-99, UW102-103, and UW102-154.

gldB and *gldC* appear to be organized in an operon, with the

translational stop codon of *gldB* overlapping the predicted translational start site of *gldC*. The 109-bp region that lies immediately upstream of *gldB* does not contain any obvious ORFs. We assume that the *gldBC* promoter lies in this region. There is currently no reliable information to allow the identification of putative *Flavobacterium* promoters by analysis of DNA sequences. The region upstream of *gldB* is relatively AT rich, however (77% A+T over the 109 nucleotides, compared to 66% A+T for the entire 3.4-kb *Xba*I-*Sal*I fragment), as is often observed in promoter regions of other bacteria (36). Immediately downstream of *gldC* is a 359-bp region that does not contain any obvious ORFs. An inverted repeat structure that is followed by a string of T's (AAAATCCCAAATTCCA ATTGCAAAACTGGAATTTGGGATTTTTTTT) is found 18 nucleotides downstream of the *gldC* stop codon. An additional inverted repeat (CATAAAAAAAAAACGCTGTAAAAC TCAAGTCTTACAGCGTTTTTTTTATG) lies 76 bp downstream of the *gldC* stop codon. These sequences may play a role in transcriptional termination or mRNA stabilization.

gldB is predicted to code for a 38.3-kDa protein. The GldB protein has two relatively hydrophobic stretches (amino acids 4 to 16 and amino acids 133 to 148). The first hydrophobic stretch has features of a signal peptide (48). Both hydrophobic regions are relatively short, but their presence tentatively suggests that GldB may be a membrane protein. GldB exhibits weak but potentially significant similarity to the *Clostridium thermocellum* β -glucanase LicA (GenBank accession no. CAA61884) (27% identical over 105 amino acids) and to the putative ADP-L-glycero-D-manno-heptose-6-epimerase *rfaD* product from *Aquifex aeolicus* (10) (23% identical over 195 amino acids). Given these weak similarities, it is difficult to predict the function of GldB in gliding motility. Since both of the proteins mentioned are thought to interact with carbohydrates, it is possible that GldB interacts with, or is involved in production of, polysaccharides or glycoproteins. Polysaccharides and glycoproteins have previously been proposed to play roles in gliding motility (15, 19, 29, 31).

gldC is predicted to code for a protein of 12.9 kDa, which appears to be relatively hydrophilic. GldC exhibits weak similarity to the fungal lectin AOL from *Arthrobotrys oligospora* (35) (34% identity over 62 amino acid residues). This level of similarity does not allow us to predict with any certainty the function of GldC but may indicate that it interacts with carbohydrates.

Analysis of the region surrounding *gldBC*. *fjo5* lies upstream of *gldB* and is oriented in the opposite direction (Fig. 2). We do not know the entire sequence of this putative gene, but analysis of the first 216 nucleotides of *fjo5* indicates that its protein product is similar to the amino terminus of a putative ATP pyrophosphatase of *Mycoplasma capricolum* (38% identity over 54 amino acid residues) (6) and to several putative NH₃-dependent NAD synthetases such as *nadE* from *Archaeoglobus fulgidus* (38% identity over 42 amino acid residues) (21). *fjo6* and *fjo7* lie downstream of *gldC*. The predicted protein products of *fjo6* and *fjo7* are not very similar to protein sequences in the databases. We do not have any evidence that *fjo5*, *fjo6*, and *fjo7* are involved in gliding motility.

Characterization of point mutations in *gldB*. To determine the nature of the mutations in the four mutants UW102-90, UW102-99, UW102-103, and UW102-154, we amplified the *gldBC* region from each mutant by PCR and determined the nucleotide sequences. Each mutant contained a single-base change within *gldB*. The mutations in UW102-90 (A612T; numbered starting from the first nucleotide of the *gldB* start codon) and UW102-154 (G375T) introduced translational stop codons at amino acid positions 205 and 126, respectively. These

mutations completely disrupted gliding motility. UW102-99 carried a mutation (C820T) resulting in a proline-to-leucine change at position 274. This mutation also completely disrupted gliding motility. The mutation in UW102-103 (T424G) resulted in a valine-to-glycine change at position 142. In the course of this analysis, we determined that UW102-103 was not completely deficient in gliding motility. Colonies of UW102-103 did not spread, but individual cells occasionally exhibited brief gliding movements over glass or agar surfaces as observed by microscopic analysis. These motile cells were probably not revertants, since we observed the same result with cells that were derived from a freshly isolated nonspreading colony of UW102-103. Apparently the valine-to-glycine mutation at position 142 results in a defective GldB protein but does not completely disrupt the ability of the protein to function in gliding motility.

Isolation of Tn4351 induced *gldB* mutants. In an independent attempt to identify gliding-motility genes, we isolated nongliding mutants following Tn4351 mutagenesis. A total of 211 mutants which formed nonspreading colonies were isolated; of these, 161 exhibited some gliding motility when movement of single cells was examined by phase-contrast microscopy. The remaining 50 mutants were nongliding (colonies failed to spread, and individual cells failed to glide over glass or agar surfaces). Of these, 30 exhibited a filamentous phenotype characteristic of a cell division defect whereas the remaining 20 had normal cell morphology. We examined these 20 nongliding mutants to determine whether any of them carried insertions in the *gldBC* region. Introduction of pDH233, which carries the wild-type *gldBC* genes, restored motility to 5 of the 20 mutants as measured by colony spreading and by observations of single-cell motility. The exact sites of insertion of Tn4351 in the five mutants were determined by cloning the disrupted genes or by amplification of the regions surrounding the transposons from chromosomal DNA and sequencing the cloned or amplified DNA. The five insertions were all unique, and all occurred within the *gldB* gene (Fig. 2). *gldB* appears to be a hot spot for transposition by Tn4351. While insertions in *gldB* accounted for 5 of the 20 Tn4351 induced nongliding mutants, insertions in *gldA* were responsible for only 2 mutations. Of the remaining 13 mutants, 2 had insertions in a gene that we refer to as *gldD* (M. J. McBride and D. W. Hunnicutt, unpublished data) and the rest had insertions that each appeared to be in other ORFs. *gldA*, *gldB*, and *gldD* do not appear to be closely linked on the *F. johnsoniae* genome.

***gldB* and *gldC* appear to be organized as an operon.** Sequence analysis suggested that *gldB* and *gldC* were arranged as an operon. To test this prediction, we constructed the *gldB* mutant CJ588 by recombining pDH227 into the chromosomal copy of *gldB* as described in Materials and Methods. Antibodies were raised against recombinant GldB and GldC proteins and were used to detect GldB and GldC in extracts of wild-type and mutant cells by Western blot analyses. Antibodies against recombinant GldB recognized GldB protein in wild-type cells but not in CJ588 cells (Fig. 3A, lanes 1 and 2). Similarly, antibodies against recombinant GldC recognized GldC protein in wild-type cells but not in CJ588 cells (Fig. 3B, lanes 1 and 2). The polarity of the insertion in *gldB* on expression of *gldC* indicates that *gldB* and *gldC* probably are cotranscribed as an operon, as predicted by sequence analysis. Surprisingly, none of the Tn4351 *gldB* mutations were polar on *gldC* (Fig. 3B, lanes 5 and 6, and data not shown). Approximately wild-type levels of GldC were produced by each of the Tn4351-induced *gldB* mutants. The mutations included examples of insertion of Tn4351 in either orientation. It has been previously reported that Tn4351 has promoters reading out of both sides that




FIG. 3. Western immunoblot detection of GldB and GldC in wild-type and mutant strains of *F. johnsoniae*. Cells were lysed in SDS-PAGE loading buffer, and proteins were separated by SDS-PAGE and transferred to PVDF membranes. GldB (A) and GldC (B) were detected using antisera raised against recombinant GldB or GldC proteins, respectively, as described in Materials and Methods. Lanes: 1, wild-type *F. johnsoniae*; 2, *gldB* insertional knockout mutant CJ588; 3, CJ767 (CJ588 complemented with pDH246, which carries *gldB*); 4, CJ594 (CJ588 complemented with pDH233, which carries *gldB* and *gldC*); 5, Tn4351-induced *gldB* mutant CJ562; 6, Tn4351-induced *gldB* mutant CJ567.

function in *Bacteroides fragilis* (41). The recognition of these promoters by *F. johnsoniae* RNA polymerase probably explains the lack of polarity of the Tn4351-induced *gldB* mutations.

GldB is required for gliding motility. Cells of wild-type *F. johnsoniae* move over glass or agar surfaces and form spreading colonies on PY2 agar (Fig. 4A). CJ588 carries an insertion of the plasmid pDH227 in the chromosomal copy of *gldB* and failed to make detectable levels of GldB or GldC protein (Fig. 3, lanes 2). Cells of CJ588 did not move over glass or agar surfaces and formed colonies that did not spread (Fig. 4B). Introduction of pDH233, which carries the *gldB* and *gldC* genes, into CJ588 restored its ability to produce the GldB and GldC proteins (Fig. 3, lanes 4) and resulted in complementation of the motility defect. Individual cells displayed gliding movements, and colonies spread nearly as well as those of wild-type cells (Fig. 4C). Introduction of pDH246, which carries only the *gldB* gene, restored production of GldB protein but not of GldC protein (Fig. 3, lanes 3) and resulted in partial restoration of gliding motility and colony spreading. Individual cells exhibited gliding movements indistinguishable from those of wild-type cells, but colonies spread less well than those of the wild type or of CJ588 complemented with pDH233 (Fig. 4D). This indicates that GldC is not absolutely required for gliding motility but that the presence of GldC enhances colony spreading.

Cellular localization of GldB and GldC. Sequence analysis suggested that GldB may be a membrane protein while GldC is probably a soluble protein. To test this, cell extracts were separated into membrane and soluble fractions by ultracentrifugation. The cell fractions were analyzed by SDS-PAGE followed by Western immunoblotting using antisera against recombinant GldB or GldC. As shown in Fig. 5, GldB localized to the membrane fraction while GldC was found in the soluble fraction, as predicted.

Differential solubilization of cytoplasmic and outer membrane components by Sarkosyl was used to determine the location of GldB. This method relies on the resistance of outer




FIG. 4. Photomicrographs of *F. johnsoniae* colonies. Colonies were grown for 2 days at 25°C on PY2 agar medium. Photomicrographs were taken with an Olympus OM-4T camera mounted on a Nikon Diaphot inverted phase-contrast microscope. (A) Wild-type *F. johnsoniae* UW101. (B) *gldBC* knockout mutant CJ588. (C) CJ594 (CJ588 complemented with pDH233, which carries *gldBC*). (D) CJ767 (CJ588 complemented with pDH246, which carries *gldB*). Bar, 1 mm.

membranes to solubilization by Sarkosyl (12). This procedure was chosen since it has proven difficult to separate the cytoplasmic and outer membranes of *F. johnsoniae* ATCC 17061 by standard gradient centrifugation methods (14). As shown in Fig. 6, Sarkosyl solubilization appears to be an effective method to fractionate cytoplasmic membrane (Sarkosyl soluble) and outer membrane (Sarkosyl insoluble) components of *F. johnsoniae*. Cytochrome *c* was used as a cytoplasmic membrane marker. *c*-type cytochromes are generally found in the cytoplasmic membrane or periplasm of respiratory gram-negative bacteria (47). *c*-type cytochromes are unusual in that they have covalently attached heme (47). They can be detected by sepa-

rating proteins by SDS-PAGE and staining for heme (34). A band corresponding to a *c*-type cytochrome was detected in the Sarkosyl-soluble (inner membrane) fraction, but none were detected in the Sarkosyl-insoluble (outer membrane) fraction (Fig. 6B). In contrast, LPS was detected only in the Sarkosyl-insoluble (outer membrane) fraction (Fig. 6C, lane 2). These results confirm that differential solubilization in Sarkosyl appears to fractionate the membranes of *F. johnsoniae* into cytoplasmic membrane and outer membrane fractions. Western blot analysis of these fractions indicated that GldB was present in the Sarkosyl-soluble fraction (Fig. 6D, lane 2). This suggests that GldB resides in the cytoplasmic membrane.

Phage resistance of *gldB* mutants. Many nongliding mutants of *F. johnsoniae* are resistant to infection by a number of *F. johnsoniae* bacteriophages (49). The reason for this pleiotropy is not known. It has been suggested that these phages are able to infect only cells which have an actively moving surface (8). The moving components of the cell surface may be needed directly for adsorption of phage or for uptake of phage nucleic acid into the cell. Alternatively, the adsorption sites on the cell surface may be covered by a layer of polysaccharide or other material in nonmotile mutants. Motile cells may keep this material moving, thus transiently exposing the sites beneath for viral adsorption. We tested the sensitivity of *F. johnsoniae* strains UW101, CJ588, and CJ767 (CJ588 complemented with pDH246) to the *F. johnsoniae* bacteriophages ϕ Cj1, ϕ Cj7, ϕ Cj13, ϕ Cj23, ϕ Cj28, ϕ Cj29, ϕ Cj42, ϕ Cj48 and ϕ Cj54. *F. johnsoniae* UW101 was readily lysed by these phages, whereas the nongliding mutant CJ588 was not lysed. Introduction of pDH246 into CJ588 restored sensitivity to each of these phages




FIG. 5. Localization of GldB and GldC. Cells were grown to mid-exponential phase, concentrated by centrifugation, and disrupted by passage through a French pressure cell. Unbroken cells and debris were removed by centrifugation at $2,500 \times g$ for 10 min. Insoluble material was pelleted from the supernatant fraction by centrifugation at $352,900 \times g$ for 30 min. The pellet was dissolved in 10 mM Tris buffer (pH 7.5), and the proteins in both the $352,900 \times g$ pellet (lanes M) and supernatant (lanes S) fractions were separated by SDS-PAGE and transferred to PVDF membranes. GldB (A) and GldC (B) were detected using antisera raised against recombinant GldB or GldC proteins, respectively.




FIG. 6. Localization of GldB to the Sarkosyl-soluble (cytoplasmic membrane) fraction of cells. Cells were disrupted and separated into soluble and membrane fractions. Membranes were fractionated further by differential solubilization in Sarkosyl as described in Materials and Methods. Equal amounts of each fraction were separated by SDS-PAGE and examined for total protein (A), cytochrome *c* (cytoplasmic membrane marker) (B), LPS (outer membrane marker) (C), and GldB protein (D). (A) Coomassie blue-stained gel to detect total protein. Lanes: 1, soluble (cytoplasmic and periplasmic) fraction; 2, Sarkosyl-soluble (cytoplasmic membrane) fraction; 3, Sarkosyl-insoluble (outer membrane) fraction. (B) Heme-stained gel to detect cytochrome *c*. Lanes: 1, soluble (cytoplasmic and periplasmic) fraction; 2, Sarkosyl-soluble (cytoplasmic membrane) fraction; 3, Sarkosyl-insoluble (outer membrane) fraction. (C) Silver-stained gel to detect LPS. Lanes: 1, Sarkosyl-soluble (cytoplasmic membrane) fraction; 2, Sarkosyl-insoluble (outer membrane) fraction. (D) Western blot analysis to detect GldB. Lanes: 1, soluble (cytoplasmic and periplasmic) fraction; 2, Sarkosyl-soluble (cytoplasmic membrane) fraction; 3, Sarkosyl-insoluble (outer membrane) fraction.

in addition to restoring gliding motility. These results are essentially identical to those observed for *gldA* mutants (2).

Movement of latex spheres by *gldB* mutant cells. Wild-type cells of *F. johnsoniae* in liquid suspension bind latex spheres on their surfaces and propel them along the length of the cells (23, 30). The spheres appear to move along multiple tracks. They may change their direction of movement, and two spheres near each other on the same cell may move in the same direction or in different directions. Spheres move at approximately the same speed as cells move over surfaces, which suggests that sphere movement is related to gliding motility.

We examined the ability of cells of *F. johnsoniae* strains UW101, CJ588, CJ767 (CJ588 complemented with pDH246), and CJ594 (CJ588 complemented with pDH233) to bind and propel latex spheres. Wild-type cells bound and propelled the spheres. Cells of the *gldB* mutant CJ588 were rarely observed to bind the spheres and were never observed to propel them. Cells of CJ767 and CJ594 bound and propelled the spheres as well as wild-type cells did. These results are essentially identical to those observed for *gldA* mutants (2).

Models to explain *F. johnsoniae* gliding motility. A number of models have been proposed to explain bacterial gliding motility (7, 19, 23, 29, 42, 51). It is possible (perhaps likely) that different mechanisms function in different groups of bacteria. *F. johnsoniae* gliding motility is thought to be powered by proton motive force (30). It is also known that exocellular or cell surface polysaccharides are important for gliding motility of *F. johnsoniae* and other unrelated bacteria (13, 15, 29, 51). These may play a passive role in mediating productive contact of *F. johnsoniae* cells with the substratum. Sulfonolipids, which are localized in the outer membrane, have also been demonstrated to play a role in *F. johnsoniae* gliding motility (1, 14). The available evidence supports a model of *F. johnsoniae* motility that involves the movement of macromolecules in or along the outer membrane along tracks that may be fixed to the peptidoglycan (23). Additional components in the periplasm and cell membrane are postulated to harvest the proton motive force and perform the work necessary to move these macromolecules. The outer membrane molecules may be proteins or glycoproteins. Others have postulated that extrusion of polysaccharide could directly propel cells (19). This would seem to

be an unlikely mechanism for *F. johnsoniae* gliding motility. Cells of *F. johnsoniae* in nonnutrient buffer remain actively motile for hours (8). It is unlikely that cells could make sufficient polysaccharide from their cellular reserves to continually propel themselves for such extensive periods by simple extrusion of polysaccharide.

The mechanism of *F. johnsoniae* gliding motility remains unclear. We previously identified one gene, *gldA*, which is required for gliding motility (2). In this paper we have identified two additional genes that are involved in gliding motility. *gldB* is required for *F. johnsoniae* gliding motility, but the exact role that GldB plays in this process is not known. GldC enhances colony spreading but is not absolutely required for gliding motility. GldB and GldC exhibit slight similarity to proteins that interact with carbohydrates. Polysaccharides and glycoproteins have been proposed to play roles in gliding motility, and so proteins required for gliding motility that are involved in polysaccharide synthesis, glycoprotein modification, or binding to carbohydrates might be expected. The weak similarities that GldB and GldC exhibit to their potential homologs prevent us from drawing strong conclusions regarding the functions of these proteins in gliding motility. The identification of components of the gliding-motility apparatus that do not exhibit strong similarity to proteins in the databases is not surprising. The mechanism of *F. johnsoniae* gliding motility is not understood, and it may be unrelated to (or distantly related to) processes that occur in other bacteria that have been extensively studied. Further studies of GldA, GldB, and GldC and of the other proteins that are involved in *F. johnsoniae* cell movement should help to clarify the mechanism of gliding motility.

ACKNOWLEDGMENTS

This research was supported by grants from the National Science Foundation (MCB-9418308 and MCB-9727825) and by a Shaw Scientist Award to M.M. from The Milwaukee Foundation.

Some of the Tn4351 induced mutants were isolated by M. Kempf and S. Agarwal. DNA sequencing was performed by the Automated DNA Sequencing Facility of the University of Wisconsin—Milwaukee Department of Biological Sciences. We thank D. Saffarini for careful reading of the manuscript.

REFERENCES

- Abbanat, D. R., E. R. Leadbetter, W. Godchaux III, and A. Escher. 1986. Sulphonolipids are molecular determinants of gliding motility. *Nature* **324**: 367–369.
- Agarwal, S., D. W. Hunnicutt, and M. J. McBride. 1997. Cloning and characterization of the *Flavobacterium johnsoniae* (*Cytophaga johnsonae*) gliding motility gene, *gldA*. *Proc. Natl. Acad. Sci. USA* **94**:12139–12144.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Bernardet, J.-F., P. Segers, M. Vancanney, F. Berthe, K. Kersters, and P. Vandamme. 1996. Cutting a gordian knot: emended classification and description of the genus *Flavobacterium*, and proposal of *Flavobacterium hydatitis* nom. nov. (basonym, *Cytophaga aquatilis* Strohl and Tait 1978). *Int. J. Syst. Bacteriol.* **46**:128–148.
- Bolivar, F., and K. Backman. 1979. Plasmids of *E. coli* as cloning vectors. *Methods Enzymol.* **68**:245–267.
- Bork, P., C. Ouzounis, G. Casari, R. Schneider, C. Sander, M. Dolan, W. Gilbert, and P. M. Gillevet. 1995. Exploring the *Mycoplasma capricolum* genome: a minimal cell reveals its physiology. *Mol. Microbiol.* **5**:955–967.
- Burchard, R. P. 1981. Gliding motility of prokaryotes: ultrastructure, physiology, and genetics. *Annu. Rev. Microbiol.* **35**:497–529.
- Chang, L. Y. E., J. L. Pate, and R. J. Betzig. 1984. Isolation and characterization of nonspreading mutants of the gliding bacterium *Cytophaga johnsonae*. *J. Bacteriol.* **159**:26–35.
- Cooper, A. J., A. P. Kalinowski, N. B. Shoemaker, and A. A. Salyers. 1997. Construction and characterization of a *Bacteroides thetaiotamicron* *recA* mutant: transfer of *Bacteroides* integrated conjugative elements is *RecA* independent. *J. Bacteriol.* **179**:6221–6227.
- Deckert, G., P. V. Warren, T. Gaasterland, W. G. Young, A. L. Lenox, D. E. Graham, R. Overbeek, M. A. Snead, M. Keller, M. Aujay, R. Huber, R. A. Feldman, J. M. Short, G. J. Olsen, and R. V. Swanson. 1998. The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*. *Nature* **392**: 353–358.
- Ferrante, A. A. 1992. Ph. D. thesis. University of Massachusetts—Amherst, Amherst.
- Filip, C., G. Fletcher, J. L. Wulff, and C. F. Earhart. 1973. Solubilization of the cytoplasmic membrane of *Escherichia coli* by the ionic detergent sodium-lauryl sarcosinate. *J. Bacteriol.* **115**:717–722.
- Godchaux, W., III, L. Gorski, and E. R. Leadbetter. 1990. Outer membrane polysaccharide deficiency in two nongliding mutants of *Cytophaga johnsonae*. *J. Bacteriol.* **172**:1250–1255.
- Godchaux, W., III, and E. R. Leadbetter. 1988. Sulfolipids are localized in the outer membrane of the gliding bacterium *Cytophaga johnsonae*. *Arch. Microbiol.* **150**:42–47.
- Godchaux, W., III, M. A. Lynes, and E. R. Leadbetter. 1991. Defects in gliding motility in mutants of *Cytophaga johnsonae* lacking a high-molecular-weight cell surface polysaccharide. *J. Bacteriol.* **173**:7607–7614.
- Guerry, P., J. van Embden, and S. Falkow. 1974. Molecular nature of two nonconjugative plasmids carrying drug resistance genes. *J. Bacteriol.* **117**: 619–630.
- Guzman, L.-M., D. Belin, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation and high-level expression by vectors containing the arabinose pBAD promoter. *J. Bacteriol.* **177**:4121–4230.
- Hedges, R. W. 1974. Transposition of ampicillin resistance from RP4 to other replicons. *Mol. Gen. Genet.* **132**:31–40.
- Hoiczky, E., and W. Baumeister. 1998. The junctional pore complex, a prokaryotic secretion organelle, is the molecular motor underlying gliding motility in cyanobacteria. *Curr. Biol.* **8**:1161–1168.
- Huang, X., and W. Miller. 1991. A time-efficient, linear-space local similarity algorithm. *Adv. Appl. Math.* **12**:337–357.
- Klenk, H.-P., R. A. Clayton, J.-F. Tomb, O. White, K. E. Nelson, K. A. Ketchum, R. J. Dodson, M. Gwinn, E. K. Hickey, J. D. Peterson, D. L. Richardson, A. R. Kerlavage, D. E. Graham, N. C. Kyrpides, R. D. Fleischmann, J. Quackenbush, N. H. Lee, G. G. Sutton, S. Gill, E. F. Kirkness, B. A. Dougherty, K. McKenney, M. D. Adams, B. Loftus, S. Peterson, C. I. Reich, L. K. McNeil, J. H. Badger, A. Glodek, L. Zhou, R. Overbeek, J. D. Gocayne, J. F. Weidman, L. McDonald, T. Utterback, M. D. Cotton, T. Spriggs, P. Atriach, B. P. Kaine, S. M. Sykes, P. W. Sadow, K. P. D'Andrea, C. Bowman, C. Fujii, S. A. Garland, T. M. Mason, G. J. Olsen, C. M. Fraser, H. O. Smith, C. R. Woese, and J. C. Venter. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**:364–370.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680–685.
- Lapidus, I. R., and H. C. Berg. 1982. Gliding motility of *Cytophaga* sp. strain U67. *J. Bacteriol.* **151**:384–398.
- Lee, C., A. Levin, and D. Branton. 1987. Copper staining: a five minute protein stain for sodium dodecyl sulfate-polyacrylamide gels. *Anal. Biochem.* **166**:308–312.
- Li, L.-Y., N. B. Shoemaker, and A. A. Salyers. 1995. Location and characterization of the transfer region of a *Bacteroides* conjugative transposon and regulation of the transfer genes. *J. Bacteriol.* **177**:4992–4999.
- McBride, M. J., and S. A. Baker. 1996. Development of techniques to genetically manipulate members of the genera *Cytophaga*, *Flavobacterium*, *Flexibacter*, and *Sporocytophaga*. *Appl. Environ. Microbiol.* **62**:3017–3022.
- McBride, M. J., and M. J. Kempf. 1996. Development of techniques for the genetic manipulation of the gliding bacterium *Cytophaga johnsonae*. *J. Bacteriol.* **178**:583–590.
- Metcalf, W. W., W. Jiang, L. L. Daniels, S.-K. Kim, A. Haldimann, and B. L. Wanner. 1996. Conditionally replicative and conjugative plasmids carrying *lacZα* for cloning, mutagenesis, and allele replacement in bacteria. *Plasmid*. **35**:1–13.
- Pate, J. L. 1988. Gliding motility. *Can. J. Microbiol.* **34**:459–465.
- Pate, J. L., and L.-Y. E. Chang. 1979. Evidence that gliding motility in prokaryotic cells is driven by rotary assemblies in the cell envelopes. *Curr. Microbiol.* **2**:59–64.
- Pate, J. L., and D. M. De Jong. 1990. Use of nonmotile mutants to identify a set of membrane proteins related to gliding motility in *Cytophaga johnsonae*. *J. Bacteriol.* **172**:3117–3124.
- Pate, J. L., S. J. Petzold, and L.-Y. E. Chang. 1979. Phages for the gliding bacterium *Cytophaga johnsonae* that infect only motile cells. *Curr. Microbiol.* **2**:257–262.
- Pearson, W. R. 1990. Rapid and sensitive sequence comparison with FASTP and FASTA. *Methods Enzymol.* **183**:63–98.
- Ritz, D., M. Bott, and H. Hennecke. 1993. Formation of several bacterial c-type cytochromes requires a novel membrane-anchored protein that faces the periplasm. *Mol. Microbiol.* **9**:729–740.
- Rosen, S., M. Kata, Y. Persson, P. H. Lipniunas, C. Wikstrom, C. Van den Hondel, J. Van den Brink, L. Rask, L. O. Heden, and A. Tunlid. 1996. Molecular characterization of a saline-soluble lectin from a parasitic fungus: extensive sequence similarities between fungal lectins. *Eur. J. Biochem.* **238**:822–829.
- Ross, W., K. K. Gosink, J. Salomon, K. Igarashi, C. Zou, A. Ishahama, K. Severinov, and R. L. Gourse. 1993. A third recognition element in bacterial promoters: DNA binding by the α subunit of RNA Polymerase. *Science* **262**:1407–1413.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Schagger, H., and G. von Jagow. 1987. Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal. Biochem.* **166**:368–379.
- Shoemaker, N. B., C. Getty, J. F. Gardner, and A. A. Salyers. 1986. Tn4351 transposes in *Bacteroides* spp. and mediates the integration of plasmid R751 into the *Bacteroides* chromosome. *J. Bacteriol.* **165**:929–936.
- Simon, R., U. Priefer, and A. Puhler. 1983. A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in Gram negative bacteria. *Bio/Technology* **2**:784–791.
- Smith, C. J., M. B. Rogers, and M. L. McKee. 1992. Heterologous gene expression in *Bacteroides fragilis*. *Plasmid* **27**:141–154.
- Spormann, A. M. 1999. Gliding motility in bacteria: Insights from studies of *Myxococcus xanthus*. *Microbiol. Mol. Biol. Rev.* **63**:621–641.
- Sprott, G. D., S. F. Koval, and C. A. Schnaitman. 1994. Cell fractionation, p. 72–103. In P. Gerhardt, R. G. E. Murray, W. A. Wood, and N. R. Krieg (ed.), *Methods for general and molecular bacteriology*. American Society for Microbiology, Washington, D.C.
- Stevens, A. M., N. B. Shoemaker, and A. A. Salyers. 1996. The region of *Bacteroides* conjugal chromosomal tetracycline resistance elements which is responsible for production of plasmidlike form from unlinked chromosomal DNA might also be involved in transfer of the element. *J. Bacteriol.* **172**: 4271–4279.
- Thaxter, R. 1892. On the Myxobacteriaceae, a new order of Schizomycetes. *Bot. Gaz.* **17**:389–406.
- Thomas, P. E. 1976. An improved staining procedure for the detection of the peroxidase activity of cytochrome P-450 on sodium dodecyl sulfate polyacrylamide gels. *Anal. Biochem.* **75**:168–176.
- Thony-Meyer, L. 1997. Biogenesis of respiratory cytochromes in bacteria. *Microbiol. Mol. Biol. Rev.* **61**:337–376.
- von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* **14**:4683–4690.
- Wolkin, R. H., and J. L. Pate. 1986. Phage adsorption and cell adherence are motility-dependent characteristics of the gliding bacterium *Cytophaga johnsonae*. *J. Gen. Microbiol.* **132**:355–367.
- Wolkin, R. H., and J. L. Pate. 1985. Selection for nonadherent or nonhydrophobic mutants co-selects for nonspreading mutants of *Cytophaga johnsonae* and other gliding bacteria. *J. Gen. Microbiol.* **131**:737–750.
- Yoderian, P. 1998. Bacterial motility: secretory secrets of gliding bacteria. *Curr. Biology*. **8**:R408–R411.