

The UspA1 Protein and a Second Type of UspA2 Protein Mediate Adherence of *Moraxella catarrhalis* to Human Epithelial Cells In Vitro

ERIC R. LAFONTAINE,¹ LESLIE D. COPE,¹ CHRISTOPH AEBI,^{1,2†} JO L. LATIMER,¹
GEORGE H. MCCrackEN, JR.,² AND ERIC J. HANSEN^{1*}

*Departments of Microbiology¹ and Pediatrics,² University of Texas Southwestern Medical Center,
Dallas, Texas 75235-9048*

Received 3 September 1999/Accepted 30 November 1999

The UspA1 and UspA2 proteins of *Moraxella catarrhalis* are structurally related, are exposed on the bacterial cell surface, and migrate as very high-molecular-weight complexes in sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Previous analysis of *uspA1* and *uspA2* mutants of *M. catarrhalis* strain 035E indicated that UspA1 was involved in adherence of this organism to Chang conjunctival epithelial cells in vitro and that expression of UspA2 was essential for resistance of this strain to killing by normal human serum (C. Aebi, E. R. Lafontaine, L. D. Cope, J. L. Latimer, S. R. Lumbley, G. H. McCracken, Jr., and E. J. Hansen, *Infect. Immun.* 66:3113–3119, 1998). In the present study, isogenic *uspA1*, *uspA2*, and *uspA1 uspA2* mutations were constructed in three additional *M. catarrhalis* strains: 012E, TTA37, and 046E. The *uspA1* mutant of strain 012E had a decreased ability to attach to Chang cells. However, inactivation of the *uspA1* gene in both strain TTA37 and strain 046E did not cause a significant decrease in attachment ability. Inactivation of the *uspA2* gene of strain TTA37 did result in a loss of attachment ability. Nucleotide sequence analysis revealed that the predicted protein encoded by the *uspA2* genes of both strains TTA37 and 046E had a N-terminal half that resembled the N-terminal half of UspA1 proteins, whereas the C-terminal half of this protein was nearly identical to those of previously characterized UspA2 proteins. The gene encoding this “hybrid” protein was designated *uspA2H*. PCR-based analysis revealed that approximately 20% of *M. catarrhalis* strains apparently possess a *uspA2H* gene instead of a *uspA2* gene. The *M. catarrhalis* *uspA1*, *uspA2*, and *uspA2H* genes were cloned and expressed in *Haemophilus influenzae* cells, which were used to prove that both the UspA1 and UspA2H proteins can function as adhesins in vitro.

Moraxella catarrhalis, an unencapsulated, gram-negative bacterium, can cause disease in both the upper and lower respiratory tracts (32). It has been estimated that approximately 20% of cases of acute bacterial otitis media in infants and young children are caused by this organism (6). *M. catarrhalis* is also associated with nearly one-third of infectious exacerbations of chronic obstructive pulmonary disease in adults (16). The ability of this organism to cause significant morbidity has resulted in increased efforts to develop an efficacious *M. catarrhalis* vaccine (35).

Outer membrane proteins have received the most attention as possible *M. catarrhalis* vaccine candidates (9, 19, 20, 31, 33, 43), and even *M. catarrhalis* lipooligosaccharide may contain potential vaccine components (15). A few of these outer membrane proteins, especially CopB (OMP B2) (4, 38), OMP CD (24), TbpA and TbpB (28), LbpA and LbpB (12), and UspA (ubiquitous surface protein A or HMW-OMP) (20, 26), which consists of two related proteins, UspA1 and UspA2 (2, 3), have been characterized in some detail. Furthermore, changes in expression of *M. catarrhalis* outer membrane proteins have been shown to affect the ability of this organism to resist clearance from the lungs of animals (27).

The UspA1 and UspA2 surface proteins of *M. catarrhalis* are structurally related but appear to mediate different biological functions. The amino acid sequences of UspA1 and UspA2 from *M. catarrhalis* strain 035E are approximately 43% identical, but each possesses an internal segment of 135 amino acids with 93% identity; this region contains an epitope which binds the monoclonal antibody (MAB) 17C7 and is present in all disease isolates of *M. catarrhalis* tested to date (20). However, these two proteins appear to have different biological functions, with UspA1 having been shown to be essential for attachment of *M. catarrhalis* strain 035E to Chang conjunctival cells in vitro, whereas UspA2 is involved directly or indirectly in serum resistance of this strain (2). Interestingly, after solubilization of *M. catarrhalis* cells at 37°C, both UspA1 and UspA2 apparently are present as oligomers or aggregates, each of which migrates in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) with an apparent molecular weight of greater than 250,000 even though their molecular masses are 88 and 62 kDa, respectively (3).

In the present study, isogenic *uspA1*, *uspA2*, and *uspA1 uspA2* double mutants were constructed in three additional strains of *M. catarrhalis*. Analysis of the ability of these mutants to adhere to Chang cells in vitro led to the discovery of a second type of UspA2 protein that is comprised of amino acid segments from both UspA1 and UspA2. Direct evidence that both UspA1 and this second type of UspA2 protein have functional activity as adhesins was obtained by expressing both of these *M. catarrhalis* proteins as recombinant molecules in *Haemophilus influenzae*.

* Corresponding author. Mailing address: Department of Microbiology, Hamon Building, NA6.200, University of Texas Southwestern Medical Center, 6000 Harry Hines Boulevard, Dallas, TX 75235-9048. Phone: (214) 648-5974. Fax: (214) 648-5905. E-mail: hansen01@utsw.swmed.edu.

† Present address: Department of Pediatrics, University of Bern, Inselspital, CH-3010 Bern, Switzerland.

MATERIALS AND METHODS

Bacterial strains, plasmids, and culture conditions. Most of the bacterial strains and plasmids used in this study are listed in Table 1. *M. catarrhalis* (3), *Escherichia coli* (11), and *H. influenzae* (18) strains were routinely cultured as described previously. Antimicrobial supplementation for *M. catarrhalis* mutants involved kanamycin (15 µg/ml), spectinomycin (15 µg/ml), or chloramphenicol (0.6 µg/ml). For bacterial adherence and serum bactericidal assays, *M. catarrhalis* strains were grown in broth without antibiotics for two to three generations. Recombinant strains of *E. coli* were selected with kanamycin (50 µg/ml), spectinomycin (150 µg/ml), or ampicillin (100 µg/ml). *H. influenzae* recombinant strains were cultured in the presence of chloramphenicol (2 µg/ml). For adherence assays, *H. influenzae* strains were grown in broth without antibiotics for two to three generations.

Recombinant DNA methods. Standard molecular biology techniques were performed as described previously (37) using *E. coli* strain DH5α or *H. influenzae* strain DB117 (39) as the host for recombinant DNA manipulations. The procedure for electroporating *M. catarrhalis* has been described in detail elsewhere (21) and was also used to electroporate *H. influenzae*.

PCR. Amplicons used for cloning of the *uspA1*, *uspA2*, and *uspA2H* genes were generated with *Pfu* DNA polymerase (Stratagene, La Jolla, Calif.). Amplification of other DNA fragments was performed either with the Gene Amp XL PCR kit (Perkin-Elmer Biosystems, Branchburg, N.J.) or with *Taq* DNA polymerase (Promega, Madison, Wis.) by the manufacturer's procedure.

Construction of isogenic mutants. The plasmid pUSPAIKAN was used to construct *uspA1* mutants of *M. catarrhalis* strains 012E and TTA37 as described previously (2). The oligonucleotide primers P1 (5'-CGGGATCCCCTCTCCCCCTAAAATCGCTG-3') and P2 (5'-AGGGATCCCGCTGTATGCCGCTACTCGCAGCT-3') (*Bam*HI restriction sites are underlined) were used in PCR to amplify a 3-kb DNA fragment containing an incomplete *uspA2* gene with a kanamycin cartridge insertion from the *M. catarrhalis uspA2* mutant 035E.2 (2). This PCR product was used to electroporate *M. catarrhalis* 012E and TTA37; kanamycin-resistant transformants were screened by PCR to identify potential *uspA2* mutants of strain 012E and *uspA2H* mutants of strain TTA37.

To construct isogenic mutants defective in expression of both UspA1 and UspA2, an incomplete *uspA2* open reading frame (ORF) was amplified from *M. catarrhalis* strain P44 by PCR using the oligonucleotide primers P1 and P2. After digestion with *Bam*HI, this 2.2-kb DNA fragment was ligated into pBS KS(+). A 0.4-kb *Bgl*II fragment was removed from the middle of this partial *uspA2* ORF and was replaced with a cartridge encoding resistance to spectinomycin (42), yielding the plasmid pELU2P44SPEC. *Eco*RI-digested pELU2P44SPEC was used to electroporate the *M. catarrhalis uspA1* mutants 012E.1 and TTA37.1. Spectinomycin-resistant transformants were screened in colony blot radioimmunoassays with the UspA1- and UspA2-reactive MAb 17C7 to identify transformants that had lost the ability to express both proteins. Linearized pELU2P44SPEC was used to electroporate *M. catarrhalis* 046E to construct an isogenic *uspA2H* mutant of this strain.

The oligonucleotide primers P3 (5'-AGGGATCCAACGACGGTCCAAGATGG-3') and P4 (5'-AGGGATCCCCTGCCACCTAAAGCCTTG-3') (underlined residues are *Bam*HI sites) were used in PCR to amplify a 3.6-kb DNA fragment from *M. catarrhalis* strain 035E.12 (2) consisting of the entire *uspA1* ORF with a chloramphenicol resistance cartridge insertion. This amplicon was introduced into *M. catarrhalis* strains 046E and 046E.2 by electroporation. Chloramphenicol-resistant colonies were screened by PCR to identify potential *uspA1* mutants.

Cloning of *M. catarrhalis* genes in *H. influenzae*. Amplicons of approximately 3 kb containing the *uspA1* genes from *M. catarrhalis* strains 035E, TTA37, 012E, 046E, and V1171 were generated by using the oligonucleotide primers P3 and P4. These PCR products were digested with *Bam*HI and ligated into the vector pACYC184 (New England Biolabs, Inc., Beverly, Mass.). These ligation mixtures were introduced into *H. influenzae* strain DB117 by electroporation. Chloramphenicol-resistant colonies were screened in colony blot radioimmunoassays with MAb 17C7 to identify recombinants expressing *M. catarrhalis* UspA1 proteins.

PCR products of approximately 3 kb containing the *M. catarrhalis* 046E, TTA37, and V1166 *uspA2H* genes were amplified with the oligonucleotide primers P5 (5'-AGGCATGCGGTTATAGCAATCCCTTG-3') and P6 (5'-GCGCATGCGCCAGCTTTATTTTATGCAGGG-3') (the underlining indicates an *Sph*I site). After digestion with *Sph*I, these amplicons were ligated into pACYC184 and used to electroporate *H. influenzae*; MAb 17C7 was used to identify transformants expressing the UspA2H protein.

The oligonucleotide primers P7 (5'-AGGCATGCTGTCCGCTGATGCTTTCTG-3') and P8 (5'-AGGCATGCGCTTTTATCCATCACTCAC-3') (the underlining indicates an *Sph*I site) were used to generate 2.0- to 2.5-kb PCR products containing the *uspA2* genes from *M. catarrhalis* 012E, 035E, and V1171. The *Sph*I-digested amplicons were ligated into pACYC184 and used to electroporate *H. influenzae*. Chloramphenicol-resistant colonies were screened as described above for expression of UspA2. Both strands of all of the cloned PCR products were sequenced.

Nucleotide sequence analysis. Nucleotide sequence analysis of recombinant plasmids and PCR amplicons was performed as described previously (10).

Characterization of protein antigens. SDS-PAGE of proteins present in whole cell lysates and Western blot analysis were performed as described previously

(10). The UspA1- and UspA2-reactive MAb 17C7 and the UspA1-specific MAb 24B5 have been described previously (2, 10). Colony blot radioimmunoassays were performed as described previously (2). The indirect antibody accessibility assay was performed as described previously (2) using MAb 17C7 to detect recombinant UspA1, UspA2, and UspA2H proteins on the surface of *H. influenzae* cells and MAb 24B5 to detect native UspA1 protein on the surface of wild-type and mutant strains of *M. catarrhalis*.

Adherence and serum resistance assays. Bacterial adherence assays were performed with Chang conjunctival epithelial cells as previously described (2). Killing of *M. catarrhalis* strains by 10% (vol/vol) normal human serum was assessed as previously described (2).

Nucleotide sequence accession numbers. The nucleotide sequences of the genes sequenced in this study have been deposited in GenBank under the following accession numbers: AF181072, *M. catarrhalis* 012E *uspA1*; AF181073, *M. catarrhalis* 012E *uspA2*; AF181076, *M. catarrhalis* TTA37 *uspA1*; AF181075, *M. catarrhalis* TTA37 *uspA2H*; U61725, *M. catarrhalis* 046E *uspA1*; and AF181074, *M. catarrhalis* 046E *uspA2H*.

RESULTS

Construction of isogenic *uspA1* and *uspA2* mutants of *M. catarrhalis* 012E, TTA37, and 046E. The *uspA1* and *uspA2* gene products of *M. catarrhalis* 035E were previously shown to be involved in adherence to human conjunctival epithelial cells and in resistance to killing by normal human serum, respectively (2). To determine whether the roles of these surface proteins are conserved among strains of *M. catarrhalis*, we constructed isogenic *uspA1*, *uspA2*, and *uspA1 uspA2* double mutants of *M. catarrhalis* strains 012E, TTA37, and 046E. (For reasons that will be explained below, the *uspA2* genes of both strains TTA37 and 046E will be designated *uspA2H* from this point forward.) PCR-based and Southern blot analyses confirmed that each mutant had the appropriate antibiotic resistance cartridge inserted into the intended target gene (data not shown).

Characterization of selected proteins expressed by wild-type and mutant *M. catarrhalis* strains. Whole cell lysates were heated at 100°C for 15 min before SDS-PAGE; this treatment allowed distinction of UspA1 and UspA2 proteins probed with a MAb (17C7) reactive with both proteins, as shown previously (10). When these preparations were probed in Western blot analysis with the UspA1-reactive MAb 24B5, it was found that the disruption of the *uspA1* ORF in *M. catarrhalis* 012E.1 (Fig. 1A, lane 4), in 012E.12 (Fig. 1A, lane 8), and in TTA37.1 and TTA37.12 (Fig. 1B, lanes 4 and 8, respectively) abolished expression of the approximately 120-kDa UspA1 protein. The *uspA2* mutation in *M. catarrhalis* 012E.2 (Fig. 1A, lane 5) and the *uspA2H* mutation in TTA37.2 (Fig. 1B, lane 5) eliminated expression of UspA2, which is present as a very high-molecular-weight complex reactive with MAb 17C7 in these whole cell lysates (Fig. 1A and B, lanes 3). The presence of both *uspA1* and *uspA2* mutations in strain 012E.12 (Fig. 1A, lanes 7 and 8) and of both *uspA1* and *uspA2H* mutations in strain TTA37.12 (Fig. 1B, lanes 7 and 8) eliminated all MAb 17C7 and MAb 24B5 reactivity from these double mutants. The *uspA1*, *uspA2H*, and *uspA1 uspA2H* mutants of strain 046E had Western blot reactivity patterns identical to those of the 012E and TTA37 mutant sets (data not shown). Taken together, these results indicate that these isogenic mutants lacked expression of the genes that were specifically targeted by our mutagenesis strategy.

Effect of *uspA1* and *uspA2* mutations on serum resistance. Inactivation of the *uspA2* gene in *M. catarrhalis* 035E renders this strain exquisitely sensitive to the bactericidal activity of normal human serum, while the lack of expression of the UspA1 protein does not affect the ability of 035E to resist killing by normal human serum (2). To determine whether the involvement of the *uspA2* gene product in serum resistance is a conserved feature of *M. catarrhalis* strains, we tested the

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Genotype or description	Reference or Source
<i>M. catarrhalis</i>		
035E	Wild-type disease isolate, attaches to Chang cells, serum resistant	2
035E.1	<i>uspA1</i> mutant of 035E, attachment deficient, serum resistant	2
035E.2	<i>uspA2</i> mutant of 035E, attaches to Chang cells, serum sensitive	2
035E.12	<i>uspA1 uspA2</i> mutant of 035E, attachment deficient, serum sensitive	2
012E	Wild-type disease isolate, attaches to Chang cells, serum resistant	1
012E.1	<i>uspA1</i> mutant of 012E, attachment deficient, serum resistant	This study
012E.2	<i>uspA2</i> mutant of 012E, attaches to Chang cells, serum sensitive	This study
012E.12	<i>uspA1 uspA2</i> mutant of 012E, attachment deficient, serum sensitive	This study
TTA37	Wild-type disease isolate, attaches to Chang cells, serum sensitive	Steven Berk
TTA37.1	<i>uspA1</i> mutant of TTA37, attaches to Chang cells, serum sensitive	This study
TTA37.2	<i>uspA2H</i> mutant of TTA37, attachment deficient, serum sensitive	This study
TTA37.12	<i>uspA1 uspA2H</i> mutant of TTA37, attachment deficient, serum sensitive	This study
046E	Wild-type disease isolate, attaches to Chang cells, serum resistant	John Nelson
046E.1	<i>uspA1</i> mutant of 046E, attaches to Chang cells, serum resistant	This study
046E.2	<i>uspA2H</i> mutant of 046E, attaches to Chang cells, serum sensitive	This study
046E.12	<i>uspA1 uspA2H</i> mutant of 046E, attachment deficient, serum sensitive	This study
TTA24	Wild-type disease isolate	10
P44	Wild-type disease isolate	25
ATCC 25240	Wild-type disease isolate	American Type Culture Collection
E22	Wild-type disease isolate	5
V1166	Wild-type isolate from nasopharynx of a healthy child	F. Henderson
V1171	Wild-type isolate from nasopharynx of a healthy child	10
<i>E. coli</i> DH5 α	Host strain for cloning experiments	37
<i>H. influenzae</i> DB117	Host strain for cloning experiments	39
Plasmids		
pBS KS(+)	Cloning vector, Amp ^r	Stratagene
pUSPA1KAN	pBS containing a truncated <i>uspA1</i> gene from <i>M. catarrhalis</i> strain 035E into which a kanamycin resistance cartridge was inserted	3
pSPEC ^r	Source of the spectinomycin resistance cartridge	42
pELU2P44SPEC	pBS containing an incomplete <i>uspA2</i> gene from <i>M. catarrhalis</i> P44 into which a spectinomycin resistance cartridge was inserted	This study
pACYC184	Cloning vector	New England Biolabs
pELU112	pACYC184 containing the <i>M. catarrhalis</i> 012E <i>uspA1</i> gene	This study
pELU212	pACYC184 containing the <i>M. catarrhalis</i> 012E <i>uspA2</i> gene	This study
pELU135	pACYC184 containing the <i>M. catarrhalis</i> 035E <i>uspA1</i> gene	This study
pELU235	pACYC184 containing the <i>M. catarrhalis</i> 035E <i>uspA2</i> gene	This study
pELU146	pACYC184 containing the <i>M. catarrhalis</i> 046E <i>uspA1</i> gene	This study
pELU246	pACYC184 containing the <i>M. catarrhalis</i> 046E <i>uspA2H</i> gene	This study
pELU137	pACYC184 containing the <i>M. catarrhalis</i> TTA37 <i>uspA1</i> gene	This study
pELU237	pACYC184 containing the <i>M. catarrhalis</i> TTA37 <i>uspA2H</i> gene	This study
pELU171	pACYC184 containing the <i>M. catarrhalis</i> V1171 <i>uspA1</i> gene	This study
pELU271	pACYC184 containing the <i>M. catarrhalis</i> V1171 <i>uspA2</i> gene	This study
pELU266	pACYC184 containing the <i>M. catarrhalis</i> V1166 <i>uspA2H</i> gene	This study




FIG. 1. Western blot analysis of whole cell lysates prepared from *M. catarrhalis* strains 012E and TTA37 and their corresponding isogenic mutants. (A) Wild-type *M. catarrhalis* 012E (lanes 1 and 2), the *uspA1* mutant 012E.1 (lanes 3 and 4), the *uspA2* mutant 012E.2 (lanes 5 and 6), and the *uspA1 uspA2* double mutant 012E.12 (lanes 7 and 8). (B) Wild-type *M. catarrhalis* TTA37 (lanes 1 and 2), the *uspA1* mutant TTA37.1 (lanes 3 and 4), the *uspA2H* mutant TTA37.2 (lanes 5 and 6), and the *uspA1 uspA2H* double mutant TTA37.12 (lanes 7 and 8). The UspA1- and UspA2-reactive MAb 17C7 was used to probe lanes 1, 3, 5, and 7 in both panels; the UspA1-specific MAb 24B5 was used to probe lanes 2, 4, 6, and 8 in both panels. Molecular mass markers are shown to the left in kilodaltons.

serum-resistant strains 012E and 046E and their isogenic mutants in a serum bactericidal assay. It should be noted that, unlike most disease isolates of *M. catarrhalis* (41), the wild-type strain TTA37 was completely killed within the first 30 min of incubation with normal human serum (data not shown). Therefore, mutants of *M. catarrhalis* TTA37 were not tested in the serum bactericidal assay.

The *M. catarrhalis uspA2* mutant 012E.2 and the *uspA1 uspA2* double mutant 012E.12, both of which lack expression of the UspA2 protein, were completely killed within 30 min of incubation with normal human serum (data not shown). In contrast, lack of expression of the UspA1 protein did not affect the ability of the *M. catarrhalis uspA1* mutant 012E.1 to resist killing by normal human serum when compared to the parent strain. The *uspA2H* mutant and *uspA1 uspA2H* double mutant of strain 046E were much more sensitive to killing by normal human serum than were the wild-type parent strain and the *uspA1* mutant of strain 046E (data not shown).

Effect of *uspA1* and *uspA2* mutations on the ability of *M. catarrhalis* strains to adhere to human epithelial cells in vitro. Lack of expression of the UspA1 protein causes an approximately 60-fold decrease in the ability of *M. catarrhalis* 035E to adhere to Chang conjunctival epithelial cells in vitro (2). To determine whether the involvement of the *uspA1* gene product in the process of adherence to human epithelial cells is a common feature of *M. catarrhalis* isolates, we tested the ability of the mutants described above to attach to Chang cells.

M. catarrhalis 012E and its *uspA2* mutant 012E.2 exhibited similar levels of attachment to Chang cells (Table 2). Lack of expression of the UspA1 protein in both the *M. catarrhalis uspA1* mutant 012E.1 and the *uspA1 uspA2* double mutant 012E.12 diminished adherence of these mutants to the mono-

TABLE 2. Adherence of wild-type and isogenic mutant strains of *M. catarrhalis* to Chang conjunctival epithelial cells in vitro

Strain	Genotype	Adherence ^a
012E	Wild type	123.0 ± 15.57
012E.1	<i>uspA1</i> mutant	9.3 ± 3.8 (0.0001)
012E.2	<i>uspA2</i> mutant	138.3 ± 19.6 (0.0088)
012E.12	<i>uspA1 uspA2</i> mutant	14.1 ± 3.8 (0.0001)
TTA37	Wild type	42.4 ± 17.3
TTA37.1	<i>uspA1</i> mutant	45.8 ± 14.7 (0.6593)
TTA37.2	<i>uspA2H</i> mutant	6.7 ± 2.1 (0.0003)
TTA37.12	<i>uspA1 uspA2H</i> mutant	4.5 ± 0.5 (0.0002)
046E	Wild type	52.1 ± 13.1
046E.1	<i>uspA1</i> mutant	55.2 ± 14.3 (0.6725)
046E.2	<i>uspA2H</i> mutant	45.4 ± 18.1 (0.4847)
046E.12	<i>uspA1 uspA2H</i> mutant	6.7 ± 2.2 (0.0001)

^a Adherence of *M. catarrhalis* to Chang cells is expressed as the mean (± standard deviation) percentage of the inoculum which adhered to the monolayers. The *P* value compared to the value for the wild-type parent strain, using a two-tailed *t* test, is given in parentheses.

layers by 1 order of magnitude (Table 2). Unexpectedly, however, *M. catarrhalis* TTA37 and its *uspA1* mutant TTA37.1 were found to attach to Chang cells at similar levels (Table 2). Disruption of the *uspA2H* ORF in the *uspA2H* mutant TTA37.2 and the *uspA1 uspA2H* mutant TTA37.12 did result in an approximately 10-fold reduction in adherence of these strains to Chang cells (Table 2). *M. catarrhalis* 046E, its *uspA1* mutant 046E.1, and the *uspA2H* mutant 046E.2 all exhibited similar levels of attachment to Chang cells (Table 2). When expression of both UspA1 and UspA2H was eliminated in the *uspA1 uspA2H* mutant 046E.12, this resulted in a 10-fold reduction in adherence (Table 2).

Nucleotide sequence analysis of the *uspA1* and *uspA2* genes of *M. catarrhalis* strains 012E, TTA37, and 046E. To investigate a possible correlation between the unexpected biological effects of the *uspA1* and *uspA2H* mutations in strains TTA37 and 046E and the structural features of the relevant proteins, we determined the nucleotide sequences of the *uspA1* and *uspA2* genes from strain 012E and those of the *uspA1* and *uspA2H* genes from strains TTA37 and 046E. The *uspA1* ORFs were predicted to encode proteins ranging in molecular weight from 91,873 to 96,959 (Table 3). The *uspA2* gene of *M. catarrhalis* 012E was predicted to encode a protein of 74,599 Da, whereas the predicted UspA2H proteins from *M. catarrhalis* 046E and TTA37 were both significantly larger than that of strain 012E (Table 3).

TABLE 3. Characteristics of the *uspA1*, *uspA2*, and *uspA2H* genes and their encoded protein products from selected *M. catarrhalis* strains

Strain	Gene	Length of ORF (nt)	Mol wt of predicted protein	No. of amino acids	Predicted start codon
012E	<i>uspA1</i>	2,766	96,959	922	ATG
	<i>uspA2</i>	2,052	74,599	684	ATG
TTA37	<i>uspA1</i>	2,619	91,873	873	ATG
	<i>uspA2H</i>	2,667	92,145	889	GTG
046E	<i>uspA1</i>	2,676	93,350	892	ATG
	<i>uspA2H</i>	2,682	94,229	894	GTG




FIG. 2. Schematic representation of the repetitive amino acid sequences and other motifs present in the UspA1 and UspA2 proteins of *M. catarrhalis* 012E and in the UspA1 and UspA2H proteins of strains TTA37 and 046E. The solid bars indicate the lengths of the proteins. The colored boxes indicate the positions and relative lengths as well as the numbers of repeats and other motifs. The GGG and FAAG motifs are indicated; the other motifs are described in detail elsewhere (10).

Approximately 130 to 140 nucleotides (nt) 5' from the predicted translational start codon of the *M. catarrhalis* 012E *uspA2* ORF were 19 repeats of the tetranucleotide AGAT (data not shown), as previously observed in the 5' regions of other *uspA2* genes (10). The nucleotide sequence upstream of the predicted translational start codon of the three *uspA1* ORFs was also found to contain a nucleotide repeat motif. As previously seen with other *uspA1* genes (10), poly(G) tracts containing 9 to 11 guanine residues were located 30 nt 5' from the ATG translational start codon of the *uspA1* ORFs in strains 012E, TTA37, and 046E (data not shown). Nucleotide sequence analysis of approximately 1.2 kb of DNA upstream of the proposed translational start codons of the *M. catarrhalis* 046E and TTA37 *uspA2H* ORFs did not detect any AGAT repeats or poly(G) tracts but did reveal that these regions were virtually identical in 046E and TTA37 (data not shown). This region did not resemble the upstream regions of either *uspA1* or *uspA2* genes from previously characterized *M. catarrhalis* strains. In addition, an ORF encoding a protein product with 26% identity to the InsB protein encoded by IS1 in the *Shigella dysenteriae* genome (34) was present in this upstream region from strains 046E and TTA37 (data not shown).

Comparison of the predicted amino acid sequences of the UspA1 proteins. Overall, the UspA1 proteins of strains 012E, TTA37, and 046E were greater than 70% identical (data not shown). As previously observed with the UspA1 proteins from four other *M. catarrhalis* strains (10), these three UspA1 proteins had virtually identical predicted signal peptides of 49 residues and contained several different amino acid repeats and other motifs (Fig. 2). Of interest are the GGG repeats, which are highly conserved in the N termini of UspA1 proteins (10) and absent from the UspA2 proteins of *M. catarrhalis* strains 035E, TTA24, ATCC 25238, and V1171 (10) and 012E (Fig. 2). The *M. catarrhalis* 012E, 046E, and TTA37 UspA1 proteins contained 9, 9, and 11 GGG repeat motifs, respectively (Fig. 2). The GGG repeat is characterized by the consensus sequence T(I/V)GGGXXNXAXGYS where the underlined residues are perfectly conserved. Following the GGG repeats, a region of 135 amino acids (designated FAAG for the first four residues present in this motif) was present in these three UspA1 proteins (Fig. 2) as well as in the four UspA1 proteins previously characterized (10). The C termini of the 012E, TTA37, and 046E UspA1 proteins were also found to contain a stretch of 193 amino acids (designated CTER1 in

Fig. 2) that were 97% identical to those observed previously within the C termini of UspA1 proteins (10).

Comparison of the predicted amino acid sequences of the UspA2 and UspA2H proteins. The UspA2 protein of *M. catarrhalis* 012E was less than 50% identical to the UspA2H proteins of strains 046E and TTA37, whereas the latter were predicted to be 62% identical. Alignment of the predicted amino acid sequences of these three proteins showed that the signal peptide of 30 residues previously seen in four other UspA2 proteins (10) was present only in the UspA2 protein from *M. catarrhalis* 012E (data not shown). However, the signal peptide of the UspA2H proteins from strains TTA37 and 046E was nearly identical to that found in the UspA1 proteins of all *M. catarrhalis* strains characterized to date (10). The C termini of both the UspA2 protein from 012E and the UspA2H proteins from strains TTA37 and 046E were also found to contain a region of 158 residues (designated CTER2 in Fig. 2) that was 87% identical among these three strains.

As before (10), several different amino acid repeats and other motifs were observed in the UspA2 and UspA2H proteins (Fig. 2). Certain amino acid repeats, such as the VEEG, NINNY, LAAY, KASS, and FET motifs, again were found to be shared by both UspA1 and the UspA2 and UspA2H proteins (Fig. 2). Interestingly, the N-terminal halves of the UspA2H proteins of *M. catarrhalis* TTA37 and 046E contained amino acid sequences that had a high degree of identity with the GGG repeats and FAAG motif present in the UspA1 protein (Fig. 2). These particular motifs were notably absent from the UspA2 proteins of *M. catarrhalis* 012E (Fig. 2) and strains 035E, TTA24, ATCC 25238, and V1171 (10). In fact, alignment of the predicted amino acid sequences of the N termini of the *M. catarrhalis* 012E, TTA37, and 046E UspA1 proteins with those of the UspA2H proteins of strains 046E and TTA37 revealed a high degree of homology (data not shown).

Analysis of *M. catarrhalis* isolates for the presence of a hybrid *uspA2* gene. Strains 046E and TTA37 both expressed a UspA2H protein which had an N-terminal half resembling that of a UspA1 protein together with a C-terminal half similar to those of other UspA2 proteins. For the sake of clarity, this type of protein has been designated UspA2H to indicate its apparent "hybrid" nature. To investigate whether other *M. catarrhalis* isolates might express a UspA2H protein, PCR was used to detect the possible presence of a *uspA2H* ORF. The oligonucleotide primer P9 (Fig. 3) corresponds to a perfectly conserved region found at the 5' ends of the *uspA1* ORFs of five *M. catarrhalis* strains (035E, TTA24, 012E, 046E, and TTA37). This same nucleotide sequence is present in the 5' ends of the *uspA2H* ORFs in strains 046E and TTA37. The oligonucleotide primer P10 (Fig. 3) was designed to bind a perfectly conserved region at the 3' ends of both the *uspA2* ORFs in strains 035E, TTA24, and 012E and the *uspA2H* ORFs in strains TTA37 and 046E. For amplification of *uspA1* ORFs, the oligonucleotide primers P9 (described above) and P12 (Fig. 3; derived from the 3' ends of the *uspA1* ORFs of strains 035E, 012E, and TTA24) were utilized. To detect the presence of *uspA2* ORFs, the oligonucleotide primers P11 (Fig. 3; derived from the 5' ends of the *uspA2* ORFs of strains 035E, 012E, and TTA24) and P10 (described above) were used in PCR.

No product was detected when primers P9 and P10 were used in PCR with chromosomal DNA from strain 035E, 012E, or TTA24; these three strains each express a UspA2 protein that has no UspA1-like sequences in its N-terminal half. In contrast, the use of these same two primers in PCR with chromosomal DNAs from strains TTA37 and 046E yielded a 2.5-kb




FIG. 3. Schematic representation of the placement of the oligonucleotide primers used in PCR to detect the presence of the *uspA1*, *uspA2*, and *uspA2H* genes in 21 *M. catarrhalis* strains.

product in both instances. When these same oligonucleotide primers (P9 and P10) were used in PCR with chromosomal DNAs from 15 additional *M. catarrhalis* strains whose *uspA2* genes had not been sequenced, amplicons of 2.3 to 2.5 kb were obtained from several strains, including ATCC 25240, E22, and V1166. A total of 11 disease isolates (035E, 012E, 046E, TTA37, TTA24, P44, P48, ATCC 25238, ATCC 25240, E22, and TTA1) and 10 normal carriage isolates (V1111, V1156, V1171, V1169, V1166, V1153, V1129, V1126, V1118, and V1121) were used in these PCRs. All 21 strains yielded a PCR product when the *uspA1*-specific primer pair P9 and P12 was used (data not shown). PCR products were obtained with the *uspA2*-specific primer pair P11 and P10 only from the 16 strains that did not possess a *uspA2H* gene (the strains other than 046E, TTA37, ATCC 25240, E22, and V1166).

Nucleotide sequence analysis of approximately 700 nt at the 3' ends of the PCR products amplified with primers P9 and P10 from *M. catarrhalis* E22, ATCC 25240, and V1166 revealed that the encoded amino acid sequence had a high level of identity with the amino acid sequence typically found in the C-terminal regions of UspA2 proteins (data not shown). The deduced amino acid sequence derived from approximately 700 nt at the 5' ends of these same amplicons, however, contained the GGG repeats and the FAAG region, which were typically present in the N termini of UspA1 proteins (data not shown).

Expression of the *M. catarrhalis* UspA1, UspA2, and UspA2H proteins in *H. influenzae*. To conclusively determine whether UspA1 and UspA2H proteins functioned as adhesins, the *uspA1*, *uspA2*, and *uspA2H* genes from various *M. catarrhalis* strains were cloned and expressed in *H. influenzae* DB117, which has been previously shown to adhere poorly to Chang cells in vitro (40).

Western blot analysis revealed that the UspA1- and UspA2-reactive MAb 17C7 bound a 120-kDa antigen in recombinant *H. influenzae* cells expressing the UspA1 protein of *M. catarrhalis* 012E, 035E, V1171, and TTA37 (data not shown). The recombinant UspA1 protein from strain 046E was not detectable in Western blot analysis despite the fact that it could be detected in colony blot radioimmunoassay (data not shown).

TABLE 4. Adherence of recombinant *H. influenzae* cells to Chang conjunctival epithelial cells in vitro and detection of recombinant proteins on the bacterial cell surface

Recombinant plasmid	<i>M. catarrhalis</i> protein ^a	Adherence ^b	Detection of protein on cell surface ^c
pACYC184 (vector)		0.2 ± 0.1	918 ± 276
pELU112	012E-UspA1	35.1 ± 8.7	13,870 ± 2,132
pELU212	012E-UspA2	0.4 ± 0.2	5,557 ± 1,546
pELU135	035E-UspA1	17.4 ± 7.3	5,050 ± 1,184
pELU235	035E-UspA2	0.2 ± 0.1	7,175 ± 3,201
pELU171	V1171-UspA1	12.3 ± 7.2	6,425 ± 1,049
pELU271	V1171-UspA2	0.4 ± 0.2	4,111 ± 1,333
pELU137	TTA37-UspA1	0.9 ± 0.4	1,286 ± 206
pELU237	TTA37-UspA2H	16.9 ± 6.3	11,630 ± 1,351
pELU146	046E-UspA1	0.8 ± 0.2	703 ± 221
pELU246	046E-UspA2H	52.6 ± 15.17	28,980 ± 1,703
pELU266	V1166-UspA2H	62.3 ± 16.6	5,632 ± 199

^a The *M. catarrhalis* protein expressed by the *H. influenzae* DB117 clone containing this recombinant plasmid.

^b Adherence of *M. catarrhalis* organisms to Chang cells is expressed as the mean (± standard deviation) percentage of the inoculum which adhered to the monolayers.

^c Counts per minute (mean ± standard deviation) of radioiodinated goat anti-mouse immunoglobulin bound to MAb 17C7 on the surface of recombinant cells. The indirect antibody-accessibility assay was performed as described previously (2).

MAb 17C7-reactive antigens with apparent molecular masses in excess of 200 kDa were detected in *H. influenzae* strains expressing the UspA2 proteins of strain 012E, 035E, and V1171. Very high-molecular-weight antigens were also detected with MAb 17C7 in whole cell lysates from *H. influenzae* cells expressing the UspA2H proteins from *M. catarrhalis* TTA37, 046E, and V1166. MAb 17C7 was used in the indirect antibody accessibility assay to detect recombinant UspA1, UspA2, and UspA2H proteins on the surface of *H. influenzae*. With the exception of the UspA1 proteins from strains TTA37 and 046E, all of these *M. catarrhalis* proteins were readily detectable on the surface of the recombinant strains (Table 4).

Expression of the *M. catarrhalis* 012E, 035E, or V1171 UspA1 protein increased the ability of *H. influenzae* DB117 to adhere to Chang cells by 2 orders of magnitude (Table 4; Fig. 4B). Similarly, expression of the *M. catarrhalis* TTA37, 046E, or V1166 UspA2H protein increased adherence by a factor of 100-fold (Table 4; Fig. 4D). In contrast, expression of the *M. catarrhalis* 012E, 035E, or V1171 UspA2 protein had little effect on the adherence of *H. influenzae* to Chang cells (Table 4; Fig. 4C). Similarly, the recombinant forms of the *M. catarrhalis* TTA37 and 046E UspA1 proteins had at best a very slight (i.e., fourfold) effect on attachment of *H. influenzae* to Chang cells (Table 4).

DISCUSSION

Despite advances made in the characterization of the biochemical and immunogenic properties of the *M. catarrhalis* UspA1 and UspA2 proteins (2, 3, 8–10, 31), information about the structure-function relationships inherent in these two macromolecules remains limited. Analysis of isogenic *uspA1* and *uspA2* mutants of *M. catarrhalis* strain 035E (2) provided the

first indication that these two proteins, which have in common a 135-amino-acid region that contains the epitope for MAb 17C7, might have different functions. Lack of expression of UspA1 severely impaired the ability of *M. catarrhalis* 035E to adhere to human epithelial cells in vitro, whereas the *uspA2* mutation rendered this strain sensitive to complement-mediated killing by normal human serum (2). In addition, the UspA1 protein purified from *M. catarrhalis* 035E binds to HEp-2 cells, whereas the UspA2 protein purified from the same strain binds these same cells very weakly (9). Furthermore, antibodies raised against purified UspA1, but not UspA2, were sufficient to significantly abrogate adherence of strain 035E to HEp-2 cells (9). It could be inferred from these findings that UspA1 and UspA2 likely had distinct biological functions and that UspA1 was directly or indirectly involved in the physical interaction of *M. catarrhalis* 035E with human epithelial cells in vitro.

Analysis of additional *uspA1* and *uspA2* mutants in the present study revealed that, at least in strain 012E, inactivation of the *uspA1* gene could be correlated with loss of adherence ability in vitro (Table 2). In contrast, inactivation of the *uspA2H* gene of strain TTA37, but not the *uspA1* gene, adversely affected the ability of this strain to adhere to Chang cells in vitro. This finding led to the discovery of a second type of UspA2 protein (UspA2H) in strain TTA37 which has a C-terminal half that is virtually identical to the C-terminal half of previously described UspA2 proteins (10) but which has an N-terminal half that is very different from that of other UspA2 proteins. More specifically, this region of the UspA2H protein contains the GGG amino acid repeats and the FAAG region, which have been observed in the N termini of all UspA1 proteins examined to date (10).

A PCR-based survey of a number of *M. catarrhalis* isolates identified additional strains which likely encoded a UspA2H protein. Determination of the complete nucleotide sequence of the *uspA2H* gene from strain 046E and partial nucleotide sequence analysis of the PCR products obtained from strains V1166, E22, and ATCC 25240 indicated that the predicted UspA2H proteins from these strains had primary amino acid sequences very similar to that of the UspA2H protein of strain TTA37.

These data derived from mutant analysis still provided only indirect evidence that UspA1 and now UspA2H were involved in the ability of *M. catarrhalis* strains to attach to Chang cells in vitro. To demonstrate directly that these *M. catarrhalis* proteins were adhesins, both the *uspA1* and *uspA2H* genes were cloned and expressed in *H. influenzae* DB117, an organism that does not normally adhere well to Chang conjunctival cells (40). *H. influenzae* recombinant clones expressing the *M. catarrhalis* 035E or 012E UspA1 protein exhibited a dramatic increase in adherence to Chang cells (Table 4; Fig. 4). These results clearly demonstrate that the UspA1 proteins from these *M. catarrhalis* isolates, whose isogenic *uspA1* mutants are deficient in attachment ability, function as an adhesin for Chang cells in vitro. Furthermore, recombinant *H. influenzae* clones expressing *M. catarrhalis* TTA37, 046E, or V1166 UspA2H protein exhibited a significant increase in adherence ability, a result which confirmed that the UspA2H protein is directly involved in the attachment ability of these *M. catarrhalis* strains. In contrast, *H. influenzae* strains expressing the *uspA2* genes from strains 035E and 012E did not attach to Chang monolayers (Table 4; Fig. 4). This latter finding can be correlated directly with the lack of effect of *uspA2* mutations on the ability of these two *M. catarrhalis* strains to attach to Chang cells (2) (Table 2). Therefore, UspA2H proteins can function as adhesins in vitro and, as such, are functionally distinct from the




FIG. 4. Light micrographs of recombinant *H. influenzae* DB117 clones incubated with Chang conjunctival epithelial cells. Adherence assays were performed as described in Materials and Methods with the following modification: after the last wash to remove nonadherent bacterial cells, the monolayers were fixed for 10 min with methanol and subsequently stained with Giemsa stain. (A) DB117 carrying the cloning vector pACYC184; (B) DB117 carrying pELU112 (expressing 012E UspA1); (C) DB117 carrying pELU212 (expressing 012E UspA2); (D) DB117 carrying pELU246 (expressing 046E UspA2H).

UspA2 proteins of previously characterized *M. catarrhalis* strains, including 035E, TTA24, ATCC 25238, and V1171 (10).

The results in Tables 2 and 4 and Fig. 4 indicate that both the UspA1 and UspA2H proteins mediate a physical interaction between *M. catarrhalis* and Chang cells in vitro. Both proteins have striking identity in their N-terminal regions (Fig. 2), and it is tempting to speculate that the region of these proteins containing the GGG repeats and FAAG motif is responsible for this interaction, although it is equally likely that another amino acid sequence(s) in the same region could be responsible for attachment to Chang cells. Both the UspA1 and UspA2 proteins have been proposed to be part of the autotransporter family of secreted proteins (22), which includes gene products that form large macromolecular complexes on the surface of bacterial cells and which often function as adherence factors (22). The N termini of some of these autotransporter proteins have been proposed to be exposed on the surface of bacteria (22). The next logical step will be to identify the region(s) of the UspA1 and UspA2H proteins responsible for conferring on *M. catarrhalis* the ability to ad-

here to human epithelial cells in vitro. Our ability to express functional, recombinant UspA1 and UspA2H proteins in *H. influenzae* should allow construction of strains expressing chimeric molecules containing different portions of UspA1 and UspA2.

It is interesting that expression of the UspA1 protein of either strain TTA37 or strain 046E in *H. influenzae* did not promote attachment to Chang cells (Table 4). Nucleotide sequence analysis of the *M. catarrhalis* DNA inserts in the respective recombinant plasmids revealed no mutations caused by PCR amplification. However, the use of MAb 17C7 in an indirect antibody accessibility assay (2) indicated that the recombinant forms of these two UspA1 proteins were not expressed on the surface of the *H. influenzae* cells (data not shown). This finding provides a physical explanation for the lack of attachment ability of the recombinant *H. influenzae* cells expressing either of these UspA1 proteins, although it is not clear why these two proteins were not translocated to the surface of *H. influenzae*.

The phenotypes of the isogenic *uspA1* and *uspA2H* mutants

of strains TTA37 and 046E must be interpreted carefully. Lack of expression of UspA1 by *M. catarrhalis* strain 046E.1 did not eliminate the ability of this *uspA1* mutant to attach to Chang cells (Table 4), because it was still expressing the UspA2H protein. Similarly, lack of expression of UspA2H by strain 046E.2 did not adversely affect attachment, because this *uspA2H* mutant still expressed a UspA1 protein. In contrast, the *uspA2H* mutant of strain TTA37 exhibited a dramatic decrease in its ability to attach to Chang cells (Table 2). UspA1, however, was found to be exposed on the surface of both the wild-type TTA37 strain and its *uspA2H* mutant in an indirect antibody accessibility assay using the UspA1-specific MAb 24B5 (data not shown). Thus, the UspA1 protein from the wild-type strain TTA37 appears to be defective and unable to confer adherence ability on this strain, because the lack of expression of the UspA2H protein is sufficient to abrogate adherence of this particular isolate to Chang cells (Table 2). Structure-function analysis of the UspA1 protein should ultimately allow determination of the region(s) in the TTA37 UspA1 protein responsible for this apparent dysfunction. Alternatively, at this time we cannot exclude the possibility that lack of expression of UspA2H in the TTA37.2 mutant somehow adversely affected the orientation and thus the function of the UspA1 protein.

Although the UspA1 and UspA2H proteins have now been shown to be adhesins which mediate attachment to Chang cells in vitro, it is likely that *M. catarrhalis* strains express other adhesins. It has been reported that the CD outer membrane protein of *M. catarrhalis* bound human mucin purified from the middle ears of children with otitis media (36). Expression of a 200-kDa protein has been found to be exclusively associated with hemagglutinating isolates of *M. catarrhalis* (13); this proposed hemagglutination factor has been recently associated with the formation of a fibrillar layer surrounding the bacteria which was observed to physically contact erythrocytes (14).

Although the UspA2 and UspA2H proteins are structurally distinct, the results of the mutant analyses indicate that both are involved in the ability of their respective *M. catarrhalis* strains to survive killing mediated by the complement components present in normal human serum. The lack of expression of UspA2 in strains 035E.2 (2) and 012E.2 and the lack of expression of UspA2H by *M. catarrhalis* 046E.2 rendered these strains exquisitely sensitive to killing by normal human serum. It is tempting to speculate that the C-terminal halves of both UspA2 and UspA2H may somehow be involved in serum resistance, since this region is common to both proteins (Fig. 2). However, the C-terminal half of the UspA2H protein of the serum-sensitive *M. catarrhalis* strain TTA37 exhibits a high degree of sequence identity to the C-terminal half of the UspA2H protein of the serum-resistant *M. catarrhalis* strain 046E. In addition, the mechanism by which many disease isolates of *M. catarrhalis* resist killing by normal human serum is unclear, although it has been proposed that binding of vitronectin to the surface of serum-resistant *M. catarrhalis* cells inhibits the formation of the membrane attack complex (41) and it has been shown that purified UspA2 binds vitronectin (9). The construction of isogenic *uspA2* mutants of various *M. catarrhalis* strains as well as the newly realized ability to express the UspA2 and UspA2H proteins in the heterologous background of *H. influenzae* DB117 may help to elucidate the involvement of these proteins in protecting *M. catarrhalis* from killing by normal human serum.

At this point, it is appropriate to address issues concerning the structures of the *uspA1*, *uspA2*, and *uspA2H* genes. The data in the present study, together with those from two previous reports (3, 10), include the nucleotide sequences of a total

of eight *uspA1* genes, six *uspA2* genes, and two *uspA2H* genes. Examination of these sequences raises two important questions: what mechanism(s) is responsible for the variability in certain regions of the *uspA1* and *uspA2* genes, and how did the *uspA2H* genes originate?

Most of the interstrain variability in the *uspA1* genes and in the *uspA2* genes is localized to the 5' half of each ORF (data not shown). In contrast, the 3' end of each *uspA1* and *uspA2* gene is highly conserved, as is the 5' end of each gene, including the beginning of the ORF. This clustering of sequence polymorphisms amidst two highly conserved regions is reminiscent of the mosaic structure described for ORFs that encode surface-exposed proteins of other bacterial pathogens, including *Neisseria meningitidis* (23, 29), *E. coli* (30), and *Neisseria gonorrhoeae* (17). In these examples, horizontal genetic exchange has been proposed as the major mechanism by which mosaic genes are generated. Similar to several of the aforementioned pathogens, *M. catarrhalis* can be transformed in vitro (7); therefore, horizontal transfer of genetic material via transformation followed by homologous recombination is a possible mechanism for generating mosaicism in the *uspA1* and *uspA2* ORFs. In addition, the presence of both a *uspA1* gene and a *uspA2* (or *uspA2H*) gene in each *M. catarrhalis* strain studied to date indicates that intrachromosomal recombination between *uspA1* and *uspA2* loci could also be involved.

The origin of the *uspA2H* genes remains open to speculation. However, it is apparent that the *uspA2H* gene was not just the result of a simple double-crossover event between a *uspA1* gene and a *uspA2* gene. The 3' end of *uspA2H* is virtually identical to that of *uspA2*, and the 5' end of the *uspA2H* ORF is clearly derived from a *uspA1* gene. However, the 1.2-kb region upstream of the translational start codon in the *uspA2H* ORFs in strains TTA37 and 046E was not derived from either *uspA1* or *uspA2*, thus posing an interesting question about the source of this nucleotide sequence. The presence in this upstream DNA of an ORF encoding a predicted protein similar to that expressed by an insertion element (i.e., *IS1*) (34) raises the possibility that several different genetic events were involved in the development of *uspA2H* genes.

In summary, we have identified two distinct yet closely related adhesins of *M. catarrhalis*. Because bacterial adherence is an important first step in the colonization of the upper respiratory tract by *M. catarrhalis*, the findings reported in this study have implications for both vaccine development and the design of therapeutic approaches which could interfere with the adherence properties conferred by UspA1 or UspA2H. It may be feasible to prevent establishment of *M. catarrhalis* in the nasopharynx, thus precluding subsequent development of otitis media caused by this organism.

ACKNOWLEDGMENTS

This study was supported by Public Health Service grant AI36344 and by Texas Advanced Technology Program Award 003660-087 to E.J.H. C.A. was supported by a research grant for young investigators from Novartis AG, Basel, Switzerland.

We thank Steven Berk, John Nelson, and Frederick Henderson for providing isolates of *M. catarrhalis*.

REFERENCES

1. Aebi, C., L. D. Cope, J. L. Latimer, S. E. Thomas, C. A. Slaughter, G. H. McCracken, Jr., and E. J. Hansen. 1998. Mapping of a protective epitope of the CopB outer membrane protein of *Moraxella catarrhalis*. *Infect. Immun.* 66:540-548.
2. Aebi, C., E. R. Lafontaine, L. D. Cope, J. L. Latimer, S. R. Lumbley, G. H. McCracken, Jr., and E. J. Hansen. 1998. Phenotypic effect of isogenic *uspA1* and *uspA2* mutations on *Moraxella catarrhalis* 035E. *Infect. Immun.* 66:3113-3119.
3. Aebi, C., I. Maciver, J. L. Latimer, L. D. Cope, M. K. Stevens, S. E. Thomas,

- G. H. McCracken, Jr., and E. J. Hansen. 1997. A protective epitope of *Moraxella catarrhalis* is encoded by two different genes. *Infect. Immun.* **65**:4367–4377.
4. Aebi, C., B. Stone, M. Beucher, L. D. Cope, I. Maciver, S. E. Thomas, G. H. McCracken, Jr., P. F. Sparling, and E. J. Hansen. 1996. Expression of the CopB outer membrane protein by *Moraxella catarrhalis* is regulated by iron and affects iron acquisition from transferrin and lactoferrin. *Infect. Immun.* **64**:2024–2030.
 5. Beaulieu, D., M. Ouellette, M. G. Bergeron, and P. H. Roy. 1988. Characterization of a plasmid isolated from *Branhamella catarrhalis* and detection of plasmid sequences within the genome of a *B. catarrhalis* strain. *Plasmid* **20**:158–162.
 6. Bluestone, C. D., J. S. Stephenson, and L. M. Martin. 1992. Ten-year review of otitis media pathogens. *Pediatr. Infect. Dis. J.* **11**:S7–S11.
 7. Catlin, B. W., and L. S. Cunningham. 1964. Genetic transformation of *Neisseria catarrhalis* by deoxyribonucleate preparations having different average base compositions. *J. Gen. Microbiol.* **37**:341–352.
 8. Chen, D., V. Barniak, K. R. VanDerMeid, and J. C. McMichael. 1999. The levels and bactericidal capacity of antibodies directed against the UspA1 and UspA2 outer membrane proteins of *Moraxella (Branhamella) catarrhalis* in adults and children. *Infect. Immun.* **67**:1310–1316.
 9. Chen, D., J. C. McMichael, K. R. van der Meid, D. Hahn, T. Mininni, J. Cowell, and J. Eldridge. 1996. Evaluation of purified UspA from *Moraxella catarrhalis* as a vaccine in a murine model after active immunization. *Infect. Immun.* **64**:1900–1905.
 10. Cope, L. D., E. R. Lafontaine, C. A. Slaughter, C. A. Hasemann, Jr., C. Aebi, F. W. Henderson, and G. H. McCracken, Jr. 1999. Characterization of the *Moraxella catarrhalis* *uspA1* and *uspA2* genes and their encoded products. *J. Bacteriol.* **181**:4026–4034.
 11. Cope, L. D., S. R. Lumbley, J. L. Latimer, J. Klesney-Tait, M. K. Stevens, L. S. Johnson, M. Purven, R. S. Munson, Jr., T. Lagergard, J. D. Radolf, and E. J. Hansen. 1997. A diffusible cytotoxin of *Haemophilus ducreyi*. *Proc. Natl. Acad. Sci. USA* **94**:4056–4061.
 12. Du, R. P., Q. J. Wang, Y.-P. Yang, A. B. Schryvers, P. Chong, M. H. Klein, and S. M. Loosmore. 1998. Cloning and expression of the *Moraxella catarrhalis* lactoferrin receptor genes. *Infect. Immun.* **66**:3656–3665.
 13. Fitzgerald, M., R. Mulcahy, S. Murphy, C. Keane, D. Coakley, and T. Scott. 1997. A 200 kDa protein is associated with haemagglutinating isolates of *Moraxella (Branhamella) catarrhalis*. *FEMS Immunol. Med. Microbiol.* **18**:209–216.
 14. Fitzgerald, M., R. Mulcahy, S. Murphy, C. Keane, D. Coakley, and T. Scott. 1999. Transmission electron microscopy studies of *Moraxella (Branhamella) catarrhalis*. *FEMS Immunol. Med. Microbiol.* **23**:57–66.
 15. Gu, X.-X., J. Chen, S. J. Barenkamp, J. B. Robbins, C.-M. Tsai, D. J. Lim, and J. Bettey. 1998. Synthesis and characterization of lipooligosaccharide-based conjugates as vaccine candidates for *Moraxella (Branhamella) catarrhalis*. *Infect. Immun.* **66**:1891–1897.
 16. Hager, H., A. Verghese, S. Alvarez, and S. L. Berk. 1987. *Branhamella catarrhalis* respiratory infections. *Rev. Infect. Dis.* **9**:1140–1149.
 17. Halter, R., J. Pohlner, and T. F. Meyer. 1989. Mosaic-like organization of IgA protease genes in *Neisseria gonorrhoeae* generated by horizontal genetic exchange *in vivo*. *EMBO J.* **8**:2737–3744.
 18. Hansen, E. J., F. R. Gonzales, N. R. Chamberlain, M. V. Norgard, E. E. Miller, L. D. Cope, S. E. Pelzel, B. Gaddy, and A. Clausell. 1988. Cloning of the gene encoding the major outer membrane protein of *Haemophilus influenzae* type b. *Infect. Immun.* **56**:2709–2716.
 19. Helminen, M. E., I. Maciver, J. L. Latimer, L. D. Cope, G. H. McCracken, Jr., and E. J. Hansen. 1993. A major outer membrane protein of *Moraxella catarrhalis* is a target for antibodies that enhance pulmonary clearance of the pathogen in an animal model. *Infect. Immun.* **61**:2003–2010.
 20. Helminen, M. E., I. Maciver, J. L. Latimer, J. Klesney-Tait, L. D. Cope, M. M. Paris, G. H. McCracken, Jr., and E. J. Hansen. 1994. A large, antigenically conserved protein on the surface of *Moraxella catarrhalis* is a target for protective antibodies. *J. Infect. Dis.* **170**:867–872.
 21. Helminen, M. E., I. Maciver, J. L. Latimer, S. R. Lumbley, L. D. Cope, G. H. McCracken, Jr., and E. J. Hansen. 1993. A mutation affecting expression of a major outer membrane protein of *Moraxella catarrhalis* alters serum resistance and survival of this organism *in vivo*. *J. Infect. Dis.* **168**:1194–1201.
 22. Henderson, I. R., F. Navarro-Garcia, and J. P. Nataro. 1998. The great escape: structure and function of the autotransporter proteins. *Trends Microbiol.* **6**:370–378.
 23. Hobbs, M. M., A. Seiler, M. Achtman, and J. G. Cannon. 1994. Microevolution within a clonal population of pathogenic bacteria: recombination, gene duplication and horizontal genetic exchange in the *opa* gene family of *Neisseria meningitidis*. *Mol. Microbiol.* **12**:171–180.
 24. Hsiao, C. B., S. Sethi, and T. F. Murphy. 1995. Outer membrane protein CD of *Branhamella catarrhalis*: sequence conservation in strains recovered from the human respiratory tract. *Microb. Pathog.* **19**:215–225.
 25. Jordan, K. L., S. H. Berk, and S. L. Berk. 1990. A comparison of serum bactericidal activity and phenotypic characteristics of bacteremic, pneumonia-causing strains, and colonizing strains of *Branhamella catarrhalis*. *Am. J. Med.* **88**(5A):28S–32S.
 26. Klingman, K. L., and T. F. Murphy. 1994. Purification and characterization of a high-molecular-weight outer membrane protein of *Moraxella (Branhamella) catarrhalis*. *Infect. Immun.* **62**:1150–1155.
 27. Kyd, J. M., A. W. Cripps, and T. F. Murphy. 1998. Outer-membrane antigen expression by *Moraxella (Branhamella) catarrhalis* influences pulmonary clearance. *J. Med. Microbiol.* **47**:159–168.
 28. Luke, N. R., T. A. Russo, N. Luther, and A. A. Campagnari. 1999. Use of an isogenic mutant constructed in *Moraxella catarrhalis* to identify a protective epitope of outer membrane B1 defined by monoclonal antibody 11C6. *Infect. Immun.* **67**:681–687.
 29. Maiden, M. C. J. 1993. Population genetics of a transformable bacterium: the influence of horizontal genetic exchange on the biology of *Neisseria meningitidis*. *FEMS Microbiol. Lett.* **112**:243–250.
 30. McGraw, E. A., J. Li, R. K. Selander, and T. S. Whittman. 1999. Molecular evolution and mosaic structure of alpha, beta and gamma intimins of pathogenic *Escherichia coli*. *Mol. Biol. Evol.* **16**:12–22.
 31. McMichael, J. C., M. J. Fiske, R. A. Fredenburg, D. N. Chakravarti, K. R. VanDerMeid, V. Barniak, J. Caplan, E. Bortell, S. Baker, R. Arumugham, and D. Chen. 1998. Isolation and characterization of two proteins from *Moraxella catarrhalis* that bear a common epitope. *Infect. Immun.* **66**:4374–4381.
 32. Murphy, T. F. 1996. *Branhamella catarrhalis*: epidemiology, surface antigenic structure, and immune response. *Microbiol. Rev.* **60**:267.
 33. Murphy, T. F., J. M. Kyd, A. John, C. Kirkham, and A. W. Cripps. 1998. Enhancement of pulmonary clearance of *Moraxella (Branhamella) catarrhalis* following immunization with outer membrane protein CD in a mouse model. *J. Infect. Dis.* **178**:1667–1675.
 34. Ohtsubo, H., K. Nyman, W. Doroszkiewicz, and E. Ohtsubo. 1981. Multiple copies of iso-insertion sequence of IS1 in *Shigella dysenteriae* chromosome. *Nature* **292**:640–643.
 35. Pelton, S. I., and J. O. Klein. 1999. The promise of immunoprophylaxis for prevention of acute otitis media. *Pediatr. Infect. Dis. J.* **18**:926–935.
 36. Reddy, M. S., T. F. Murphy, H. S. Faden, and J. M. Bernstein. 1997. Middle ear mucin glycoprotein: purification and interaction with nontypeable *Haemophilus influenzae* and *Moraxella catarrhalis*. *Otolaryngol. Head Neck Surg.* **116**:175–180.
 37. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 38. Sethi, S., J. M. Surface, and T. F. Murphy. 1997. Antigenic heterogeneity and molecular analysis of CopB of *Moraxella (Branhamella) catarrhalis*. *Infect. Immun.* **65**:3666–3671.
 39. Setlow, J. K., D. C. Brown, M. E. Boling, A. Mattingly, and M. P. Gordon. 1968. Repair of deoxyribonucleic acid in *Haemophilus influenzae*. *J. Bacteriol.* **95**:546–558.
 40. St. Geme, J. W., III, M. L. de la Morena, and S. Falkow. 1994. A *Haemophilus influenzae* IgA protease-like protein promotes intimate interaction with human epithelial cells. *Mol. Microbiol.* **14**:217–233.
 41. Verduin, C. M., M. Jansze, C. Hol, T. E. Molles, J. Verhoef, and H. van Dijk. 1994. Differences in complement activation between complement-resistant and complement-sensitive *Moraxella (Branhamella) catarrhalis* strains occur at the level of membrane attack complex formation. *Infect. Immun.* **62**:589–595.
 42. Whithy, P. W., D. J. Morton, and T. L. Stull. 1998. Construction of antibiotic resistance cassettes with multiple paired restriction sites for insertional mutagenesis of *Haemophilus influenzae*. *FEMS Microbiol. Lett.* **158**:57–60.
 43. Yang, Y.-P., L. E. Myers, U. McGuinness, P. Chong, Y. Kwok, M. H. Klein, and R. E. Harkness. 1997. The major outer membrane protein, CD, extracted from *Moraxella (Branhamella) catarrhalis* is a potential vaccine antigen that induces bactericidal antibodies. *FEMS Immunol. Med. Microbiol.* **17**:187–199.