

Identification of SopE2, a *Salmonella* Secreted Protein Which Is Highly Homologous to SopE and Involved in Bacterial Invasion of Epithelial Cells

C. S. BAKSHI,^{1,2} V. P. SINGH,^{1,2} M. W. WOOD,¹ P. W. JONES,¹ T. S. WALLIS,¹ AND E. E. GALYOV^{1*}

Institute for Animal Health, Compton, Berkshire RG20 7NN, United Kingdom,¹ and Indian Veterinary Research Institute, Izatnagar, Bareilly 243122, India²

Received 12 November 1999/Accepted 31 January 2000

Type III secreted Sop protein effectors are delivered into target eukaryotic cells and elicit cellular responses underlying *Salmonella* pathogenicity. In this work, we have identified another secreted protein, SopE2, and showed that SopE2 is an important invasion-associated effector. SopE2 is encoded by the *sopE2* gene which is present and conserved in pathogenic strains of *Salmonella*. SopE2 is highly homologous to SopE, a protein encoded by a gene within a temperate bacteriophage and present in only some pathogenic strains.

A variety of gram-negative pathogenic bacteria possess a dedicated protein secretion system, denoted type III, which is of primary importance for the successful engagement of eukaryotic host cells (for a recent review, see reference 8). Type III secretion systems play a central role in virulence, directing secretion and translocation of several bacterial effector proteins into the cytoplasm of host cells. It is becoming apparent that the mechanism of type III secretion and translocation is conserved in a variety of bacteria, although each pathogen appears to have a specific set of secreted protein effectors. Upon translocation, the bacterial protein effectors affect different host cell functions and elicit a variety of responses. This, in turn, has a major impact on the development and progress of infection.

Recent work by our group and others has identified a number of protein effectors (3, 4, 5, 9, 12, 17) secreted and translocated into eukaryotic cells by the *Salmonella* Inv/Spa type III secretion system, which is one of two type III systems present in salmonellae (for a review, see reference 2). The Inv/Spa system is encoded within a pathogenicity island, SPI-1 (13), and is required for eliciting host cell responses which in turn result in host cell cytoskeleton rearrangements which assist bacterial entry into nonphagocytic epithelial cells and aid in the production of proinflammatory signals (for a recent review, see reference 1). Four of the key proteins in these processes are Sip A, B, C, and D. All these proteins are encoded by a single polycistronic operon located adjacent to the *inv/spa* loci (7, 10, 11, 16). At least some of the Sip proteins appear to have multiple activities and perform several functions, but it is believed that the major function of Sip B, C, and D is to execute the translocation of a set of specific effector proteins into eukaryotic cells. One such translocated effector is SopE (5, 17). SopE contributes to the expression of *Salmonella* invasion by stimulating membrane ruffling via guanidine nucleotide exchange on Rho GTPases CDC42 and Rac (6). The *sopE* gene is located outside SPI-1 and is encoded within a temperate bacteriophage (5, 15). Interestingly, not all strains of *Salmonella* carry the *sopE* gene (15), suggesting either that the SopE function is redundant or that there is another protein with

functions similar to those of SopE, in at least some *Salmonella* strains. We have investigated this suggestion.

SopE was initially identified as a component of protein aggregates accumulated in the culture media of *Salmonella enterica* serovar Dublin B1 (*sipB* mutant) (17). This suggested that *sipB* mutants of different *Salmonella* strains may have different profiles of Sop proteins deposited into filamentous aggregates. In order to assess the Sop protein profiles from *Salmonella* strains with different host specificities, we first transduced the polar *sipB* mutation from serovar Dublin B1 into a variety of other *Salmonella* strains (virulent field isolates from the IAH strain collection) by using P22 transduction. The correct mutations in Cm^r transfectants (one for each strain) were confirmed by PCR (data not shown). Different *sipB* mutant strains were then grown overnight at 25°C in Luria-Bertani medium, were diluted 10-fold in fresh LB medium, and were incubated for 5 h at 37°C. All *Salmonella* strains tested produced filaments similar to those produced by serovar Dublin B1. The protein filament aggregates accumulated in each flask were collected and washed in phosphate-buffered saline, essentially as described earlier (17). Filaments were dissolved in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) loading buffer, and proteins were separated by SDS-12% PAGE. The gel was stained with Coomassie brilliant blue (Fig. 1A) or was blotted onto a nitrocellulose membrane and probed with an anti-SopE monoclonal antibody (Fig. 1B). The analysis of the Coomassie-stained gel revealed that filament aggregates isolated from each *Salmonella* strain were composed of a similar set of proteins, migrating in each case with a profile closely resembling that of serovar Dublin (Fig. 1A). The monoclonal anti-SopE antibodies (17), however, did not recognize the ~30-kDa protein from *Salmonella enterica* serovar Typhimurium F98 isolated from a chicken and serovar Typhimurium ST5306 isolated from a pig (Fig. 1B). In both cases, however, this protein appeared to migrate with the same apparent molecular weight as SopE from serovar Dublin. We then attempted to amplify a *sopE*-specific DNA fragment by PCR by using genomic DNA from different *Salmonella* strains as a template with SEup (5'-ACCGAGAAAAGATCTTTAG CAAAAA-3') and SEdown (5'-CAAGATCAGCTCACACT-3') primers, based on the known DNA sequence of *sopE* from serovars Dublin and Typhimurium. No DNA fragment was amplified when DNA from serovar Typhimurium F98 or ST5306 was used as a template, but a *sopE*-specific DNA

* Corresponding author. Mailing address: Institute for Animal Health, Compton, Berkshire RG20 7NN, United Kingdom. Phone: (44) 1635 577291. Fax: (44) 1635 577243. E-mail: edouard.galyov@bbsrc.ac.uk.




FIG. 1. Analysis of the proteins secreted by different *Salmonella* strains (*sipB* mutants). The secreted proteins deposited by each strain into proteinaceous filaments were isolated as described earlier (17). The protein samples from serovar Dublin B1 (lane 1), *Salmonella enterica* serovar Gallinarum S9.B1 (lane 2), *Salmonella enterica* serovar Enteritidis S13.B1 (lane 3), serovar Typhimurium F98.B1 (lane 4), serovar Typhimurium ST4/74.B1 (lane 5), serovar Typhimurium ST5306.B1 (lane 6), and *Salmonella enterica* serovar Choleraesuis A57.B1 (lane 7) were separated by SDS-12% PAGE and were stained with Coomassie brilliant blue (A) or were transferred onto nitrocellulose membrane and probed with an anti-SopE monoclonal antibody (B). Molecular mass is shown on the left.

fragment of the expected size was amplified in all other samples (data not shown). This further suggested that serovar Typhimurium F98 and ST5306 did not possess *sopE* and that the ~30-kDa protein from these two strains may be a different protein. To investigate this protein further, we extracted the 30-kDa protein produced by serovar Typhimurium F98, subjected it to N-terminal amino acid sequencing, and used it for immunization of mice to raise monoclonal antibodies. The sequence of 15 amino acids was determined to be X-N-I-T-L-S-T-Q-H-Y-R-I-H-R-S. This sequence was then analyzed by using the BLAST search program against a partial genomic sequence of serovar Typhimurium and *Salmonella enterica* serovar Paratyphi available from the Genome Sequencing Center at Washington University School of Medicine, St. Louis, Mo. (<http://genome.wustl.edu/gsc/bacterial/salmonella.shtml>) and against that of serovar Typhi available from Sanger Center (http://www.sanger.ac.uk/Projects/S_typhi/blast_server.shtml). This analysis revealed a match with the product of an open reading frame (ORF) present in all three *Salmonella* serotypes. (In the available nucleotide sequence from serovar Typhi there is a deletion of one nucleotide at position 73 within this ORF leading to a frame shift. It is unclear at this moment if this is due to a possible sequencing mistake.) The deduced amino acid sequence of the protein product of this ORF was highly homologous to SopE (see below). We therefore designated this protein SopE2, and the ORF was designated *sopE2*. DNA fragments containing a complete *sopE2* gene and upstream areas from serovars Dublin 2229 and Typhimurium F98 were amplified by PCR by using an upstream primer SE3 (5'-TTA

```

SopE: 1 MTKITLFFHNFRIQKQEAETPLKEKSTERNLAKSILAVKNHFILKLSKLS 50
      | | | | | : : : : : | : : : : : | : : : : : | : : : : : |
SopE2: 1 MTNITLSTQHYRIHRSDVEVPVKEKTEKDI FAKSITAVRNSFISLSTLSLS 50

SopE: 51 ERFISHKNTSSATHFHRGSASEGRAVLTKNKVKNFMQLTLHDIDIRGSA 100
      : | | : : : : : | : : : : : | : : : : : | : : : : : |
SopE2: 51 DRFSLHQQTIDIPTHFHRGNASEGRAVLTSKTVKDEMLQLKNSLDIKGNA 100

SopE: 101 SKDPAYASQTREATLSAVYSKYKDYCNLLISKGIDIAPELKEIGEAAQN 150
      | | | | | | | : | : | | | | | | | : | : | | | | | | |
SopE2: 101 SKDPAYARQTCDAMLSAVYSNNKHCCCKLLISKGVSIITPFLKEIGEAAQN 150

SopE: 151 AGLPGATKNDVFSFSGAGANPFITPLITSAYSKYPHMFTSQHQKASFNLY 200
      | | | | | | | : | : | | | | | | | : | | | | | | | | |
SopE2: 151 AGLPGEIKNGVFTPGGAGANPFVVPLIAAASIKYPHMFINHQVSVFKAY 200

SopE: 201 AEKIIMTEVVPVLFNECAMPTPQQFQOILENIANKYIQNTPT 240
      | | | | | | | | | | | | | | | | | | | | | | |
SopE2: 201 AEKIVMKEVTPLEFNKGTMPPTPQQFQOLTENIANKYLQNAS 240

```

FIG. 2. Sequence alignment of serovars Dublin SopE and Typhimurium F98 SopE2. Lines indicate identical amino acids, and colons and periods indicate conservative amino acid substitutions.

AGCATAAGCTTAATTCCATTGTT-3') and a downstream primer SE4 (5'-CTGATAAATGAATTCAGGCCGC ATC-3') based on the available sequence from serovar Typhi and modified to include restriction sites. Chromosomal DNA from serovars Dublin 2229 and Typhimurium F98 were used as templates. The resulting DNA fragments were cloned into pBluescript to yield pSopE2sd and pSopE2st, respectively. The sequence of the cloned DNA in both plasmids was identical to that of an ORF obtained from the serovar Typhimurium genome database. The deduced amino acid sequence of the SopE2 protein was 64% identical to that of SopE (Fig. 2). The SopE2 sequence was found to be remarkably conserved in different *Salmonella* serotypes, with only six amino acid substitutions in SopE2 in serovar Paratyphi compared to that in serovars Typhimurium and Dublin. We next designed two primers, SE2up (5'-TCGTGGGAGCGGATCCGAGGGTAG GGCAGTAT-3') and SE2down (5'-TGCGCAGCCTCGAGT ATCTCTTTTCAGAAA-3'), from an internal part of *sopE2* and used them in PCR with DNA from different *Salmonella* strains as template. The *sopE2*-specific DNA was present in all *Salmonella* strains analyzed, including strains with different host specificities (data not shown). Together, these data suggest that the *sopE2* sequence is present and conserved in different *S. enterica* strains.

A DNA fragment amplified by using serovar Typhimurium DNA was cloned into the pDM4 suicide plasmid (14) to yield pSB1. *sopE2* mutant serovar Typhimurium SE2.1 was constructed by crossing pSB1 from *Escherichia coli* S17.1 λ pir into parental serovar Typhimurium F98, and correct insertion of the suicide plasmid was confirmed by PCR. The mutation was trans complemented by introducing the pSopE2st plasmid into serovar Typhimurium SE2.1. The growth characteristics of the *sopE2* mutant and the trans complemented strains in LB medium at 37°C were undistinguishable from those of the wild-type strain (data not shown). The proteins produced and secreted by the different *Salmonella* strains in LB medium at 37°C were isolated as described earlier (17) and were analyzed by SDS-PAGE and Western blotting by using monoclonal anti-SopE2 antibodies. The SDS-PAGE analysis revealed that the *sopE2* mutation did not cause any apparent effect on the expression and secretion of secreted proteins other than a 30-kDa protein (data not shown). The Western blot analysis revealed that a faint protein band at 30 kDa, present in the wild-type strain, was missing in the *sopE2* mutant but reappeared in the trans complemented strain (Fig. 3), suggesting that the observed 30-kDa protein is SopE2.

The wild-type, mutant, and trans complemented strains were




FIG. 3. Western blot analysis of the secreted proteins of the wild-type serovar Typhimurium F98 (lane 2), serovar Typhimurium SE2.1 *sopE2* mutant (lane 3), and serovar Typhimurium SE2.1/pSopE2st transcomplemented *sopE2* mutant (lane 4) strains. Culture supernatant proteins from different *Salmonella* strains were prepared by precipitation with 10% trichloroacetic acid as previously described (17), were separated by SDS-PAGE, were transferred to nitrocellulose membrane, and were probed with anti-SopE2 monoclonal antibody.

then assessed in a cultured cell invasion assay, essentially as described earlier (17). The results of this experiment clearly showed that serovar Typhimurium SE2.1 had a reduced ability to invade HeLa cells (Fig. 4). The invasion phenotype was partially restored by complementing the mutations in trans, demonstrating that the invasion defect in the mutant strain is due to inactivation of *sopE2*.

Thus, we have identified SopE2, an invasion-associated secreted protein of *Salmonella*. Structurally, *sopE2* is very similar to *sopE*, suggesting that these two genes have the same evolu-




FIG. 4. Invasion of HeLa cells by the wild-type serovar Typhimurium F98, serovar Typhimurium SE2.1 *sopE2* mutant, and serovar Typhimurium SE2.1/pSopE2st transcomplemented *sopE2* mutant strains. All strains were grown overnight at 26°C, then diluted 10-fold and grown at 37°C to an optical density at 600 nm of 0.5. The cells were infected with bacteria at an approximate ratio of 1:10. HeLa cells were infected for 1 h and then washed and exposed to gentamicin (100 µg/ml) treatment for 1 h. The cells were lysed with 1% Triton X-100, and the numbers of bacteria surviving gentamicin treatment were calculated by viable count. The experiments were carried out in triplicate. Each bar is derived from the mean of three samples per strain. The error bars indicate standard errors of the means calculated from triplicate plating.

tionary origin and/or that one of these genes is a product of gene duplication. The high structural similarity between SopE2 and SopE and the similar phenotypic effects caused by inactivation of SopE- and SopE2-encoding genes suggest that these two proteins are likely to have similar mechanisms of action. Unlike *sopE*, *sopE2* appears to be broadly distributed in salmonellae, suggesting that conservation of this gene may be more important than that of *sopE*. Thus, it appears that at least some *Salmonella* strains possess two functional genes, encoding virulence-associated type III secreted proteins highly similar to each other. This finding prompted us to compare sequences of other known secreted proteins with sequences deposited in the *Salmonella* genome database. No homologues were found for SopB (3), AvrA (4), or SptP (12). However, our analysis revealed that there is a gene encoding a SopD (9) homologue. The sequence of this protein is 41% identical to that of SopD (data not shown).

The type III secretion/translocation mechanism appears to be conserved in many gram-negative bacteria; however, each pathogen has a specific set of type III secreted effector proteins. An ability to expand the repertoire of such effectors without the risk of losing essential functions may allow pathogens more flexibility in adapting within different host environments. It is interesting to note a remarkable conservation of the SopE2 sequence in different *Salmonella* serotypes, as this is in contrast to the conservation of SopE. SopE proteins from serovars Dublin and Typhimurium, for example, show only 90% overall identity, indicating a rapid evolution of the *sopE* gene in *Salmonella*. It is also worthy to note that the *sopE* gene is carried by a bacteriophage (5, 15). Together, these data suggest that *Salmonella* appears to possess a mechanism which enables the preservation of one copy of some genes encoding effector proteins while allowing simultaneous rapid evolution of the gene sequence. Those mutated gene variants that provide selective advantage can then be disseminated throughout the population via mobile genetic elements.

Nucleotide sequence accession number. The sequence of the DNA fragment from serovar Typhimurium F98 identical to that from serovar Dublin 2229 was deposited in the EMBL database under accession no. AF200952.

The work presented here was supported by the Biotechnology and Biological Sciences Research Council of the United Kingdom; the United Kingdom Ministry of Agriculture, Fisheries and Food; the India/UK TOMBIT project financed by the United Kingdom Department for International Development; and the Indian Council for Agricultural Research.

C.S.B. and V.P.S. contributed equally to this work.

REFERENCES

1. Darwin, K. H., and V. L. Miller. 1999. Molecular basis of the interaction of *Salmonella* with the intestinal mucosa. *Clin. Microbiol. Rev.* **12**:405–428.
2. Galán, J. E. 1999. Interaction of *Salmonella* with host cells through the centisome 63 type III secretion system. *Curr. Opin. Microbiol.* **2**:46–50.
3. Galyov, E. E., M. W. Wood, R. Rosqvist, P. B. Mullan, P. R. Watson, S. Hedges, and T. S. Wallis. 1997. A secreted effector protein of *Salmonella dublin* is translocated into eukaryotic cells and mediates inflammation and fluid secretion in infected ileal mucosa. *Mol. Microbiol.* **25**:903–912.
4. Hardt, W.-D., and J. E. Galán. 1997. A secreted *Salmonella* protein with homology to a virulence determinant of plant pathogenic bacteria. *Proc. Natl. Acad. Sci. USA* **94**:9887–9892.
5. Hardt, W.-D., H. Urlaub, and J. E. Galán. 1998. A target of the centisome 63 type III protein secretion system of *Salmonella typhimurium* is encoded by a cryptic bacteriophage. *Proc. Natl. Acad. Sci. USA* **95**:2574–2579.
6. Hardt, W.-D., L.-M. Chen, K. E. Shuebel, X. R. Bustello, and J. E. Galán. 1998. *Salmonella typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell* **93**:815–826.
7. Hermant, D., R. Menard, N. Arricau, C. Parsot, and M. Y. Popoff. 1995. Functional conservation of the *Salmonella* and *Shigella* effectors of entry into epithelial cells. *Mol. Microbiol.* **17**:781–789.

8. Hueck, C. J. 1998. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* **62**:379–433.
9. Jones, M. A., M. W. Wood, P. B. Mullan, P. R. Watson, T. S. Wallis, and E. E. Galyov. 1998. Secreted effector proteins of *Salmonella dublin* act in concert to induce enteritis. *Infect. Immun.* **66**:5799–5804.
10. Kaniga, K., D. Trollinger, and J. E. Galán. 1995. Identification of two targets of the type III secretion system encoded in the *inv* and *spa* loci of *Salmonella typhimurium* that share homology to IpaD and IpaA proteins. *J. Bacteriol.* **177**:7078–7085.
11. Kaniga, K., S. G. Tucker, D. Trollinger, and J. E. Galán. 1995. Homologues of the *Shigella* invasins IpaB and IpaC are required for *Salmonella typhimurium* entry into cultured cells. *J. Bacteriol.* **177**:3965–3971.
12. Kaniga, K., J. Uralil, J. B. Bliska, and J. E. Galán. 1996. A secreted tyrosine phosphatase with modular effector domains encoded by the bacterial pathogen *Salmonella typhimurium*. *Mol. Microbiol.* **21**:633–641.
13. Mills, D. M., V. Bajaj, and C. A. Lee. 1995. A 40 kb chromosomal fragment encoding *Salmonella typhimurium* invasion genes is absent from the corresponding region of the *Escherichia coli* K-12 chromosome. *Mol. Microbiol.* **15**:749–759.
14. Milton, D. L., R. O'Toole, P. Hörstedt, and H. Wolf-Watz. 1996. Flagellin A is essential for the virulence of *Vibrio anguillarum*. *J. Bacteriol.* **178**:1310–1319.
15. Mirold, S., W. Rabsch, M. Rohde, S. Stender, H. Tschäpe, H. Rüssmann, E. Igwe, and W.-D. Hardt. 1999. Isolation of a temperate bacteriophage encoding the type III effector protein SopE from epidemic *Salmonella typhimurium* strain. *Proc. Natl. Acad. Sci. USA* **96**:9845–9850.
16. Pegues, D. A., M. J. Hantman, I. Behlau, and S. I. Miller. 1995. PhoP/PhoQ transcriptional repression of *Salmonella typhimurium* invasion genes: evidence for a role in protein secretion. *Mol. Microbiol.* **17**:169–181.
17. Wood, M. W., R. Rosqvist, P. B. Mullan, M. H. Edwards, and E. E. Galyov. 1996. SopE, a secreted protein of *Salmonella dublin*, is translocated into the target eukaryotic cell via a sip-dependent mechanism and promotes bacterial entry. *Mol. Microbiol.* **22**:327–338.