

Differential Roles of the *Pseudomonas aeruginosa* PA14 *rpoN* Gene in Pathogenicity in Plants, Nematodes, Insects, and Mice

ERIK L. HENDRICKSON,^{1,2†} JOULIA PLOTNIKOVA,^{1,2} SHALINA MAHAJAN-MIKLOS,^{1,2‡}
LAURENCE G. RAHME,^{3,4} AND FREDERICK M. AUSUBEL^{1,2*}

Department of Genetics¹ and Department of Surgery,³ Harvard Medical School, and Department of Molecular Biology²
and Shriners' Burns Institute,⁴ Massachusetts General Hospital, Boston, Massachusetts 02114

Received 4 June 2001/Accepted 18 September 2001

We cloned the *rpoN* (*ntrA*, *glnF*) gene encoding the alternate sigma factor σ^{54} from the opportunistic multihost pathogen *Pseudomonas aeruginosa* strain PA14. A marker exchange protocol was used to construct the PA14 *rpoN* insertional mutation *rpoN::Gen^r*. PA14 *rpoN::Gen^r* synthesized reduced levels of pyocyanin and displayed a variety of phenotypes typical of *rpoN* mutants, including a lack of motility and the failure to grow on nitrate, glutamate, or histidine as the sole nitrogen source. Compared to wild-type PA14, *rpoN::Gen^r* was ca. 100-fold less virulent in a mouse thermal injury model and was significantly impaired in its ability to kill the nematode *Caenorhabditis elegans*. In an *Arabidopsis thaliana* leaf infectivity assay, although *rpoN::Gen^r* exhibited significantly reduced attachment to trichomes, stomata, and the epidermal cell surface, did not attach perpendicularly to or perforate mesophyll cell walls, and proliferated less rapidly in *Arabidopsis* leaves, it nevertheless elicited similar disease symptoms to wild-type *P. aeruginosa* PA14 at later stages of infection. *rpoN::Gen^r* was not impaired in virulence in a *Galleria mellonella* (greater wax moth) pathogenicity model. These data indicate that *rpoN* does not regulate the expression of any genes that encode virulence factors universally required for *P. aeruginosa* pathogenicity in diverse hosts.

In gram-negative bacteria, the alternate sigma factor σ^{54} , working in concert with a transcriptional activator that belongs to the NtrC superfamily, activates a variety of genes that are regulated in response to external stimuli (1). For example, in various bacteria, σ^{54} is required for expression of the enzymatic pathways responsible for nitrogen utilization, dicarboxylate transport, xylene degradation, and hydrogen utilization (6, 32, 39, 41, 61). σ^{54} is also involved in the regulation of virulence-related factors in both plant and animal pathogens, including pilin, flagellin, and alginate synthesis in *Pseudomonas aeruginosa* (19, 58, 60); capsular expression in *Klebsiella pneumoniae* (3); and regulation of *hrp* gene expression and coronatine biosynthesis in *Pseudomonas syringae* (23, 24).

Our laboratory has developed a bacterial pathogenicity model that utilizes a clinical isolate of *P. aeruginosa* (strain UCBPP-PA14 [referred to here as PA14]) that elicits severe soft-rot-like symptoms and proliferates when infiltrated into *Arabidopsis* leaves (52), kills the larvae of the wax moth caterpillar *Galleria mellonella* (30), causes lethal sepsis in a mouse full-skin-thickness burn model (52), and kills the nematode *Caenorhabditis elegans* (40, 56, 57). Interestingly, there is significant overlap among the PA14 virulence factors required for pathogenesis in plants, nematodes, insects, and mice. For example, among 21 genes identified as being involved in pathogenesis by screening transposon-induced PA14 mutants in plants and nematodes, 18, 17, 19, and 21 of these genes were

required for pathogenicity in *Arabidopsis*, nematodes, wax moths, and mice, respectively (30, 40, 53, 57).

In other studies, we showed that *rpoN* is a key virulence factor for the plant pathogen *P. syringae* (23, 24). Specifically, molecular and genetic analysis showed that the *P. syringae rpoN* gene is required for expression of the *P. syringae hrp* gene cluster, a block of contiguous genes, some of which encode components of a type III secretory system (2, 18, 26, 45, 46, 59).

Given the facts that RpoN activates the expression of a wide variety of environmentally regulated genes and is required for virulence in a variety of pathogens, we hypothesized that RpoN would play a central role in the evolution of *P. aeruginosa's* ability to be a pathogen of evolutionarily disparate hosts. In this study we describe the results of experiments that involved the construction of a *P. aeruginosa* PA14 *rpoN* mutant to study the role of σ^{54} in *P. aeruginosa* pathogenesis in a variety of plant and animal hosts. Surprisingly, we report that the *P. aeruginosa rpoN* gene is not a universal virulence factor required for multihost pathogenesis.

MATERIALS AND METHODS

Bacterial strains, plasmids, and culture conditions. The bacterial strains and plasmids used and constructed in this study are listed in Table 1. *Escherichia coli* and *P. aeruginosa* strains were grown at 37°C in L broth, King's A (KA), King's B (KB) (33), or M9 minimal salts media. Nitrogen source utilization tests for PA14 *rpoN* mutants were performed in M9 salts minimal medium by replacing ammonium chloride with an alternative nitrogen source at 5 mM when required. Bacterial motility was tested on "swarm plates" (35). Pyocyanin assays (17) were carried out in KA broth containing 100 μ M FeCl₃ (17, 33). Pyoverdinin was assayed on KB plates as described previously (57). *C. elegans* killing assays were carried out on NG agar ("slow killing" [56]) or PGS agar ("fast killing" [56]) as described elsewhere. Antibiotic concentrations for *E. coli* strains were as follows: streptomycin, 150 μ g/ml; kanamycin, 25 μ g/ml; tetracycline, 12 μ g/ml; gentamicin, 5 to 10 μ g/ml; and spectinomycin, 20 μ g/ml. Antibiotic concentrations for *P. aeruginosa* were as follows: streptomycin, 200 μ g/ml; kanamycin, 200 μ g/ml;

* Corresponding author. Mailing address: Department of Molecular Biology, Wellman 10, Massachusetts General Hospital, Boston, MA 02114. Phone: (617) 726-5969. Fax: (617) 726-5949. E-mail: ausubel@molbio.mgh.harvard.edu.

† Present address: Department of Microbiology, University of Washington, Seattle, WA 98195.

‡ Present address: Microbia, Inc., Cambridge, MA 02139.

TABLE 1. Bacterial strains and plasmids

Strain or plasmid	Relevant genotype, phenotype, or role ^a	Source or reference
<i>P. aeruginosa</i>		
PAK N1	<i>rpoN</i> ::Gen ^r	S. Lory (28)
PA14	Wild type	M. Schroth (52)
PA14 <i>rpoN</i> ::Gen ^r	Contains Gen ^r cassette inserted into <i>rpoN</i>	This study
PA14(pSMC21)	PA14 expressing GFP	56
PA14 <i>rpoN</i> ::Gen ^r (pSMC21)	<i>rpoN</i> ::Gen ^r expressing GFP	This study
<i>E. coli</i>		
DH5a	F ⁻ <i>lacZ</i> ΔM15 <i>endA1 recA1 hsdR17 supE44 thi-1 gyrA relA1</i> λ ⁻ ; host for cosmid library and other plasmids	Bethesda Research Laboratories (20)
MM294(pRK2013)	Donor of transfer functions for triparental crosses	14
Plasmids		
pJSR1	Ap ^r , cosmid cloning vector	52
pBSK(+)	Ap ^r , cloning vector	Stratagene, Inc.
pBR322	Ap ^r Tet ^r , cloning vector	New England Biolabs, Inc.
pKI11	Source of <i>P. aeruginosa</i> PAK <i>rpoN</i> gene	S. Lory (28)
pPAR4	4-kb fragment containing PA14 <i>rpoN</i> gene in pBSK(+)	This study
pPAR4SR	4-kb fragment containing PA14 <i>rpoN</i> gene in pJSR1	This study
pRPONgent	PA14 <i>rpoN</i> ::Gen ^r in pBR322	This study
pRPON10	PA14 <i>rpoN</i> gene in pJSR1	This study
pSMC21	Derivative of plasmid pSMC2; carries the <i>A. victoria</i> GFP	G. A. O'Toole (7)

^a Ap^r, ampicillin resistance; Tet^r, tetracycline resistance.

tetracycline, 75 μg/ml; gentamicin, 30 μg/ml; nalidixic acid, 50 μg/ml; rifampin, 100 μg/ml, and carbenicillin, 300 μg/ml.

Bacterial genetics. pJSR1 derivatives were introduced into *Pseudomonas* strains via triparental matings with MM294/pRK2013 as the donor of transfer functions as described previously (14). Plasmid pSMC21 (7) containing the *Aequorea victoria* green fluorescent protein (GFP) was introduced into PA14 *rpoN*::Gen^r by electroporation.

Pyocyanin assays. Pyocyanin was measured at 520 nm in acidic solution by using a modified version of a previously described method (17). Cultures were grown from a 100-fold dilution of a log-phase culture in KA broth modified with 100 μM FeCl₃. After 20 h, 1 ml of the culture was extracted with 2 ml of chloroform and centrifuged for 5 min. The blue chloroform solution was transferred into a new tube containing 1 ml of 0.2 N HCl to extract pyocyanin into the acidic solution. The concentration was determined by measuring the optical density at 520 nm (OD₅₂₀).

Plant material and growth of plants. *Arabidopsis* ecotypes Llagostera (LI-0) and Landsberg *erecta* (La-er) were obtained from the *Arabidopsis* Biological Resource Center, Columbus, Ohio. *Arabidopsis* plants were grown in Metro-Mix 2000 in either a climate-controlled greenhouse at 19°C under a 12-h light-dark cycle with supplemental fluorescent illumination or in a Percival AR-60L growth chamber at 20°C and 50% relative humidity.

***Arabidopsis* pathogenicity assays.** Six- to eight-week-old intact or detached *Arabidopsis* rosette leaves were used for pathogenicity assays. The pathogenicity of PA14 strains was tested by placing detached LI-0 leaves on a 1.5% water agar surface with their petioles embedded into the agar and inoculating the leaves by growing lawns of PA14 or PA14 *rpoN*::Gen^r overnight on LB agar medium containing rifampin (PA14) and gentamicin (*rpoN*::Gen^r), cutting 3-mm-diameter agar cylinders from these plates, and placing the cylinders bacterial side down on one or both sides of the central vein in the top half of a leaf. The growth of PA14 or PA14 *rpoN*::Gen^r in *Arabidopsis* leaves was determined by infiltrating the leaves of intact La-er plants with 5 × 10⁴ CFU/cm² of leaf area as described previously (52). The growth of PA14 or PA14 *rpoN*::Gen^r in detached *Arabidopsis* leaves was determined by infiltrating LI-0 leaves with a bacterial suspension in 10 mM MgSO₄ at a density of 5 × 10⁴ CFU/ml for 1 h under a slight vacuum at room temperature in the wells of a six-well microtiter plate. At days 0, 1, 2, 3, and 4, the titer of the bacteria in each of two punches in each of three leaves was determined as described previously (15).

Infection of *G. mellonella* larvae. The 50% lethal dose (LD₅₀) of PA14 strains in *G. mellonella* larvae was determined as described previously (30). In brief, overnight cultures grown in KB medium were diluted 1:100, allowed to grow until they reached an OD₆₀₀ of 0.3 to 0.4, and resuspended in 10 mM MgSO₄. After dilution to an OD₆₀₀ of 0.1 with 10 mM MgSO₄, serial 10-fold dilutions were

made in 10 mM MgSO₄ containing 1 mg of rifampin/ml and 10 mg of carbenicillin/ml. A 10-μl Hamilton syringe was used to inject 5-μl aliquots into the hindmost left proleg of fifth-instar *G. mellonella* larvae purchased from Van der Horst Wholesale, St. Mary's, Ohio. Groups of 10 larvae infected with the same dose of bacteria were placed in petri dishes and incubated at 25°C for 60 h. Larvae were scored as dead when they no longer moved upon shaking of the petri dish or poking with a pipette tip.

Mouse full-skin-thickness burn model. Inbred AKR mice that had been subjected to a thermal burn injury were infected with PA14 strains as described previously (54). In brief, 6-week-old male mice were anesthetized by the injection of phenobarbital. The animals were then shaved, and a ventral skin fold was elevated. Two brass blocks preheated to 92 to 95°C were applied to the skin fold for 5 s to deliver a full-skin-thickness burn covering ca. 5% of the body surface area. The burn eschar was injected with 100 μl of bacterial suspension at a titer of 5 × 10⁴ or 5 × 10⁶ bacteria per ml. Bacteria for the inoculation were grown overnight in LB, diluted 1:100, allowed to grow until they reached an OD₆₀₀ of 1.6 to 1.7, pelleted by centrifugation, and resuspended and diluted in 10 mM MgSO₄. The mouse protocol was reviewed and approved by the Animal Care Committee of the Massachusetts General Hospital.

***C. elegans* killing assay.** The killing kinetics of *C. elegans* strain Bristol N2 by PA14 strains were determined on PGS agar (fast-killing assay) or NG agar (slow-killing assay) as described previously (56).

Nucleic acid manipulations. Routine DNA manipulations such as DNA blots and plasmid DNA isolation were performed as described previously (4). Restriction enzymes, T4 DNA ligase, and calf intestine phosphatase were purchased from Boehringer Mannheim and New England BioLabs and used according to the manufacturers' specifications.

Cloning the *P. aeruginosa* PA14 *rpoN* gene and construction of a PA14 *rpoN* mutant. A cosmid clone, pRPON10, containing a presumptive *P. aeruginosa* PA14 *rpoN* gene was identified in a pJSR1 cosmid library (52) by colony hybridization with *P. aeruginosa* strain PAK *rpoN* gene on plasmid pKI11 as a hybridization probe. The presumptive PA14 *rpoN* gene was mapped to a 4.0-kb *Xho*I-*Eco*RI fragment and was subcloned into pBSK(+) to produce pPAR4. DNA sequence analysis showed 95% identity out of 141 nucleotides sequenced between the PA14 *rpoN* gene and the *P. aeruginosa* PAO1 *rpoN* gene (reference 31 and data not shown).

A mutant derivative of the PA14 *rpoN* gene was constructed by inserting a DNA cassette conferring gentamicin resistance (Gen^r) into the *Cl*aI site of the PA14 *rpoN* gene in pPAR4 to create *rpoN*::Gen^r and then excising a 5.5-kb *Xho*I-*Spe*I fragment containing *rpoN*::Gen^r and ligating it into the *Eco*RI site of pBR322 to create pRPONgent. pRPONgent (containing *rpoN*::Gen^r) was conjugated into PA14 via a triparental mating by using pRK2013 (14), and

TABLE 2. Nitrogen utilization of *P. aeruginosa* strains

Nitrogen source	Nitrogen utilization of strain:				
	PA14			PAK	
	Wild type (plates ^a)	<i>rpoN</i> mutant		Wild type (plates ^a)	<i>rpoN</i> mutant (plates ^a)
		Plates ^a	Liquid ^b		
Ammonia	2	4	+	2	4
Arginine	2	3	+	2	3
Glutamate	2	—	—	2	—
Glutamine	2	2	+	2	2
Histidine	2	—	—	2	—
Nitrate	2	—	—	2	—

^a Measured as the number of days until the appearance of colonies. — indicates that only tiny colonies grew.

^b Growth (+) or absence of growth (—) in liquid culture. Growth experiments were conducted twice with identical results

rpoN::Gen^r was marker exchanged into the PA14 genome by first selecting for gentamicin resistance and then screening for carbenicillin sensitivity.

Scanning electron and confocal microscopy. To monitor the attachment of PA14 GFP-labeled strains to the epidermal cell walls of LI-0 leaves and to LI-0 trichomes by confocal laser microscopy, 4-mm-diameter leaf disks were immersed in bacterial suspensions at a density of ca. 10⁹ cells/ml under weak vacuum to remove possible air bubbles from the surface of the leaf disks and make the leaf surface more accessible to bacteria. Subsequently, leaf disks were incubated for 24 h in the bacterial suspensions with shaking under normal pressure and room temperature. The leaf disks were examined with a confocal laser spectrophotometer (Leica TCS NT) by excitation at 488 nm and by monitoring the emission intensity at 511 nm. To monitor the development of PA14 strains in LI-0 parenchyma cells by scanning electron microscopy, 4-mm-diameter leaf disks were cut from the central portion of a leaf from either side of the central vein with a cork borer and then immersed in a PA14 or PA14 *rpoN::Gen^r* suspension at an OD₆₀₀ of 0.02 in 10 mM MgSO₄ in the wells of a 24-well microtiter dish, followed by incubation at room temperature. Leaf disks were fixed at 1, 2, 3, and 4 days postinfection (dpi) in 4% paraformaldehyde, passed through an ethanol series (30, 50, 70, 96, and 100%), and freeze fractured in liquid nitrogen. The plant material was dried in a critical-point drying apparatus (Samdri-PVT-3B; Tousimis), mounted on stubs, coated with a 20- to 25-μm layer of gold-palladium in a Hummer II Sputter Coater (Technics), and studied by using an AMRAY 1000 scanning electron microscope.

RESULTS AND DISCUSSION

PA14 *rpoN::Gen^r* exhibits a variety of characteristic RpoN⁻ phenotypes. As described in Materials and Methods, an interspecies hybridization method was used to identify and clone the *P. aeruginosa* PA14 *rpoN* gene. The PA14 *rpoN* gene was partially sequenced and over this region showed 95% identity to both *P. aeruginosa* PAO1 and PAK *rpoN* sequences (data not shown). Also, as described in Materials and Methods, a PA14 *rpoN* mutant (*rpoN::Gen^r*) was constructed by inserting a DNA cassette conferring gentamicin resistance into the *Clal* site of the PA14 *rpoN* gene and transferring the disrupted gene into the PA14 genome by homologous recombination. The *Gen^r* cassette is inserted within the N-terminal 33% of *rpoN*, before the highly conserved carboxy-terminal region (31, 41).

Unlike wild-type PA14, *rpoN::Gen^r* exhibited a variety of characteristic RpoN⁻ phenotypes. Similar to a previously described *P. aeruginosa* PAK *rpoN* mutant (strain PAK N1) (58), PA14 *rpoN::Gen^r* was nonmotile and did not grow well on glutamate, histidine, or nitrate as the sole nitrogen source (Table 2). Electron microscopic examination showed that PA14 *rpoN::Gen^r* cells were nonflagellated (data not shown). In contrast to a previous report that *P. aeruginosa* PAK *rpoN*

(strain N1) is a glutamine auxotroph (58), we found that both PAK *rpoN* (strain N1) and PA14 *rpoN::Gen^r* grew slowly in the absence of glutamine. Restricted nitrogen utilization, rather than glutamine auxotrophy, is also observed with *rpoN* mutants of *P. syringae* and *P. putida* (36).

To confirm that the RpoN⁻ phenotypes of PA14 *rpoN::Gen^r* were due to the *rpoN::Gen^r* insertion, a 4.0-kb DNA fragment containing a wild-type copy of PA14 *rpoN* on plasmid pPAR4SR was introduced into the PA14 *rpoN::Gen^r* strain. This plasmid restored motility and wild-type growth on all of the nitrogen sources listed in Table 2. Although this complementation experiment did not rule out the possibility that the *rpoN::Gen^r* insertion has a polar effect on downstream genes, it appears likely that the RpoN⁻ phenotypes of PA14 *rpoN::Gen^r* are primarily due to the disruption of the *rpoN* gene. This conclusion is based on the conservation of the genomic region surrounding *rpoN* in *P. aeruginosa* strain PAO1 in comparison to the analogous genomic regions in several bacterial species (29, 31, 42, 44, 50, 55). Upstream of the PAO1 *rpoN* gene is an open reading frame (ORF) with homology to the ATP-binding component of an ABC transporter (21). Downstream of *rpoN* in PAO1 are four ORFs (55). Homologous downstream ORFs are found in several species (29, 31, 42, 44, 50, 55). The second ORF is homologous to the enzyme IIA domains of several proteins of the bacterial sugar phosphotransferase system (PTS) (31). The fourth ORF is homologous to the gene for the HPr protein, which is also a component of the PTS (50). ORF1 and -3 encode peptides that have no homology to proteins of known function. These downstream ORFs appear to play a role in modifying RpoN activity, although the effects are relatively minor. Disruption of the second downstream ORF in *P. aeruginosa* strain PAK reduced the ability of PAK to grow without glutamine, but overexpression of ORF1 or ORF2 by the *tac* promoter had no measurable effect (31). Mutation of either the first or second downstream ORFs in *K. pneumoniae* increased expression of a number of *rpoN*-regulated genes (42). In *Caulobacter crescentus* and *Rhizobium etli*, mutations of the downstream ORFs had only a minor effect on one *rpoN*-regulated gene, *flkK* (29), or had no measurable effect (44), respectively.

PA14 *rpoN::Gen^r* synthesizes reduced levels of pyocyanin. *P. aeruginosa* secretes a variety of compounds under nutrient-limiting conditions, including the yellow-green siderophore pyoverdinin and the blue microbial toxin pyocyanin (10, 43). Pyoverdinin may be a virulence factor in burn wound infections (5, 43), and pyocyanin has been implicated in pulmonary artery injury (9, 22). KA Fe³⁺ medium or swarm plates turned noticeably blue when PA14 was grown to stationary phase,

TABLE 3. Lethality of PA14 in a mouse burn model

<i>P. aeruginosa</i> strain	Mortality ratio (no. of animals that died/total no. of animals [%]) at 10 days post burn	
	Low dose ^a	High dose ^b
PA14	7/7 (100)	8/8 (100)
PA14 <i>rpoN::Gen^r</i>	1/7 (14)	5/8 (62)

^a Six-week-old male AKR/J mice (Charles River Laboratories) were injected with 5 × 10³ cells as described previously (54).

^b Injected with 5 × 10⁵ cells.




FIG. 1. *P. aeruginosa* adheres to leaf surfaces. Confocal scanning microscopy of PA14 or *rpoN::Gen^f* expressing GFP in association with *Arabidopsis* ecotype LI-O. (A) PA14(pSCM21) adheres to the entire surface of the leaf, including the stomatal (S) openings. Bar, 20 μ m. (B) PA14 *rpoN::Gen^f*(pSCM21) cells primarily attached to the grooves formed at cell junctions; very few of the mutant bacteria were located above the stomata (S). Bar, 20 μ m. (C) PA14(pSCM21) (b) formed large multilayered clusters on trichomes (T). Bar, 10 μ m. (D) PA14 *rpoN-GFP::Gen^f*(pSCM21) did not attach to the trichome (T) surface and concentrated loosely around them. Bar, 10 μ m. Leaf disks were incubated with PA14/pSCM21 (A and C) or PA14 *rpoN::Gen^f*/pSCM21 (B and D) in bacterial suspensions at a density of $\sim 10^9$ CFU/ml under slight vacuum for 1 h, followed by incubation for 24 h in an orbital shaker at room temperature, and then examined by confocal laser microscopy as described in Materials and Methods.

whereas no blue color was observed after the growth of PA14 *rpoN*::Gen^r. Quantitative measurement showed that PA14 *rpoN*::Gen^r produced 56% ± 13% less pyocyanin than did the wild type and that pyocyanin production was restored to wild-type levels in PA14 *rpoN*::Gen^r(pPAR4SR). Pyoverdinin production appeared to be unaffected in PA14 *rpoN*::Gen^r. PA14 colonies grown on KB plates express pyoverdinin, which can be detected by fluorescence when the colonies are exposed to long-wavelength UV light. Visual inspection revealed no noticeable difference between wild-type and PA14 *rpoN*::Gen^r colonies.

PA14 *rpoN*::Gen^r exhibits reduced pathogenicity in mice. As shown in Table 3, PA14 *rpoN*::Gen^r was significantly less pathogenic than wild-type PA14 in a mouse burn model. A high percentage of lethality was observed with PA14 at both a relatively high dose of 5×10^5 cells (100% lethality) and a relatively low dose of 5×10^3 cells (~80% lethality). In contrast, only five of eight and one of seven animals died when inoculated with 5×10^5 and 5×10^3 PA14 *rpoN*::Gen^r, respectively. The use of two different doses allows a better estimation of the mutation's effects on virulence. In the case of *rpoN*, even the high dose was not as lethal as wild type at a 100-fold-lower dose, demonstrating a significant reduction in virulence. These burn model data are consistent with previous reports demonstrating that *P. aeruginosa rpoN* mutants or *P. aeruginosa* mutants which contain lesions in RpoN-regulated genes exhibit reduced colonization and virulence in a number of other model systems. *rpoN* mutants showed a reduced ability to colonize in a chronic murine intestinal mucosal model (48) or in human respiratory epithelial xenografts (11) and exhibited reduced virulence in a murine corneal scratch model (51) or in a murine model of acute pneumonia (12). One limitation of these studies, as well as of our own, however, is the fact that *rpoN* mutants are nonmotile, and it is not known to what extent the nonmotile phenotype of the *rpoN* mutants contributes to the loss of virulence. Nonmotile mutants of *P. aeruginosa* strains M2, PA01, and MT1200 exhibited significant reduction in virulence in a mouse burn model (16). The reduction was at the same level or greater than that seen with the PA14 *rpoN* mutant. Thus, the entire effect on virulence of the *rpoN* mutant in *P. aeruginosa* PA14 in the mouse burn model could be due to the loss of motility. RpoN is also an important virulence factor for virulence of *Vibrio anguillarum* in fish (47) and *Vibrio cholerae* in an infant mouse colonization model (13, 34). In the case of *V. cholerae*, it appears that unknown *rpoN*-regulated genes, in addition to the flagellum genes, are required for colonization.

PA14 *rpoN*::Gen^r exhibits reduced pathogenicity in *Arabidopsis*. We have developed several assays to assess the pathogenicity of *P. aeruginosa* in *Arabidopsis* leaves, comparing attachment to plant leaf surfaces and monitoring growth rate and symptom development. We previously showed (49) that when *Arabidopsis* ecotype LI-0 leaf disks are incubated in a dense bacterial suspension (i.e., $\sim 10^9$ cells/ml) for 24 h, wild-type PA14 cells attached to the entire leaf epidermal cell surface, congregated above most of the stomata, and formed large, multilayered clusters on the trichomes (Fig. 1A and C). In contrast, as shown in Fig. 1B and D, PA14 *rpoN*::Gen^r cells primarily attached to the grooves formed at cell junctions, and




FIG. 2. Growth of PA14 and PA14 *rpoN*::Gen^r in *Arabidopsis* leaves. Detached LI-0 leaves were vacuum infiltrated with bacterial suspensions as described in Materials and Methods. The initial titers were ca. 5×10^4 CFU/cm² leaf area. Bacterial titers in the infiltrated leaves were measured immediately after infiltration and at 1, 2, 3, and 4 days after infiltration as described in Materials and Methods. Each value represents the average of three replicates. Similar results were obtained when the leaves of intact 6-week-old *La-er* plants were infiltrated at a titer of 5×10^4 CFU/cm² of leaf area and incubated as described in Materials and Methods (not shown). Symbols: ●, PA14; ■, PA14 *rpoN*::Gen^r.

very few of the mutant bacteria were located above the stomata or associated with trichomes.

As shown in Fig. 2, PA14 *rpoN*::Gen^r showed an ~10-fold decrease in its ability to proliferate in detached *Arabidopsis* LI-0 leaves compared to PA14 after gentle vacuum infiltration of a relatively low inoculum of bacteria. Similar results were obtained when intact *La-er Arabidopsis* plants were hand infiltrated with a 1-ml syringe without a needle (data not shown). With either mode of infection, PA14 *rpoN*::Gen^r ultimately elicited severe soft-rot symptoms.

We also tested the ability of PA14 *rpoN*::Gen^r to infect LI-0 leaves and elicit disease symptoms without being forced into the leaf by a syringe or vacuum infiltration. Lawns of PA14 or PA14 *rpoN*::Gen^r were grown on LB-rifampin agar. Agar cylinders 3 mm in diameter were cut and placed bacterial side down on detached *Arabidopsis* LI-0 leaves as described in Materials and Methods. Both PA14 and PA14 *rpoN*::Gen^r formed lesions under these assay conditions, but at relatively early stages of infection (4 days postinfection [dpi]) the wild-type strain produced significantly larger lesions than PA14 *rpoN*::Gen^r. However, by 7 dpi, we did not find any significant differences in the symptoms produced by PA14 and PA14 *rpoN*::Gen^r. At 4 dpi, the PA14 lesions were, on average, 6.3 ± 1.45 mm in diameter, whereas the diameter of the PA14 *rpoN*::Gen^r-elicited lesions were 4.4 ± 1 mm. At 7 dpi, the mean diameters of the PA14 and PA14 *rpoN*::Gen^r lesions were 7.3 ± 0.85 and 6.91 ± 1.14 mm, respectively. Interestingly, chlorotic rings around the lesions formed by the less-virulent PA14 *rpoN*::Gen^r mutant were significantly larger than around PA14 lesions (data not shown).

Finally, we showed in a recent publication that one of the hallmarks of PA14 pathogenesis of *Arabidopsis* is perpendicular attachment of PA14 cells to the leaf epidermis and to mesophyll cell walls (49). Moreover, as shown in Fig. 3B and C,




FIG. 3. Scanning electron micrographic visualization of freeze-fractured infected *Arabidopsis* LI-0 leaves depicting the failure of *P. aeruginosa* PA14 *rpoN::Gen^f* cells to attach perpendicularly to and penetrate the cell walls of vessel parenchyma cells. The leaves of intact LI-0 plants were infiltrated under vacuum, the infected plants were incubated for 72 h, and then examined by scanning electron microscopy as described in Materials and Methods. *b*, bacterium, *W*, plant wall; *WC*, wall convolution. (A) PA14 *rpoN::Gen^f* cells parallel to the surface of a smooth cell wall of a parenchyma vessel cell. Bar, 1 μ m. (B and C) PA14 cells attached perpendicularly to the surface of a highly convoluted cell wall of a parenchyma vessel cell. Holes (arrows) in the plant cell walls with diameters similar to that of the bacteria are readily visible. Bars, 1 μ m.

PA14 causes the convolution of and the formation of small holes in mesophyll cell walls. The diameter of these holes is approximately the same as the diameter of the bacteria. In contrast to the wild type, PA14 *rpoN*::Gen^r cells did not attach perpendicularly under the conditions examined to the leaf epidermal surface 24 h postinfection (data not shown) and, as shown in Fig. 3A, *rpoN*::Gen^r cells oriented themselves parallel to mesophyll cell walls and did not cause the formation of the small holes. Thus, although we found that PA14 *rpoN*::Gen^r has diminished ability to attach to plant cell surfaces, it still has the ability to cause soft-rot symptoms. In contrast, we have shown that the *P. syringae rpoN* gene is absolutely required for full pathogenicity in *Arabidopsis*, most likely because it is required for *hrp* gene expression (23, 24). The *hrp* genes encode the components of a type III secretory system related to the *Yersinia* species Yop secretion system (37). *P. aeruginosa* has recently been shown to contain a type III secretion system as well, which is required for the export of exotoxin S (62). Our data indicate that the *P. aeruginosa* type III system is either *rpoN* independent or that a type III secretion system is not absolutely required for *P. aeruginosa* pathogenicity in plants.

PA14 *rpoN*::Gen^r exhibits reduced *C. elegans* killing. When the nematode *C. elegans* is fed a lawn of PA14 grown on solid medium, the nematodes die in two characteristic ways depending on the medium on which PA14 is grown (56). On a high-osmolarity but low-phosphate medium (e.g., PGS), the nematodes die rapidly (fast killing) over the course of 24 h. In contrast, nematodes that are fed PA14 grown on a low-osmolarity but high-phosphate medium (NGM) are killed at a slower rate (slow killing), dying after 2 to 3 days. Previous studies have shown that the killing of *C. elegans* by PA14 under these two different conditions is mediated by distinct molecular mechanisms (40, 56, 57). As shown in Fig. 4, there was significantly less killing of *C. elegans* by PA14 *rpoN*::Gen^r compared to killing by wild-type PA14 under both the fast- and slow-killing conditions. The killing of PA14 *rpoN*::Gen^r in the fast-killing assay was restored to wild-type levels by pPAR4SR, which carries wild-type *rpoN*, indicating that the loss of pathogenicity phenotype of PA14 *rpoN*::Gen^r was due to the disruption of the *rpoN* gene (data not shown). The decrease in fast killing by PA14 *rpoN*::Gen^r is consistent with the results described above showing that PA14 *rpoN*::Gen^r synthesizes reduced levels of pyocyanin. Previous results from our laboratory showed that pyocyanin is an important toxin mediating the fast-killing process (40).

PA14 *rpoN*::Gen^r does not exhibit reduced pathogenicity in *G. mellonella*. A single PA14 cell is sufficient to kill a greater wax moth caterpillar when PA14 is injected into the hemolymph (30). We found that PA14 and PA14 *rpoN*::Gen^r were indistinguishable in their ability to kill fifth-instar wax moth larvae (data not shown). The LD₅₀ in both cases was approximately one bacterial cell. This is interesting in light of the decreased virulence of PA14 *rpoN*::Gen^r in the mouse burn model. Microbial defense in insects shows interesting parallels to innate immunity in mammals and plants (8, 25, 27). Members of the cecropin class of antibacterial peptides have been found in both mammals and insects, and cecropin A in *Drosophila melanogaster* is regulated by a cascade similar to the one involving NF- κ B activation in the mammalian inflammatory response (38). The reduction in virulence of PA14




FIG. 4. PA14 *rpoN*::Gen^r killing of *C. elegans* under fast- and slow-killing conditions. (A) Fast killing. L4 stage worms were seeded onto bacterial lawns on PGS medium, and mortality was recorded as described earlier (40). Each value is the mean for three replicates. Symbols: ■, PA14; ◆, PA14 *rpoN*::Gen^r. (B) Slow killing. L4 stage worms were seeded onto bacterial lawns on NGM media, and mortality was recorded as described previously (56). Each value is the mean for six replicates. Symbols: ●, PA14; ◆, *rpoN*::Gen^r.

rpoN::Gen^r in mice but not in wax moths suggests that *rpoN* does not play a significant role in the regulation of virulence factors that affect components of the host innate immune system conserved between insects and mammals.

Conclusions. Previous work with PA14 in our laboratory has shown that PA14 utilizes a common set of virulence factors in evolutionarily disparate hosts (30, 40, 53, 57). Given this and the highly pleiotropic nature of *rpoN* mutants, we expected that disruption of *rpoN* would result in a significant impairment in plant, nematode, insect, and mouse pathogenicity. Surprisingly, a major reduction in pathogenicity of the *P. aeruginosa* PA14 *rpoN* mutant was observed only in *C. elegans* killing and in the elicitation of sepsis in the mouse burn model. Although decreased lesion size and a 10-fold reduction in in planta growth was observed in *Arabidopsis* at early stages of an infection (3 to 4 dpi), at later stages (7 dpi) the *P. aeruginosa rpoN* mutant elicited disease symptoms that were indistinguishable from those caused by the wild type. This contrasts with the absolute requirement for RpoN function in *P. syringae* for pathogenicity (23, 24). No effect of the *rpoN* mutation was observed in *G. mellonella* killing. Thus, although it has been previously reported that *P. aeruginosa rpoN* mutants exhibit

impaired virulence in mice, the results reported here provide a more complete perspective concerning the role of RpoN in the diverse pathogenic interactions that *P. aeruginosa* has with several evolutionarily disparate hosts. The major conclusion is that, in contrast to our expectations, *rpoN* does not appear to regulate any genes that are likely to encode virulence factors universally required for pathogenicity irrespective of the host.

ACKNOWLEDGMENTS

This work was supported by NIH grant GM48707 to F.M.A. and by grants from Hoechst AG and Aventis SA to Massachusetts General Hospital.

REFERENCES

- Albright, L. M., E. Huala, and F. M. Ausubel. 1989. Prokaryotic signal transduction mediated by sensor and regulator protein pairs. *Annu. Rev. Genet.* **23**:311–336.
- Alfano, J. R., and A. Collmer. 1997. The type III (Hrp) secretion pathway of plant pathogenic bacteria: trafficking harpins, Avr proteins, and death. *J. Bacteriol.* **179**:5655–5662.
- Arakawa, Y., R. Wacharotayankun, T. Nagatsuka, H. Ito, N. Kato, and M. Ohta. 1995. Genomic organization of the *Klebsiella pneumoniae* *cps* region responsible for serotype K2 capsular polysaccharide synthesis in the virulent strain Chedid. *J. Bacteriol.* **177**:1788–1796.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 2001. *Current protocols in molecular biology*. John Wiley & Sons, New York, N.Y.
- Barton, H. A., Z. Johnson, C. D. Cox, A. I. Vasil, and M. L. Vasil. 1996. Ferric uptake regulator mutants of *Pseudomonas aeruginosa* with distinct alterations in the iron-dependent repression of exotoxin A and siderophores in aerobic and microaerobic environments. *Mol. Microbiol.* **21**:1001–1017.
- Black, L. K., and R. J. Maier. 1995. IHF- and RpoN-dependent regulation of hydrogenase expression in *Bradyrhizobium japonicum*. *Mol. Microbiol.* **16**:405–413.
- Bloemberg, G. V., G. A. O'Toole, B. J. Lugtenberg, and R. Kolter. 1997. Green fluorescent protein as a marker for *Pseudomonas* spp. *Appl. Environ. Microbiol.* **63**:4543–4551.
- Boman, H. G. 1995. Peptide antibiotics and their role in innate immunity. *Annu. Rev. Immunol.* **13**:61–92.
- Britigan, B. E., G. T. Rasmussen, and C. D. Cox. 1997. Augmentation of oxidant injury to human pulmonary epithelial cells by the *Pseudomonas aeruginosa* siderophore pyochelin. *Infect. Immun.* **65**:1071–1076.
- Byng, G. S., D. C. Eustice, and R. A. Jensen. 1979. Biosynthesis of phenazine pigments in mutant and wild-type cultures of *Pseudomonas aeruginosa*. *J. Bacteriol.* **138**:846–852.
- Cohn, L. A., A. Weber, T. Phillips, S. Lory, M. Kaplan, and A. Smith. 2001. *Pseudomonas aeruginosa* infection of respiratory epithelium in a cystic fibrosis xenograft model. *J. Infect. Dis.* **183**:919–927.
- Comolli, J. C., A. R. Hauser, L. Waite, C. B. Whitechurch, J. S. Mattick, and J. N. Engel. 1999. *Pseudomonas aeruginosa* gene products PilT and PilU are required for cytotoxicity in vitro and virulence in a mouse model of acute pneumonia. *Infect. Immun.* **67**:3625–3630.
- Correa, N. E., C. M. Lauriano, R. McGee, and K. E. Klose. 2000. Phosphorylation of the flagellar regulatory protein FlrC is necessary for *Vibrio cholerae* motility and enhanced colonization. *Mol. Microbiol.* **35**:743–755.
- Ditta, G., S. Stanfield, D. Corbin, and D. R. Helinski. 1980. Broad host-range DNA cloning system for gram-negative bacteria: construction of a gene bank of *Rhizobium meliloti*. *Proc. Natl. Acad. Sci. USA* **77**:7347–7351.
- Dong, X., M. Mindrinos, K. R. Davis, and F. M. Ausubel. 1991. Induction of *Arabidopsis* defense genes by virulent and avirulent *Pseudomonas syringae* strains and by a cloned avirulence gene. *Plant Cell* **3**:61–72.
- Drake, D., and T. C. Montie. 1988. Flagella motility and invasive virulence of *Pseudomonas aeruginosa*. *J. Gen. Microbiol.* **124**:43–52.
- Essar, D. W., L. Eberly, A. Hadero, and I. P. Crawford. 1990. Identification and characterization of genes for a second anthranilate synthase in *Pseudomonas aeruginosa*: interchangeability of the two anthranilate synthases and evolutionary implications. *J. Bacteriol.* **172**:884–900.
- Galan, J., and A. Collmer. 1999. Type III secretion machines: bacterial devices for protein delivery into host cells. *Science* **284**:1322–1328.
- Goldberg, J. B., and T. Dahnke. 1992. *Pseudomonas aeruginosa* AlgB, which modulates the expression of alginate, is a member of the NtrC subclass of prokaryotic regulators. *Mol. Microbiol.* **6**:59–66.
- Hanahan, D., and M. Meselson. 1983. Plasmid screening at high colony density. *Methods Enzymol.* **100**:333–342.
- Hartig, E., and W. G. Zumft. 1998. The requirement of RpoN (sigma factor σ^{54}) in denitrification by *Pseudomonas stutzeri* is indirect and restricted to the reduction of nitrite and nitric oxide. *Appl. Environ. Microbiol.* **64**:3092–3095.
- Hassan, H. M., and I. Fridovich. 1980. Mechanism of the antibiotic action of pyocyanine. *J. Bacteriol.* **141**:1556–1563.
- Hendrickson, E. L., P. Guevera, and F. M. Ausubel. 2000. The alternative sigma factor RpoN is required for *hrp* activity in *Pseudomonas syringae* pv. *maculicola* and acts at the level of *hrpL* transcription. *J. Bacteriol.* **182**:3508–3516.
- Hendrickson, E. L., P. Guevera, A. Penaloza-Vazquez, J. Shao, C. Bender, and F. M. Ausubel. 2000. Virulence of the phytopathogen *Pseudomonas syringae* pv. *maculicola* is *rpoN* dependent. *J. Bacteriol.* **182**:3498–3507.
- Hoffmann, J. A. 1995. Innate immunity of insects. *Curr. Opin. Immunol.* **7**:4–10.
- Huang, H.-C., R.-H. Lin, C.-J. Chang, A. Collmer, and W.-L. Deng. 1995. The complete *hrp* gene cluster of *Pseudomonas syringae* pv. *syringae* 61 includes two blocks of genes required for hairpin Pss secretion that are arranged colinearly with *Yersinia ysc* homologs. *Mol. Plant-Microbe Interact.* **8**:733–746.
- Hultmark, D. 1993. Immune reactions in *Drosophila* and other insects: a model for innate immunity. *Trends Genet.* **9**:178–183.
- Ishimoto, K. S., and S. Lory. 1989. Formation of pilin in *Pseudomonas aeruginosa* requires the alternative sigma factor (RpoN) of RNA polymerase. *Proc. Natl. Acad. Sci. USA* **86**:1954–1957.
- Janakiraman, R. S., and Y. V. Brun. 1997. Transcriptional and mutational analyses of the *rpoN* operon in *Caulobacter crescentus*. *J. Bacteriol.* **179**:5138–5147.
- Jander, G., L. G. Rahme, and F. M. Ausubel. 2000. Positive correlation between virulence of *Pseudomonas aeruginosa* mutants in mice and insects. *J. Bacteriol.* **182**:3843–3845.
- Jin, S., K. Ishimoto, and S. Lory. 1994. Nucleotide sequence of the *rpoN* gene and characterization of two downstream open reading frames in *Pseudomonas aeruginosa*. *J. Bacteriol.* **176**:1316–1322.
- Kessler, B., S. Marqués, T. Köhler, J. L. Ramos, K. N. Timmis, and V. de Lorenzo. 1994. Cross talk between catabolic pathways in *Pseudomonas putida*: XylS-dependent and-independent activation of the *TOL meta* operon requires the same *cis*-acting sequences within the *Pm* promoter. *J. Bacteriol.* **176**:5578–5582.
- King, E. O., M. K. Ward, and D. E. Raney. 1954. Two simple media for the demonstration of phycocyanin and fluorescein. *J. Lab. Clin. Med.* **44**:301–307.
- Klose, K. E., and J. J. Mekalanos. 1998. Distinct roles of an alternative sigma factor during both free-swimming and colonizing phases of the *Vibrio cholerae* pathogenic cycle. *Mol. Microbiol.* **28**:501–520.
- Kohler, T., S. Harayama, J. L. Ramos, and K. N. Timmis. 1989. Involvement of *Pseudomonas putida* RpoN σ factor in regulation of various metabolic functions. *J. Bacteriol.* **171**:4326–4333.
- Kustu, S., E. Santero, J. Keener, D. Popham, and D. Weiss. 1989. Expression of sigma 54 (*ntrA*)-dependent genes is probably united by a common mechanism. *Microbiol. Rev.* **53**:367–376.
- Lee, C. A. 1997. Type III secretion systems: machines to deliver bacterial proteins into eukaryotic cells? *Trends Microbiol.* **5**:148–156.
- Lemaitre, B., E. Nicolas, L. Michaut, J.-M. Reichhart, and J. A. Hoffmann. 1996. The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* **86**:973–983.
- Macaluso, A., E. A. Best, and R. A. Bender. 1990. Role of the *nac* gene product in the nitrogen regulation of some NTR-regulated operons of *Klebsiella aerogenes*. *J. Bacteriol.* **172**:7249–7255.
- Mahajan-Miklos, S., M. W. Tan, L. G. Rahme, and F. M. Ausubel. 1999. Molecular mechanisms of bacterial virulence elucidated using a *Pseudomonas aeruginosa*-*Caenorhabditis elegans* pathogenesis model. *Cell* **96**:47–56.
- Merrick, M. J. 1993. In a class of its own—the RNA polymerase sigma factor σ^{54} (σ^N). *Mol. Microbiol.* **10**:903–909.
- Merrick, M. J., and J. R. Coppard. 1989. Mutations in genes downstream of the *rpoN* gene (encoding σ^{54}) of *Klebsiella pneumoniae* affect repression from σ^{54} -dependent promoters. *Mol. Microbiol.* **3**:1765–1775.
- Meyer, J., A. Neely, A. Stintzi, C. Georges, and I. A. Holder. 1996. Pyoverdinin is essential for virulence of *Pseudomonas aeruginosa*. *Infect. Immun.* **64**:518–523.
- Michaels, J., T. Van Soom, I. D'hooghe, B. Dombrecht, T. Benhassine, P. de Wilde, and J. Vanderleyden. 1998. The *Rhizobium etli* *tpoN* locus: DNA sequence analysis and phenotypic characterization of *rpoN*, *ptsN*, and *ptsA* mutants. *J. Bacteriol.* **180**:1729–1740.
- Mudgett, M., and B. Staskawicz. 1999. Characterization of the *Pseudomonas syringae* pv. *tomato AvrRpt2* protein: demonstration of secretion and processing during bacterial pathogenesis. *Mol. Microbiol.* **32**:927–941.
- Mudgett, M., and B. Staskawicz. 1998. Protein signaling via type III secretion pathways in phytopathogenic bacteria. *Curr. Opin. Microbiol.* **1**:109–114.
- O'Toole, R., D. L. Milton, P. Horstedt, and H. Wolf-Watz. 1997. RpoN of the fish pathogen *Vibrio (Listonella) anguillarum* is essential for flagellum production and virulence by the water-borne but not intraperitoneal route of inoculation. *Microbiology* **143**:3849–3859.
- Pier, G. B., G. Meluleni, and E. Neuger. 1992. A murine model of chronic mucosal colonization by *Pseudomonas aeruginosa*. *Infect. Immun.* **60**:4768–4776.
- Plotnikova, J. M., L. G. Rahme, and F. M. Ausubel. 2000. Pathogenesis of the human opportunistic pathogen *Pseudomonas aeruginosa* PA14 in *Arabi-*

- dopsis*. *Plant. Physiol.* **124**:1766–1774.
50. Powell, B. S., D. L. Court, T. Inada, Y. Nakamura, V. Michotey, X. Cui, A. Reizer, M. H. Saier, and J. Reizer. 1995. Novel proteins of the phosphotransferase system encoded within the *rpoN* operon of *Escherichia coli*. Enzyme IIANtr affects growth on organic nitrogen and the conditional lethality of an *erats* mutant. *J. Biol. Chem.* **270**:4822–4839.
 51. Preston, M. J., S. M. Fleiszig, T. S. Zaidi, J. B. Goldberg, V. D. Shortridge, M. L. Vasil, and G. B. Pier. 1995. Rapid and sensitive method for evaluating *Pseudomonas aeruginosa* virulence factors during corneal infections in mice. *Infect. Immun.* **63**:3497–3501.
 52. Rahme, L. G., E. J. Stevens, S. F. Wolfort, J. Shao, R. G. Tompkins, and F. M. Ausubel. 1995. Common virulence factors for bacterial pathogenicity in plants and animals. *Science* **268**:1899–1902.
 53. Rahme, L. G., M. W. Tan, L. Le, S. M. Wong, R. G. Tompkins, S. B. Calderwood, and F. M. Ausubel. 1997. Use of model plant hosts to identify *Pseudomonas aeruginosa* virulence factors. *Proc. Natl. Acad. Sci. USA* **94**:13245–13250.
 54. Stevens, E. J., C. M. Ryan, J. S. Friedberg, R. L. Barnhill, M. L. Yarmush, and R. G. Tompkins. 1994. A quantitative model of *Pseudomonas aeruginosa* infection in injury. *J. Burn Care Rehabil.* **15**:232–235.
 55. Stover, C. K., X. Q. Pham, A. L. Erwin, S. D. Mizoguchi, P. Warriner, M. J. Hickey, F. S. Brinkman, W. O. Hufnagle, D. J. Kowalik, M. Lagrou, R. L. Garber, L. Goltry, E. Tolentino, S. Westbrook-Wadman, Y. Yuan, L. L. Brody, S. N. Coulter, K. R. Folger, A. Kas, K. Larbig, R. Lim, K. Smith, D. Spencer, G. K. Wong, Z. Wu, and I. T. Paulsen. 2000. Complete genome sequence of *Pseudomonas aeruginosa* PA01, an opportunistic pathogen. *Nature* **406**:959–964.
 56. Tan, M. W., S. Mahajan-Miklos, and F. M. Ausubel. 1999. Killing of *Caenorhabditis elegans* by *Pseudomonas aeruginosa* used to model mammalian bacterial pathogenesis. *Proc. Natl. Acad. Sci. USA* **96**:715–720.
 57. Tan, M. W., L. G. Rahme, J. A. Sternberg, R. G. Tompkins, and F. M. Ausubel. 1999. *Pseudomonas aeruginosa* killing of *Caenorhabditis elegans* used to identify *P. aeruginosa* virulence factors. *Proc. Natl. Acad. Sci. USA* **96**:2408–2413.
 58. Totten, P. A., J. C. Lara, and S. Lory. 1990. The *rpoN* gene product of *Pseudomonas aeruginosa* is required for expression of diverse genes, including the flagellin gene. *J. Bacteriol.* **172**:389–396.
 59. van Dijk, K., D. E. Fouts, A. H. Rehm, A. R. Hill, A. Collmer, and J. R. Alfano. 1999. The Avr (effector) proteins HrmA (HopPsyA) and AvrPto are secreted in culture from *Pseudomonas syringae* pathovars via the Hrp (type III) protein secretion system in a temperature- and pH-sensitive manner. *J. Bacteriol.* **181**:4790–4797.
 60. Woods, D. E., D. C. Straus, W. G. Johanson, Jr., V. K. Berry, and J. A. Bass. 1980. Role of pili in adherence of *Pseudomonas aeruginosa* to mammalian buccal epithelial cells. *Infect. Immun.* **29**:1146–1151.
 61. Wu, Z.-L., T. C. Charles, H. Wang, and E. W. Nester. 1992. The *ntrA* gene of *Agrobacterium tumefaciens*: identification, cloning, and phenotype of a site-directed mutant. *J. Bacteriol.* **174**:2720–2723.
 62. Yahr, T. L., J. Goranson, and D. W. Frank. 1996. Exoenzyme S of *Pseudomonas aeruginosa* is secreted by a type III pathway. *Mol. Microbiol.* **22**:991–1003.