

Enterococcus faecalis 3-Hydroxy-3-Methylglutaryl Coenzyme A Synthase, an Enzyme of Isopentenyl Diphosphate Biosynthesis†

Autumn Sutherland,¹ Matija Hedl,¹ Barbara Sanchez-Neri,¹ John W. Burgner II,²
Cynthia V. Stauffacher,² and Victor W. Rodwell^{1*}

Departments of Biochemistry¹ and Biological Sciences,² Purdue University, West Lafayette, Indiana 47907

Received 28 February 2002/Accepted 29 April 2002

Biosynthesis of the isoprenoid precursor isopentenyl diphosphate (IPP) proceeds via two distinct pathways. Sequence comparisons and microbiological data suggest that multidrug-resistant strains of gram-positive cocci employ exclusively the mevalonate pathway for IPP biosynthesis. Bacterial mevalonate pathway enzymes therefore offer potential targets for development of active site-directed inhibitors for use as antibiotics. We used the PCR and *Enterococcus faecalis* genomic DNA to isolate the *mvaS* gene that encodes 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) synthase, the second enzyme of the mevalonate pathway. *mvaS* was expressed in *Escherichia coli* from a pET28 vector with an attached N-terminal histidine tag. The expressed enzyme was purified by affinity chromatography on Ni²⁺-agarose to apparent homogeneity and a specific activity of 10 $\mu\text{mol}/\text{min}/\text{mg}$. Analytical ultracentrifugation showed that the enzyme is a dimer (mass, 83.9 kDa; $s_{20,w}$, 5.3). Optimal activity occurred in 2.0 mM MgCl₂ at 37°C. The ΔH_a was 6,000 cal. The pH activity profile, optimum activity at pH 9.8, yielded a pK_a of 8.8 for a dissociating group, presumably Glu78. The stoichiometry per monomer of acetyl-CoA binding was 1.2 ± 0.2 and that of covalent acetylation was 0.60 ± 0.02 . The K_m for the hydrolysis of acetyl-CoA was 10 μM . Coupled conversion of acetyl-CoA to mevalonate was demonstrated by using HMG-CoA synthase and acetoacetyl-CoA thiolase/HMG-CoA reductase from *E. faecalis*.

Two distinct pathways, the mevalonate pathway (Fig. 1) and the nonmevalonate pathway (20), give rise to isopentenyl diphosphate (IPP), the monomer unit for isoprenoid biosynthesis. Mammals and the archaea studied use exclusively the mevalonate pathway (3), whereas plants employ both pathways (1). Inspection of the sequences of microbial genomes has revealed that, whereas many gram-negative eubacteria contain genes that encode enzymes of the nonmevalonate pathway, the gram-positive cocci, *Borrelia burgdorferi* (27), and *Streptomyces* species (10, 24) possess genes that appear to encode the enzymes of the mevalonate pathway.

The mevalonate pathway is essential for the survival of representative gram-positive cocci. Wilding et al. (28) established that the survival of knockout mutants of the genes that encode 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase (EC 1.1.1.34) of *Streptococcus pneumoniae* and *Staphylococcus aureus* depended on externally supplied mevalonate. While the mevalonate pathway is also essential for human subjects, there are two classes of HMG-CoA reductase that differ both with respect to their structure and their inhibition characteristics (4). In addition, there may be significant differences between the bacterial and human forms of other enzymes of the pathway. The enzymes of the mevalonate pathway may therefore represent potential targets for the development of new antibiotics.

An unusual feature of the mevalonate pathway of *Enterococcus faecalis* is that a dual-function polypeptide, acetoacetyl-

CoA thiolase/HMG-CoA reductase (11, 27), catalyzes the first and third reactions of the pathway (Fig. 1). The second reaction of the mevalonate pathway, the functionally irreversible condensation of acetoacetyl-CoA and acetyl-CoA to form HMG-CoA, is catalyzed by HMG-CoA synthase (EC 4.1.3.5). Our interest in *E. faecalis* HMG-CoA synthase, which appears to link the activities of the dual-function *E. faecalis* enzyme, reflects both the potential of the mevalonate pathway enzymes of enterococci as targets for antibiotics and the fact that no bacterial HMG-CoA synthase has previously been characterized.

MATERIALS AND METHODS

Reagents. Purchased reagents and kits included T4 DNA ligase, Vent DNA polymerase, and restriction enzymes (New England Biolabs); deoxynucleoside triphosphates (Invitrogen); 40% acrylamide (Amresco); Bradford reagent and MicroSpin BioP-30 columns (Bio-Rad); IPTG (isopropyl- β -D-thiogalactopyranoside; Gibco-BRL); glass fiber filters (Millipore); [¹⁴C]acetyl-CoA (Amersham-Pharmacia); and nickel-nitrilotriacetic acid (Ni-NTA) agarose, a QIAprep spin miniprep kit, and a gel extraction kit (Qiagen). *E. faecalis* chromosomal DNA was a gift of Michael Gwynn at GlaxoSmithKline. Synthetic oligonucleotides were prepared by Integrated DNA Technologies, Coralville, Iowa. *E. faecalis* acetoacetyl-CoA thiolase/HMG-CoA reductase was prepared as previously described (11). Unless otherwise specified, all other reagents were from Sigma.

Construction of the expression plasmid. The *mvaS* gene was PCR amplified from *E. faecalis* chromosomal DNA with primers that introduced an *Nde*I site (5'-CGTAAAGGAGTTAAACATATGACAATTGGG-3') and a *Bam*HI site (5'-GAATCGGGGGATCCAAATACTTAGTTTCG-3') at the ends of the amplified fragment. The 1,176-bp PCR fragment was cloned into *Nde*I/*Bam*HI-cut pET28 in frame with the N-terminal histidine tag and thrombin cleavage site to give the expression plasmid MvaS-pET28-6H. The insert was sequenced by the Iowa State University DNA Sequencing Facility, Ames, to verify that no mutations had been introduced during PCR amplification.

Expression and purification of the gene product. *Escherichia coli* BL21(DE3) cells transformed with MvaS-pET28-6H were grown at 37°C, with shaking, on Luria-Bertani broth (22) containing 10 μg of kanamycin per ml. Cells harvested by centrifugation were washed with 0.9% saline, suspended in Buffer A (10 mM

* Corresponding author. Mailing address: Department of Biochemistry, Purdue University, West Lafayette, IN 47907-1153. Phone: (765) 494-1608. Fax: (765) 494-7897. E-mail: vrodwell@purdue.edu.

† Journal paper 16732 from the Purdue Agricultural Experiment Station.




FIG. 1. Intermediates and enzymes of the mevalonate pathway for IPP biosynthesis.

imidazole, 300 mM NaCl, 1 mM phenylmethylsulfonyl fluoride, 1 mM dithiothreitol, 20 mM HEPES; pH 8.0), and lysed in a French press. The supernatant liquid (cytosol) obtained after centrifugation of the cell lysate in a Beckman L8-70 ultracentrifuge (30,000 rpm, 60 min, 4°C) was applied to an Ni-NTA column. The column was washed with Buffer A and then eluted successively with 50 and 100 mM imidazole in Buffer A. Fractions with high activity were combined and stored in liquid nitrogen. Protein concentration was determined by the Bradford method (5).

Analytical ultracentrifugation. *E. faecalis* HMG-CoA synthase, i.e., 430 μ l of a 1-mg/ml solution in 100 mM NaCl–10 mM Tris (pH 8.0), was loaded in a Beckman analytical ultracentrifuge cell with sapphire windows and an aluminum-filled epoxy centerpiece. The cell was placed in a Beckman Model XL-I centrifuge and allowed to come to thermal equilibrium at 20°C for 1 h. The sample was then spun at 50,000 rpm for 4 h. Rayleigh interference scans were taken at 1-min intervals.

HMG-CoA synthase activity. Determination of HMG-CoA synthase activity employed a Hewlett-Packard model 8452 diode array spectrophotometer with the cell compartment maintained at 37°C to monitor the change in absorbance at 302 nm that accompanies the acetyl-CoA-dependent disappearance of the enolate form of acetoacetyl-CoA (21). The standard assay medium included 500 μ M acetyl-CoA, 20 μ M acetoacetyl-CoA, 5.0 mM $MgCl_2$, and 50 mM Tris (pH 9.75). Assays employed a final volume of 220 μ l. One enzyme unit (eu) represents the disappearance in 1 min of 1 μ mol of acetoacetyl-CoA. The reported data represent mean values for at least triplicate determinations.

Acetyl-CoA hydrolase activity. Acetyl-CoA hydrolase activity was determined by measuring the release of coenzyme A with DTNB [5,5'-dithiobis(2-nitroben-




FIG. 2. SDS-PAGE of fractions obtained during purification of *E. faecalis* HMG-CoA synthase. Lane 1, prestained standards of the indicated molecular weight; lane 2, cell extract; lane 3, cytosol; lane 4, Ni-NTA fraction.

zoic acid)]. The assay employed a Hewlett-Packard model 8452 diode array spectrophotometer with the cell compartment maintained at 37°C to monitor the change in absorbance at 412 nm that accompanies the reaction of CoA with DTNB. The assay employed modifications of the procedures of Weitzman (26) and Mizioroko et al. (19). Standard assay conditions consisted of 1 mM DTNB, 20 μ M acetyl-CoA, and 50 mM Tris (pH 9.8) in a final volume of 220 μ l. One eu represents the release in 1 min of 1 μ mol of CoA. The reported data represent mean values for at least triplicate determinations.

RESULTS

Expression and purification of the gene product of *E. faecalis mvaS*. The *mvaS* gene of *E. faecalis*, which encodes HMG-CoA synthase, was cloned into pET28 in frame with an N-terminal histidine tag. A soluble protein was expressed in *E. coli* to high levels at 37°C and was readily purified 20-fold on a nickel-chelating column. The yield was 4 mg per liter of >98% homogeneous protein, as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (Fig. 2).

Sequence alignments. Animal HMG-CoA synthases are cytosolic (cholesterogenic) or mitochondrial (ketogenic). CLUSTALW (www.expasy.ch) was used to align the amino acid sequences of HMG-CoA synthases from known animal and putative bacterial forms of the enzyme. The cysteine (17), histidine (16), and glutamate (8) implicated as catalytic residues in the avian enzyme are conserved across all forms of the enzyme (Table 1). The sequence identities were 75 and 80% for the mitochondrial and cytosolic isozymes, respectively, and ca. 50% across the two classes. The bacterial HMG-CoA synthases share >20% identity among themselves but only 10% identity with the animal enzymes. The relatively low identity between the bacterial and animal forms might indicate differences in their active sites that might ultimately be exploited for antibiotic design.

Multimeric state. The multimeric state of *E. faecalis* HMG-CoA synthase was investigated by analytical ultracentrifugation and gel filtration chromatography. Approximately 90% of the

TABLE 1. Selected sequences and conserved residues of animal and bacterial HMG-CoA synthases^a

Sequence set and source ^b	Sequence (residue position)
Set 1	(95)
Animal	L E V G T E T I I D K S K x V K
Bacterial	V I V A T E S x I D x x K A x x
Set 2	(129)
Animal	D T T N A C Y G G T
Bacterial	E x x x A C Y x A T
Set 3	(264)
Animal	M I F H x P x x K
Bacterial	x x F H x P x C K

^a Capital letters indicate conserved residues. Residues in boldface are conserved in both animal and bacterial HMG-CoA synthases. An "x" indicates a nonconserved residue. Glu95 (8), Cys129 (17), and His264 (16) have been identified as catalytic residues in the avian enzyme. For *E. faecalis* HMG-CoA synthase these residues are Glu79, Cys111, and His233.

^b Animal cytosolic HMG-CoA synthases aligned are from rats, hamsters, humans, and chickens. Animal mitochondrial HMG-CoA synthases aligned are from mouse, rat, human, and pig. Aligned bacterial HMG-CoA synthase sequences are from *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Enterococcus faecium*, *Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus aureus*, and a *Streptomyces* sp.

total protein had a sedimentation constant of ca. 5.3 S (Fig. 3), which is consistent with a molecular mass of 83.9 kDa. The dimer mass calculated from the amino acid sequence is 84.3 kDa. Material migrating at 3.6 S probably represents the monomer, although the calculated mass of 45 kDa is somewhat larger than the predicted size of 42 kDa. Gel filtration on a Superdex 200 column confirmed these results (data not shown).




FIG. 3. Sedimentation velocity ultracentrifugation of *E. faecalis* HMG-CoA synthase. (A) Sedimentation boundaries measured by using Rayleigh interference optics plotted against radial position. The data shown are for 25-min intervals and represent one-fifth of the data used in the analysis. The jagged lines represent the observed fringes, and the smooth lines represent the calculated best-fit distribution calculated by using Sedfit 8.3 and the Lamm equation model (23). (B) Best-fit $c(s)$ sedimentation coefficient distribution, allowing for systematic time-invariant noise. The uncorrected values for the sedimentation coefficients for the minor and major peaks are 3.6 and 5.3 S, respectively.

Effect of temperature, magnesium ion concentration, and hydrogen ion concentration. Enzymes of the mevalonate pathway are present in mesophiles, thermophiles, and halophiles. We therefore investigated the effect of temperature, magnesium ion concentration, and hydrogen ion concentration on *E. faecalis* HMG-CoA synthase activity. Optimum activity occurred at ca. 37°C (Fig. 4, top). A ΔH_a of 6,000 cal was calculated from the Arrhenius plot (Fig. 4, top [inset]). Activity was optimal at 2 mM $MgCl_2$ (Fig. 4, middle). Neither 5 to 10 mM EDTA nor 2 to 10 mM KCl affected activity. Optimal activity occurred at pH 9.8 (Fig. 4, bottom). Modeling of the data for pH values from pH 8.5 to 9.25 gave an estimated value for the pK_a of a dissociating group of 8.8.

Kinetic parameters. For synthesis of HMG-CoA, the K_m for acetyl-CoA was 350 μM . Since acetoacetyl-CoA competes with acetyl-CoA for the catalytic cysteine, only a K_m^{app} can be determined (14). At a concentration of 500 μM acetyl-CoA, the K_m^{app} for acetoacetyl-CoA was 10 μM . Table 2 summarizes these kinetic parameters.

Detection of an acetyl-S-enzyme intermediate. Formation of HMG-CoA proceeds in three stages (Fig. 5), the first and third of which can be studied experimentally. Detection of covalently bound acetyl-CoA documents the formation of an acetyl-S-enzyme intermediate in stage 1. Determination of the stoichiometry of acetyl-CoA bound indicates the number of active sites. Covalent acetylation of HMG-CoA synthase was determined by the procedure of Mizioroko et al. (19), and acetyl-CoA binding was determined by the procedure of Vollmer et al. (25). *E. faecalis* HMG-CoA synthase bound 1.2 ± 0.2 mol of acetyl-CoA per mol of active sites, 60% of which were covalently associated with the enzyme (Table 3). All of the active sites appear to be associated with acetyl-CoA, and at least half of the sites contain an acetyl-S-intermediate.

Hydrolytic release of enzyme-bound acetyl-CoA. The hydrolytic release of enzyme-bound HMG-CoA cannot readily be measured directly because the overall reaction is irreversible. Stage 3 can, however, be modeled by the hydrolysis of the acetylated enzyme, albeit at only 1% the rate of the overall reaction (7, 19). A K_m of 10 μM for acetyl-CoA and a V_{max} of 2×10^{-2} $\mu mol/min/mg$ were determined for the hydrolysis of acetyl-CoA. These parameters approximate those previously reported for avian HMG-CoA synthase (Table 3).

Coupled conversion of acetyl-CoA to mevalonate. The first and third reactions of the mevalonate pathway in *E. faecalis*, the synthesis of acetoacetyl-CoA and the formation of mevalonate, are catalyzed by a single bifunctional enzyme (11). Since HMG-CoA synthase catalyzes the second reaction of the pathway, we investigated the coupled conversion of acetyl-CoA to mevalonate. The pH optima for HMG-CoA synthase, acetoacetyl-CoA thiolase, and HMG-CoA reductase range from pH 6.5 to 10.5. We therefore studied the coupled conversion of acetyl-CoA to mevalonate at several pH values. Coupled conversion was indeed observed. Optimal synthesis of mevalonate from acetyl-CoA occurred at pH 8.5 (Fig. 6).

DISCUSSION

Only recently has it been possible to deduce from inspection of the sequences of bacterial genomes that gram-positive bacteria employ the mevalonate pathway to synthesize IPP (27).




FIG. 4. Effect of temperature, MgCl_2 concentration, and hydrogen ion concentration. (Top) Temperature. Assays were conducted at the indicated temperatures under otherwise standard conditions. The inset shows selected data shown as an Arrhenius plot. (Middle) MgCl_2 .

TABLE 2. Kinetic constants for selected HMG-CoA synthases

Source (reference)	K_m for acetyl-CoA (μM)	K_m^{app} for acetoacetyl-CoA (μM)	V_{max} ($\mu\text{mol}/\text{min}/\text{mg}$)
<i>E. faecalis</i>	350	10	10
Avian liver cytosol (7)	270	1.2	4.4
<i>Blattella germanica</i> (6)	1,500	<0.5	66

Of the enzymes of the mevalonate pathway in enterococci, only *S. aureus* HMG-CoA reductase (28) and the acetoacetyl-CoA thiolase/HMG-CoA reductase of *E. faecalis* (11) have been expressed and characterized. We report here the cloning, expression, purification, and characterization of the first bacterial HMG-CoA synthase.

Analytical ultracentrifugation and gel filtration revealed that *E. faecalis* HMG-CoA synthase is a dimer and thus has the same multimeric state as other HMG-CoA synthases (14). Optimal activity occurred at 37°C in the presence of 2.0 mM MgCl_2 (Fig. 4), which stabilizes acetoacetyl-CoA (9). The free energy of activation, ΔH_a , was 6,000 cal. A pK_a of 8.8 was estimated from the effect of hydrogen ion concentration on activity (Fig. 4). This value is similar to the pK_a of 8.6 reported for Glu95 and to its presumably hydrophobic environment (8). Glu95, and by inference its conjugate residue Glu78 of the *E. faecalis* enzyme, acts as a general acid during the condensation stage of the HMG-CoA synthase reaction (Fig. 5).

As for the avian enzyme (18), the stoichiometry of acetyl-CoA binding was 1.2 ± 0.2 per monomer and of covalent acetylation was 0.60 ± 0.02 per monomer (Table 3). The difference in stoichiometry may indicate that the presence of an acetyl-S-enzyme intermediate at one active site may induce conformational changes that preclude simultaneous formation of an acetyl-S-enzyme intermediate at the other site. K_m for the hydrolysis of acetyl-CoA, a reaction that models the third or hydrolytic stage of the overall reaction, was 10 μM . The V_{max} of 10 eu/mg and the specific activity of 6 $\mu\text{mol}/\text{min}/\text{mg}$ of protein for *E. faecalis* HMG-CoA synthase are only twofold greater than those for the avian enzyme (Table 2). However, the 10 μM K_m^{app} for acetoacetyl-CoA, which is almost an order of magnitude higher than that for the avian enzyme, may be indicative of subtle differences at their active sites.

The overall conversion of acetyl-CoA to mevalonate comprises the first three reactions of the mevalonate pathway (Fig. 1). The coupled conversion of acetyl-CoA to mevalonate was demonstrated by using the dual enzyme acetoacetyl-CoA thiolase/HMG-CoA reductase and HMG-CoA synthase of *E. faecalis*. Optimal activity for the coupled reactions occurred at pH 8.5.

The enzymes of the mevalonate pathway of IPP biosynthesis represent potential targets for metabolic intervention. The mevalonate pathway is essential for the survival of *S. aureus* and,

Assays were conducted at the indicated concentrations of MgCl_2 under otherwise standard conditions. (Bottom) Hydrogen ion concentration. Assays were conducted in 50 mM sodium acetate, 50 mM glycine, 50 mM Tris, and 50 mM 2-(*N*-morpholino)ethanesulfonic acid at the indicated pH under otherwise standard conditions.




FIG. 5. Proposed mechanism for catalysis of the HMG-CoA synthase reaction. Active site residues of the avian mitochondrial enzyme include Glu95, Cys129, and His264. During the acetylation step, Cys129 attacks the carbonyl group of acetyl-CoA, forming an acetyl-S-intermediate (17). After the addition and condensation of acetoacetyl-CoA, HMG-CoA is released by hydrolysis. Glu95 acts as the general acid in the condensation step (8), and His264 anchors binding of acetoacetyl-CoA (16).

by inference, for other gram-positive bacteria (28). Although the mevalonate pathway is also essential for human subjects, there are two classes of HMG-CoA reductase that differ both with respect to their structure and their inhibition characteristics (4). The K_i values for statin drug inhibition of the class I HMG-CoA reductases of eukaryotes are nanomolar (2) but are millimolar for the class II HMG-CoA reductases of *S. aureus* (28) and *E. faecalis* (11). Significant differences also characterize the crystal structures of the class I human enzyme

(12) and the class II enzyme of *Pseudomonas mevalonii* (13). It thus may be possible to develop inhibitors, i.e., "class II statins," specific to class II HMG-CoA reductases. Furthermore, the existence of two classes of one enzyme of the mevalonate

TABLE 3. Parameters for the partial reactions that model the first and third stages of the overall reaction catalyzed by HMG-CoA synthase

Source (reference)	Acetyl-CoA bound (mol/mol of monomer) ^a	Acetyl-CoA covalently bound (mol/mol of monomer) ^b	Hydrolysis of acetyl-CoA	
			V_{\max} (eu/mg)	K_m (μM)
<i>E. faecalis</i>	1.2 ± 0.2^c	0.60 ± 0.02^c	0.02	8.5
Avian liver cytosol (7, 8)	1.1	0.63	0.016	11

^a To determine the stoichiometry of acetyl-CoA binding, 100 μg of HMG-CoA synthase and 0.5 mM or 1 mM [^{14}C]acetyl-CoA (specific activity, 150 cpm/nmol) in 50 mM Tris (pH 9.75) were combined in a volume 70 μl and incubated at room temperature for 5 min. The entire reaction mixture was then applied to a MicroSpin BioP-30 gel filtration column and centrifuged for 4 min at 1,000 \times g. The filtrate was then assayed for protein concentration (5) and radioactivity. The stoichiometry was determined by dividing the moles of acetyl-CoA by the moles HMG-CoA synthase monomer present.

^b The stoichiometry of covalently bound acetyl-CoA is presented. HMG-CoA synthase (100 μg), 200 μg of bovine serum albumin as a carrier protein, and 0.5, 1.0, 2.0, or 3.0 mM [^{14}C]acetyl-CoA (specific activity, 150 cpm/nmol) in 50 mM Tris buffer (pH 9.75) at a final volume of 150 μl were mixed on ice. After 2 min, 1 ml of 10% trichloroacetic acid was added, and each incubation mixture was applied to a glass fiber filter. Filters were washed with 10% trichloroacetic acid and then with 1 ml of ice-cold absolute ethanol, allowed to dry, and counted. The stoichiometry of covalently bound acetyl-CoA was determined by dividing the moles of acetyl-CoA by the moles HMG-CoA synthase monomer present.

^c Mean \pm the standard deviation.




FIG. 6. Coupled conversion of acetyl-CoA to mevalonate catalyzed by *E. faecalis* acetoacetyl-CoA thiolase/HMG-CoA reductase and HMG-CoA synthase. Shown is the effect of hydrogen ion concentration on the conversion of acetyl-CoA to acetoacetyl-CoA (\circ), of HMG-CoA to mevalonate (\square), and of acetyl-CoA to mevalonate (\bullet). All assays employed 50 mM sodium acetate, 50 mM glycine, 50 mM Tris, and 50 mM 2-(*N*-morpholino)ethanesulfonic acid at the indicated pH. Assays of acetoacetyl-CoA thiolase and HMG-CoA reductase activity were conducted essentially as previously described (11), but under the above conditions. For the conversion of acetyl-CoA to mevalonate, the additions were: 1 mM acetyl-CoA, 0.4 mM NADPH, 9 nM *E. faecalis* acetoacetyl-CoA thiolase/HMG-CoA reductase subunit, and 16 nM *E. faecalis* HMG-CoA synthase.

pathway suggests that significant differences may also characterize the bacterial and human forms of additional enzymes of the pathway. Although the glutamate (8), cysteine (17), and histidine (15) implicated as participating in catalysis in the avian form of the enzyme are conserved in *E. faecalis* HMG-CoA synthase (Table 1), the bacterial HMG-CoA synthases and their animal counterparts exhibit only ca. 10% overall sequence identity. The sequences of bacterial HMG-CoA synthases also cluster away from those of the eukaryotic synthases (27). Although low sequence identity and remote clustering of the enzymes from different kingdoms are indicative of different classes of HMG-CoA synthase, this inference must await confirmation by detailed structural investigations.

ACKNOWLEDGMENTS

This research was funded by American Heart Association grant 0150503N (V.W.R.) and by National Institutes of Health grants HL 47113 (V.W.R.) and 52115 (C.V.S.).

We thank Michael Gwynn and Imogen Wilding of Glaxo SmithKline for the gift of *E. faecalis* genomic DNA and Kevin Lehnbeuter for technical assistance.

REFERENCES

- Bach, T. J. 1995. Some new aspects of isoprenoid biosynthesis in plants: a review. *Lipids* **30**:191–202.
- Bischoff, K. M., and V. W. Rodwell. 1996. 3-Hydroxy-3-methylglutaryl-CoA reductase from *Haloflex volcanii*: purification, characterization, and expression in *Escherichia coli*. *J. Bacteriol.* **178**:19–23.
- Bochar, D. A., J. A. Friesen, C. V. Stauffacher, and V. W. Rodwell. 1999. Biosynthesis of mevalonic acid from acetyl-CoA, p. 15–44. In David Cane (ed.), *Isoprenoids including carotenoids and steroids*. Pergamon Press, Oxford, United Kingdom.
- Bochar, D. A., C. V. Stauffacher, and V. W. Rodwell. 1999. Sequence comparisons reveal two classes of 3-hydroxy-3-methylglutaryl coenzyme A reductase. *Mol. Genet. Metab.* **66**:122–127.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Cabano, J., C. Buesa, F. G. Hegardt, and P. F. Marrero. 1997. Catalytic properties of recombinant 3-hydroxy-3-methylglutaryl coenzyme A synthase-1 from *Blattella germanica*. *Insect Biochem. Mol. Biol.* **27**:499–505.
- Chun, K. Y., D. A. Vinarov, and H. M. Miziorko. 2000. 3-Hydroxy-3-methylglutaryl-CoA synthase: participation of invariant acidic residues in formation of the acetyl-S-enzyme reaction intermediate. *Biochemistry* **39**:14670–14681.
- Chun, K. Y., D. A. Vinarov, J. Zajicek, and H. M. Miziorko. 2000. 3-Hydroxy-3-methylglutaryl-CoA synthase: a role for glutamate 95 in general acid/base catalysis of C-C bond formation. *J. Biol. Chem.* **275**:17946–17953.
- Clinkenbeard, K. D., T. Sugiyama, and M. D. Lane. 1975. Cystolic 3-hydroxy-3-methylglutaryl-CoA synthase from chicken liver. *Methods Enzymol.* **35**:160–167.
- Hamano, T., T. Dairi, M. Yamamoto, T. Kawasaki, H. Kaneda, T. Kuzuyama, N. Itoh, and H. Seto. 2001. Cloning of a gene cluster encoding enzymes responsible for the mevalonate pathway from a terpenoid antibiotic-producing *Streptomyces* strain. *Biosci. Biotechnol. Biochem.* **65**:1627–1635.
- Hedl, M., A. Sutherland, E. I. Wilding, M. Mazzulla, D. McDevitt, P. Lane, J. W. Burgner II, K. R. Lehnbeuter, C. V. Stauffacher, M. N. Gwynn, and V. W. Rodwell. 2002. *Enterococcus faecalis* acetoacetyl-CoA thiolase/3-hydroxy-3-methylglutaryl coenzyme A reductase, a dual-function protein of isopentenyl diphosphate biosynthesis. *J. Bacteriol.* **184**:2116–2122.
- Istvan, E. S., M. Palnitkar, S. K. Buchanan, and J. Deisenhofer. 2000. Crystal structure of the catalytic portion of human HMG-CoA reductase: insights into regulation of activity and catalysis. *EMBO J.* **19**:819–830.
- Lawrence, C. M., V. W. Rodwell, and C. V. Stauffacher. 1995. Crystal structure of *Pseudomonas mevalonii* HMG-CoA reductase at 3.0 angstrom resolution. *Science* **268**:1758–1762.
- Lowe, D. M., and P. K. Tubbs. 1985. 3-Hydroxy-3-methylglutaryl-coenzyme A synthase from ox liver. *Biochem. J.* **227**:591–599.
- Misra, I., H. A. Charlier, Jr., and H. M. Miziorko. 1995. Avian cytosolic 3-hydroxy-3-methylglutaryl-CoA synthase: evaluation of the role of cysteines in reaction chemistry. *Biochim. Biophys. Acta* **1247**:253–259.
- Misra, I., and H. M. Miziorko. 1996. Evidence for the interaction of avian 3-hydroxy-3-methylglutaryl-CoA synthase histidine 264 with acetoacetyl-CoA. *Biochemistry* **35**:9610–9616.
- Misra, I., C. Narasimhan, and H. M. Miziorko. 1993. Avian 3-hydroxy-3-methylglutaryl-CoA synthase: characterization of a recombinant cholesterogenic isozyme and demonstration of the requirement for a sulfhydryl functionality in formation of the acetyl-enzyme reaction intermediate. *J. Biol. Chem.* **268**:12129–12135.
- Miziorko, H. M., and M. D. Lane. 1977. 3-Hydroxy-3-methylglutaryl-CoA synthase. Participation of acetyl-S-enzyme and enzyme-S-hydroxymethylglutaryl-S-CoA intermediates in the reaction. *J. Biol. Chem.* **252**:1414–1420.
- Miziorko, H. M., K. D. Clinkenbeard, W. D. Reed, and M. D. Lane. 1975. 3-Hydroxy-3-methylglutaryl-CoA synthase: evidence for an acetyl-S-enzyme intermediate and identification of a cysteinyl sulfhydryl as the site of acetylation. *J. Biol. Chem.* **250**:5768–5773.
- Rohmer, M. 1999. A mevalonate-independent root to isopentenyl diphosphate, p. 45–67. In David Cane (ed.), *Isoprenoids including carotenoids and steroids*. Pergamon Press, Oxford, United Kingdom.
- Rokosz, L. L., D. A. Boulton, E. A. Butkiwicz, G. Sanyal, M. A. Cueto, P. A. Lachance, and J. D. Hermes. 1994. Human cytoplasmic 3-hydroxy-3-methylglutaryl coenzyme A synthase: expression, purification, and characterization of recombinant wild-type and Cys¹²⁹ mutant enzymes. *Arch. Biochem. Biophys.* **312**:1–13.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Schuck, P., M. A. Perugini, N. R. Gonzales, G. J. Howlett, and D. Schubert. 2002. Size-distribution analysis of proteins by analytical ultracentrifugation: strategies and application to model systems. *Biophys. J.* **82**:1096–1111.
- Takagi, M., T. Kuzuyama, S. Takahashi, and H. Seto. 2000. A gene cluster for the mevalonate pathway from *Streptomyces* sp. strain CL190. *J. Bacteriol.* **182**:4153–4157.
- Vollmer, S. H., L. M. Mende-Mueller, and H. M. Miziorko. 1988. Identification of the site of acetyl-S-enzyme formation on avian liver mitochondrial 3-hydroxy-3-methylglutaryl-CoA synthase. *Biochemistry* **27**:4288–4292.
- Weitzman, P. D. J. 1969. Citrate synthase from *Escherichia coli*. *Methods Enzymol.* **13**:22–26.
- Wilding, E. I., J. R. Brown, A. P. Bryant, A. F. Chalker, D. J. Holmes, K. A. Ingraham, S. Iordanescu, C. Y. So, M. Rosenberg, and M. N. Gwynn. 2000. Identification, evolution, and essentiality of the mevalonate pathway for isopentenyl diphosphate biosynthesis in gram-positive cocci. *J. Bacteriol.* **182**:4319–4327.
- Wilding, E. I., D.-Y. Kim, A. P. Bryant, M. N. Gwynn, R. D. Lunsford, D. McDevitt, J. E. Myers, Jr., M. Rosenberg, D. Sylvester, C. V. Stauffacher, and V. W. Rodwell. 2000. Essentiality, expression and characterization of the class II 3-hydroxy-3-methylglutaryl coenzyme A reductase of *Staphylococcus aureus*. *J. Bacteriol.* **182**:5147–5152.