

# Functions Required for Extracellular Quinolone Signaling by *Pseudomonas aeruginosa*

Larry A. Gallagher,<sup>1</sup> Susan L. McKnight,<sup>2</sup> Marina S. Kuznetsova,<sup>1</sup>  
Everett C. Pesci,<sup>2</sup> and Colin Manoil<sup>1\*</sup>

Department of Genome Sciences, University of Washington, Seattle, Washington 98195,<sup>1</sup> and Department of Microbiology and Immunology, East Carolina University, Greenville, North Carolina 27858<sup>2</sup>

Received 22 May 2002/Accepted 28 August 2002

**A set of 30 mutants exhibiting reduced production of the phenazine poison pyocyanin were isolated following transposon mutagenesis of *Pseudomonas aeruginosa* PAO1. The mutants could be subdivided into those with defects in the primary phenazine biosynthetic pathway and those with more pleiotropic defects. The largest set of pleiotropic mutations blocked the production of the extracellular *Pseudomonas* quinolone signal (PQS), a molecule required for the synthesis of secondary metabolites and extracellular enzymes. Most of these *pqs* mutations affected genes which appear to encode PQS biosynthetic functions, although a transcriptional regulator and an apparent response effector were also represented. Two of the genes required for PQS synthesis (*phnA* and *phnB*) had previously been assumed to encode phenazine biosynthetic functions. The transcription of one of the genes required for PQS synthesis (PA2587/*pqsH*) was regulated by the LasI/R quorum-sensing system, thereby linking quorum sensing and PQS regulation. Others of the pleiotropic phenazine-minus mutations appear to inactivate novel components of the quorum-sensing regulatory network, including one regulator (np20) previously shown to be required for virulence in neutropenic mice.**

A complex network of regulatory factors governs the production of secondary metabolites and other virulence factors in the opportunistic pathogen *Pseudomonas aeruginosa*. This network regulates gene expression in response to stimuli such as growth phase, culture density, and oxygen and iron availability (12, 26, 28, 37). Central components of the network are the *las* and *rhl* quorum-sensing systems, which activate gene expression in response to culture density (13). Each system is made up of two genes, one encoding an enzyme which produces a specific acylated homoserine lactone autoinducer (*lasI/rhlI*), and a second encoding a transcriptional activator that binds the corresponding autoinducer (*lasR/rhlR*). The *las* system directs expression of virulence factors such as elastases A and B and alkaline protease (16, 25). The *rhl* system directs expression of rhamnolipid biosynthesis enzymes, pyocyanin biosynthesis enzymes, and hydrogen cyanide synthase (3, 24, 28). In addition, LasI/R regulates expression of both itself and *rhlI* (1, 26). The *las* and *rhl* systems together have been shown to influence the expression of over two hundred genes (36).

Recently, a third signaling system based on 2-heptyl-3-hydroxy-4-quinolone, designated the *Pseudomonas* quinolone signal (PQS), has been shown to be a part of the quorum-sensing regulatory network in *P. aeruginosa* (27). The production of PQS depends on *lasR* (27), and exogenous PQS strongly induces expression of elastase B and *rhlI* in a *lasR* mutant background (22). These results place PQS between the *las* and *rhl* quorum-sensing systems in the quorum-sensing regulatory network (22).

We have described a process (“paralytic killing”) in which *P.*

*aeruginosa* PAO1 rapidly kills the nematode *Caenorhabditis elegans* by cyanide poisoning (8, 14). Previous studies of a different *P. aeruginosa* strain (PA14) had implicated a different poison, the phenazine pyocyanin, in nematode killing (20). Pyocyanin is a redox cycling agent synthesized as a secondary metabolite from chorismate by the *phz* gene products (21). Phenazines have been used as electron acceptors for cyanide production in vitro (2, 6), and it appeared possible that they could be required for cyanide production and nematode killing by strain PAO1. To help address this issue, we isolated and characterized PAO1 mutants defective in phenazine production. Unexpectedly, the majority of the mutants we found appear to be defective in the regulation of phenazine synthesis rather than in the biosynthetic pathway itself. This study presents an analysis of the regulatory mutants and shows that the most common class is defective in PQS signaling.

## MATERIALS AND METHODS

**Strains, plasmids, growth media, and culture conditions.** The *P. aeruginosa* strains used were PAO1 (17) from L. Passador and B. Iglewski (MPAO1) and S. Lory (PAO1seq) and those listed in Tables 1 and 2. Chromosomal transposon insertion mutations were transferred between strains by transformation using a technique to be described in detail elsewhere (N. Benkers and C. Manoil, unpublished studies). Briefly, recipient strains carrying a plasmid encoding phage lambda *red* recombination functions were electroporated with chromosomal DNA isolated from donor strains, with selection for transposon antibiotic-resistant transformants. Transformants were cured of the plasmid carrying lambda *red* prior to further analysis. The *lacZ* transcriptional gene fusion alleles used to construct the strains for Tables 2 and 3 were isolated by Whiteley et al. (36) as transposon insertions in genes *qsc101*, *qsc118*, *qsc128*, *qsc105*, *qsc131*, and *qsc135*. The *Escherichia coli* strains used were DH5 $\alpha$  (30) for plasmid construction and SM10 $\lambda$ pir (32) for conjugal suicide plasmid delivery. The growth media used were brain heart infusion (BHI) agar (Difco), low-phosphate succinate minimal medium (LPSM) (7), L agar (30), and L broth. For visual analysis of  $\beta$ -galactosidase activity, L agar was supplemented with 50- $\mu$ g/ml 5-bromo-4-chloro-3-indolyl galactoside (X-Gal). Plasmids were maintained in *P. aeruginosa* in medium supplemented with 200- $\mu$ g/ml carbenicillin and in *E. coli* in medium

\* Corresponding author. Mailing address: Department of Genome Sciences, Box 357730, University of Washington, Seattle, WA 98195. Phone: (206) 543-7800. Fax: (206) 543-0754. E-mail: manoil@u.washington.edu.

TABLE 1. Mutants defective in pyocyanin production<sup>a</sup>

Strain	Parent	Insert site	Gene	Description	Source or reference
<i>phzI</i> region					
MP601	MPAO1	4,712,812	<i>phzM</i>	Pyocyanin biosynthesis	This study
MP651	PAO1seq	4,714,312	<i>phzA1</i>	Core phenazine biosynthetic operon	This study
*MP705	MPAO1	4,714,312	<i>phzA1</i>	Core phenazine biosynthetic operon	This study <sup>b</sup>
MP602	MPAO1	4,715,531	<i>phzC1</i>	Core phenazine biosynthetic operon	This study
MP652	PAO1seq	4,718,219	<i>phzE1</i>	Core phenazine biosynthetic operon	This study
*MP706	MPAO1	4,718,219	<i>phzE1</i>	Core phenazine biosynthetic operon	This study <sup>b</sup>
MP653	PAO1seq	4,720,699	<i>phzS</i>	Pyocyanin biosynthesis	This study
*MP702	MPAO1	4,720,699	<i>phzS</i>	Pyocyanin biosynthesis	This study <sup>b</sup>
<i>phnAB</i> region					
MP654	PAO1seq	1,081,164	PA0998 ( <i>pqsC</i> )	β-Ketoacyl-acyl protein synthase	This study
*MP703	MPAO1	1,081,164	PA0998 ( <i>pqsC</i> )	β-Ketoacyl-acyl protein synthase	This study <sup>b</sup>
MP603	MPAO1	1,081,418	PA0998 ( <i>pqsC</i> )	β-Ketoacyl-acyl protein synthase	This study
MP604	MPAO1	1,081,470	PA0998 ( <i>pqsC</i> )	β-Ketoacyl-acyl protein synthase	This study
MP655	PAO1seq	1,082,732	PA0999 ( <i>pqsD</i> )	β-Ketoacyl-acyl protein synthase	This study
*MP704	MPAO1	1,082,732	PA0999 ( <i>pqsD</i> )	β-Ketoacyl-acyl protein synthase	This study <sup>b</sup>
MP605	MPAO1	1,082,972	PA1000 ( <i>pqsE</i> )	Hypothetical	This study
MP710	MPAO1	ND <sup>c</sup>	<i>phnA</i>	Anthranilate synthase.	This study
MP551	MPAO1	1,086,674	PA1003 ( <i>pqsR</i> )	<i>lysR</i> -like transcriptional regulator	14
PA2587 region					
MP562	MPAO1	2,927,500	PA2587 ( <i>pqsH</i> )	FAD-dependent monooxygenase	14
MP606	MPAO1	2,928,247	PA2587/PA2588	Between genes	This study
Known regulators					
MP701	MPAO1	ND	<i>lasR</i>	<i>lasR</i> quorum-sensing regulator	15; this study
MP607	MPAO1	3,889,853	<i>rhlI/R</i>	<i>rhl</i> quorum-sensing regulatory locus	This study
MP608	MPAO1	1,013,341	<i>gacS</i>	Global regulator	This study
MP502	MPAO1	1,015,249	<i>gacS</i>	Global regulator	14
Putative regulators					
MP501	MPAO1	4,423,808	PA3946	Similar to <i>B. pertussis</i> virulence gene regulator <i>bvgS</i>	14
MP609	MPAO1	5,302,623	<i>dkxA</i>	Repressor of <i>rhl</i>	This study
MP553	MPAO1	5,304,386	PA4725	Putative two-component sensor kinase	14
MP552	MPAO1	5,304,811	PA4725	Putative two-component sensor kinase	14
MP656	PAO1seq	5,306,302	PA4725	Putative two-component sensor kinase	This study
MP610	MPAO1	5,482,018	PA4886	Putative two-component sensor kinase.	This study
MP611	MPAO1	6,192,554	np20	Similar to <i>E. coli</i> zinc uptake regulator Zur	This study
Miscellaneous					
MP612	MPAO1	447,569	PA0406	Hypothetical protein	This study
MP613	MPAO1	448,033	PA0406	Hypothetical protein	This study
MP614	MPAO1	812,055	PA0744	Probable enoyl-CoA hydratase/isomerase	This study
MP504	MPAO1	812,969	PA0745	Probable enoyl-CoA hydratase/isomerase	14
MP615	MPAO1	3,395,202	PA3031/PA3032	Between genes	This study

<sup>a</sup> All mutants correspond to insertions of IS*phoA*/hah-Tc except the *phnA* and *lasR* mutants, which were constructed by in vitro methods.

<sup>b</sup> Strain was derived by transferring into the chromosome of MPAO1 the insertion allele from the strain listed immediately above in the list.

<sup>c</sup> ND, not determined

supplemented with 100 μg/ml ampicillin. To construct plasmid pLG10, cosmid 122 (containing the *phnAB* genomic region) from the laboratory of S. Lory was first digested with *Hind*III and *Nhe*I. The 10,301-bp, *phnAB*-containing fragment from this digestion was gel purified (QIAGEN kit) and DNA ligase joined to the gel-purified 4,532-bp fragment from pUCP18 (31) cleaved with *Hind*III and *Xba*I. To construct plasmid pLG12, the 6,084-bp *Pst*I fragment from pLG10 was cloned into the *Pst*I site of pUCP18. To construct plasmid pLG14, pLG12 was digested with *Asc*I and *Hind*III and treated with Klenow enzyme (New England Biolabs). The larger fragment was gel purified and self-ligated. To construct plasmid pLG16, pLG10 was digested with *Sac*I and *Asc*I and treated with mung bean nuclease (New England Biolabs). The larger fragment was then gel purified

and self-ligated. All constructs were confirmed by restriction analysis. Standard molecular biology protocols were used (30).

**Transposon mutagenesis and mutant screening.** Transposon IS*phoA*/hah-Tc mutagenesis was carried out as described previously (14). Lawns of mutants derived from strain MPAO1 were screened visually for pigmentation defects after 24 h of growth at 37°C on 3.5-cm-diameter BHI agar plates. Small cultures (200 μl) of mutants of PAO1seq were screened visually for pigmentation defects after 2 days of growth at room temperature in a 96-well format in LPSM.

**DNA sequencing.** The chromosomal DNA flanking the transposon insertions was sequenced as described previously (14), except that primer T*phoA*-II (5'-GTGCAGTAATATCGCCCTGAGCA-3') replaced primer MTN51.1, primer

TABLE 2.  $\beta$ -Galactosidase activities in strains carrying chromosomal *lacZ* reporter fusions to quorum-sensing-controlled genes<sup>a</sup>

Mutant gene	<i>phzC1-lacZ</i>			<i>hcnB-lacZ</i>			<i>pqsH-lacZ</i>			<i>phzD2-lacZ</i>			<i>rhlI-lacZ</i>			<i>cytC-lacZ</i>		
	% Activity	(SEM)	n <sup>c</sup>	% Activity	(SEM)	n	% Activity	(SEM)	n	% Activity	(SEM)	n	% Activity	(SEM)	n	% Activity	(SEM)	n
w-t	(100)	(8)	13 <sup>b</sup>	(100)	(8)	12 <sup>b</sup>	(100)	(3)	10 <sup>b</sup>	(100)	(6)	12 <sup>b</sup>	(100)	(2)	9 <sup>b</sup>	(100)	(8)	10 <sup>b</sup>
<i>pqsC</i>	16	(3)	3	43	(4)	3	85	(4)	2	117	(16)	2	91	(5)	2	71	(13)	2
<i>pqsD</i>	4	(0)	3	30	(5)	2	82	(1)	2	140	(9)	2	84	(7)	2	65	(10)	2
<i>pqsE</i>	3	(0)	3	30	(3)	2	84	(2)	2	121	(11)	2	88	(4)	4 <sup>b</sup>	74	(12)	2
<i>phnA</i>	14	(2)	3	38	(3)	2	83	(4)	2	108	(20)	2	89	(6)	2	75	(17)	2
<i>pqsR</i>	5	(1)	8 <sup>b</sup>	27	(0)	5 <sup>b</sup>	80	(2)	5 <sup>b</sup>	148	(14)	4 <sup>b</sup>	88	(3)	5 <sup>b</sup>	76	(9)	3
<i>pqsH</i>	8	(2)	5 <sup>b</sup>	32	(1)	3	ND <sup>d</sup>			128	(14)	5 <sup>b</sup>	84	(4)	4 <sup>b</sup>	72	(6)	5 <sup>b</sup>
<i>lasR</i>	1	(0)	3	7	(2)	8 <sup>b</sup>	9	(2)	3	5	(1)	4 <sup>b</sup>	5	(0)	4 <sup>b</sup>	49	(4)	4 <sup>b</sup>
np20	ND			6	(1)	6 <sup>b</sup>	6	(0)	6 <sup>b</sup>	4	(0)	7 <sup>b</sup>	5	(0)	6 <sup>b</sup>	61	(2)	7 <sup>b</sup>
PA3946	162	(17)	8 <sup>b</sup>	20	(1)	5 <sup>b</sup>	47	(1)	4 <sup>b</sup>	85	(4)	7 <sup>b</sup>	58	(1)	4 <sup>b</sup>	64	(5)	3

<sup>a</sup> The chromosomal *qsc*-gene-*lacZ* reporter alleles tested are listed at the top of each data column. The gene mutated in each strain is indicated in the far left column.  $\beta$ -Galactosidase activity is shown for each reporter allele as a percentage of the average activity in the wild-type background. 100%  $\beta$ -galactosidase activity represents, for *phzC1-lacZ*, 813 Miller units; for *hcnB-lacZ*, 534 units; for *pqsH-lacZ*, 268 units; for *phzD2-lacZ*, 77 units; for *rhlI-lacZ*, 2,000 units; and for *cytC-lacZ*, 95 units.

<sup>b</sup> Value determined from analysis of duplicate constructed strains.

<sup>c</sup> n, number of independent assays performed.

<sup>d</sup> ND, not done.

HAH-1 (5'-ATCCCCCTGGATGGAAAACGG-3') replaced primer MTN50.1, and primer HAH-2 (5'-AAACGGGAAAGGTTCCGTTCCA-3') replaced primer MTN55.1. Chromosomal locations were determined by BLAST analysis of the transposon-adjacent chromosomal DNA sequences compared with the complete strain PAO1 genome sequence (obtained at [www.pseudomonas.com](http://www.pseudomonas.com)).

**Nematode killing,  $\beta$ -galactosidase, pyocyanin and PQS assays.** Nematode killing assays were carried out as described previously (14).  $\beta$ -Galactosidase assays for Table 2 were carried out as described previously (23) after growth with aeration at 37°C of 5 ml cultures in L broth. All cultures were inoculated from fresh colonies and adjusted to an optical density at 600 nm (OD<sub>600</sub>) of 0.02 before incubation. To assess the optimal growth point for comparing gene expression levels,  $\beta$ -galactosidase levels were monitored over 15 h of growth. For all six reporter alleles in the wild-type background, the  $\beta$ -galactosidase levels reached a linear rate of increase by early stationary phase (~10 h of incubation), and the slopes remained constant, even after 15 h of incubation (data not shown). The results were reproducible in multiple trials, and all subsequent  $\beta$ -galactosidase assays were carried out at the 13-h time point. For extracellular complementation analysis, strains were patched directly next to one another on L agar containing 50- $\mu$ g/ml X-Gal. Increased  $\beta$ -galactosidase activity from the complementation was scored visually at adjacent regions of growth after 2 days of incubation at 37°C. For Table 4, pyocyanin extracted from agar medium was assayed as previously described (14). For Table 5, samples of culture supernatants (4 ml) from 5-ml cultures grown with aeration for 16 h at 37°C from an initial inoculum at OD<sub>600</sub> of 0.02 were extracted with 3 ml of chloroform; 2.5 ml of the chloroform phase was then further extracted with 0.5 ml of 0.2 N HCl, and the OD<sub>520</sub> of the aqueous phase was measured (14). For PQS assays, bacteria from plates grown overnight were inoculated into 5 ml of Luria-Bertani (LB)

broth and cultures were incubated at 37°C with shaking (260 rpm) for approximately 6 h. Cells were then subcultured into 1 ml of LB broth at an A<sub>660</sub> of 0.02 and grown for 24 h at 37°C with shaking (260 rpm). After growth, 300  $\mu$ l of culture was extracted twice with 900  $\mu$ l of acidified ethyl acetate (27) by vigorously vortexing for 30 s, followed by centrifugation at 16,000  $\times$  g for 5 min. An 800- $\mu$ l aliquot of the upper organic layer from each extraction was transferred to a single microcentrifuge tube and allowed to dry overnight at room temperature. The following day, dried extracts were either analyzed by thin-layer chromatography (TLC) or stored at -20°C until analysis. For TLC analysis, dried extracts were resuspended in 50  $\mu$ l of a 1:1 acidified ethyl acetate-acetonitrile mixture by vortexing and pulse spinning multiple times. Ten microliters of each extract was loaded onto a Silica Gel 60 F<sub>254</sub> (10 by 20 cm; EM Science), along with synthetic PQS. Chromatography was performed with a solvent mixture containing a 17:2:1 ratio of methylene chloride-acetonitrile-1,4-dioxane. When the solvent front neared the top of the plates, the plates were photographed under long-wave UV light (365 nm) using a Polaroid camera with ISO 3000 black-and-white film.

## RESULTS

**Isolation of *P. aeruginosa* mutants deficient in pyocyanin production.** We identified transposon insertion mutants of *P. aeruginosa* strain PAO1 defective in production of the blue phenazine pigment pyocyanin by visual screening (Materials and Methods). In the process of carrying out the screen, we observed that different isolates of wild-type PAO1 behaved

TABLE 3. Induction of *phzC1-lacZ* expression by extracellular complementation

Adjacent strain <sup>a</sup>	<i>phzC1-lacZ</i> fusion strain <sup>a,b</sup>						
	MP882.1 ( <i>pqsC</i> )	MP883.1 ( <i>pqsD</i> )	MP884.1 ( <i>pqsE</i> )	MP894.1 ( <i>phnA</i> )	MP885.2 ( <i>pqsR</i> )	MP886.1 ( <i>pqsH</i> )	MP892.1 ( <i>lasR</i> )
MPAO1 (wild-type)	+	++	-	+	+/-	+++	++++
MP703 ( <i>pqsC</i> )	-	-	-	-	-	-	-
MP704 ( <i>pqsD</i> )	-	-	-	-	-	+	-
MP605 ( <i>pqsE</i> )	+	++	-	+	+/-	+++	++++
MP710 ( <i>phnA</i> )	-	-	-	-	-	-	-
MP551 ( <i>pqsR</i> )	-	-	-	-	-	-	+/-
MP562 ( <i>pqsH</i> )	+	+	-	+/-	+/-	-	+++
MP701 ( <i>lasR</i> )	+/-	+	-	-	-	-	-
MP607 ( <i>rhlR/I</i> )	+/-	++	-	-	-	+++	++++
MP611 (np20)	-	-	-	-	-	-	-

<sup>a</sup> The gene mutated in each strain is indicated parenthetically

<sup>b</sup> Increased  $\beta$ -galactosidase activity over background levels as measured by X-Gal hydrolysis was assessed visually and ranged from slight (+/-) to extensive (++++).

TABLE 4. Pyocyanin production, nematode killing, and PQS production by *P. aeruginosa* mutants

Strain	Mutant gene	Pyocyanin <sup>a</sup>	% Nematode killing <sup>a</sup>	PQS
MPAO1		(1.00)	98 (2)	+++
<i>phz1</i> cluster				
MP601	<i>phzM</i>	0.01 (0.00)	100 (0)	+++
MP705	<i>phzA1</i>	0.04 (0.01)	99 (1)	+++
MP706	<i>phzE1</i>	0.17 (0.02)	99 (1)	+++
MP702	<i>phzS</i>	0.02 (0.00)	78 (11)	+++
<i>phnAB</i> region				
MP703	PA0998 ( <i>pqsC</i> )	0.04 (0.00)	36 (8)	–
MP704	PA0999 ( <i>pqsD</i> )	0.02 (0.01)	14 (2)	–
MP605	PA1000 ( <i>pqsE</i> )	0.01 (0.01)	25 (8)	+++
MP710	<i>phnA</i>	0.04 (0.02)	37 (8)	–
MP551	PA1003 ( <i>pqsR</i> )	0.01 (0.00)	12 (10)	–
PA2587 region (MP562)	PA2587 ( <i>pqsH</i> )	0.02 (0.01)	9 (4)	–
Known regulators				
MP701	<i>lasR</i>	0.01 (0.00)	2 (1)	+/-
MP607	<i>rhII/R</i>	0.02 (0.00)	11 (8)	+++
MP502	<i>gacS</i>	0.10 (0.01)	50 (5)	+++
Putative regulators				
MP501	PA3946	0.31 (0.02)	22 (7)	+++
MP552	PA4725	0.19 (0.04)	19 (8)	+++
MP610	PA4886	0.29 (0.03)	40 (6)	+++
MP611	np20	0.00 (0.00)	4 (3)	+/-
Miscellaneous				
MP613	PA0406	0.07 (0.00)	48 (17)	+
MP504	PA0745	0.03 (0.01)	50 (13)	+
MP615	PA3031/PA3032	0.03 (0.01)	1 (1)	+

<sup>a</sup> Values listed are averages from three separate assays. Pyocyanin recoveries relative to the parent strain (MPAO1) are presented. Numbers in parentheses indicate standard errors of the mean.

differently with respect to pyocyanin production. During liquid growth in LPSM, a medium that induces strong pyocyanin production (7), the PAO1 strain whose genome had been sequenced (designated PAO1seq) (33) produced significantly more pyocyanin than did our laboratory strain of PAO1 (designated MPAO1, obtained from L. Passador and B. Iglewski) (data not shown). When grown on BHI agar, MPAO1 produced much more pyocyanin than did PAO1seq. We initially screened for mutants in the PAO1seq strain (2,800 IS*phoA*/hah-Tc transposon insertion mutants screened) and switched to screening in our laboratory strain once the conditions promoting its pyocyanin production were identified (2,100 IS*phoA*/hah-Tc transposon insertion mutants screened) (Materials and Methods).

Twenty-one pyocyanin-deficient mutants which formed normal-size colonies on L agar were isolated (Table 1). Seven additional pyocyanin-deficient mutants obtained in a previous screen (14) and two inactivating mutations constructed in vitro (in *lasR* and *phnA*) were also represented in the set of mutants analyzed (Table 1). We determined the transposon insertion sites for the new mutants by PCR amplification and sequencing of the genomic DNA flanking each transposon insertion, followed by BLAST analysis against the completed PAO1 genome sequence.

A variety of genes were affected in the mutant set, including clusters in the *phz1* and *phnAB* regions (Table 1). The *phz1* region (genes *phzM*, *phzA1* through *phzG1*, and *phzS*) encodes pyocyanin biosynthetic enzymes (21). The *phnA* and *phnB*

TABLE 5. Pyocyanin production by *phnAB*-region mutants carrying plasmids

Plasmid	<i>pqs</i> genes on plasmid	Strain <sup>a</sup>					
		MPAO1	MP703 ( <i>pqsC</i> )	MP704 ( <i>pqsD</i> )	MP605 ( <i>pqsE</i> )	MP710 ( <i>phnA</i> )	MP551 ( <i>pqsR/mvfR</i> )
pUCP18		0.29 (0.05)	0.02 (0.00)	0.00 (0.00)	0.00 (0.00)	0.01 (0.00)	0.01 (0.00)
pLG10	<i>pqsABCDE</i>	0.93 (0.10)	0.78 (0.04)	0.82 (0.06)	0.80 (0.09)	0.77 (0.05)	0.00 (0.00)
pLG12	<i>pqsABCD</i>	0.13 (0.02)	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)	0.01 (0.00)	0.00 (0.00)
pLG14	<i>pqsABC</i>	0.17 (0.04)	0.01 (0.00)	0.01 (0.01)	0.02 (0.01)	0.02	0.01 (0.00)
pLG16	<i>pqsABC</i>	0.35	0.02	0.01	0.01	0.02	0.01

<sup>a</sup> The gene mutated in each strain is indicated. Pyocyanin was assayed as described previously (14), and its recovery is expressed in absorbance units. Standard errors of the mean are shown where multiple independent assays were performed.



genes encode an anthranilate synthase homologue originally proposed to encode a phenazine biosynthetic function (10), a role which has been questioned in a recent report (21). Numerous regulatory genes were also identified in the mutant set, including three (*lasR*, *rhlI*, and *gacS*) known to be a part of the quorum-sensing regulatory network. Four more novel putative regulatory genes were also identified, including a homologue of the *E. coli* zinc uptake regulator Zur (np20) and three putative two-component regulatory genes (PA3946, PA4725, and PA4886). (PA gene numbers refer to ORF designations assigned to unnamed genes in the PAO1 genome sequence [www.pseudomonas.com].) Additional potential regulators were also identified (Table 1).

Since the two supposedly identical parent strains (MPAO1 and PAO1seq) displayed different pyocyanin production characteristics, it was essential for further analysis that the new mutations be characterized in an isogenic strain background. The transposon insertion alleles generated in PAO1seq were therefore transferred into the chromosome of strain MPAO1 by transformation (Materials and Methods). The resulting MPAO1 derivatives are marked with asterisks (Table 1). The inactivating alleles of *lasR* (from strain PAO-R1 [15]) and *phnA* (S. L. McKnight and E. C. Pesci, unpublished data) were also transferred into the MPAO1 strain background (Table 1).

**Pyocyanin is not required for nematode killing.** We measured pyocyanin production and nematode killing for a subset of the pyocyanin-deficient mutants, representing insertions in most of the genes identified in the mutant set (Table 4). Previous studies had shown that hydrogen cyanide is necessary and sufficient for the killing (14). Strains carrying mutations in genes of the core phenazine biosynthetic locus (*phzM*, *phzA1*, *phzE1*, and *phzS*) (21) killed worms efficiently in spite of severe defects in pyocyanin production (Table 4), implying that pyocyanin (and probably other phenazines) is not essential for cyanide production in *P. aeruginosa*. Similar results were obtained in studies of several phenazine-deficient mutants of other *Pseudomonas* species (C. Cosma, D. Mavrodi, C. Manoil, and L. Thomashow, unpublished results). Remarkably, all of the pyocyanin-deficient mutants affecting genes outside of the *phz1* locus failed to kill worms efficiently and are thus apparently pleiotropically defective in both cyanide and pyocyanin production. Direct assays of HCN production (14) and expression of HCN biosynthetic gene-*lacZ* fusions (see below) provide additional evidence of this pleiotropy.

**Identification of genes required for PQS production.** The recently identified PQS is known to be required for phenazine production in *P. aeruginosa* (21, 22). Since one of the genes we identified, PA2587, had been found to be required for PQS production (E. C. Pesci, unpublished), we examined production of PQS by the other pyocyanin-deficient mutants. PQS production was assayed using TLC under conditions which distinguish PQS from the acylated homoserine lactone autoinducers (Fig. 1 and Table 4). Although most of the strains produced PQS at levels comparable to the wild-type strain MPAO1, seven were negative, including most of the strains with mutations in the *phnAB* region (in genes PA0998, PA0999, *phnA*, and PA1003/*mvfR*), PA2587, *lasR*, and np20. Detectable spots with an  $R_f$  similar to PQS were observed for the *lasR* and np20 mutants, but these spots were purple as




FIG. 1. PQS production by *P. aeruginosa* strains. PQS samples extracted from 24-h cultures were analyzed by TLC. (A) Lanes: 1, 50 ng of PQS; 2, strain MP701 (*lasR*); 3, strain MP607 (*rhlR/I*); 4, strain MP502 (*gacS*); 5, strain MP501 (PA3946); 6, strain MP552 (PA4725); 7, strain MP610 (PA4886); 8, strain MP611 (np20); 9, strain MP613 (PA0406); 10, strain MP504 (PA0745); 11, strain MP615 (PA3031/3032); 12, strain MPAO1. (B) Lanes: 1, 50 ng of PQS; 2, strain MP703 (*pqsC*); 3, strain MP704 (*pqsD*); 4, strain MP605 (*pqsE*); 5, strain MP710 (*phnA*); 6, strain MP551 (*pqsR/mvfR*); 7, strain MP562 (*pqsH*); 8, strain MP601 (*phzM*); 9, strain MP705 (*phzA1*); 10, strain MP706 (*phzE1*); 11, strain MP702 (*phzS*); 12, strain MPAO1. The arrowhead in each panel indicates the position of PQS.

opposed to the fluorescent blue of authentic PQS (data not shown).

The finding that mutations in several genes in the *phnAB* region prevented PQS production suggested that this region may encode PQS biosynthetic functions. The genomic organization of the region suggests a possible five-gene operon made up of genes PA0996 to PA1000 (Fig. 2). Based on the results presented here and in an independent study (9) confirming that these genes play essential roles in PQS biosynthesis and signaling, we propose naming the genes *pqsABCDE*. We pro-




FIG. 2. Map of the *phnAB* genomic region and plasmids used for complementation analysis. The vertical arrows in the map of the *pqs* and *phn* genes indicate the locations of insertions in strains MP703, MP603, MP604, MP704, MP605, MP710, MP551, and MP562, respectively (reading left to right). The *phnA* mutation in MP710 is a constructed deletion carrying a Tc<sup>r</sup> cassette. The chromosomal regions present in recombinant plasmids pLG10, pLG12, pLG14, and pLG16 are shown, with the open triangle indicating the orientation of the *P<sub>lac</sub>* promoter in the pUCP18 vector.

pose the designations *pqsH* for the unlinked gene PA2587 and *pqsR* for PA1003, which encodes a member of the LysR family of transcriptional regulators corresponding to strain PA14 Mvfr (5).

Although most of the mutations in the *phnAB* region prevented PQS production, the insertion in *pqsE* produced a substance comigrating with PQS (Fig. 1). This result suggests that *pqsE* is not required for PQS synthesis, even though it is required for phenazine and cyanide production. The PQS-deficient and pyocyanin-deficient phenotypes cotransformed with the *pqsE* insertion mutation (data not shown), implying that they were not due to an undetected secondary mutation in the original mutant. While *pqsE* may play a role in pyocyanin production that is unrelated to PQS signaling, it appears more likely, given its genetic context, that *pqsE* is either responsible for a modification in the PQS biosynthetic pathway that is not distinguishable by our assay or that *pqsE* is required for the cellular response to PQS. Studies presented below support the second of these alternatives.

**Complementation analysis.** We carried out complementation studies to investigate whether polar effects were responsible for the *phnAB*-region mutant phenotypes. Four plasmids were constructed (Fig. 2): pLG10 (which carries *pqsABCDE*), pLG12 (*pqsABCD*), pLG14 (*pqsABC*) and pLG16 (*pqsABC*). Selected *phnAB* region mutants carrying these plasmids were examined for PQS and pyocyanin production. Plasmids pLG10 (*pqsABCDE*) and pLG12 (*pqsABCD*) fully restored PQS production to all the mutants tested except the *pqsR* (*mvfr*) mutant (Fig. 3). These results demonstrate that polar effects on *phnA* and *phnB* were not responsible for the PQS defects observed in the *pqsC* and *pqsD* mutants and confirm that the *pqsABCDE* operon is itself required for PQS production. Plasmids pLG14 (*pqsABC*) and pLG16 (*pqsABC*) restored PQS production to the *pqsC* mutant but not to the *pqsD* mutant (Fig. 3), indicating that *pqsC* and *pqsD* are both needed for PQS biosynthesis.

The pattern of complementation of the pyocyanin production defects differed from that of the PQS defects (Table 5). Although, as expected, plasmid pLG10 restored pyocyanin production to all the mutants tested except the *pqsR* (*mvfr*) mutant, none of the other plasmids restored pyocyanin production to any of the mutants. Since none of the other plasmids carry the *pqsE* gene (Fig. 2A), the results imply that *pqsE* is required for pyocyanin (but not PQS) production and that the mutations in *pqsC* and *pqsD* are polar on *pqsE*.

Surprisingly, although none of the complementing plasmids carried the *phnA* gene, they all complemented the PQS defect (and pLG10 complemented the pyocyanin defect) of a *phnA* mutant (Fig. 3 and Table 5). *phnA* and *phnB* encode an anthranilate synthase (10), and anthranilate is a precursor of PQS (4). Since there are five anthranilate synthase homologues in addition to PhnAB encoded in the *P. aeruginosa* genome (10, 11; Pesci, unpublished), it is possible that multicopy expression of the plasmid-borne *pqsA-D* leads to PQS production using anthranilate synthesized by one of the other enzymes. Recruitment of anthranilate produced by another anthranilate synthetase may also account for why the presence of pLG10 (which lacks *phnAB*) leads to significantly increased production of PQS and pyocyanin over wild-type in the strains examined (including the *phnA* mutant) (Table 5 and Fig. 3). An-




FIG. 3. Complementation of *Pqs*<sup>-</sup> mutants. PQS samples extracted from 24-h cultures of strains carrying different *pqs* genes in plasmids were analyzed by TLC. (A) Lane 1, strain MPAO1 grown without a plasmid; lanes 2 to 7, strains MPAO1, MP703 (*pqsC*), MP704 (*pqsD*), MP605 (*pqsE*), MP710 (*phnA*), and MP551 (*mvfr*), respectively, carrying plasmid pUCP18; lanes 8 to 13, the same strains as the previous six lanes, but carrying plasmid pLG10; lane 14, 50 ng of PQS. (B) Lane 1, strain MPAO1 grown without a plasmid; lanes 2 to 7, strains MPAO1, MP703 (*pqsC*), MP704 (*pqsD*), MP605 (*pqsE*), MP710 (*phnA*), and MP551 (*pqsR*), respectively, carrying plasmid pLG12; lanes 8 to 10, strains MPAO1, MP703 (*pqsC*), and MP704 (*pqsD*), respectively, carrying plasmid pLG14; lanes 11 to 14, strains MPAO1, MP703 (*pqsC*), MP704 (*pqsD*), and MP605 (*pqsE*), respectively, carrying plasmid pLG16; lane 15, 50 ng of PQS. The arrowhead indicates the position of PQS. (C) Summary of complementation results shown in panels A and B.

thraniolate synthase redundancy could also help account for the finding that the effect of *phnA* inactivation on PQS production depends critically on strain background (Pesci, unpublished).

**Transcriptional analysis of mutants.** We examined the effects of *pqs* and other pleiotropic mutations on the expression of six quorum sensing-controlled (*qsc*) *lacZ* transcriptional gene fusions isolated by Whiteley et al. (36). The reporter gene fusions were transferred into the chromosomes of the wild-type (MPAO1) and pyocyanin-deficient mutant strains by transformation (Materials and Methods). The six fusions used for the analysis (Table 2, top row) represent three of the four major classes of quorum sensing-regulated genes identified in the earlier study (36) and include *phzC1* (phenazine biosynthesis), *hcnB* (cyanide biosynthesis), and *pqsH* (PQS synthesis).

We assessed the optimal growth phase at which to compare the *qsc* gene expression levels in the various mutants by monitoring expression of each of the six fusion alleles in the wild-type (MPAO1) strain background during growth in liquid culture. By mid-stationary phase, all six strains showed significant and reproducible  $\beta$ -galactosidase activity (data not shown). We therefore chose a time point in mid-stationary phase for analyzing expression levels in the various mutant backgrounds (Materials and Methods). For most combinations, two independent constructions were assayed. The  $\beta$ -galactosidase levels measured in the different reporter strains are shown in

Table 2, with the values of the mutant strains expressed relative to the corresponding wild-type values.

The mutations in genes implicated in PQS signaling (*pqsCDE*, *phnA*, *pqsR*, and *pqsH*) all reduced transcription of both *phzC1* (more than sixfold) and *hcnB* (about threefold), results compatible with their decreased pyocyanin production and nematode killing (Table 4). The *pqs* mutations did not significantly reduce expression of the other four *lacZ* fusions examined. Thus, for all six fusions, the effects on expression were similar for mutations in the different *pqs* genes, a finding compatible with the hypothesis that all participate in a common regulatory process.

We were surprised that none of the mutations in the genes required for PQS synthesis significantly reduced *rhII* transcription (Table 2). This finding appears to contradict the previous finding that PQS supplied exogenously to a *lasR* mutant greatly enhanced *rhII* transcription from a multicopy plasmid (22). Since the latter result was obtained at a later growth phase than that examined here, the two findings could reflect growth-stage-dependent differences in *rhII* transcription. To test this possibility, we examined *rhII* transcription in some of our mutant strains at the same late growth point that was used for the plasmid-based assay (18 h of growth). However, we observed that even at the later growth point, *rhII* transcription in the *pqsC*, *pqsD*, *pqsE*, *phnA*, *pqsR*, and *pqsH* mutant backgrounds was at 85% (2%), 94% (0%), 88%, 92%, 91% (0%), and 89% (8%) of wild-type levels, respectively (standard errors given in parentheses). Our results thus indicate that loss of PQS does not prevent *rhII* transcription.

The *lasR* and *np20* mutations were highly pleiotropic, causing greater than 10-fold reductions in the activities of the majority of the reporter fusions analyzed, including the *pqsH-lacZ* fusion. The *lasR* and *np20* mutant defects in producing PQS can thus be at least partially explained by reduced *pqsH* expression. Of the three novel two-component regulators (Table 1), only the mutation in PA3946 showed significant reductions in *qsc* fusion expression, reducing transcription of *hcnB* fivefold and that of *pqsH* twofold (Table 2 and data not shown).

**Extracellular complementation.** To further investigate the roles that the various *pqs* genes play in PQS signaling, we tested whether diffusible substances (e.g., PQS) produced by the mutants could induce transcription of phenazine biosynthetic genes in other mutant strains. Wild-type MPAO1 or PQS-defective mutants were grown adjacent to mutants carrying the chromosomal *phzC1-lacZ* transcriptional reporter on LB agar containing X-Gal indicator, and expression of the gene fusion was scored by visual inspection (Table 3).

Most of the findings were in agreement with the results of the PQS assays presented above. Strains shown to produce PQS (wild-type, *pqsE*, and *rhII/R*) increased  $\beta$ -galactosidase expression in nearly all of the *phzC1-lacZ* fusion strains except that carrying the *pqsE* mutation. The responses of the *pqsH* and *lasR* mutant strains were particularly strong. The failure of the *pqsE* mutant strain to be induced is compatible with a role of PqsE in the response to PQS, as suggested above. (The findings also indicate that the polarity of the *pqsC* and *pqsD* mutations on *pqsE* [see above] may be incomplete.) The failure of most of the PQS-nonproducing mutants (*pqsC*, *pqsD*, *phnA*, and *pqsR*) to significantly complement any of the *phzC1-lacZ*

fusion strains is also in accord with expectations. On the other hand, the relatively strong extracellular complementation by the *pqsH* mutant of several of the PQS-deficient fusion strains is not simply explained and suggests the action of a diffusible PQS intermediate (i.e., analogous to auxotroph cross-feeding).

The behavior of several regulatory mutations was also in accord with the PQS assays. Both *lasR* and *np20* mutations greatly reduced extracellular complementation of the *phzC1-lacZ* fusion strains, whereas the *rhII/R* mutation showed little reduction relative to wild type.

## DISCUSSION

This report presents a genetic analysis of pyocyanin production in *P. aeruginosa*. Pyocyanin is a redox cycling phenazine poison produced as a secondary metabolite from chorismate by a multistep pathway (21). Pyocyanin production is highly regulated, being subject to control by two quorum-sensing systems (Las and Rhl), a novel extracellular quinolone signal (PQS), and other components of the quorum-sensing regulatory network (3, 19, 22, 29, 36).

We initially set out to determine whether pyocyanin or other phenazine compounds were required for cyanide-mediated nematode killing by *P. aeruginosa* PAO1. Although our previous experiments had shown that hydrogen cyanide is sufficient for the killing (14), studies with a different strain of *P. aeruginosa* (PA14) had revealed a partial correlation between killing of nematodes and production of pyocyanin (20). Biochemical studies had also shown that under some conditions phenazines participate in the synthesis of cyanide (2, 6). We therefore identified PAO1 mutants showing reduced production of pyocyanin and tested them for killing. Several of the mutants with severe defects in pyocyanin production due to mutations inactivating the biosynthetic pathway showed normal nematode killing, indicating that pyocyanin is not required for nematode killing by *P. aeruginosa* PAO1.

Remarkably, the majority of the mutations leading to pyocyanin defects did not alter biosynthetic pathway genes and were pleiotropically defective in both phenazine production and nematode killing (cyanide production). Three mutations (in *lasR*, *rhII/R*, and *gacS*) inactivated components of the quorum-sensing regulatory network already known to be required for production of the two substances. However, the largest class of pleiotropic mutations was found to cause dramatic reductions in the production of the quinolone signal (PQS). These genes included a cluster in the *phnAB* region (PA0998, PA0999, and PA1003) and an unlinked gene (PA2587). A *phnA* deletion mutant constructed *in vitro* was also defective in PQS synthesis. A different study has demonstrated that two other *phnAB* region genes (PA0996 and PA0997) are also required for PQS synthesis (9). Another gene in the *phnAB* cluster (PA1000) was not required for PQS production, but may participate in the cellular response to PQS (see below). The new genes required for the production and action of PQS were named *pqsA-R* (Fig. 2 and 4).

PQS biosynthetic functions of several of the *pqs* products are suggested by their sequence homologies (Fig. 4). PhnA and PhnB presumably synthesize the anthranilate precursor of PQS from chorismate (4, 10). Gene *pqsA* encodes a product homologous to benzoate coenzyme A ligase, which may be involved






FIG. 4. PQS synthesis and regulation. The genomic organization of the *pqs* genes and a model for PQS biosynthesis and regulation are shown. We propose that PQS is synthesized from chorismate via anthranilate by the *phnAB*, *pqsABCD*, and *pqsH* gene products as indicated. Points of regulation by PqsR, LasI/R, np20, and PqsE based on the results presented in this and a previous report (5) are shown. The PqsR (MvfR) regulation of *pqsABCDE* is suggested by the finding that a plasmid carrying *pqsABCDE* complements a *phnA* mutant but not a *pqsR* mutant for PQS production (Fig. 3C).

in activating anthranilate for PQS synthesis. Genes *pqsB*, *pqsC*, and *pqsD* encode proteins homologous to  $\beta$ -keto-acyl-acyl carrier protein synthases and are presumably involved in the production of a long chain hydrocarbon which reacts with anthranilate in the PQS biosynthetic pathway (4). Gene *pqsE* is not homologous to any defined proteins and our results indicate that it is not required for PQS synthesis. Gene *pqsH* encodes a putative FAD-dependent monooxygenase that may be responsible for the addition of the hydroxyl group to PQS.

Although *phnA* and *phnB* were originally assumed to encode an anthranilate synthetase comprising part of the phenazine biosynthetic pathway (10), recent studies by Mavrodi et al. (21) imply that they are unlikely to participate directly in the formation of the phenazine nucleus. The requirement of *phnAB* for PQS biosynthesis indicates that rather than playing a direct role in phenazine biosynthesis, the defect of *phnAB* mutants in pyocyanin production is most likely due to the highly pleiotropic phenotype caused by reduced PQS signaling. The observation that the *lysR* regulator encoded by *mvfR* (*pqsR*) in strain PA14 is required for the expression of *phnAB* also accounts directly for why the gene is required for the production of multiple virulence factors (5). An earlier study showed that PA14 *mvfR* was required for production of the Las system autoinducer and/or PQS (the assay used did not distinguish between the two substances), although the nature of the requirement was not clear (5).

To further characterize the Pqs proteins and other functions needed for pyocyanin production, we examined the effects of the corresponding mutations on the expression of several quorum sensing-regulated *lacZ* transcriptional gene fusions described by Whiteley et al. (36). Several conclusions emerged from this analysis. First, mutations in the genes required for PQS production (*pqsC*, *pqsD*, *phnA*, *pqsR*, and *pqsH*) and re-

sponse (*pqsE*) reduced expression of pyocyanin and hydrogen cyanide biosynthetic functions (*phzC1* and *hcnB*), indicating that reduced transcription accounts for the defects in the production of both substances. These findings are compatible with models proposing that PQS signaling controls expression of genes regulated by the *rhl* quorum-sensing system (22). Second, transcription of *pqsH* was severely reduced in the *lasR* mutant background. Since *pqsH* is required for PQS synthesis, this finding accounts for a previous observation that PQS production requires LasR function (27) and provides a specific link between the Las and PQS regulatory systems. Third, mutations in the putative regulators np20 and PA3946 reduced transcription of a number of quorum sensing-controlled genes, showing that these functions participate in the quorum-sensing regulatory network. The np20 mutant exhibited a particularly strong phenotype, suggesting that it functions above PQS signaling in the network. np20 is a homologue of the *E. coli* zinc uptake regulator Zur, whose expression is induced by respiratory mucus from cystic fibrosis patients and which is required for virulence in neutropenic mice (34, 35). PA3946 is homologous to *Bordetella pertussis* *bvgS*, a primary regulator of virulence genes (18).

The role of *pqsE* differs from that of the other *pqs* genes. Loss of *pqsE* function causes defects in pyocyanin production, worm killing, and expression of quorum sensing-regulated *lacZ* gene fusions, defects which parallel those observed for the other *pqs* mutants. However, *pqsE* is not required for PQS biosynthesis as assessed by chromatography and extracellular complementation. The findings suggest a role for PqsE in the cellular response to PQS. Such a role is supported by the finding that *pqsE* mutants are not complemented extracellularly by wild-type and other strains for *phzC1-lacZ* expression.

In conclusion, this study has helped specify functions required for the action of the novel extracellular quinolone signaling system in *P. aeruginosa*. In addition, this work has identified several new regulators which appear to belong to the quorum-sensing regulatory network. Although the biological meaning of the formidable complexity of this network remains elusive, specifying the components which make it up constitutes an important step in approaching this goal.

#### ACKNOWLEDGMENTS

We thank Marvin Whiteley and E. Peter Greenberg for providing reporter strains, Matt Wolfgang and Steve Lory for cosmids, Linda Thomashow and Dmitri Mavrodi for helpful discussions and strains, Chris Cosma for sharing unpublished data, and David D'Argenio for helpful discussions.

This work was supported in part by a Cystic Fibrosis Research Development Program postdoctoral fellowship (to L.A.G.), by National Institutes of Health grant R01-AI46682 (to E.C.P.), and by a grant from the Cystic Fibrosis Foundation (to C.M.).

#### REFERENCES

1. Albus, A. M., E. C. Pesci, L. J. Runyen-Janecky, S. E. West, and B. H. Iglewski. 1997. Vfr controls quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* **179**:3928–3935.
2. Blumer, C., and D. Haas. 2000. Mechanism, regulation, and ecological role of bacterial cyanide biosynthesis. *Arch. Microbiol.* **173**:170–177.
3. Britz, J. M., and D. E. Ohman. 1995. Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhlI, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *J. Bacteriol.* **177**:7155–7163.
4. Calfee, M. W., J. P. Coleman, and E. C. Pesci. 2001. Interference with *Pseudomonas* quinolone signal synthesis inhibits virulence factor expression by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **98**:11633–11637.



5. Cao, H., G. Krishnan, B. Goumnerov, J. Tsongalis, R. Tompkins, and L. G. Rahme. 2001. A quorum sensing-associated virulence gene of *Pseudomonas aeruginosa* encodes a LysR-like transcription regulator with a unique self-regulatory mechanism. *Proc. Natl. Acad. Sci. USA* **98**:14613–14618.
6. Castric, C. A. 1981. The metabolism of hydrogen cyanide by bacteria, p. 233–261. In B. Vennesland, E. E. Conn, C. J. Knowles, J. Westley, and F. Wissing (ed.), *Cyanide in biology*. Academic Press, New York, N.Y.
7. Cox, C. D. 1986. Role of pyocyanin in the acquisition of iron from transferrin. *Infect. Immun.* **52**:263–270.
8. Darby, C., C. L. Cosma, J. H. Thomas, and C. Manoil. 1999. Lethal paralysis of *Caenorhabditis elegans* by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **96**:15202–15207.
9. D'Argenio, D., M. W. Calfee, P. B. Rainey, and E. C. Pesci. 2002. Autolysis and autoaggregation in *Pseudomonas aeruginosa* colony morphology mutants. *J. Bacteriol.* **184**:6481–6489.
10. Essar, D. W., L. Eberly, A. Hadero, and I. P. Crawford. 1990. Identification and characterization of genes for a second anthranilate synthase in *Pseudomonas aeruginosa*: interchangeability of the two anthranilate synthases and evolutionary implications. *J. Bacteriol.* **172**:884–900.
11. Essar, D. W., L. Eberly, C. Y. Han, and I. P. Crawford. 1990. DNA sequences and characterization of four early genes of the tryptophan pathway in *Pseudomonas aeruginosa*. *J. Bacteriol.* **172**:853–866.
12. Fuqua, C., M. R. Parsek, and E. P. Greenberg. 2001. Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu. Rev. Genet.* **35**:439–468.
13. Fuqua, C., S. C. Winans, and E. P. Greenberg. 1996. Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu. Rev. Microbiol.* **50**:727–751.
14. Gallagher, L. A., and C. Manoil. 2001. *Pseudomonas aeruginosa* PAO1 kills *Caenorhabditis elegans* by cyanide poisoning. *J. Bacteriol.* **183**:6207–6214.
15. Gambello, M. J., and B. H. Iglewski. 1991. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *J. Bacteriol.* **173**:3000–3009.
16. Gambello, M. J., S. Kaye, and B. H. Iglewski. 1993. *LasR* of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (*apr*) and an enhancer of exotoxin A expression. *Infect. Immun.* **61**:1180–1184.
17. Holloway, B. W., V. Krishnapillai, and A. F. Morgan. 1979. Chromosomal genetics of *Pseudomonas*. *Microbiol. Rev.* **43**:73–102.
18. Kinnear, S. M., R. R. Marques, and N. H. Carbonetti. 2001. Differential regulation of Bvg-activated virulence factors plays a role in *Bordetella pertussis* pathogenicity. *Infect. Immun.* **69**:1983–1993.
19. Latifi, A., M. K. Winson, M. Foglino, B. W. Bycroft, G. S. Stewart, A. Lazdunski, and P. Williams. 1995. Multiple homologues of LuxR and LuxI control expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Mol. Microbiol.* **17**:333–343.
20. Mahajan-Miklos, S., M. W. Tan, L. G. Rahme, and F. M. Ausubel. 1999. Molecular mechanisms of bacterial virulence elucidated using a *Pseudomonas aeruginosa*-*Caenorhabditis elegans* pathogenesis model. *Cell* **96**:47–56.
21. Mavrodi, D. V., R. F. Bonsall, S. M. Delaney, M. J. Soule, G. Phillips, and L. S. Thomashow. 2001. Functional analysis of genes for biosynthesis of pyocyanin and phenazine-1-carboxamide from *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **183**:6454–6465.
22. McKnight, S. L., B. H. Iglewski, and E. C. Pesci. 2000. The *Pseudomonas* quinolone signal regulates *rhl* quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* **182**:2702–2708.
23. Miller, J. F. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
24. Ochsner, U. A., and J. Reiser. 1995. Autoinducer-mediated regulation of rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **92**:6424–6428.
25. Passador, L., J. M. Cook, M. J. Gambello, L. Rust, and B. H. Iglewski. 1993. Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science* **260**:1127–1130.
26. Pesci, E. C., and B. H. Iglewski. 1997. The chain of command in *Pseudomonas* quorum sensing. *Trends Microbiol.* **5**:132–135.
27. Pesci, E. C., J. B. Milbank, J. P. Pearson, S. McKnight, A. S. Kende, E. P. Greenberg, and B. H. Iglewski. 1999. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **96**:11229–11234.
28. Pessi, G., and D. Haas. 2000. Transcriptional control of the hydrogen cyanide biosynthetic genes *hcnABC* by the anaerobic regulator ANR and the quorum-sensing regulators LasR and RhIR in *Pseudomonas aeruginosa*. *J. Bacteriol.* **182**:6940–6949.
29. Reimann, C., M. Beyeler, A. Latifi, H. Winteler, M. Foglino, A. Lazdunski, and D. Haas. 1997. The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer N-butyryl-homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol. Microbiol.* **24**:309–319.
30. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
31. Schweizer, H. P. 1991. Escherichia-Pseudomonas shuttle vectors derived from pUC18/19. *Gene* **97**:109–121.
32. Silhavy, T. J., M. L. Berman, and L. W. Enquist. 1984. Experiments with gene fusions. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
33. Stover, C. K., X. Q. Pham, A. L. Erwin, S. D. Mizoguchi, P. Warrenner, M. J. Hickey, F. S. Brinkman, W. O. Hufnagle, D. J. Kowalik, M. Lagrou, R. L. Garber, L. Goltry, E. Tolentino, S. Westbrook-Wadman, Y. Yuan, L. L. Brody, S. N. Coulter, K. R. Folger, A. Kas, K. Larbig, R. Lim, K. Smith, D. Spencer, G. K. Wong, Z. Wu, I. T. Paulsen, J. Reizer, M. H. Saier, R. E. W. Hancock, S. Lory, and M. V. Olson. 2000. Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* **406**:959–964.
34. Wang, J., S. Lory, R. Ramphal, and S. Jin. 1996. Isolation and characterization of *Pseudomonas aeruginosa* genes inducible by respiratory mucus derived from cystic fibrosis patients. *Mol. Microbiol.* **22**:1005–1012.
35. Wang, J., A. Mushegian, S. Lory, and S. Jin. 1996. Large-scale isolation of candidate virulence genes of *Pseudomonas aeruginosa* by in vivo selection. *Proc. Natl. Acad. Sci. USA* **93**:10434–10439.
36. Whiteley, M., K. M. Lee, and E. P. Greenberg. 1999. Identification of genes controlled by quorum sensing in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **96**:13904–13909.
37. Whiteley, M., M. R. Parsek, and E. P. Greenberg. 2000. Regulation of quorum sensing by RpoS in *Pseudomonas aeruginosa*. *J. Bacteriol.* **182**:4356–4360.