

## *Ralstonia eutropha* H16 Encodes Two and Possibly Three Intracellular Poly[D-(–)-3-Hydroxybutyrate] Depolymerase Genes

Gregory M. York,<sup>1†</sup> Joachim Lupberger,<sup>1‡</sup> Jiamin Tian,<sup>2</sup> Adam G. Lawrence,<sup>1</sup>  
JoAnne Stubbe,<sup>1,2\*</sup> and Anthony J. Sinskey<sup>1\*</sup>

Departments of Biology<sup>1</sup> and Chemistry,<sup>2</sup> Massachusetts Institute of Technology,  
Cambridge, Massachusetts 02139

Received 14 January 2003/Accepted 16 April 2003

Intracellular poly[D-(–)-3-hydroxybutyrate] (PHB) depolymerases degrade PHB granules to oligomers and monomers of 3-hydroxybutyric acid. Recently an intracellular PHB depolymerase gene (*phaZ1*) from *Ralstonia eutropha* was identified. We now report identification of candidate PHB depolymerase genes from *R. eutropha*, namely, *phaZ2* and *phaZ3*, and their characterization in vivo. *phaZ1* was used to identify two candidate depolymerase genes in the genome of *Ralstonia metallidurans*. *phaZ1* and these genes were then used to design degenerate primers. These primers and PCR methods on the *R. eutropha* genome were used to identify two new candidate depolymerase genes in *R. eutropha*: *phaZ2* and *phaZ3*. Inverse PCR methods were used to obtain the complete sequence of *phaZ3*, and library screening was used to obtain the complete sequence of *phaZ2*. *PhaZ1*, *PhaZ2*, and *PhaZ3* share ~30% sequence identity. The function of *PhaZ2* and *PhaZ3* was examined by generating *R. eutropha* H16 deletion strains ( $\Delta$ *phaZ1*,  $\Delta$ *phaZ2*,  $\Delta$ *phaZ3*,  $\Delta$ *phaZ1* $\Delta$ *phaZ2*,  $\Delta$ *phaZ1* $\Delta$ *phaZ3*,  $\Delta$ *phaZ2* $\Delta$ *phaZ3*, and  $\Delta$ *phaZ1* $\Delta$ *phaZ2* $\Delta$ *phaZ3*). These strains were analyzed for PHB production and utilization under two sets of conditions. When cells were grown in rich medium, *PhaZ1* was sufficient to account for intracellular PHB degradation. When cells that had accumulated ~80% (cell dry weight) PHB were subjected to PHB utilization conditions, *PhaZ1* and *PhaZ2* were sufficient to account for PHB degradation. *PhaZ2* is thus suggested to be an intracellular depolymerase. The role of *PhaZ3* remains to be established.

Polyhydroxyalkanoates (PHAs) are polyoxoesters produced by a wide range of bacteria when they find themselves in an environment with an available carbon source but limited in additional nutrient(s) required for growth (9). The short-chain-length PHAs, where R is a methyl or ethyl, have properties of thermoplastics and are biodegradable (Fig. 1). Much effort has focused on understanding the biology of PHA homeostasis for several reasons. First, this understanding could lead to expression of the appropriate gene set in heterologous systems to make PHA production economically competitive with oil-based polymers. Second, understanding PHA homeostasis serves as a paradigm for understanding the mechanism of homopolymerization reactions in which the product undergoes a phase transition during its formation, generating insoluble inclusions (granules). The intracellular PHAs can be degraded when the bacteria require carbon but are in otherwise nutrient-replete conditions, and the monomers and energy released can be reused to allow the bacteria to grow (17). The insoluble PHA granules must, therefore, be biosynthesized in a controlled fashion to facilitate enzymatic degradation. A variety of proteins associated with PHA homeostasis have been identified and are being characterized (3, 8, 10, 12, 20, 22). As part

of our research to understand the mechanisms that control polymer size and reuse, we have been interested in identifying the intracellular depolymerases that degrade poly[D-(–)-3-hydroxybutyrate] (PHB) within the granules. The cloning, sequencing, and characterization in vivo of two new putative intracellular PHB depolymerases from *Ralstonia eutropha* are reported.

Extracellular depolymerases, in contrast to the intracellular depolymerases, have been extensively characterized over the last decade (5). These enzymes are secreted into the environment to degrade PHA released from dead bacteria. These proteins in general contain a N-terminal signal peptide of 25 to 38 amino acids. The peptide is cleaved during passage of the protein out of the cytosol. These depolymerases contain a large N-terminal catalytic domain with a lipase box (GX SXG), a C-terminal PHB binding domain, and a linker region that connects the N- and C-terminal domains (for an excellent review, see reference 6). Several organisms contain extracellular depolymerase isozymes whose functions have not yet been clearly delineated.

The first sequence of an intracellular depolymerase was reported by Saegusa et al. (GenBank accession no. AB017612) (14) and was somewhat surprising, as the protein had no sequence similarity to the extracellular depolymerases. Recently this intracellular depolymerase, *PhaZ* (here designated *PhaZ1*), from *R. eutropha* was expressed, and its properties were examined in crude extracts. *PhaZ1* is 47 kDa and has neither a lipase box nor an identifiable PHB binding domain. The depolymerase was shown to work best on amorphous PHB as found in the granules inside the cell. Furthermore, the enzyme was shown to convert the polymer into oligomers and

\* Corresponding author. Mailing address for JoAnne Stubbe: Department of Chemistry, Bldg. 68-438, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA 02139. Phone: (617) 253-1814. Fax: (617) 258-7247. E-mail: stubbe@mit.edu. Mailing address for Anthony Sinskey: Department of Biology, Bldg. 68-370, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA 02139. Phone: (617) 253-6721. Fax: (617) 253-8550. E-mail: asinskey@mit.edu.

† Present address: USB Corporation, Cleveland, OH 44128.

‡ Present address: Feldmoos 5, D-88273 Fronreute, Germany.




FIG. 1. Thermoplastic short-chain-length PHAs are synthesized from monomers with short side chains.

monomers of hydroxybutyrate by the end of the degradation process (14). The striking differences between the sequences of the extracellular and intracellular depolymerases make the intracellular depolymerases interesting subjects for mechanistic and biological studies.

The function of PhaZ1 *in vivo* has recently been addressed by two groups independently (4, 14). Each group generated a  $\Delta$ *phaZ1* in *R. eutropha* H16 and then examined the fate of accumulated PHB under different growth conditions. The authors concluded from their studies that additional depolymerase(s) must be present to account for the observed phenotypes.

The availability of the *R. eutropha phaZ1* sequence and the *Ralstonia metallidurans* genome sequence allowed us to successfully devise a strategy to find two additional candidate depolymerase genes in *R. eutropha*, designated *phaZ2* and *phaZ3*. To understand the function of PhaZ2 and PhaZ3, a set of *R. eutropha* H16 *phaZ* gene deletion strains have been generated ( $\Delta$ *phaZ1*,  $\Delta$ *phaZ2*,  $\Delta$ *phaZ3*,  $\Delta$ *phaZ1\Delta**phaZ2*,  $\Delta$ *phaZ1\Delta**phaZ3*,  $\Delta$ *phaZ2\Delta**phaZ3*, and  $\Delta$ *phaZ1\Delta**phaZ2\Delta**phaZ3*). The effect of these *phaZ* gene deletions on PHB accumulation and utilization when different *R. eutropha* strains were grown in rich medium and in PHB utilization medium are reported. These studies reveal that PhaZ2 is an intracellular depolymerase. The function of PhaZ3 remains to be established.

## MATERIALS AND METHODS

**Strains and plasmids.** All strains and plasmids used in this study are listed in Table 1. The oligonucleotides used in this study are listed in Table 2.

**Media and growth conditions.** All *Escherichia coli* strains were grown aerobically at 37°C in Luria-Bertani (LB) medium or solid LB agar (1.2%). All *R. eutropha* strains were cultivated aerobically at 30°C in LB medium or Tryptic soy broth-dextrose free (TSB) medium (Becton Dickinson Microbiology Systems, Cockeysville, Md.). The mating of *Escherichia coli* S17-1 with *R. eutropha* was performed at 30°C on LB agar supplemented with thiamine (100 µg/liter), magnesium sulfate (1 mM), and calcium chloride (0.25 mM). The *sacB* gene selection was performed on sodium chloride-free LB agar supplemented with 5% sucrose and 0.2% fructose at 30°C. PHB production was examined in minimal medium (11) supplemented with 1% fructose and 0.01% ammonium chloride (PHB high); PHB utilization was examined in minimal medium supplemented with 0.5% ammonium chloride (PHB no carbon). Both processes were also studied in TSB rich medium. For *E. coli* strains, typical concentrations of antibiotics used were: ampicillin, 100 µg/ml; kanamycin, 25 µg/ml; or gentamicin, 10 µg/ml. For *R. eutropha* strains, the same concentrations were applied except for kanamycin, which was used at 270 µg/ml.

**Chemicals and enzymes.** All restriction endonucleases, DNA markers, and phage T4 DNA ligase (EC 6.5.1.1) were obtained from New England Biolabs, Beverly, Mass. *Thermus aquaticus* (*Taq*) DNA polymerase (EC 2.7.7.7) and shrimp alkaline phosphatase (EC 3.1.3.1) were obtained from Roche Molecular Biochemicals, Indianapolis, Ind. All antibiotics and fine chemicals were obtained from Sigma Biochemicals, St. Louis, Mo. All DNA oligonucleotides were synthesized by MWG Biotech Inc., High Point, N.C.

**DNA preparation and manipulation.** Standard approaches were used for DNA preparation and manipulation (1). Genomic DNA was extracted from *R. eutropha* with modifications previously described (23). PCR products were routinely cloned into pCR2.1-TOPO using a TOPO TA cloning kit (Invitrogen Corp., Carlsbad, Calif.).

**Design of probes to identify intracellular depolymerase genes.** Candidate PHB depolymerase genes were identified by a BLAST comparison using *phaZ1* and the recently sequenced *R. metallidurans* CH34 genome: [http://jgi.doe.gov/JGI\\_microbial/html/ralstonia/ralston\\_homepage.html](http://jgi.doe.gov/JGI_microbial/html/ralstonia/ralston_homepage.html). Analysis yielded two open reading frames (ORF), *phaZ1*<sub>CH34</sub> and *phaZ2*<sub>CH34</sub>, with high homology to *R. eutropha phaZ1*. A codon table was generated for *R. eutropha* based on the sequences of 17 *R. eutropha* H16 genes. The deduced amino acid sequences of

TABLE 1. Strains and plasmids used in this study<sup>a</sup>

Strain or plasmid	Description <sup>b</sup>	Reference or source
<i>R. eutropha</i> strains		
Ae H16	Wild-type; Gm resistant	ATCC17699
Re1097	$\Delta$ <i>phaZ1</i> strain, derived from Ae H16/pGY96	This study
Re1107	$\Delta$ <i>phaZ3</i> strain, derived from Ae H16/pPK51	This study
Re1108	$\Delta$ <i>phaZ1\Delta</i> <i>phaZ3</i> strain, derived from Re1097/pPK51	This study
Re1110	$\Delta$ <i>phaZ2</i> strain, derived from AeH16/pJOE39	This study
Re1111	$\Delta$ <i>phaZ1\Delta</i> <i>phaZ2\Delta</i> <i>phaZ3</i> strain, derived from Re1108/pJOE39	This study
Re1112	$\Delta$ <i>phaZ1\Delta</i> <i>phaZ2</i> strain, derived from Re1097/pJOE39	This study
Re1113	$\Delta$ <i>phaZ2\Delta</i> <i>phaZ3</i> strain, derived from Re1107/pJOE39	This study
<i>E. coli</i> strains		
DH5 $\alpha$	Strain for ligation, cloning, and heterologous expression of PHA genes	New England Biolabs
DH5 $\alpha$ F'	F'/endA1 hsdR17 glnV44 thi-1 recA1 gyrA relA1DU169 deoR	19
S17-1	recA pro hsdR RP4-2-Tc::Mu-Km::Tn7	ATCC47055
Plasmids		
PBluescriptIIKS	Cloning vector; <i>LacZa</i> Ap resistance	Stratagene
pCR2.1-TOPO	High-copy-number plasmid used for cloning; confers Ap and Km resistance	Invitrogen
pJQ200mp 18Km	Derivative of pJQ200mp 18; Gm resistance gene disrupted; confers Km resistance	21
pGY96	$\Delta$ <i>phaZ1</i> gene replacement plasmid; confers Km resistance	This study
pJOE39	$\Delta$ <i>phaZ2</i> gene replacement plasmid; confers Km resistance	This study
pPK51	$\Delta$ <i>phaZ3</i> gene replacement plasmid; confers Km resistance	This study
pWt-250-2	~240-bp PCR product corresponding to <i>phaZ1</i> ; cloned in pCR2.1-TOPO	This study
pZ15	~240-bp PCR product corresponding to <i>phaZ2</i> ; cloned in pCR2.1-TOPO	This study
pZ31	~240-bp PCR product corresponding to <i>phaZ3</i> ; cloned in pCR2.1-TOPO	This study

<sup>a</sup> Plasmids constructed in this study that were used only as intermediates for construction of other plasmids are described only in Materials and Methods.

<sup>b</sup> Abbreviations: Ap, ampicillin; Gm, gentamicin; Km, kanamycin.

TABLE 2. Oligonucleotides used in this study

Purpose and oligonucleotides	Sequence <sup>a</sup>	Location and orientation <sup>b</sup>
Probing intracellular depolymerases		
phaZB-3	GTSTAYRTBACXGAYTGG	Degenerate oligonucleotides (degeneracy: 192)
phaZB-4	GCRTCGATBGGVCCXSCSAT	Degenerate oligonucleotides (degeneracy: 288)
Inverse PCR		
phaZC1	AAGTGCCCGGCGTCAAGCG	5' <i>phaZ3</i> core sequence
phaZC2	GCATATCGTGGCGGTTTGCC	3' <i>phaZ3</i> core sequence
Library screening		
phaZB6	GCCGATCAGGTGCTGCACG	5' <i>phaZ2</i> core sequence
phaZB10	GCGGGATCCCAAATCCAGGTCCGGTGG	3' <i>phaZ2</i> core sequence
Generation of deletion strains		
phaZ6	CCGGAGGATCCCGCAACAGGTGGCGAC	5' end of region upstream of <i>phaZ1</i> ORF (+)
phaZ7	CGCAATCGCGGGCGTTTTTCGCCTTTTCTGCCTGGGTCTA	Fusion upstream and downstream of <i>phaZ1</i> ORF (-)
phaZ8	TAGACCCAGGCAGAAAAGGCGAAAACGCCCGCATTGCC	Fusion upstream and downstream of <i>phaZ1</i> ORF (+)
phaZ9	GCCGAGGATCCGCTGATCAACCCGGTGGTGG	3' end of region downstream of <i>phaZ1</i> ORF (-)
phaZD1b	CGTGCCAGGCATAAACTGATGGCCCCGGCAGCCGCCAGC	Fusion upstream and downstream of <i>phaZ2</i> ORF (-)
phaZD2c	AAAGGATCCCGAAGACAAAGCAAAGGGGTAG	3' end of region downstream of <i>phaZ2</i> ORF (-)
phaZD3b	GCTGGCGGCTGCCGGGGCCATCAGTTTATGCCTGGCAGC	Fusion upstream and downstream of <i>phaZ2</i> ORF (+)
phaZD4b	TTTGGATCCAGCCTTGGGGTGGATTTCATTC	5' end of region upstream of <i>phaZ2</i> ORF (+)
phaZC8	GCCAAGGCGACGGAGCGCTGCGATTCCCGCCTTTTGG	Fusion upstream and downstream of <i>phaZ3</i> ORF (-)
phaZC9	CAAAAAGCGGGGAATCGCAGCGCTCCGTCGCCTTGGC	Fusion upstream and downstream of <i>phaZ3</i> ORF (+)
phaZC10	GGCCGGATCCACTTGTATTGCAAGCTGCTCC	5' end of region upstream of <i>phaZ3</i> ORF (+)
phaZC11	GGCCGGATCCTTATTCGAGCACAAAGTGCG	3' end of region downstream of <i>phaZ3</i> ORF (-)
Deletion strain verification (additional oligonucleotides)		
phaZ2as	AATGGCATGTTGATCGTTGGTG	Upstream of <i>phaZ2</i> ORF (-)
phaZ2se	CCTCCTTACTGCTTGTGGCCG	Downstream of <i>phaZ2</i> ORF (+)

<sup>a</sup> Restriction site (*Bam*HI) engineered into sequences is indicated in bold lettering. Degenerate oligonucleotide code: B = T, C, G; R = A, G; S = C, G; V = A, G, C; X = T, C, A, G; Y = T, C.

<sup>b</sup> Forward (+) or reverse (-) orientation relative to ORF is indicated.

*phaZ1*, *phaZ1*<sub>CH34</sub>, and *phaZ2*<sub>CH34</sub> were aligned using the Clustal W algorithm (18). The alignment revealed two conserved regions of six and seven amino acids that were used for design of the degenerate oligonucleotides phaZB-3 and phaZB-4 (2). The oligonucleotides were used for amplification of ~250-bp fragments of potential PHB depolymerase gene sequences from the Re1097 ( $\Delta$ *phaZ1*) strain. Two candidate fragments with high homology to PHB depolymerase genes were identified and designated *phaZ2* and *phaZ3*.

**Construction of *phaZ1* precise deletion gene replacement plasmid pGY96.** The 294- and 295-bp fragments of *R. eutropha* DNA corresponding to the regions immediately upstream and downstream of the *phaZ1* ORF were amplified by PCR with the oligonucleotides phaZ6/phaZ7 and phaZ8/phaZ9. *Bam*HI sites were introduced into both products by phaZ6 and phaZ9, and an overlapping segment of 39 nucleotides was introduced into the two products by phaZ7 and phaZ8. The two PCR products were combined and amplified with phaZ6 and phaZ9 to yield a 0.6-kb PCR product with sequence upstream and downstream of the *phaZ1* ORF, flanked by *Bam*HI sites, and lacking the *phaZ1* gene. The PCR product was cloned into the *Afl*III site of pCR2.1-TOPO to yield an intermediate plasmid. The 0.6-kb *Bam*HI fragment of this intermediate plasmid was then cloned into the *Bam*HI site of pJQ200mp18Km to yield pGY96.

**Inverse PCR to obtain the complete *phaZ3* sequence.** Genomic DNA from *R. eutropha* RE1097 ( $\Delta$ *phaZ1*) was isolated and partially digested with *Sau*3AI (62 mU/ $\mu$ g of DNA, 80 min) in high dilution. The resulting DNA was self ligated. Inverse PCR (16) was performed on the resulting circular chromosomal fragments using the primer set phaZC1-phaZC2, designed within the known region of *phaZ3* so that the primer 3' ends faced towards the start codon and stop codon of the designated *phaZ3* ORF. The unknown sequence portions of *phaZ3* were amplified and flanked now by the inverted priming sites of phaZC1 and phaZC2. The resulting PCR products were cloned into pCR2.1-TOPO and sequenced.

**Library generation and screening to obtain the complete *phaZ2* sequence.** Inverse PCR was unsuccessful at obtaining the full sequence of *phaZ2*. A *R. eutropha* library was generated from the partially digested genomic DNA of

Re1108. The digestion reaction (*Sau*3AI: 62 mU/ $\mu$ g of DNA; 80 min) was stopped with 0.1% sodium dodecyl sulfate (wt/vol). Fragments between 1 and 10 kb were cloned into the vector pBluescriptIIKS. Successfully transformed *E. coli* DH5 $\alpha$ F' clones were cultivated and stored (10% dimethyl sulfoxide; -80°C) in 96-well tissue culture plates (VWR International Inc., Buffalo Grove, Ill.). The library was screened by PCR in a 96-well multiblock PCR cyclor (Primus 96 Plus, MWG Biotech Inc., High Point, N.C.). The presence of a 388-bp *phaZ2* fragment was detected by the primer set phaZB6-phaZB10. Specificity was verified by restriction analysis of the PCR products with *Bgl*I. Positive candidates were sequenced, and the complete sequence was assembled using Contig Express (InforMax Inc., Annapolis, Md.).

**Construction of *phaZ3* and *phaZ2* precise deletion gene replacement plasmids pPK51 and pJOE39, respectively.** The protocol was identical to that described above for pGY96. The 410- and 306-bp fragments immediately upstream and downstream of the *phaZ3* ORF were generated using PhaZC8-PhaZC10 and PhaZC9-PhaZC11. The resulting PCR product was placed in pJQ200mp18Km to produce pPK51. Fragments upstream (455-bp) and downstream (487-bp) of the *phaZ2* ORF were generated by using PhaZD3b-PhaZD4b and PhaZD1b-PhaZD2c. The resulting PCR product was used to generate pJOE39.

**Construction of depolymerase deletion strains in *R. eutropha* H16.** The deletion strains in *R. eutropha* H16 were created by homologous recombination by a standard procedure (13, 21). The *phaZ* replacement plasmids pGY96, pJOE39, and pPK51 were introduced into the appropriate *R. eutropha* strains to generate  $\Delta$ *phaZ1*,  $\Delta$ *phaZ2*,  $\Delta$ *phaZ3*,  $\Delta$ *phaZ1\Delta**phaZ2*,  $\Delta$ *phaZ1\Delta**phaZ3*,  $\Delta$ *phaZ2\Delta**phaZ3*, and  $\Delta$ *phaZ1\Delta**phaZ2\Delta**phaZ3* by conjugation from donor strain *E. coli* S17-1. Construction of the desired deletions was confirmed by PCR with the oligonucleotides in Table 2. Oligonucleotides PhaZ6-PhaZ9 were used to confirm  $\Delta$ *phaZ1* strains; PhaZas-PhaZse were used to verify  $\Delta$ *phaZ2* strains; and PhaZC10-PhaZC11 were used to confirm  $\Delta$ *phaZ3* strains. Southern blot analyses were also carried out to verify the strain construction.






FIG. 3. PHB production and utilization in TSB medium. (A) Results for wt (○),  $\Delta phaZ1$  (◇),  $\Delta phaZ2$  (□), and  $\Delta phaZ3$  (×) *R. eutropha* strains. (B) Results for wt (○),  $\Delta phaZ1\Delta phaZ2$  (◇),  $\Delta phaZ1\Delta phaZ3$  (□), and  $\Delta phaZ1\Delta phaZ2\Delta phaZ3$  (×) *R. eutropha* strains.

of these strains was examined for PHB when grown in rich medium (TSB) over 72 h and when grown in nutrient-limited medium (PHB high) for 72 h followed by transfer to PHB utilization medium (PHB no carbon) for 48 or 72 h. The results from the experiments in TSB are shown in Fig. 3A and B. They revealed a peak of PHB production between 8 and 12 h. In the wt, Re110 ( $\Delta phaZ2$ ), and Re1107 ( $\Delta phaZ3$ ) strains, all of the PHB was removed by 48 h. In all of the other strains, each of which has *phaZ1* deleted, the PHB levels remain unchanged after 12 h. At present, it is not clear if the differences in PHB observed for different strains during the first 12 h are significant. The results in TSB medium therefore provided no evidence that PhaZ2 or PhaZ3 can function as PHB depolymerases in vivo.

The results from experiments in which PHB was accumulated to ~80% cell dry weight in PHB high medium and the cells were then switched into PHB no-carbon medium are shown in Fig. 4 and Fig. 5A and B, respectively. During the 72-h cultivation period in PHB (high), the PHB accumulation pattern of each *phaZ* deletion strain was very similar to that of the wt strain, all approaching 80% cell dry weight. When each strain was switched into PHB utilization medium, the wt strain had utilized half of its PHB content after 48 h. The  $\Delta phaZ1\Delta phaZ2\Delta phaZ3$  strain, on the other hand, did not lose any PHB content within the same period, suggesting that either PhaZ2 or PhaZ3 was involved in PHB degradation. The effect of single and double deletion of the three putative genes was further examined in an effort to understand the function of the newly identified genes. The  $\Delta phaZ1$  strain utilized only ~20% of its PHB content, similar to previous results reported by Saito. The most interesting observations were that the  $\Delta phaZ1\Delta phaZ2$  strain did not lose any significant amount of PHB, while the  $\Delta phaZ2$  strain behaved like the wt strain. The result for the  $\Delta phaZ1\Delta phaZ2$  strain, in comparison to those for the  $\Delta phaZ1$  and  $\Delta phaZ2$  strains, suggested that PhaZ2 is an intracellular depolymerase and may act upon the product(s) generated by PhaZ1. The PHB degradation in  $\Delta phaZ1\Delta phaZ3$  in several experiments appeared to be less than that in the  $\Delta phaZ1$  strain. However, at present, growth conditions have not yet been found to demonstrate unambiguously the function of PhaZ3 as a depolymerase.

## DISCUSSION

An understanding of PHA homeostasis requires identification of all the proteins involved in polymer biosynthesis, degradation, and their regulation. A number of proposals concerning the role of intracellular depolymerases have been advanced, with the most recent studies suggesting that there are multiple depolymerases in *R. eutropha* and that one might be constitutively active and a second might be inducible (4, 14). A test of the proposals requires identification of all of the proteins involved in PHB degradation. We now report the identification of two new candidate PHA depolymerase genes, *phaZ2* and *phaZ3*, in *R. eutropha*. A sequence comparison (Fig. 2) reveals a high level of sequence identity. Given that PhaZ1 has been shown in vitro to degrade amorphous PHB granules to small oligomers, it is likely that PhaZ2 and PhaZ3 will possess similar activity. Thus far, no lab has purified the intracellular depolymerase to homogeneity or analyzed the products produced in the early stages of the depolymerase-catalyzed reactions. Thus differentiation between depolymerase




FIG. 4. PHB production in PHB(high) medium by wt (○),  $\Delta phaZ1$  (◇),  $\Delta phaZ1\Delta phaZ3$  (□), and  $\Delta phaZ1\Delta phaZ2$  (△) *R. eutropha* strains. The production pattern is the same in all deletion strains (additional data not shown).




FIG. 5. PHB utilization in PHB (no carbon) medium. (A) Results for wt (○),  $\Delta$ phaZ1 (◇),  $\Delta$ phaZ2 (+),  $\Delta$ phaZ1 $\Delta$ phaZ3 (□),  $\Delta$ phaZ2 $\Delta$ phaZ3 (×), and  $\Delta$ phaZ1 $\Delta$ phaZ2 (△) *R. eutropha* strains. (B) Results for wt (○),  $\Delta$ phaZ1 (◇),  $\Delta$ phaZ3 (□),  $\Delta$ phaZ1 $\Delta$ phaZ3 (×), and  $\Delta$ phaZ1 $\Delta$ phaZ2 $\Delta$ phaZ3 (△) *R. eutropha* strains. These results are from an experiment independent of that corresponding to the results shown in panel A.

functions and the requirement for isozymes requires further investigation.

To obtain evidence that PhaZ2 and PhaZ3 are actually PHB depolymerases, the role of these proteins in a variety of *R. eutropha* deletion strains has been examined under rich medium growth conditions and conditions for PHB utilization. Comparison of the results from the wt strain with those of various deletion strains after 72 h of cultivation under conditions in which PHB is utilized to provide a carbon source for growth establishes that PhaZ1 and PhaZ2 are involved in PHB degradation. PhaZ1, identified by Saito et al., was observed to have the largest effect. Our results provide the first direct evidence that PhaZ2 is an active PHB depolymerase.

Studies of the deletion strains relative to the wt strain when grown on rich TSB medium present a different picture. Under these conditions, the data exhibited a higher degree of variation between individual cultures. However, despite these variations, PhaZ1 is consistently the only protein involved in PHB degradation.

To understand the function of each of the depolymerases now requires analysis of mRNA levels, protein presence, and granule structure as a function of growth conditions. In addition, purification of the proteins and development of assays in the early stages of PHB degradation is essential to our understanding of and differentiation between the functions of the different depolymerases. These studies are in progress.

#### ACKNOWLEDGMENTS

This work was supported by NIH grant GM49171 to JoAnne Stubbe and Anthony J. Sinskey, by a DAAD grant "Kurzstipendien für Abschlussarbeiten" for Joachim Lupberger, and by NIH training grant 5T32GM08334 to Jiamin Tian.

Plasmid pPK51 was constructed by Peter Kok.

#### REFERENCES

- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1994. Current protocols in molecular biology. John Wiley & Sons, Inc., New York, N.Y.
- Compton, T. 1990. Degenerate primers for DNA amplification, p. 39-45. In M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White (ed.), PCR protocols: a guide to methods and applications. Academic Press, San Diego, Calif.
- Gerngross, T. U., K. D. Snell, O. P. Peoples, A. J. Sinskey, E. Cushai, S. Masamune, and J. Stubbe. 1994. Overexpression and purification of the soluble polyhydroxyalkanoate synthase from *Alcaligenes eutrophus*: evidence for a required posttranslational modification for catalytic activity. *Biochemistry* **33**:9311-9320.
- Handrick, R., S. Reinhardt, and D. Jendrossek. 2000. Mobilization of poly(3-hydroxybutyrate) in *Ralstonia eutropha*. *J. Bacteriol.* **182**:5916-5918.
- Jendrossek, D., and R. Handrick. 2002. Microbial degradation of polyhydroxyalkanoates. *Annu. Rev. Microbiol.* **56**:403-432.
- Jendrossek, D. 2001. Microbial degradation of polyesters. *Adv. Biochem. Eng. Biotechnol.* **71**:293-325.
- Karr, D. B., J. K. Waters, and D. W. Emerich. 1983. Analysis of poly-beta-hydroxybutyrate in *Rhizobium japonicum* bacteroids by ion-exclusion high-pressure liquid-chromatography UV detection. *Appl. Environ. Microbiol.* **46**:1339-1344.
- Liebergessel, M., K. Sonomoto, M. Madkour, F. Mayer, and A. Steinbüchel. 1994. Purification and characterization of the poly(hydroxyalkanoic acid) synthase from *Chromatium vinosum* and localization of the enzyme at the surface of poly(hydroxyalkanoic acid) granules. *Eur. J. Biochem.* **226**:71-80.
- Madison, L. L., and G. W. Huisman. 1999. Metabolic engineering of poly(3-hydroxyalkanoates): from DNA to plastic. *Microbiol. Mol. Biol. Rev.* **63**:21-53.
- Maehara, A., Y. Doi, T. Nishiyama, Y. Takagi, S. Ueda, H. Nakano, and T. Yamane. 2001. PhaR, a protein of unknown function conserved among short-chain-length polyhydroxyalkanoic acids producing bacteria, is a DNA-binding protein and represses *Paracoccus denitrificans* phaP expression *in vitro*. *FEMS Microbiol. Lett.* **200**:9-15.
- Peoples, O. P., and A. J. Sinskey. 1989. Poly- $\beta$ -hydroxybutyrate (PHB) biosynthesis in *Alcaligenes eutrophus* H16: identification and characterization of the PHB polymerase gene (*phbC*). *J. Biol. Chem.* **264**:15298-15303.
- Pieper-Fürst, U., M. H. Madkour, F. Mayer, and A. Steinbüchel. 1994. Purification and characterization of a 14-kilodalton protein that is bound to the surface of polyhydroxyalkanoic acid granules in *Rhodococcus ruber*. *J. Bacteriol.* **176**:4328-4337.
- Quandt, J., and M. F. Hynes. 1993. Versatile suicide vectors which allow direct selection for gene replacement in gram-negative bacteria. *Gene* **127**:15-21.
- Saegusa, H., M. Shiraki, C. Kanai, and T. Saito. 2001. Cloning of an intracellular poly[D(-)-3-hydroxybutyrate] depolymerase gene from *Ralstonia eutropha* H16 and characterization of the gene product. *J. Bacteriol.* **183**:94-100.
- Saito, T., K. Takizawa, and H. Saegusa. 1995. Intracellular poly(3-hydroxybutyrate) depolymerase in *Alcaligenes eutrophus*. *Can. J. Microbiol.* **41**(suppl. 1):187-191.
- Silver, J. 1991. Inverse polymerase chain reaction, p. 137-146. In M. J. McPherson, P. Quirke, and G. R. Taylor (ed.), PCR: a practical approach. Oxford University Press, New York, N.Y..
- Steinbüchel, A., and S. Hein. 2001. Biochemical and molecular basis of microbial synthesis of polyhydroxyalkanoates in microorganisms. *Adv. Biochem. Eng. Biotechnol.* **71**:81-123.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. Clustal W: improving the sensitivity of progressive multiple sequence alignment through se-

- quence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
19. **Whittaker, P. A., A. J. Campbell, E. M. Southern, and N. E. Murray.** 1988. Enhanced recovery and restriction mapping of DNA fragments cloned in a new lambda vector. *Nucleic Acids Res.* **16**:6725–6736.
  20. **Wieczorek, R., A. Pries, A. Steinbüchel, and F. Mayer.** 1995. Analysis of a 24-kilodalton protein associated with the polyhydroxyalkanoic acid granules in *Alcaligenes eutrophus*. *J. Bacteriol.* **177**:2425–2435.
  21. **York, G. M., J. Stubbe, and A. J. Sinskey.** 2001. New insight into the role of the PhaP phasin of *Ralstonia eutropha* in polyhydroxybutyrate production. *J. Bacteriol.* **183**:2394–2397.
  22. **York, G. M., J. Stubbe, and A. J. Sinskey.** 2002. The *Ralstonia eutropha* PhaR protein couples synthesis of the PhaP phasin to the presence of polyhydroxybutyrate in cells and promotes polyhydroxybutyrate production. *J. Bacteriol.* **184**:59–66.
  23. **York, G. M., B. H. Junker, J. A. Stubbe, and A. J. Sinskey.** 2001. Accumulation of the PhaP phasin of *Ralstonia eutropha* is dependent on production of polyhydroxybutyrate in cells. *J. Bacteriol.* **183**:4217–4226.