

GacA Regulates Symbiotic Colonization Traits of *Vibrio fischeri* and Facilitates a Beneficial Association with an Animal Host

Cheryl A. Whistler and Edward G. Ruby*

Pacific Biomedical Research Center, University of Hawaii, Honolulu, Hawaii 96813

Received 3 July 2003/Accepted 23 September 2003

The GacS/GacA two-component system regulates the expression of bacterial traits during host association. Although the importance of GacS/GacA as a regulator of virulence is well established, its role in benign associations is not clear, as mutations in either the *gacS* or *gacA* gene have little impact on the success of colonization in nonpathogenic associations studied thus far. Using as a model the symbiotic association of the bioluminescent marine bacterium *Vibrio fischeri* with its animal host, the Hawaiian bobtail squid, *Euprymna scolopes*, we investigated the role of GacA in this beneficial animal-microbe interaction. When grown in culture, *gacA* mutants were defective in several traits important for symbiosis, including luminescence, growth in defined media, growth yield, siderophore activity, and motility. However, *gacA* mutants were not deficient in production of acylated homoserine lactone signals or catalase activity. The ability of the *gacA* mutants to initiate squid colonization was impaired but not abolished, and they reached lower-than-wild-type population densities within the host light organ. In contrast to their dark phenotype in culture, *gacA* mutants that reached population densities above the luminescence detection limit had normal levels of luminescence per bacterial cell in squid light organs, indicating that GacA is not required for light production within the host. The *gacA* mutants were impaired at competitive colonization and could only successfully cocolonize squid light organs when present in the seawater at higher inoculum densities than wild-type bacteria. Although severely impaired during colonization initiation, *gacA* mutants were not displaced by the wild-type strain in light organs that were colonized with both strains. This study establishes the role of GacA as a regulator of a beneficial animal-microbe association and indicates that GacA regulates utilization of growth substrates as well as other colonization traits.

During colonization of animal or plant tissue, bacteria must adapt to the requirements of these environments and prevail over host defenses. There is great interest in understanding how signaling between beneficial bacteria and their hosts is initiated and a stable association is permitted, while at the same time detrimental pathogens are detected and infection is prevented. Although long-term, benign bacterial associations with animals are ubiquitous, studies of these associations are often confounded by the complexity of consortial populations and our inability to culture obligate symbionts. Elucidating the mechanisms underlying recognition and persistence by bacteria in beneficial associations with animal hosts will both add to our understanding of health and aid in the successful treatment of disease.

Vibrio fischeri is a bioluminescent marine bacterium that forms long-term, beneficial associations with certain fishes and sepiolid squid, such as the Hawaiian bobtail squid, *Euprymna scolopes*. In this association, newly hatched squid acquire *V. fischeri* from the surrounding seawater in which they are present at a few hundred CFU per ml (30) in a total background of about 10^6 other marine bacteria per ml. Only *V. fischeri* colonizes the nascent light-emitting organs of the juvenile squid, forming an essentially monospecific culture (44). The specificity of the association suggests that specialized col-

onization mechanisms in the bacterial symbiont have co-evolved with cognate recognition mechanisms in the squid host (53).

Squid colonization is both spatially and temporally dynamic. Only motile *V. fischeri* cells can migrate specifically through ducts before they reach the crypt spaces of the light organ (18). During colonization, bacterial symbionts that reach the crypt grow to a population density of 10^5 to 10^6 CFU, using as growth substrates host-derived nutrients, including small peptides (20). The increase in population size allows the density-dependent induction of luminescence gene (*lux*) expression via the accumulation of acylated homoserine lactone (acyl-HSL) quorum-sensing molecules (7). Two different acyl-HSL signals, *N*-(3-oxohexanoyl) homoserine lactone (C_6 -HSL) and *N*-octanoyl homoserine lactone (C_8 -HSL), work in concert to activate the *lux* operon (32), which contains both the structural genes for luciferase and the aldehyde synthetase genes (34). Although the squid host expels an estimated 95% of the bacterial contents of its light organ daily (29), creating a new level of selective pressure, regrowth of the remaining bacterial cells results in their persistent association with the host. Mutants defective or reduced in luminescence effectively initiate colonization and grow to a normal cell density but are impaired at longer-term host association during these subsequent regrowth periods (32, 52).

Bacteria often coordinately express multiple traits that are generally important for host association, including motility, attachment, and stress defense, together with other traits that are important to their interaction with specific hosts or host

* Corresponding author. Mailing address: Pacific Biomedical Research Center, University of Hawaii, 41 Ahui St., Honolulu, HI 96813. Phone: (808) 539-7309. Fax: (808) 599-4817. E-mail: eruby@hawaii.edu.

tissues, such as the production and secretion of effectors, secondary metabolites, or virulence determinants. For instance, in the genus *Pseudomonas*, both host association traits and virulence are globally controlled by a two-component regulatory system composed of the sensor kinase GacS and the response regulator GacA (22). Early studies with plant-pathogenic *Pseudomonas* spp. revealed that GacS/GacA controls the production of exoenzymes and is required for virulence in a number of host-microbe systems (22). In the opportunistic pathogen *Pseudomonas aeruginosa*, the production of acyl-HSL signal molecules and factors necessary for virulence on both plant and animal hosts are GacA controlled (22). More recent studies have also linked the GacA homologs ExpA, SirA, UvrY, VarA, and LetA, respectively, to (i) the regulation of extracellular enzymes and acyl-HSL signals in *Erwinia carotovora*, (ii) motility and invasion gene expression in *Salmonella* spp., (iii) colonization and stress response of *Escherichia coli*, (iv) colonization and virulence of *Vibrio cholerae*, and (v) motility, transmission, and stress response in *Legionella pneumophila* (21, 22, 40, 49).

The GacS/GacA system also regulates the expression of traits during benign host association. For example, in biocontrol *Pseudomonas* spp., GacA controls the production of anti-fungal secondary metabolites that contribute to the health of their host plants, although the production of these compounds has little impact on host-microbe association (22). Additionally, GacS/GacA mutants of these biocontrol bacteria have an enhanced fluorescence typical of overproduction of fluorescent siderophore compounds that function in iron sequestration and bacterial competition (9–11, 55). Thus, GacS/GacA regulation of host association in these superficial and nonspecific benign associations appears to function indirectly and predominantly influences microbe-microbe interactions.

We cloned the *gacA* gene from *V. fischeri* and investigated its role in global regulation of symbiotic colonization of the squid *E. scolopes* to establish what role GacA plays in colonization in this intimate and specific bacterium-host association. Our studies showed that GacA controlled multiple traits important for successful squid colonization, including motility, growth-substrate utilization, and luminescence. However, the effect of *gacA* on luminescence was not accomplished through a deficiency in acyl-HSL accumulation but instead involved depression of luminescence via an undefined mechanism. Animal studies with the *gacA* mutant alone, or in competition with the wild-type strain, revealed that GacA facilitates but is not required for host colonization. Although GacA was required for luminescence in culture, *gacA* strain-colonized squid became luminous, demonstrating that GacA effects differed in the host and in culture. The results of this study indicate that GacA is a global regulator of colonization traits and that mutations result in defects in symbiotic colonization, most notably during colonization initiation and growth within the light organ.

MATERIALS AND METHODS

Bacterial strains, plasmids, and culture conditions. Bacteria and plasmids are listed in Table 1. Chemicals used for culture media preparation were from Sigma (St. Louis, Mo.) unless otherwise noted. Wild-type *V. fischeri* strain ES114, isolated from an adult specimen of *E. scolopes* (6), and its derivatives were routinely grown at 28°C in either liquid seawater-tryptone broth (SWT) (6) with shaking at 200 rpm or on nutrient agar medium with added salt (LBS) (18). *V.*

fischeri was also grown on minimal agar plates supplemented with either 24 mM ribose (20), 50 mM fumarate, or 0.5% Casamino Acids (CA; Difco) as a carbon source. *E. coli* strains were routinely grown in Luria-Bertani (LB) broth (45) or in brain heart infusion medium (Difco). When required, media were supplemented with antibiotics at the following concentrations: for *V. fischeri*, kanamycin at 50 µg/ml, chloramphenicol (Ch) at 5 µg/ml for multiple copies of the resistance gene in plasmids and at 2.5 µg/ml for a single copy on the chromosome, and erythromycin at 5 µg/ml; for *E. coli*, kanamycin at 50 µg/ml, Ch at 25 µg/ml, ampicillin at 100 µg/ml, and erythromycin at 150 µg/ml. Plates were supplemented with 40 mg of 5-bromo-4-chloro-3-indolyl-β-galactopyranoside/ml for visualization of β-galactosidase activity. Where appropriate, C₆-HSL and C₈-HSL (Aurora Biosciences, San Diego, Calif.) were added to media at 120 nM before inoculation with bacteria as previously described (32). Conditioned broth was prepared by combining fresh broth with spent broth at a ratio of 1:1. Spent broth was prepared by growing bacterial cultures to a final optical density at 600 nm (OD₆₀₀) of 1.8, pelleting cells by centrifugation at 12,000 × g, and filter sterilizing the cleared broth by passage through a 0.2-µm-pore-sized filter.

Recombinant DNA techniques. Standard molecular methods were used for transformations, restriction enzyme digestions, gel electrophoresis, Southern analysis, and PCR (45). Restriction enzymes were from New England BioLabs (Beverly, Mass.). Gel purification of restriction enzyme-digested DNA was performed using the QiaQuick gel extraction kit (Qiagen, Valencia, Calif.). Plasmids for laboratory procedures were purified using the Qiaprep Spin Miniprep kit (Qiagen). Plasmid DNA for sequence analysis was prepared using the Perfect Prep plasmid mini kit (Eppendorf Scientific Inc., Westbury, N.Y.). Ligations were performed by the thermal cycling method (31). Genomic DNA was isolated by a cetyltrimethylammonium bromide method (3). Digoxigenin-11-dUTP-labeled probes for Southern blotting were generated by PCR using materials and protocols supplied by the manufacturer (Boehringer Mannheim Corporation, Indianapolis, Ind.). Oligonucleotide primers were synthesized by Integrated DNA Technologies (Coralville, Iowa).

Cloning and sequence analysis of *gacA*. Alignments of the predicted amino acid sequence from various GacA homologs were used to identify conserved regions to which degenerate primers were designed. The forward (5'-GARGC NGAYWSNNGGNGARGA-3') and reverse (5'-NARYTTYTCNGTRTCNARD ATNCC-3') primers were generated and used to amplify a 550-bp fragment from the genome of *V. fischeri* strain ES114 by PCR. After confirmation that the sequence identity was similar to that of the GacA homolog VarA from *V. cholerae* (56) by directly sequencing the PCR product, gene-specific primers (forward primer Vgac2, ATGAGTTAATTCAACGTCTCAC; reverse primer Vgac3, TTATGGATATGAACATGCCTGG) were designed. These primers were used to amplify an internal fragment of GacA (525 bp in length) which served as a gene-specific probe to identify and isolate a genomic clone containing the intact gene. Southern analysis of genomic fragments from *V. fischeri* generated by digestion with various restriction enzymes revealed that the gene was contained on a 4.0-kb *Hind*III fragment. A genomic sublibrary of 4.0-kb *Hind*III-digested fragments of ES114 was generated by gel purification of the restriction fragments and subsequent ligation of ES114 DNA to *Hind*III-digested pEV579 plasmid DNA. The library DNA was transformed into *E. coli* cells which were plated on LB with Ch and 5-bromo-4-chloro-3-indolyl-β-galactopyranosidase. White colonies containing plasmids with inserted DNA were directly screened by PCR using the Vgac2 and Vgac3 primers. From a single clone, pVCW1A7, we amplified the predicted 525-bp fragment. Sequencing of the entire *gacA* gene was performed at the Biotechnology Molecular Biology Instrumentation Facility, University of Hawaii, using primers designed to the sequence obtained from the PCR-amplified fragment. Comparisons of the cloned DNA with sequences in GenBank were performed using the BLAST software package (2).

Generation of mutants. Plasmid pVCW1A7 that contained the wild-type *gacA* region was mutagenized using the EZ::TN<KAN-2> insertion kit following protocols supplied by the manufacturer (Epicentre, Madison, Wis.). Insertions in *gacA* were identified by PCR amplification and confirmed to be in the *gacA* open reading frame (ORF) by sequence analysis of the clones. One random EZ::TN<KAN-2> insertion was identified within the sequence encoding amino acid residue P58 of GacA. The mutagenized gene was recombined with the chromosomal copy of *gacA* by marker exchange mutagenesis as described previously (47). A single colony, designated strain VCW2A1, was confirmed to have the *gacA*::EZ::TN<KAN-2> mutation by Southern analysis. A random EZ::TN<KAN-2> insertion in the *uvrC* gene was generated by a similar approach, creating strain VCW2E1.

To make an in-frame deletion of *gacA*, we created an *Aat*II restriction site within the *gacA* ORF. A single nucleotide conversion from T to C at position 600 in the nucleotide sequence was generated using the QuikChange site-directed mutagenesis kit and protocols supplied by the manufacturer (Stratagene, La

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Description ^a	Source or reference
<i>E. coli</i> strains		
DH5 α	F ⁻ <i>recA1 endA1 hsdR17 supE44 thi-1 gyrA96 relA1</i> Δ (<i>argF-lacZYA</i>) U169 ϕ 80 <i>lacZ</i> Δ M15 λ ⁻	Gibco-BRL, Inc.
CC118 λ <i>pir</i>	Δ (<i>arg-leu</i>) <i>araD</i> Δ <i>lacX74 galE galK phoA20 thi-1 rpsE rpoB argE</i> (Am) <i>recA1</i> , lysogenized with <i>lambda pir dam dcm</i>	24
<i>V. fischeri</i> strains		
ES114	Wild-type <i>E. scolopes</i> light organ isolate	6
KV433	ES114 Rf ⁺ derivative ESR1; <i>katA::erm</i>	54
SP301	ES114 Rf ⁺ derivative ESR1; <i>glnD::mini-Tn5Cm</i>	19
DM66	Spontaneously arising, hyperflagellated, hyperswimmer variant of ES114	36
KV495	ES114 Rf ⁺ derivative ESR1; <i>iucA::mini-Tn10</i>	50
JRM200	Ch ⁺ derivative of ES114	33
VCW2A1	ES114 derivative with a random EZ::TN<KAN-2> insertion in the <i>gacA</i> gene disrupting the codon for P58; Km ^r ; <i>gacA::EZ::TN<KAN-2></i>	This study
VCW2E1	ES114 derivative with a random EZ::TN<KAN-2> insertion in <i>uvrC</i> ; Km ^r ; <i>uvrC</i>	This study
VCW2F5	ES114 derivative with an internal in-frame <i>AatII</i> deletion resulting in loss of coding region between I27 and G194; Δ <i>gacA</i>	This study
VCW2H7	ES114 derivative with a frameshift mutation in <i>luxI</i> ; C6-HSL ⁻ ; <i>luxI</i>	32
Plasmids		
pVO8	pACYC, Ch ⁺ , Em ^r ; multicopy vector stably maintained by <i>V. fischeri</i>	51
pEVS79	pBCSK derivative, Mob ⁺ , Ch ⁺	47
pEVS104	R6K γ derivative of pRK2013; Δ ColE1, <i>oriT tra trb</i> Δ Tn903 Km ^r	47
pHV200I ⁻	Entire <i>lux</i> operon from <i>V. fischeri</i> ES114 with a frameshift mutation in the <i>luxI</i> gene cloned into pBR322; Ap ^r	39
p395B	<i>aidA::lacZ</i> fusion; Sp ^r Tc ^r	16
pVCW1A7	pEVS79, containing a 4.0-kb <i>HindIII</i> genomic insert from <i>V. fischeri</i> strain ES114 including the intact <i>gacA</i> gene and partial <i>uvrC</i> gene; Ch ⁺	This study
pVCW1C7	Random EZ::TN<KAN> insertion in the <i>gacA</i> gene in pVCW1A7; Ch ⁺ Km ^r	This study
pVCW1D7	Random EZ::TN<KAN> insertion in the <i>uvrC</i> gene in pVCW1A7; Ch ⁺ Km ^r	This study
pVCW2A5	Plasmid pVCW1A7 with a single nucleotide substitution from T to C, resulting in the generation of an <i>AatII</i> restriction site; Ch ⁺	This study
pVCW2D5	Plasmid pVCW2A5 with a 519-bp <i>AatII</i> deletion within the <i>gacA</i> gene; Ch ⁺	This study
pVCW2A6	Plasmid pEVS79 with an 8-kb <i>SalI</i> fragment from KV29 containing frameshifted <i>luxI</i> ; Ch ⁺	32
pVCW3C3	Plasmid pVO8 with a 4.0-kb fragment containing wild-type <i>gacA</i> from pVCW1A7; Em ^r	This study

^a Ap, ampicillin; Ch, chloramphenicol; Em, erythromycin; Km, kanamycin; Rf, rifampin; Sp, spectinomycin; Tc, tetracycline.

Jolla, Calif.). Two overlapping primers, VGA-AATF (3'-AAGTGGAGACGTC GAATTAACATCATCTAGCTATTCGTC-5') and VGA-AATR (5'-ATCAGCA ATAGCTAGATGAGTTAATTCGACGTCCTCCAC-3'), were used for mutagenesis. An in-frame deletion of 89% of the *gacA* ORF was generated within the predicted protein sequence from amino acid residue I27 to G194 by digesting the resulting plasmid, pVCW2A5, with *AatII* and self-ligating it, creating plasmid pVCW2D5. The mutation was exchanged with the wild-type *gacA* gene by marker-exchange mutagenesis, creating strain VCW2F5, and gene replacement was confirmed by Southern analysis.

A *LuxI* mutation in strain ES114 constructed for these studies was generated similarly to that mutation in *V. fischeri* strain ESR1 (52) and was described in a previous publication (32).

Luminescence detection. A 1-ml aliquot of *V. fischeri* cells grown in broth culture was removed at regular intervals to determine the luminescence and optical density (OD₆₀₀). Luminescence levels were measured with a Turner 20/20 luminometer (Turner Designs, Sunnyvale, Calif.) calibrated with a light standard. Where appropriate, decanal suspended in 95% ethyl alcohol was added at a final concentration of 0.01% to an aliquot of culture immediately prior to measurement of luminescence (32).

Luminescence of *V. fischeri* within colonized squid was routinely measured at 24, 48, and 72 h postinoculation. The luminescence detection limit was determined on squid monitored continuously between 7 and 9 h postinoculation with the wild-type strain ES114. Squid with a luminescence level between 1 and 5 luminescence units (LU) were immediately frozen, and the number of bacteria contributing to luminescence was quantified subsequently by homogenization of the squid in seawater (SW) that was formerly sterilized by autoclaving, plating the contents onto LBS agar plates, and enumerating the colonies of *V. fischeri* that grew following overnight incubation. The experiment was done twice with

similar results, and the data from both experiments were combined and reported as the mean CFU.

Quantification of acyl-HSLs. Published methods for the purification and quantification of C₆-HSL and C₈-HSL were used (46). Briefly, acyl-HSLs were extracted twice with an equal volume of acidified ethyl acetate from cell-free supernatants of ES114 and derivative cultures grown in SWT broth to a final OD₆₀₀ of 1.6. The samples were concentrated by evaporation before analysis. Because C₈-HSL is produced at a higher level (micromolar) than C₆-HSL (nanomolar) in strain ES114 (A. Schaefer, personal communication), C₆-HSL cross-reaction with the biological reporter strain for C₈-HSL does not interfere with quantification of C₈-HSL; therefore, C₈-HSL was directly quantified from extracts of 15-ml cultures. However, C₈-HSL cross-reaction with the biological reporter strains for C₆-HSL could interfere with its quantification. Therefore, we first purified C₆-HSL from 500-ml cultures using a C₁₈ reverse-phase high-performance liquid chromatography column and a linear, 10-to-100% (vol/vol) methanol-water gradient at 0.5 ml/min. The elution profile of synthetic C₆-HSL was determined to identify which fractions contained activity, and these and flanking 1-ml fractions from extracts were assayed for activity. C₆-HSL was quantified using the reporter strain *E. coli* VJS533 harboring plasmid pHV200I⁻, which does not produce its own acyl-HSL but which responds to C₆-HSL by producing luminescence (39). C₈-HSL was quantified using the reporter strain *Ralstonia solanacearum* AW1-A18 harboring plasmid p395B, which expresses *lacZ* in response to exogenous C₈-HSL (16). *LacZ* activity was measured by a standard assay (35). The amounts of C₆-HSL and C₈-HSL were determined by comparing the activity measured from a dilution series of the extracted and purified samples to the linear range of each standard curve.

Siderophore and catalase activities. The ability of mutant strains to produce siderophores was assessed qualitatively with chrom-azurolo S indicator (CAS

agar plates and compared to that of ES114, which produces an orange halo around bacterial colonies indicative of sequestration of the iron from CAS, and two strains defective in siderophore production, KV495 and SP301, which produce no halo indicative of the absence of siderophore secretion or activity, as negative controls. The CAS was added to artificial seawater medium (6) supplemented with 0.3% CA and buffered with piperazine-*N,N'*-bis(ethanesulfonic acid) (pH 6.8) as previously described (19, 28) to make CAS agar plates.

Published methods for quantification of catalase activity were used without modifications, using a strain that is defective in catalase production, KV433, as a control (4, 54). Protein concentrations were determined using the Bio-Rad DC protein assay kit with protocols supplied by the manufacturer (Bio-Rad, Hercules, Calif.). The experiment was repeated with similar results.

Motility, flagellation, and chemotaxis. Motility of exponentially growing (OD_{600} of 0.2 to 0.4) cells of *V. fischeri* in liquid cultures was assessed by light microscopy. Flagella were examined and the number of flagella per cell was determined from a total of 75 cells per treatment from three separate experiments by transmission electron microscopy (36). The data were combined and reported as the mean number of flagella \pm the standard error (SE).

Swimming motility in soft agar was determined using SWT containing concentrations between 0.3 and 0.7% of Bacto Agar (Difco, Detroit, Mich.). At these agar concentrations, the polarly flagellated *V. fischeri* ES114 is presumed to swim, as the pattern of movement is not typical of the swarming motility seen for peritrichously flagellated *Vibrio* spp. (48). Three microliters of an exponentially growing culture (OD_{600} of 0.4) was spotted on the surface of duplicate agar plates, and the movement of the cells in the agar as a concentric circle away from the spotted culture was periodically measured at the leading edge. The ability of strains to move toward attractants was assessed by spotting 10 μ l of an exponentially growing culture (OD_{600} of 0.4) on soft agar plates made with 0.25% Bacto Agar in 70% artificial seawater, 1% tryptone (Difco) with or without the addition of 0.5% CA or 1.6 mM serine and observing the bands of cells migrating up a concentration gradient created by the degradation of each band's attractant (13). The experiments were repeated with similar results, and the data from one representative experiment are reported.

Animal colonization. The ability of *V. fischeri* strains to colonize juvenile *E. scolopes* squid was determined as previously described (36, 44) with the following modifications. Exponentially growing bacteria (OD_{600} between 0.2 and 0.4) from cultures grown with shaking at 200 rpm were suspended in a volume of between 50 and 250 ml of filtered-sterilized seawater (FSW) at a final concentration of between 110 and 20,000 CFU/ml. Squid were placed collectively into bowls in a volume that allowed a minimum of 2 ml of SW/squid for either 3 h or overnight and then transferred to fresh FSW before being placed in individual vials containing 4 ml of FSW. Each morning, squid were aseptically transferred to fresh vials containing 4 ml of FSW. Colonization of squid light organs based on bacterial cell counts recovered from squid was routinely assessed at 24, 48, and 72 h postinoculation by rinsing squid in FSW and then freezing animals at -70°C before homogenizing, serially diluting, and plating the homogenate on LBS agar plates to determine the number of CFU of *V. fischeri*/light organ. Aposymbiotic animals placed in SW without bacteria and otherwise treated identically were also plated to confirm the absence of contaminating *V. fischeri* bacteria. These experiments were repeated a minimum of two times with both strains VCW2A1 and VCW2F5, which were comparable to each other, and one representative experiment with VCW2F5 was reported.

The ability of bacteria expelled from luminous squid to colonize previously uncolonized squid was also determined. Previously uncolonized juveniles were placed in vials containing serial dilutions of FSW that contained bacteria expelled from luminous VCW2F5- or ES114-colonized animals. The number of *V. fischeri* cells in the SW could not be determined by direct plating due to a high background of other bacteria. After 3 h, squid were placed in fresh vials and luminescence and colonization were determined at 30 h postcolonization.

Cocolonization experiments were performed by placing squid overnight in FSW containing both VCW2F5 and a wild-type ES114 derivative, JRM200 (33), containing a Ch resistance gene inserted in the genome in single copy, at various concentrations. In cocolonization experiments, the identity of the light organ symbionts plated on LBS agar was assessed by replica plating colonies onto LBS agar containing antibiotic (Ch) selection and by visual assessment of colony morphology.

The ability to complement the colonization defects of VCW2F5 was tested with the *gacA*-containing plasmid, pVCW3C3, or with a vector control, pVO8. Squid inoculated with bacteria at 3,000 CFU/ml of SW were subsequently maintained in FSW containing Ch (2 $\mu\text{g}/\text{ml}$) to select for the plasmids. At 24 h postinoculation, luminescence and colonization levels were determined.

Nucleotide sequence accession number. The nucleotide sequence of the *gacA* gene from *V. fischeri* along with flanking DNA has been submitted to the GenBank databases under accession number AY377390.

RESULTS

Cloning, characterization, and mutagenesis of the *gacA* gene in *V. fischeri*. Amplification of ES114 (wild-type) genomic DNA using fully degenerate primers to various *gacA* homologs (see Materials and Methods) generated a PCR product with high sequence similarity to the genes encoding GacA homologs. Subsequently, a genomic clone containing the intact *gacA* gene was isolated and sequenced. Sequence analysis of the clone revealed an ORF that was 642 bp in length, encoding a predicted protein of 214 amino acids with the alternative start codon GTG. The predicted sequence of the protein was 85% identical to VarA from *V. cholerae* (56). Within the amino-terminal receiver domain, between amino acids 1 and 123, we identified the putative phosphate-accepting aspartate residue (D54) involved in phosphorelay and, located in the carboxy-terminal region of the protein, between amino acids 146 and 203, was a conserved helix-turn-helix domain. A partial ORF 276 bp downstream of the *gacA* ORF, also beginning with GTG, was homologous to UvrC. The *uvrC* gene is also located downstream of the *gacA* genes in several other bacterial species.

Three mutants were generated to study the role of GacA in *V. fischeri* (Table 1). Two *gacA* mutants included a marked-insertion mutant strain, VCW2A1 (*gacA*::EZ::TN<KAN-2>), and an unmarked, in-frame deletion mutant strain, VCW2F5 (Δ *gacA*). In other bacterial species, *gacA* and the downstream gene *uvrC* are cotranscribed; thus, insertions can cause polar loss of UvrC. To control for potential polar effects of the insertion on UvrC, a third mutant harboring an insertion in the *uvrC* gene was generated in strain VCW2E1 (*uvrC*). When grown on LBS agar plates, the two *gacA* mutants had colonies that were smaller, less yellow, and had a translucent morphology when compared to the wild-type strain; however, the *uvrC* mutant strain colony morphology was indistinguishable from that of the wild type, suggesting colony morphology was affected by GacA and not polar loss of UvrC.

The *gacA* mutants have a growth yield defect that is relieved by the addition of Casamino Acids. The small colony size of *gacA* mutants led us to evaluate whether growth was influenced by GacA. The exponential growth rates of *V. fischeri* mutant and wild-type strains in SWT broth did not differ (Fig. 1). However, the *gacA* mutants had a growth yield defect (Fig. 1) and reached a lower final cell density (OD_{600} of 1.8) compared to the wild type (OD_{600} of >5.0). No growth yield defect was observed in the *uvrC* mutant, which reached a final cell density similar to the wild type. When cultured on minimal agar plates with either ribose or fumarate as the sole carbon source, the Δ *gacA* mutant did not grow, although the wild type grew on these media, indicating that the Δ *gacA* mutant was unable to adapt to the metabolic requirements of prototrophic growth on these sole carbon compounds. The addition of 0.5% CA, which can serve as a source of nitrogen, carbon, and vitamins, improved but did not restore the growth of the mutant to the level seen for the wild type, indicating that an amino acid auxotrophy alone could not account for the growth defect.




FIG. 1. Growth of wild-type *V. fischeri* and derivatives in culture. The optical densities (OD₆₀₀) of wild-type (■), *luxI* (▲), and $\Delta gacA$ (●) cultures grown in SWT were determined throughout the growth cycle. One representative experiment is presented.

To assess whether the growth yield defect of the $\Delta gacA$ mutant was caused by either insufficient nutrients or accumulation of repressive compounds in the supernatant, we measured the growth rate and final growth yield of the wild type and the $\Delta gacA$ mutant in complex media of different composition. The media included (i) SWT diluted to different extents with 70% SW, (ii) SWT supplemented with CA, or (iii) SWT conditioned with an equal volume of cell-free supernatants from either the wild-type or $\Delta gacA$ strain or both strains grown to a final OD₆₀₀ of 1.8 (Fig. 2). In all media tested, the $\Delta gacA$ mutant attained the same exponential growth rate as the wild type (data not shown), but it reached a lower cell density than the wild type (Fig. 2). Although the addition of 0.5% CA (the same amount that improved but did not restore growth of the mutant in minimal medium) to SWT did not substantially improve the growth of either strain, the addition of 2.5% CA

to SWT increased the yield of both strains. Additionally, the $\Delta gacA$ mutant reached the same final cell density in 0.5× SWT as it did in SWT conditioned with $\Delta gacA$ broth. This cell density was lower than the final cell density that the mutant reached in broth conditioned with either a mixture of both wild type and $\Delta gacA$ supernatants at a 1:1 ratio or wild type alone. In contrast, the wild type reached the same cell density in all conditioned broths. These data support the hypothesis that the growth yield defect of the mutant was caused by a limitation of growth substrates rather than the generation of growth-restrictive compounds by the $\Delta gacA$ mutant. The growth yield of the mutant was fully restored by carrying the *gacA* gene in *trans* on pVCW3C3 (data not shown).

The *gacA* mutants have a luminescence defect that is not complemented by the addition of acyl-HSLs or aldehyde. Luminescence was not detected from broth cultures either of the *gacA*::EZ::TN<KAN-2> mutant (data not shown) or of the $\Delta gacA$ mutant (Fig. 3) unless the *gacA* gene was restored in *trans* on pVCW3C3 (data not shown). In contrast, luminescence was detected from both the *uvrC* mutant (data not shown) and the *luxI* mutant (Fig. 3), which harbors a mutation in the C₆-HSL synthetase and therefore is defective in the production of one of the two activating signals of the *lux* biosynthetic operon. Although the *luxI* mutant produces less luminescence than the wild type, the *luxI* mutant was significantly more luminous than the $\Delta gacA$ mutant. This finding implies that GacA did not simply affect LuxI activity, but instead influenced luminescence by another mechanism. One hypothesis is that the *gacA* mutation affected C₈-HSL production, as mutants in its synthetase, AinS, produce no luminescence in culture (32).

To determine whether the $\Delta gacA$ mutant was defective in the synthesis of C₈-HSL, which is produced at a relatively high level in culture by the wild type (32), we tested the ability of excess amounts of either acyl-HSL or decanal, a substrate of the luciferase reaction that is limiting in culture, to complement the mutant's luminescence defect. Exponentially growing




FIG. 2. Growth yield of wild-type *V. fischeri* (black bars) and the $\Delta gacA$ mutant (gray bars) in various diluted and amended complex media (SWT) after 18 h of incubation. Conditioned broth was prepared by mixing SWT with cell-free supernatants of either the wild type or the $\Delta gacA$ mutant at a ratio of 1:1, or by combining SWT with cell-free supernatants of both the wild type and the $\Delta gacA$ mutant at a ratio of 2:1:1. Bars indicate the SE.




FIG. 3. Luminescence of wild-type *V. fischeri* and derivatives in culture. The luminescence of wild-type (■), *luxI* (▲), and Δ *gacA* (●) cultures grown in SWT was determined throughout the growth cycle. One representative experiment is presented. Bars indicate the SE and are sometimes obscured by the symbols.

culture of the wild type or the *luxI* or Δ *gacA* mutant that produced no detectable light became luminous after the addition of decanal or either of the acyl HSLs (Table 2). The wild type and the *luxI* mutant were similar in their luminescence response to all three substances (Table 2). However, with the addition of C₆-HSL or decanal, the Δ *gacA* mutant produced only about 20% of the luminescence of either the wild type or the *luxI* mutant. With C₈-HSL, the Δ *gacA* mutant produced only 2% of the luminescence of the other two strains (Table 2). The inability of exogenous acyl-HSL to complement its luminescence defect suggests that repression of luminescence in the *gacA* mutants is not caused solely by an acyl-HSL or decanal deficiency but is effected by another mechanism.

Since the Δ *gacA* mutant was minimally responsive to addition of excess acyl-HSLs, it was still unclear whether the strain produced these compounds. Quantification of acyl-HSLs revealed the Δ *gacA* mutant produced both C₆-HSL (0.07 nM) and C₈-HSL (1.2 μ M) at the same molarity as the wild type (0.15 nM and 1.3 μ M, respectively). The *luxI* mutant also

TABLE 2. Luminescence response to acyl-HSL and decanal

Strain	Specific luminescence (LU/OD) ^a			
	No additions	C ₆ -HSL ^b	C ₈ -HSL ^b	Decanal ^c
Wild type	<0.02	252	3.5	20
<i>luxI</i> mutant	<0.02	203	2.7	16
Δ <i>gacA</i> mutant	<0.01	43	0.069	3.5

^a One LU = 1.3×10^7 quanta/s. Luminescence was determined on aliquots of exponentially growing cultures at an OD₆₀₀ of 0.25 to 0.35 from which no luminescence was detected prior to additions. One representative time point is shown.

^b Cells were grown with shaking in the presence of 120 nM C₆-HSL or C₈-HSL.

^c Decanal (0.01%) was added to an aliquot of culture prior to measuring luminescence.




FIG. 4. Effect of medium viscosity on the motility of wild-type *V. fischeri* and derivatives. The extent of movement of duplicate samples of wild-type (gray bars), Δ *gacA* (black bars), *gacA::EZ::TN<KAN>* (white bars), and hyperswimmer strain DM66 (hatched bars) cells was measured over time. Average values (± 1 standard deviation) were normalized to the wild-type rates at each viscosity. The absence of error bars indicates no variability within treatment.

produced C₈-HSL at a similar concentration (1.7 μ M); however, as expected, no C₆-HSL was detectable (<0.005 nM).

Additional colonization traits are affected by the GacA mutations in culture. Previous studies have identified additional traits of *V. fischeri* important during host colonization. These include the production of catalase (32, 54) and siderophore (19), as well as motility (18, 36), all of which have been shown to be regulated by GacA in other bacterial species (22). Therefore, we determined whether GacA from *V. fischeri* globally controls these colonization phenotypes in culture.

The *gacA::EZ::TN<KAN-2>* mutant was not defective in catalase activity, as culture extracts were comparable to the wild type in the degradation of hydrogen peroxide. However, both the *gacA* mutants, but not the *uvrC* mutant, were defective at siderophore-mediated iron sequestration on CAS agar plates. CAS agar, which is a defined, low-iron medium containing 0.3% CA, did sustain growth of the *gacA* mutants, although they grew more slowly than other mutants that are also defective in siderophore activity (see Materials and Methods), indicating that iron limitation alone did not cause the *gacA* growth defect observed on minimal agar plates. An intact *gacA* gene supplied in *trans* on pVCW3C3 restored siderophore production to the *gacA* mutants.

GacA also regulates motility behavior in *V. fischeri*, but its impact was complex. Exponentially growing *gacA* mutants of *V. fischeri* from liquid cultures were motile; however, their ability to swim through various concentrations of soft agar was altered. Although other *Vibrio* spp. exhibit swarming behavior on higher concentrations of agar due to a lateral flagellar gene system distinct from the polar flagella used for swimming (48), *V. fischeri* ES114 swims on agar concentrations between 0.25 and 0.7% and has not been reported or observed to be peritrichously flagellated or to exhibit swarming motility. At a relatively low viscosity (0.3% agar), both *gacA* mutants swam faster than the wild type and were similar to a hyperswimmer strain of *V. fischeri*, DM66 (Fig. 4). However, at a higher viscosity (0.7% agar), the *gacA* mutants swam more slowly than




FIG. 5. Patterns of chemotaxis in soft agar by hyperswimmer derivatives of *V. fischeri*. The relative migration rates towards serine (inner ring) or nucleosides (outer ring) are indicated by the ring diameters. Shown are the patterns of strain DM66 (A and C), which is the same as the wild-type pattern, and the $\Delta gacA$ mutant (B and D) in medium without (A and B) or with (C and D) the addition of 1.6 mM serine.

the wild type, which swam more slowly than DM66 (Fig. 4). The motility of the *gacA* mutants when grown at an intermediate viscosity (0.5% agar) (Fig. 4) and at all agar concentrations tested when *gacA* was supplied in *trans* on pVCW3C3 (data not shown) was indistinguishable from that of the wild type. The *uvrC* mutation had no detectable effect on motility (data not shown). Examination by transmission electron microscopy of the $\Delta gacA$ mutant grown in broth cultures revealed no apparent differences in flagellum length, width, or appearance; however, it was slightly hyperflagellated (5.2 ± 0.3 flagella per cell) compared to the wild type (3.1 ± 0.2 flagella per cell).

Differences in chemotaxis between *gacA* mutants and the wild type were also observed. At the leading edge of the migrating front, both the wild type (data not shown) and the hyperswimmer strain DM66 (Fig. 5A) created two distinct concentric bands, representing chemotaxis up a gradient created by degradation of each of two attractants, nucleosides and serine (13). In contrast, the concentric bands of the $\Delta gacA$ mutant migrated more closely to each other (Fig. 5B) and often appeared as one diffuse band. With the addition of either 0.5% CA (data not shown) or 1.6 mM serine to the medium, which slows the migration of the inner band of both wild-type (data not shown) and DM66 (Fig. 5C) cells, the two bands generated by migration of the *gacA* mutant became distinctly separated (Fig. 5D). This observation is consistent with the hypothesis that the *gacA* mutant depleted serine from the medium more rapidly than the wild type did.

Symbiotic *gacA* mutants are impaired in host colonization and growth, but not in luminescence. Because *gacA* influenced traits in culture that could affect (i) colonization initiation, e.g., motility (18), (ii) growth within the light organ, e.g., substrate utilization (20), and (iii) persistent association, e.g., lumines-

TABLE 3. Luminescence characteristics of colonized squid

Strain	% Colonized squid that were detectably luminous at:			LU ^a	CFU ^b	Specific luminescence ^c
	24 h	48 h	72 h			
Wild type	100	100	100	247 ± 42	(1.2 ± 0.1) × 10 ⁵	5.8 ± 3
$\Delta gacA$ mutant	10	11	14	16.4 ± 8.7	(1.0 ± 0.4) × 10 ⁴	5.0 ± 2

^a One LU = 1.3×10^4 quanta/s and is reported as the mean ± SE. Luminescence is reported from 8 of 72 $\Delta gacA$ mutant-colonized squid and 32 wild-type-colonized squid that were detectably luminous on the day cell counts were determined.

^b Mean CFU of only luminous squid reported ± SE.

^c The specific luminescence was calculated as the mean of the sum of individual specific luminescence values (LU/10³ CFU from each luminous individual) reported ± SE and does not represent the average LU/average CFU.

cence (52) and siderophore production (19), we tested the ability of the $\Delta gacA$ mutant to colonize, grow within, and maintain an association with juvenile squid. When newly hatched juvenile squid were placed for 3 h in SW containing 2×10^3 wild-type cells/ml, 100% of the animals became colonized; however, only 51% of animals exposed to the same concentration of $\Delta gacA$ cells were successfully colonized. A colonization efficiency of 100% was achieved by *gacA* mutants only after a 14- to 18-h inoculation with $\geq 10^4$ CFU/ml, a level that is at least 50-fold higher than that required by the wild type.

Although most $\Delta gacA$ mutant-colonized squid produced no detectable luminescence, a subset was luminous (Table 3). These data contrast with what we observed with the *gacA* mutant grown in culture, which never produced detectable luminescence without the addition of acyl-HSL or decanal (Fig. 3 and Table 2). We confirmed that bacteria isolated from these luminous squid were $\Delta gacA$ based on both their colony morphology on LBS and CAS agar and their luminescence and growth yield phenotypes in culture (data not shown); however, the possibility remained that a mutation had occurred that suppressed squid phenotypes or that the strains had adapted in some other way to the light organ environment. We confirmed that $\Delta gacA$ mutants from luminous animals had not acquired a mutation that suppressed *gacA* colonization phenotypes, because such squid isolates retained a comparably low efficiency of colonization and a low proportion of luminescence (10%) characteristic of the original $\Delta gacA$ mutant inoculum. Similarly, bacteria directly expelled from luminous *gacA* mutant-colonized squid and not cultured in medium prior to a subsequent exposure to squid were characteristically impaired at colonizing juveniles, whereas expelled, wild-type bacteria were not impaired, even when diluted 100-fold (data not shown). Thus, there was no evidence that $\Delta gacA$ symbionts in luminous animals had adapted to the host, improving their ability to reinfect squid.

Enumeration of bacteria from the light organs of colonized squid provided insight into why there were differences in luminescence levels between various $\Delta gacA$ mutant-colonized squid. Wild-type bacterial populations averaged 1.2×10^5 CFU/squid, whereas $\Delta gacA$ populations averaged only 1.5×10^3 CFU/squid, and this density was maintained over several




FIG. 6. Colonization levels of wild-type *V. fischeri* and its $\Delta gacA$ derivative. The mean number (\pm SE) of symbiotic bacteria per colonized squid for each treatment was determined by plating light organ contents at different times following colonization with the wild type (gray bars) or the $\Delta gacA$ mutant (black bars). The colonization levels of the subsets of $\Delta gacA$ mutant-colonized animals that were either detectably luminous (white bars) or not (hatched bars) are also plotted separately. The dashed line represents the average CFU at the luminescence detection limit for wild-type-colonized squid. The mean CFU level in nonluminous wild-type-colonized squid was below this detection limit.

days (Fig. 6), whereas previously characterized derivatives of *V. fischeri* in which luminescence has been either abolished or reduced do not maintain their initial population levels and their population diminishes by 48 h postinoculation (32). Squid colonized by the wild type became detectably luminous between 7 and 9 h postinoculation, when their populations reached an average of 8×10^3 CFU/squid (Fig. 6). On average, $\Delta gacA$ mutant-colonized animals that were dark contained populations of only 4×10^2 cells and, thus, were below this minimum level of luminescence detection. The $\Delta gacA$ mutant-colonized animals that were detectably luminous had larger symbiont populations (Table 3) and were above the detection limit (Fig. 6). Therefore, the inability of the $\Delta gacA$ mutant to reach a normal colonization level in the light organ most likely prevented the induction of detectable levels of luminescence in these animals. Because a percentage of $\Delta gacA$ mutant-colonized animals was detectably luminous and their symbionts did not differ from the wild type in their luminescence per bacterial cell (Table 3), we inferred that GacA was not required to achieve light emission in the squid. Normal (100%) colonization efficiency (data not shown), colonization level, and luminescence of the mutant were fully restored by carrying the *gacA* gene in *trans* on pVCW3C3 (Table 4).

TABLE 4. Complementation of $\Delta gacA$ symbiotic defects by *gacA*

Strain	Plasmid	LU ^a	CFU ^b
Wild type	pVO8	175 ± 54	$(4.9 \pm 1.2) \times 10^5$
$\Delta gacA$ mutant	pVO8	10 ± 8	$(1.3 \pm 0.4) \times 10^3$
$\Delta gacA$ mutant	pVCW3C3	117 ± 22	$(3.4 \pm 0.6) \times 10^5$

^a One LU = 1.3×10^4 quanta/s and is reported as the mean \pm SE.

^b Mean CFU per light organ is reported \pm SE.




FIG. 7. Colonization of squid by wild-type *V. fischeri* and its $\Delta gacA$ derivative in mixed bacterial inoculations. (A and B) The proportion of wild-type-colonized (light gray), $\Delta gacA$ mutant-colonized (black), cocolonized (dark gray), or uncolonized (white) squid after inoculation with either 6,000 CFU of each strain by itself or 12,000 CFU of both strains combined at a 1:1 ratio (wild type/ $\Delta gacA$ mutant) ($n = 20$ for each treatment) (A), or with a total of 110 CFU of the wild type by itself ($n = 20$), 2,500 CFU of the $\Delta gacA$ mutant by itself ($n = 20$), or 2,600 CFU of both strains combined at a 1:23 ratio (wild type/ $\Delta gacA$) ($n = 70$) (B). (C) The mean number (\pm SE) of wild-type (light gray) and $\Delta gacA$ (black) CFU per light organ at 48 h postinoculation is shown from the three subgroups that resulted from the combined treatment presented in panel B ($n = 46$). For wild-type-colonized or cocolonized squid a minimum of 200 colonies were identified, but for $\Delta gacA$ mutant-colonized squid frequently fewer than 100 bacterial colonies were available for assessment due to this mutant's lower colonization level.

***gacA* mutants are severely impaired at competitively initiating colonization but are not displaced by the wild type in cocolonized light organs.** Determining the relative effectiveness of mutant and wild-type bacteria during coinoculation experiments can help elucidate interactions between the strains as they initiate association with the host, because a direct competition can accentuate defects and reveal otherwise subtle differences between strains. Thus, we used such competition experiments to determine (i) whether the wild type either complemented or exacerbated the association defects of the mutant and, conversely, (ii) whether the mutant interfered with colonization by the wild type.

When squid were coinoculated for 3 h with both the $\Delta gacA$ mutant and the wild type, each at 6×10^3 CFU/ml, less than 0.05% of the cells present in the symbiotic population at either 24 or 48 h were $\Delta gacA$ (Fig. 7A). This result indicated that the $\Delta gacA$ mutant was at a competitive disadvantage in colonization in the presence of wild-type cells. Since at an inoculum of 6×10^3 CFU/ml the wild type by itself colonized 100% of the squid, whereas the $\Delta gacA$ mutant by itself colonized only 65% of the squid (Fig. 7A), we hypothesized that the mutant's defect was expressed during the initiation of the colonization rather than during competitive growth within the light organ. To test this hypothesis, we adjusted the inoculum so that it would give the two strains an equal chance at initiating symbiosis. To equalize the strains' colonization efficiencies, we combined $\Delta gacA$ cells at a concentration of 2.5×10^3 CFU/ml, which by itself resulted in colonization of 55% of the squid, with wild-type cells at a concentration of only 1.1×10^2 CFU/ml, which by itself resulted in colonization of 65% of the squid (Fig. 7B). With this 23:1 advantage, the $\Delta gacA$ mutant initiated the cocolonization process with the efficiency expected and successfully cocolonized squid with the wild type (Fig. 7B). In the mixed symbiotic populations that resulted, the mutant and the wild type attained essentially the same levels as they did when they colonized in the absence of the other strain (Fig. 7C). Thus, it appears that the $\Delta gacA$ competition defect is important primarily during initiation. Even after initiation, wild-type cells neither complemented the growth defect of the $\Delta gacA$ mutant nor, surprisingly, displaced the $\Delta gacA$ population after 48 h postinoculation. Similarly, there was no indication that the presence of the $\Delta gacA$ mutant affected the ability of the wild type to attain and maintain its normal level of colonization.

DISCUSSION

We show here that in *V. fischeri* GacA globally controls multiple phenotypes, several of which are generally important to bacteria during host colonization, and at least one of which, luminescence, plays a specific role in this symbiotic association (52). This study also establishes that growth substrate utilization is coregulated by GacA along with other colonization traits. The contribution of each phenotype to the observed colonization defects remains speculative. However, this study confirms the conservation of GacA as a bacterial colonization regulator and indicates that GacA regulates not only pathogenic associations but also a specific, beneficial animal association.

Substrate utilization and growth. Although GacS/GacA has long been associated with the regulation of secondary metabolism (22), until recently the implication that GacA is an important regulator of growth had been largely overlooked. In several bacterial species, including *Pseudomonas fluorescens* (55), *Azotobacter vinelandii* (8), and *E. coli* (37), GacA controls the production of RpoS, an alternative sigma factor associated with the transition from rapid growth rates to slow or non-growing states. Consistent with reductions in RpoS, spontaneous *gacS* and *gacA* mutants in *P. fluorescens* are frequently isolated from and can overtake stationary-phase cultures (14), as altered RpoS function can confer a growth advantage in stationary-phase (GASP) phenotype (57, 58). Recent work has

confirmed the role of GacS/GacA in growth and substrate utilization, as mutants in the GacS/GacA homologs of *E. coli* are similar to CsrA mutants and show preference for growth on gluconeogenic substrates, such as amino acids, but not glycolytic growth substrates (41). Enhanced ability to catabolize amino acids can also confer a GASP phenotype (59). In several bacterial species, including *P. fluorescens* (1, 5, 23), *E. carotovora* (12, 25), and *E. coli* (49), GacA antagonizes the repressive activity of CsrA homologs via positive regulation of small regulatory RNA paralogs of *csrB*. These examples demonstrate that there is a strong link between GacA and growth.

Several phenotypes of the *gacA* mutants of *V. fischeri* substantiate the role of *gacA* in primary metabolism, including (i) their inability to grow on a minimal medium with simple sugars as a carbon source, (ii) their low growth yield in rich medium (Fig. 1), and (iii) a growth yield defect in squid light organs (Table 3; Fig. 6). Further results suggest that as with the homologous mutant of *E. coli* (41), the *gacA* mutant of *V. fischeri* preferentially utilized amino acids as growth substrates. These include (i) growth on minimal medium with CA as a sole carbon source, (ii) improved growth yield in rich medium when supplemented with CA (Fig. 2), and (iii) enhanced chemotaxis toward serine consistent with a more rapid utilization and depletion of this amino acid (Fig. 5). Interestingly, the wild type did not preferentially deplete the substrates that are growth limiting for the *gacA* mutant; in fact, the *gacA* mutant reached a higher cell density in wild-type-conditioned medium than in *gacA*-conditioned medium, whereas the wild type reached the same cell density in both conditioned media (Fig. 2). Such differences between the wild type and the *gacA* mutant in growth substrate utilization could allow the strains to occupy different nutritional niches during early stages of growth in cocolonized light organs, allowing the *gacA* mutant to maintain its minority population despite the abundance of competitors (Fig. 7C). Although the extent of the metabolic defects of the *gacA* mutant of *V. fischeri* remains unknown, the inability of the mutant to grow on the gluconeogenic substrate fumarate implies that regulation by GacA in *V. fischeri* may be more complex than a defect in switching between gluconeogenesis and glycolysis, as has been observed with *E. coli* (41).

The growth defects described both in culture and during symbiotic association imply that the *gacA* mutation interfered with the ability of *V. fischeri* to sense and adapt to the nutrient conditions of the light organ. For instance, the limited availability of amino acids in the light organ could underlie the restricted growth of the *gacA* mutant much as it does for amino acid auxotrophs (20). Recently, it has been reported that pathogenic *Salmonella enterica* serovar Typhimurium recognizes its location within the enteric tract by sensing the presence of intestinal short-chain fatty acids and, in response, induces invasion genes through a process mediated by the GacS/GacA homologs SirA/BarA (27). A similar inability to respond appropriately to a light organ signal could impair the *gacA* mutant not only during growth in the light organ (Table 3; Fig. 6) but also during initiation (Fig. 7A and B). Further characterization of the *gacA* mutant may elucidate which nutrient resources serve as host-specific signals during symbiotic association.

Luminescence regulation. One of the most striking phenotypes of the GacA mutants in culture was their inability to

produce luminescence, a trait that is specifically important for the squid-*V. fischeri* association (52). Due to the dependence of squid luminescence on acyl-HSL signal accumulation (32, 52) and the linkage of GacS/GacA to acyl-HSL expression in other host-associated bacteria (9, 15, 43), we suspected that the dark phenotype of the GacA mutant of *V. fischeri* resulted from a deficiency in acyl-HSL synthesis or accumulation. However, the GacA mutant produced typical levels of both C₆-HSL and C₈-HSL in culture and responded only partially to the addition of excess acyl-HSLs (Table 2), suggesting that luminescence expression could be blocked in the absence of GacA. Surprisingly, whereas GacA was required for luminescence in culture, it was not required for characteristic induced levels of luminescence per bacterial cell in the host light organ (Table 3). In contrast *luxI* mutants, which are luminous in culture (Fig. 3), are not luminous in the light organ even though they initially reach populations similar to the wild-type strain (32, 52). Although other factors may play a role, these results demonstrate that C₆-HSL-mediated induction of luminescence and not GacA is the dominant activating pathway in the squid host.

Other symbiosis-related phenotypes. The appropriate expression of motility behavior, which is regulated by GacA in other bacteria (17, 26, 56), is critical during early stages of squid-host association (18, 36). Although the hyperflagellation of *gacA* mutants may explain their hyperswimmer phenotype in low-viscosity medium, it is unclear why they appear less motile than the wild type in high-viscosity medium, since other hyperflagellated strains swim faster than the wild type at all medium viscosities (Fig. 4) (36). Furthermore, because *V. fischeri* ES114 does not exhibit the swarming motility that other *Vibrio* spp. exhibit (48), such differences cannot be explained as a defect in lateral flagella. Since nonmotile *V. fischeri* strains cannot initiate colonization (18), this study implies that the *gacA* mutants were motile during squid association; however, their hyperflagellation phenotype could lead to a delay in colonization (36).

We investigated two additional colonization traits that are often present in GacA regulons. Catalase production was identified as an important bacterial factor during growth in the squid light organ (54) and indicated that the oxidative environment of the light organ may restrict the growth of certain bacteria. Although bacterial defenses to oxidative damage are controlled by GacA in other bacteria (37, 55), *V. fischeri* did not require GacA for normal catalase activity in culture. In contrast, GacA was required for the production of another colonization factor, siderophore. The production of siderophores by pathogenic bacteria can contribute to virulence by mediating iron acquisition from host sources, but they can also contribute to protection from oxidative damage by preventing the Fe²⁺-catalyzed generation of free radicals (42). A recent study determined that the siderophore biosynthetic gene, *iucA*, is induced by *V. fischeri* cells within squid light organs (50), supporting the importance of iron sequestration during persistent host association (19).

GacA and symbiont specificity. This study demonstrated that *gacA* mutants were not only defective in reaching normal colonization levels but also were severely impaired during initiation and early colonization phases of symbiosis, suggesting that GacA may coregulate defense and communication activities along with nutrient acquisition. During host association, it is postulated that a selective winnowing occurs that eventually

allows colonization only by *V. fischeri* (53). Indeed, while other bacterium species can participate in the initial stages of association, even at these early stages *V. fischeri* exhibits dominance (38). Such a selection process is likely to involve not only symbiont defense traits but also reciprocal bacterium-host signaling and recognition. Analysis of the GacA regulon in *V. fischeri* is ultimately aimed at discovering such traits that may elucidate how bacteria colonize and maintain beneficial associations with animals.

ACKNOWLEDGMENTS

We thank C. DeLoney-Marino, K. Visick, and J. Graf for protocols and helpful suggestions and A. Schaefer for assistance with acyl-HSL purification and quantification. We further thank A. Schaefer, D. Millikan, C. Lupp, J. McCann, J. Graber and L. Sycuro for technical assistance, helpful conversations, and comments on earlier versions of the manuscript and D. K. Willis for guidance in the initiation of these studies.

This work was supported in part by a postdoctoral fellowship in microbial biology from the National Science Foundation to C.A.W., by National Institutes of Health grant RR12294 to E.G.R. and M. McFall-Ngai, by National Science Foundation grant IBN0211673 to M. McFall-Ngai and E.G.R., and by a W. M. Keck Foundation grant to E.G.R. and others.

REFERENCES

1. Aarons, S., A. Abbas, C. Adams, A. Fenton, and F. O'Gara. 2000. A regulatory RNA (prfB RNA) modulates expression of secondary metabolite genes in *Pseudomonas fluorescens* F113. *J. Bacteriol.* **182**:3913–3919.
2. Altschul, S., T. Madden, A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
3. Ausubel, F., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1990. *Current protocols in molecular biology*. Wiley and Sons, Inc., New York, N.Y.
4. Beers, R. F., Jr., and I. W. Sizer. 1952. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.* **195**:133–140.
5. Blumer, C., S. Heeb, B. Pessi, and D. Haas. 1999. Global GacA-steered control of cyanide and exoprotease production in *Pseudomonas fluorescens* involves specific ribosome binding sites. *Proc. Natl. Acad. Sci. USA* **96**:14073–14078.
6. Boettcher, K. J., and E. G. Ruby. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the sepiolid squid *Euprymna scolopes*. *J. Bacteriol.* **172**:1053–1058.
7. Boettcher, K. J., and E. G. Ruby. 1995. Detection and quantification of *Vibrio fischeri* autoinducer from symbiotic squid light organs. *J. Bacteriol.* **177**:1053–1058.
8. Castaneda, M., J. Sanchez, S. Moreno, C. Nunez, and G. Espin. 2001. The global regulators GacA and sigma S form part of a cascade that controls alginate production in *Azotobacter vinelandii*. *J. Bacteriol.* **183**:6787–6793.
9. Chancey, S. T., D. W. Wood, and L. S. Pierson III. 1999. Two-component transcriptional regulation of *N*-acyl-homoserine lactone production in *Pseudomonas aureofaciens*. *Appl. Environ. Microbiol.* **65**:2294–2299.
10. Chancey, S. T., D. W. Wood, E. A. Pierson, and L. S. Pierson III. 2002. Survival of GacS/GacA mutants of the biological control bacterium *Pseudomonas aureofaciens* 30–84 in the wheat rhizosphere. *Appl. Environ. Microbiol.* **68**:3308–3314.
11. Corbell, N. A., and J. E. Loper. 1995. A global regulator of secondary metabolite production in *Pseudomonas fluorescens* PF-5. *J. Bacteriol.* **177**:6230–6236.
12. Cui, Y., A. Chatterjee, and A. K. Chatterjee. 2001. Effects of the two-component system comprising GacA and GacS of *Erwinia carotovora* subsp. *carotovora* on the production of global regulatory rsmB RNA, extracellular enzymes, and HarpinEcc. *Mol. Plant Microbe Interact.* **14**:516–526.
13. DeLoney-Marino, C. R., C. R. Wolfe, and K. L. Visick. Chemoattraction of *Vibrio fischeri* to serine, nucleosides, and *N*-acetylneuraminic acid, a component of squid light-organ mucus. *Appl. Environ. Microbiol.*, in press.
14. Duffy, V. K., and G. Defago. 2000. Controlling instability in *gacS-gacA* regulatory genes during inoculate production of *Pseudomonas fluorescens* bio-control strains. *Mol. Plant Microbe Interact.* **66**:3142–3150.
15. Eriksson, A. R. B., R. A. Andersson, M. Pirhonen, and E. T. Palva. 1998. Two-component regulators involved in the global control of virulence in *Erwinia carotovora* subsp. *carotovora*. *Mol. Plant Microbe Interact.* **8**:743–752.

16. Flavier, A. B., L. M. Ganova-Raeva, M. A. Schell, and T. P. Denny. 1997. Hierarchical autoinduction in *Ralstonia solanacearum*: control of acyl-homoserine lactone production by a novel autoregulatory system responsive to 3-hydroxypalmitic acid methyl ester. *J. Bacteriol.* **179**:7089–7097.
17. Goodier, R. I., and B. M. M. Ahmer. 2001. SirA orthologs affect both motility and virulence. *J. Bacteriol.* **183**:2249–2258.
18. Graf, J., P. V. Dunlap, and E. G. Ruby. 1994. Effect of transposon-induced motility mutations on colonization of the host light organ by *Vibrio fischeri*. *J. Bacteriol.* **176**:6986–6991.
19. Graf, J., and E. G. Ruby. 2000. Novel effects of a transposon insertion in the *Vibrio fischeri glnD* gene: defects in iron uptake and symbiotic persistence in addition to nitrogen utilization. *Mol. Microbiol.* **37**:168–179.
20. Graf, J., and E. G. Ruby. 1998. Host-derived amino acids support the proliferation of symbiotic bacteria. *Proc. Natl. Acad. Sci. USA* **95**:1818–1822.
21. Hammer, B. K., E. S. Tateda, and M. S. Swanson. 2002. A two-component regulator induces the transmission phenotype of stationary-phase *Legionella pneumophila*. *Mol. Microbiol.* **44**:107–118.
22. Heeb, S., and D. Haas. 2001. Regulatory roles of the GacS/GacA two-component system in plant-associated and other gram-negative bacteria. *Mol. Plant Microbe Interact.* **14**:1351–1363.
23. Heeb, S., C. Blumer, and D. Haas. 2002. Regulatory RNA as mediator in GacA/RsmA-dependent global control of exoproduct formation in *Pseudomonas fluorescens* CHA0. *J. Bacteriol.* **184**:1046–1056.
24. Herrero, M., V. de Lorenzo, and K. N. Timmis. 1990. Transposon vectors containing non-antibiotic resistance selection markers for cloning and stable chromosomal insertion of foreign genes in gram-negative bacteria. *J. Bacteriol.* **172**:6557–6567.
25. Hyttiainen, H., M. Montesano, and E. T. Palva. 2001. Global regulators ExpA (GacA) and Kdgr modulate extracellular enzyme gene expression through the RsmA-RsmB system in *Erwinia carotovora* subsp. *carotovora*. *Mol. Plant Microbe Interact.* **14**:931–938.
26. Kinscherf, T. G., and D. K. Willis. 1999. Swarming by *Pseudomonas syringae* B728a requires *gacS* (*lemA*) and *gacA* but not the acyl-homoserine lactone biosynthetic gene *ahlI*. *J. Bacteriol.* **181**:4133–4136.
27. Lawhon, S. D., R. Maurer, M. Suyemoto, and C. Altier. 2002. Intestinal short-chain fatty acids alter *Salmonella typhimurium* invasion gene expression and virulence through BarA/SirA. *Mol. Microbiol.* **46**:1451–1464.
28. Lee, K. H., and E. G. Ruby. 1994. Competition between *Vibrio fischeri* strains during initiation and maintenance of a light organ symbiosis. *J. Bacteriol.* **176**:1985–1991.
29. Lee, K. H., and E. G. Ruby. 1994. Effect of the squid host on the abundance and distribution of symbiotic *Vibrio fischeri* in nature. *Appl. Environ. Microbiol.* **60**:1565–1571.
30. Lee, K. H., and E. G. Ruby. 1995. Symbiotic role of the viable but nonculturable state of *Vibrio fischeri* in Hawaiian coastal seawater. *Appl. Environ. Microbiol.* **61**:278–283.
31. Lund, A. H., M. Dutch, and F. S. Pedersen. 1996. Increased cloning efficiency by temperature-cycle ligation. *Nucleic Acids Res.* **24**:800–801.
32. Lupp, C., M. Urbanowski, E. P. Greenberg, and E. G. Ruby. 2003. The *Vibrio fischeri* quorum-sensing systems *ain* and *lux* sequentially induce luminescence gene expression and are important for persistence in the squid host. *Mol. Microbiol.* **50**:319–331.
33. McCann, J., E. V. Stabb, D. S. Millikan, and E. G. Ruby. 2003. Population dynamics of *Vibrio fischeri* during infection of *Euprymna scolopes*. *Appl. Environ. Microbiol.* **69**:5928–5934.
34. Meighen, E. A., and P. V. Dunlap. 1993. Physiological, biochemical and genetic control of bacterial bioluminescence. *Adv. Microb. Physiol.* **34**:1–67.
35. Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
36. Millikan, D. S., and E. G. Ruby. 2002. Alterations in *Vibrio fischeri* motility correlate with a delay in symbiosis initiation and are associated with additional symbiotic colonization defects. *Appl. Environ. Microbiol.* **68**:2519–2528.
37. Mukhopadhyay, S., J. P. Audia, R. N. Roy, and H. E. Schellhorn. 2000. Transcriptional induction of the conserved alternative sigma factor RpoS in *Escherichia coli* is dependent on BarA, a probable two-component regulator. *Mol. Microbiol.* **37**:371–381.
38. Nyholm, S. V., B. Deplancke, H. R. Gaskins, M. A. Apicella, and M. J. McFall-Ngai. 2002. Roles of *Vibrio fischeri* and nonsymbiotic bacteria in the dynamics of mucus secretion during symbiont colonization of the *Euprymna scolopes* light organ. *Appl. Environ. Microbiol.* **68**:5113–5122.
39. Pearson, J. P., K. M. Gray, L. Passador, K. D. Tucker, A. Eberhard, B. H. Iglewski, and E. P. Greenberg. 1994. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc. Natl. Acad. Sci. USA* **91**:197–201.
40. Pernestig, A. K., S. J. Normark, D. M. Georgellis, and O. Melefors. 2000. The role of the AirS two-component system in uropathogenic *Escherichia coli*. *Adv. Exp. Med. Biol.* **485**:137–142.
41. Pernestig, A. K., D. Georgellis, T. Romeo, K. Suzuki, H. Tomenius, S. Normark, and O. Melefors. 2003. The *Escherichia coli* BarA-UvrY two-component system is needed for efficient switching between glycolytic and gluconeogenic carbon sources. *J. Bacteriol.* **185**:843–853.
42. Ratledge, C., and L. G. Dover. 2000. Iron metabolism in pathogenic bacteria. *Annu. Rev. Microbiol.* **54**:881–941.
43. Reimann, C., M. Beyeler, M. Latifi, H. Winteler, M. Foglino, A. Lazdunski, and D. Hass. 1997. The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer *N*-butyryl-homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol. Microbiol.* **24**:309–319.
44. Ruby, E. G., and L. M. Asato. 1993. Growth and flagellation of *Vibrio fischeri* during initiation of the sepiolid squid light organ symbioses. *Arch. Microbiol.* **159**:160–167.
45. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
46. Schaefer, A. L., B. L. Hanselka, M. R. Parsek, and E. P. Greenberg. 2000. Detection, purification, and structural elucidation of the acylhomoserine lactone inducer of *Vibrio fischeri* luminescence and other related molecules. *Methods Enzymol.* **305**:288–301.
47. Stabb, E. V., and E. G. Ruby. 2002. RP4-based plasmids for conjugation between *Escherichia coli* and members of the *Vibrionaceae*. *Methods Enzymol.* **358**:413–426.
48. Stewart, B. J., and L. L. McCarter. 2003. Lateral flagellar gene systems of *Vibrio parahaemolyticus*. *J. Bacteriol.* **185**:4508–4518.
49. Suzuki, K., X. Wang, T. Weilbacher, A.-K. Pernestig, O. Melefors, D. Georgellis, P. Babitzke, and T. Romeo. 2002. Regulatory circuitry of the CsrA/CsrB and BarA/UvrY systems of *Escherichia coli*. *J. Bacteriol.* **184**:5130–5140.
50. Visick, K. L., and E. G. Ruby. 1997. New genetic tools for use in the marine bioluminescent bacterium *Vibrio fischeri*, p. 119–122. In J. W. Hastings, L. J. Kricka, and P. E. Stanley (ed.), *Bioluminescence and chemiluminescence*. John Wiley & Sons, New York, N.Y.
51. Visick, K. L., and E. G. Ruby. *TnluxAB* insertion mutants of *Vibrio fischeri* with symbiosis-regulated phenotypes. *Nova Acta Leopold.*, in press.
52. Visick, K. L., J. Foster, J. Doimo, M. McFall-Ngai, and E. G. Ruby. 2000. *Vibrio fischeri lux* genes play an important role in colonization and development of the host light organ. *J. Bacteriol.* **182**:4578–4586.
53. Visick, K. L., and M. J. McFall-Ngai. 2000. An exclusive contract: specificity in the *Vibrio fischeri-Euprymna scolopes* partnership. *J. Bacteriol.* **182**:1779–1787.
54. Visick, K. L., and E. G. Ruby. 1998. The periplasmic group III catalase of *Vibrio fischeri* is required for normal symbiotic competence and is induced both by oxidative stress and by approach to stationary phase. *J. Bacteriol.* **180**:2087–2092.
55. Whistler, C. A., N. A. Corbell, A. Sarniguet, W. Ream, and J. E. Loper. 1998. The two-component regulators GacS and GacA influence accumulation of the stationary-phase sigma factor σ^{54} and the stress response in *Pseudomonas fluorescens* Pf-5. *J. Bacteriol.* **180**:6635–6641.
56. Wong, S. M., P. A. Carroll, L. G. Rahme, F. M. Ausubel, and S. B. Calderwood. 1998. Modulation of expression of the ToxR regulon in *Vibrio cholerae* by a member of the two-component family of response regulators. *Infect. Immun.* **66**:5854–5861.
57. Zambrano, M. M., and R. Kolter. 1996. GASping for life in stationary phase. *Cell* **86**:181–184.
58. Zambrano, M. M., D. A. Siegle, M. Almiron, A. Tormo, and R. Kolter. 1993. Microbial competition: *Escherichia coli* mutants that take over stationary phase cultures. *Science* **259**:1757–1760.
59. Zinser, E. R., and R. Kolter. 1999. Mutations enhancing amino acid catabolism confer a growth advantage in stationary phase. *J. Bacteriol.* **181**:5800–5807.