

FlhF, the Third Signal Recognition Particle-GTPase of *Bacillus subtilis*, Is Dispensable for Protein Secretion

Geeske Zanen,¹ Haike Antelmann,² Helga Westers,^{1†} Michael Hecker,² Jan Maarten van Dijl,^{1*} and Wim J. Quax¹

Department of Pharmaceutical Biology, University of Groningen, Groningen, The Netherlands,¹ and Institut für Mikrobiologie und Molekularbiologie, Ernst-Moritz-Arndt-Universität Greifswald, Greifswald, Germany²

Received 11 March 2004/Accepted 8 June 2004

***Bacillus subtilis* contains three proteins of the signal recognition particle-GTPase family known as Ffh, FtsY, and FlhF. Here we show that FlhF is dispensable for protein secretion, unlike Ffh and FtsY. Although *flhF* is located in the *fla/che* operon, *B. subtilis* 168 *flhF* mutant cells assemble flagella and are motile.**

In eukaryotes, prokaryotes, and archaea, a large number of proteins is transported across membranes in order to fulfill their biological function. Complex and well-organized protein transport systems have evolved for membrane translocation of these proteins. Most proteins that play a role outside the cytoplasm contain a signal peptide, which directs the (pre)protein to its final destination (2, 23, 28, 29). Chaperones and targeting factors recognize this signal peptide and keep preproteins in an export-competent state before targeting to the translocation machinery in the membrane. The major machinery for protein transport is the Sec translocase, which handles preproteins in an unfolded state (7).

On the basis of proteomic studies, it has been proposed that the majority of secretory proteins of the gram-positive bacterium *Bacillus subtilis* are targeted to the Sec translocase by the so-called signal recognition particle (SRP) (9). This SRP seems to be involved in preprotein targeting to membranes of organisms belonging to all three domains of life. The *B. subtilis* SRP complex consists of the Ffh (Fifty-four homolog) protein (10), a small cytoplasmic RNA (scRNA) (15, 16), and a histone-like protein (HBSu) (17). Preprotein targeting by this SRP complex presumably involves the presence of the SRP receptor-like protein FtsY (18). Both Ffh and FtsY belong to the widely conserved family of SRP-GTPases (8). Interestingly, *B. subtilis* and several other bacterial species (but not *Escherichia coli*) contain a third gene encoding a protein belonging to the SRP-GTPase family. In *B. subtilis*, this paralogue of Ffh and FtsY was named FlhF (flagellum-associated protein) because it appeared to be required for the flagellar assembly and motility of this bacterium (5). Specifically, the *B. subtilis* FlhF protein has 46% identical residues and conservative replacements in a stretch of 175 residues with *B. subtilis* Ffh and 37% identical residues and conservative replacements in a stretch of 318 residues with *B. subtilis* FtsY. As shown by sequence alignments and domain searches, FlhF contains the conserved N

and G domains of the SRP-like GTPases (Fig. 1). However, it lacks the so-called M domain typical for the C termini of Ffh-like proteins and contains a basic B domain instead of the acidic A domain of FtsY-like proteins of bacteria and yeasts. Notably, the mammalian SRP receptor SR α contains a more basic N-terminal domain, like FlhF of *B. subtilis*. Consistent with its proposed function, the *flhF* gene is located within the *che/fla* operon, which encodes the majority of the chemotaxis and flagellar proteins (11). Pandza and coworkers (20) showed that the FlhF homologue of *Pseudomonas putida* has a role in polar flagellar placement and in induction of the general stress response.

On the basis of the similarity between FlhF and Ffh/FtsY, Carpenter et al. (5) proposed that FlhF might be involved in protein secretion. Notably, however, FlhF is dispensable for growth and viability, whereas Ffh and FtsY are essential, like the key components SecA, SecY, and SecE of the Sec translocase (12). This raised the questions of whether and, if so, to what extent FlhF is involved in protein secretion.

Construction of a *B. subtilis* 168 *flhF* mutant. Since all of our previous studies on protein secretion by *B. subtilis* were performed with sequenced strain 168 (13), a *B. subtilis* 168 *flhF::cat* mutant strain was constructed by transforming *B. subtilis* 168 with chromosomal DNA of *flhF* mutant strain OI2735, which was constructed by Carpenter et al. (5) (Table 1). *B. subtilis* 168 was transformed as previously described (22). Chloramphenicol-resistant transformants were screened by PCR with primers cat1 (5'-GAT TTA GAC AAT TGG AAG) and cat2 (5'-GAC AAT TCC TGA ATA GAG) to show the presence of the *cat* gene (data not shown). PCR was carried out with the *Pwo* DNA polymerase (Roche) as described previously (26).

FlhF is not required for protein secretion by *B. subtilis* strains 168 and DB430. To investigate the involvement of FlhF in protein secretion, the composition of the extracellular proteome of *B. subtilis* 168 *flhF::cat* was analyzed and compared to that of parental strain 168. In addition, similar experiments were performed with protease-deficient strain DB430 and a DB430 *flhF::cat* derivative that was obtained by transformation of strain DB430 with chromosomal DNA of strain OI2735. For analysis of their extracellular proteomes, all strains were grown at 37°C under vigorous agitation in rich medium. After 1 h of

* Corresponding author. Present address: Department of Molecular Bacteriology, University of Groningen, P.O. Box 30001, 9700 RB Groningen, The Netherlands. Phone: 31-50-3633079. Fax: 31-50-3633528. E-mail: J.M.van.Dijl@med.rug.nl.

† Present address: Department of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, 9751 NN Haren, The Netherlands.




FIG. 1. Conserved domains in proteins of the SRP-GTPase family. The SRP-GTPase family members of yeast (SRP54, SR α), *E. coli* (P48, FtsY_Ec), and *B. subtilis* (Ffh, FlhF, FtsY_Bs) are represented schematically. Different domains that can be distinguished are the acidic A domain; the basic B domain, the conserved N domain, the M domain involved in RNA and preprotein binding, and the GTP-binding G domain. The five conserved boxes, G1 to G5, in the G domain, as defined by Eichler and Moll (8), are shown.

postexponential growth, cells were separated from the growth medium by centrifugation and proteins secreted into the growth medium were concentrated by trichloroacetic acid precipitation. The resulting samples were used for two-dimensional gel electrophoresis, and protein spots were identified by matrix-assisted laser desorption ionization–time of flight mass spectrometry and/or N-terminal sequencing as previously described (1). After dual-channel imaging to visualize possible changes in the extracellular protein composition, no major differences were observed between the extracellular proteomes of the *flhF::cat* mutants and the respective parental strains, 168

and DB430 (Fig. 2A). Under the conditions tested, some fluctuation in the levels of prophage-encoded proteins XkdK, XkdG, XkdM, and YolA was observed in the growth media of *flhF* mutant strains, as well as in the media of parental strains 168 and DB430 (Fig. 2B). Most likely, this reflects fluctuations in the expression of genes located on the PBSX prophage (*xkdK*, *xkdG*, and *xkdM*) and the SP β prophage (*yolA*). Unexpectedly, the extracellular accumulation of proteins known to be required for cell motility, such as FlgK (flagellar hook-associated protein 1), FliD (flagellar hook-associated protein 2), and the flagellin Hag, was not affected by the absence of FlhF.

It was previously shown that the absence of some components of the Sec machinery of *B. subtilis*, such as SecDF (4) and SecG (27), has no detectable effect on protein secretion unless the secretion machinery is challenged with overproduced secretory proteins. For example, this was shown by high-level expression of the α -amylase AmyQ of *Bacillus amyloliquefaciens* (4) with plasmid pKTH10 (Table 1). To study the importance of FlhF for AmyQ secretion at high levels, the *flhF* mutant strain and parental strain 168 were transformed with pKTH10. After overnight growth in TY medium (1% Bacto Tryptone, 0.5% Bacto Yeast Extract, 1% NaCl) supplemented with kanamycin, cells and medium fractions were separated by centrifugation (2 min, 16,000 \times g, room temperature). Next, protein samples for sodium dodecyl sulfate-polyacrylamide gel electrophoresis were prepared as described previously (25). After separation by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, proteins were transferred to a Protran nitrocellulose transfer membrane (Schleicher & Schuell) as described by Kyhse-Andersen (14). AmyQ was visualized with specific antibodies and horseradish peroxidase-goat anti-rabbit immunoglobulin G conjugates (BioSource International). As shown in Fig. 3A, disruption of the *flhF* gene affects neither the amounts of AmyQ secreted into the growth medium nor the amounts of AmyQ present in the cells. Taken together, these findings demonstrate that FlhF is dispensable for protein secretion, even when the secretion machinery of *B. subtilis* is challenged by the high-level production of a secretory protein.

Since FlhF is a paralogue of Ffh and FtsY, it is conceivable that the *B. subtilis* cell can suppress the effects of the absence of FlhF by production of Ffh or FtsY at increased levels. To study possible changes in the levels of Ffh and FtsY in the absence of FlhF, Western blotting experiments were performed. For this purpose, overnight cultures of *B. subtilis* 168

TABLE 1. Plasmid and bacterial strains used in this study

Plasmid or strain	Relevant properties ^a	Reference
pKTH10	Encodes α -amylase AmyQ of <i>B. amyloliquefaciens</i> , 6.8 kb; Km ^r	19
<i>B. subtilis</i>		
168	<i>trpC2</i>	13
O12735	<i>flhF::cat</i> Cm ^r	5
168 <i>flhF::cat</i>	Like 168; <i>flhF::cat</i> Cm ^r	This paper
BFA2616	Like 168; <i>ylxH::pMutin2</i> Em ^r	12
BFA2616 <i>flhF::cat</i>	Like BFA2616; <i>flhF::cat</i> Cm ^r Em ^r	This paper
DB430	Like 168; <i>nprE aprE bpf isp1</i>	6
DB430 <i>flhF::cat</i>	Like DB430; <i>flhF::cat</i> Cm ^r	This paper

^a Km^r, kanamycin resistance (20 μ g/ml); Cm^r, chloramphenicol resistance (5 μ g/ml); Em^r, erythromycin resistance (2 μ g/ml).




FIG. 2. Extracellular proteome of *B. subtilis flhF::cat*. The extracellular proteins of the *flhF::cat* mutant strains and the respective parental strains 168 and DB430 were separated by two-dimensional gel electrophoresis, after which dual-channel fluorescence imaging was used to visualize possible changes in extracellular protein composition (3). Protein spots identified by mass spectrometry and/or N-terminal sequencing are indicated. Green protein spots are predominantly present in the image of the extracellular proteins of the parental strain, red protein spots are predominantly present in the image of the extracellular proteins of the *flhF* mutant strain, and yellow protein spots are present in similar amounts in both images. (A) Extracellular proteomes of *B. subtilis* DB430 and DB430 *flhF::cat*. (B) Variable extracellular levels of prophage-encoded proteins YolA, XkdM, XkdG, and XkdK (top to bottom).




FIG. 3. Absence of FlhF has no impact on secretion of AmyQ and cellular levels of Ffh and FtsY. The secretion of overproduced AmyQ (A) and the intracellular levels of Ffh and FtsY (B) were analyzed by Western blotting with cellular (c) and/or growth medium (m) fractions of *B. subtilis* 168 *flhF::cat* and parental strain 168. d, degradation products of AmyQ.

flhF::cat and parental strain 168 were diluted to an optical density at 600 nm of 0.05 and grown until 1 h after the transition between exponential and postexponential growth. Subsequently, cells were collected by centrifugation and prepared for Western blotting as indicated above. Ffh and FtsY were visualized with specific antibodies and alkaline phosphatase-conjugated anti-rabbit immunoglobulin G (BioSource International) and a standard nitroblue tetrazolium-5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside reaction (21). The results demonstrate that the intracellular levels of Ffh and FtsY are not affected by disruption of the *flhF* gene (Fig. 3B). Consequently, it can be concluded that *B. subtilis* cells do not compensate for the absence of FlhF by upregulation of the production of Ffh and/or FtsY. However, complementation of the *flhF* mutation by the production of Ffh and/or FtsY at normal levels cannot be ruled out.

FlhF has a minor role in the motility of strain 168. As the *flhF* mutation in *B. subtilis* O12735 was shown to result in nonmotility, we verified whether the same would be true for cells of *B. subtilis* 168 *flhF::cat* by using a motility plate assay. *B. subtilis* cultures were grown overnight at 37°C in TY medium. Next, the optical density at 600 nm was measured and adjusted to 1.0 with fresh TY medium. Subsequently, an aliquot of 2 μ l was spotted onto TY plates containing 0.27% agar (supplemented with 1 mM isopropyl- β -D-thiogalactopyranoside [IPTG] when appropriate). Finally, after incubation for 12 h at 37°C, the swarming distances of the different strains




FIG. 4. Motility assays. (A) Comparison of the motilities of *B. subtilis* strains 168, OI2735, 168 *flhF::cat*, BFA2616 (grown in the absence or presence of 1 mM IPTG), and BFA2616 *flhF::cat* (grown in the absence or presence of 1 mM IPTG) after 12 h of incubation on 0.27% agar plates at 37°C. (B) Schematic representation of *ylxH* gene disruption by pMutin2 via Campbell-type integration. *lacI*, *E. coli lacI* gene; *ori* pBR322, origin of replication of plasmid pBR322; *ApR*, ampicillin resistance marker; *EmR*, erythromycin resistance marker; t1 t2, transcriptional terminators on pMutin2; *Pspac*, IPTG-dependent promoter; *Pfla/che*, promoter of the *fla/che* operon; *ylxH'*, 3'-truncated *ylxH* gene; *ylxH*, 5'-truncated *ylxH* gene; *flhF*, *flhF* gene; *cheB*, *cheB* gene.

were compared (Fig. 4A). Consistent with the report of Carpenter et al. (5), *B. subtilis flhF* mutant strain OI2735 is non-motile. Remarkably, however, disruption of the *flhF* gene in *B. subtilis* 168 has no effect on cell motility (Fig. 4A). Moreover, scanning electron microscopy shows that *B. subtilis* 168 *flhF::cat* produces apparently intact flagella (data not shown). These observations imply that there are substantial differences in the genetic backgrounds of the two *flhF* mutant strains. To investigate whether genes downstream of *flhF* might be involved in this phenomenon, the motility of *B. subtilis* 168 with an integrated copy of the pMutin2 plasmid in the *ylxH* gene (strain BFA2616; Fig. 4B) was tested. As shown in Fig. 4A, the motility of strain BFA2616 was significantly reduced compared to that of *B. subtilis* 168. However, induction of the transcription of genes downstream of *ylxH* by activation of the pMutin2-derived *Pspac* promoter with IPTG resulted in less severely impaired motility of the cells. This shows that the YlxH protein has a role in cell motility. In addition to YlxH, proteins encoded by genes downstream of the *ylxH* gene are required for motility. To further investigate a possible role of FlhF in this process, the BFA2616 *flhF::cat* double mutant was constructed by transformation of strain BFA2616 with chromosomal DNA of strain OI2735. Irrespective of the presence of IPTG to induce transcription of genes downstream of *ylxH*, the motility of cells of the double-mutant strain was more severely affected than that of strain BFA2616 cells. It has to be noted, however, that upon incubation of the swarming plates for 24 h some motility of the BFA2616 *flhF::cat* double mutant was observed when it was grown in the presence of IPTG. In contrast, strain OI2735 displayed no motility at all. The fact that the *flhF* single mutation had no effect on motility while the combined *flhF* and *ylxH* mutations affected motility more severely than the *ylxH* single mutation indicates that the functions of FlhF and YlxH overlap at least partly. In this respect, it is interesting that YlxH has a putative nucleotide binding site, like FlhF. If FlhF and YlxH act cooperatively, the function of FlhF can be taken over by YlxH in *flhF* mutant cells, but the opposite seems not to occur. Remarkably, integration of the pMutin2 plasmid into the

ylxH gene appears to result in a polar effect on the expression of downstream genes, while there is no evidence for polar effects upon integration of the *cat* gene into *flhF*. Nevertheless, transcriptome analyses with *B. subtilis* strain 168, as documented on the JAFAN website (<http://bacillus.genome.jp/>), show that the expression profiles of *flhF* and *ylxH*, as well as the surrounding genes, are highly similar under the 10 different growth conditions tested. This strongly suggests that these genes are part of one operon or regulon. Moreover, studies by West and coworkers (30) support the idea that the expression of *flhF* and *ylxH* is controlled by one promoter. Taken together, our observations demonstrate that FlhF has a minor role in the motility of *B. subtilis* 168 cells. It is not clear why disruption of *flhF* in strain OI2735 results in a complete block of motility (5).

In conclusion, our combined proteomic and biochemical analyses of sequenced *B. subtilis* strain 168 demonstrate that FlhF, the third SRP-GTPase of this organism, is dispensable for protein secretion and has a minor role in cell motility. It remains to be investigated whether FlhF has a role in the biogenesis of membrane proteins, as shown for its homologues Ffh and FtsY in *E. coli* (24) and proposed for gram-positive bacteria (28).

We thank George Ordal for providing *B. subtilis* strain OI2735 and Marc Kolkman from Genencor International for providing anti-Ffh and anti-FtsY antibodies. Jan Jongbloed, Joen Luirink, Rob Meima, and Bauke Oudega are thanked for helpful discussions.

G.Z. and W.J.Q. were supported by the Stichting Technische Wetenschappen (BVI.4837), H.W. and J.M.V.D. were supported by the CEU (BIO4-CT98-0250, QLK3-CT-1999-00413, and QLK3-CT-1999-00917), and H.A. and M.H. were supported by the Deutsche Forschungsgemeinschaft, the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie, and the Fonds der Chemischen Industrie.

REFERENCES

- Antelmann, H., H. Tjalsma, B. Voigt, S. Ohlmeier, S. Bron, J. M. van Dijk, and M. Hecker. 2001. A proteomic view on genome-based signal peptide predictions. *Genome Res.* **11**:1484–1502.
- Blobel, G., and B. Dobberstein. 1975. Transfer to proteins across membranes. II. Reconstitution of functional rough microsomes from heterologous components. *J. Cell Biol.* **67**:852–862.

3. Blum, H., H. Beier, and H. J. Gross. 1987. Improved silver staining of plant-proteins, RNA and DNA polyacrylamide gels. *Electrophoresis* **8**:93–99.
4. Bolhuis, A., C. P. Broekhuizen, A. Sorokin, M. L. van Roosmalen, G. Venema, S. Bron, W. J. Quax, J. M. van Dijk. 1998. SecDF of *Bacillus subtilis*, a molecular Siamese twin required for the efficient secretion of proteins. *J. Biol. Chem.* **273**:21217–21224.
5. Carpenter, P. B., D. W. Hanlon, and G. W. Ordal. 1992. *flhF*, a *Bacillus subtilis* flagellar gene that encodes a putative GTP-binding protein. *Mol. Microbiol.* **6**:2705–2713.
6. Doi, R. H., S. L. Wong, and F. Kawamura. 1986. Potential use of *Bacillus subtilis* for secretion and production of foreign proteins. *Trends Biotechnol.* **4**:232–235.
7. Driessen, A. J., E. H. Manting, and C. van der Does. 2001. The structural basis of protein targeting and translocation in bacteria. *Nat. Struct. Biol.* **8**:492–498.
8. Eichler, J., and R. Moll. 2001. The signal recognition particle of Archaea. *Trends Microbiol.* **9**:130–136.
9. Hirose, I., K. Sano, I. Shioda, M. Kumano, K. Nakamura, and K. Yamane. 2000. Proteome analysis of *Bacillus subtilis* extracellular proteins: a two-dimensional protein electrophoretic study. *Microbiology* **146**:65–75.
10. Honda, K., K. Nakamura, M. Nishiguchi, and K. Yamane. 1993. Cloning and characterization of a *Bacillus subtilis* gene encoding a homolog of the 54-kilodalton subunit of mammalian signal recognition particle and *Escherichia coli* Ffh. *J. Bacteriol.* **175**:4885–4894.
11. Kirsch, M. L., P. B. Carpenter, and G. W. Ordal. 1994. A putative ATP-binding protein from the *che/fla* locus of *Bacillus subtilis*. *DNA Sequence* **4**:271–275.
12. Kobayashi, K., S. D. Ehrlich, A. Albertini, G. Amati, K. K. Andersen, M. Arnaud, K. Asai, S. Ashikaga, S. Aymerich, P. Bessieres, F. Boland, S. C. Brignell, S. Bron, K. Bunai, J. Chapuis, L. C. Christiansen, A. Danchin, M. Debarbouille, E. Dervyn, E. Deuerling, K. Devine, S. K. Devine, O. Dreesen, J. Errington, S. Fillinger, et al. 2003. Essential *Bacillus subtilis* genes. *Proc. Natl. Acad. Sci. USA* **100**:4678–4683.
13. Kunst, F., N. Ogasawara, I. Moszer, A. M. Albertini, G. Alloni, V. Azevedo, M. G. Bertero, P. Bessieres, A. Bolotin, S. Borchert, R. Borriss, L. Boursier, A. Brans, M. Braun, S. C. Brignell, S. Bron, S. Brouillet, C. V. Bruschi, B. Caldwell, V. Capuano, N. M. Carter, S. K. Choi, J. J. Codani, I. F. Conner-ton, A. Danchin, et al. 1997. The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249–256.
14. Kyhse-Andersen, J. 1984. Electroblotting of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose. *J. Biochem. Biophys. Methods* **10**:203–209.
15. Nakamura, K., Y. Imai, A. Nakamura, and K. Yamane. 1992. Small cytoplasmic RNA of *Bacillus subtilis*: functional relationship with human signal recognition particle 7S RNA and *Escherichia coli* 4.5S RNA. *J. Bacteriol.* **174**:2185–2192.
16. Nakamura, K., M. Nishiguchi, K. Honda, and K. Yamane. 1994. The *Bacillus subtilis* SRP54 homologue, Ffh, has an intrinsic GTPase activity and forms a ribonucleoprotein complex with small cytoplasmic RNA *in vivo*. *Biochem. Biophys. Res. Commun.* **199**:1394–1399.
17. Nakamura, K., S. Yahagi, R. Yamazaki, and K. Yamane. 1999. *Bacillus subtilis* histone-like protein, HBSu, is an integral component of a SRP-like particle that can bind the Alu domain of small cytoplasmic RNA. *J. Biol. Chem.* **274**:13569–13576.
18. Ogura, A., H. Kakeshita, K. Honda, H. Takamatsu, K. Nakamura, and K. Yamane. 1995. *srb*: a *Bacillus subtilis* gene encoding a homologue of the alpha-subunit of the mammalian signal recognition particle receptor. *DNA Res.* **2**:95–100.
19. Palva, I. 1982. Molecular cloning of α -amylase gene from *Bacillus amyloliquefaciens* and its expression in *Bacillus subtilis*. *Gene* **19**:81–87.
20. Pandza, S., M. Baetens, C. H. Park, T. Au, M. Keyhan, and A. Martin. 2000. The G-protein FlhF has a role in polar flagellar placement and general stress response induction in *Pseudomonas putida*. *Mol. Microbiol.* **36**:414–423.
21. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
22. Tjalsma, H., A. Bolhuis, M. L. van Roosmalen, T. Wiegert, W. Schumann, C. P. Broekhuizen, W. J. Quax, G. Venema, S. Bron, and J. M. van Dijk. 1998. Functional analysis of the secretory precursor processing machinery of *Bacillus subtilis*: identification of a eubacterial homolog of archaeal and eukaryotic signal peptidases. *Genes Dev.* **12**:2318–2331.
23. Tjalsma, H., A. Bolhuis, J. D. H. Jongbloed, S. Bron, and J. M. van Dijk. 2000. Signal peptide-dependent protein transport in *Bacillus subtilis*: a genome-based survey of the secretome. *Microbiol. Mol. Biol. Rev.* **64**:515–547.
24. Valent, Q. A., J. W. de Gier, G. von Heijne, D. A. Kendall, C. M. ten Hagen-Jongman, B. Oudega, and J. Luirink. 1997. Nascent membrane and presecretory proteins synthesized in *Escherichia coli* associate with signal recognition particle and trigger factor. *Mol. Microbiol.* **25**:53–64.
25. van Dijk, J. M., A. de Jong, H. Smith, S. Bron, and G. Venema. 1991. Non-functional expression of *Escherichia coli* signal peptidase I in *Bacillus subtilis*. *J. Gen. Microbiol.* **137**:2073–2083.
26. van Dijk, J. M., A. de Jong, G. Venema, and S. Bron. 1995. Identification of the potential active site of the signal peptidase SipS of *Bacillus subtilis*: structural and functional similarities with LexA-like proteases. *J. Biol. Chem.* **270**:3611–3618.
27. van Wely, K. H. M., J. Swaving, C. P. Broekhuizen, M. Rose, W. J. Quax, and A. J. M. Driessen. 1999. Functional identification of the product of the *Bacillus subtilis* *yvaL* gene as a SecG homologue. *J. Bacteriol.* **181**:1786–1792.
28. van Wely, K. H., J. Swaving, R. Freudl, and A. J. Driessen. 2001. Translocation of proteins across the cell envelope of gram-positive bacteria. *FEMS Microbiol. Rev.* **25**:437–454.
29. von Heijne, G. 1998. Life and death of a signal peptide. *Nature* **396**:111–113.
30. West, J. T., W. Estacio, and L. Márquez-Magaña. 2000. Relative roles of the *fla/che* P_A, P_{D-3}, and P_{sigD} promoters in regulating motility and *sigD* expression in *Bacillus subtilis*. *J. Bacteriol.* **182**:4841–4848.