

Analysis of Gene Islands Involved in Methanopterin-Linked C₁ Transfer Reactions Reveals New Functions and Provides Evolutionary Insights

Marina G. Kalyuzhnaya,¹ Natalia Korotkova,¹† Gregory Crowther,¹ Christopher J. Marx,¹‡
Mary E. Lidstrom,^{1,2} and Ludmila Chistoserdova^{1*}

Department of Chemical Engineering¹ and Department of Microbiology,²
University of Washington, Seattle, Washington 98195

Received 24 January 2005/Accepted 15 March 2005

In this study, the occurrence and chromosomal clustering of genes encoding C₁ transfer reactions linked to tetrahydromethanopterin (H₄MPT) were analyzed in a variety of proteobacteria and in representatives of the Planctomycetes via genomic analysis or via partial sequencing by cosmid walking. Although a tendency for clustering was found common for the genes of interest, significant variations in gene order and the degree of clustering were uncovered both between and within different groups of Proteobacteria and between Proteobacteria and Planctomycetes. Phylogenetic analyses suggested that the evolution of genes encoding H₄MPT-linked reactions in Proteobacteria involved lateral transfers within Proteobacteria and possibly between Proteobacteria and other phyla. Gene cluster comparisons revealed a number of novel genes potentially involved in the C₁ transfer reactions, and these were analyzed by mutation and expression analyses. Four genes, a homolog of *pabB*, and three genes conserved between methanogenic Archaea and Bacteria possessing H₄MPT-linked functions, *orfY*, *orf1*, and *afpA* were shown to be involved in formaldehyde oxidation/detoxification, as judged by specific mutant phenotypes. In particular, *pabB* contributes to the biosynthesis of *para*-aminobenzoic acid, a precursor of both tetrahydrofolate and H₄MPT, and *afpA* apparently encodes a novel dihydromethanopterin reductase, based on mutant complementation experiments.

One of the major breakthroughs in the understanding of methylotrophy in *Bacteria* during the recent decade, the recognition of the tetrahydromethanopterin (H₄MPT)-linked pathway for formaldehyde oxidation as a major C₁ oxidation pathway, was due to a serendipitous discovery of a cluster of genes in *Methylobacterium extorquens* AM1 that are homologous to the genes involved in methanogenesis in *Archaea* (7). It has since been demonstrated that this pathway is nearly ubiquitous in gram-negative methylotrophs (26). More recently, the pathway's presence has been expanded beyond methylotrophs (20), and even beyond *Proteobacteria* into the *Planctomycetes* (5, 10). Although phylogenetic analysis has argued against recent lateral transfer of the genes in question between methanogenic *Archaea* and *Proteobacteria* (5), the history of these genes in *Bacteria* remains poorly understood. Although some congruence has been observed between the phylogenetic positions of the respective species within *Proteobacteria* and phylogenies of genes involved in H₄MPT-linked reactions (13, 15), the relationships of the latter are not always well resolved and sometimes are complicated by the presence of multiple gene copies. In the present study we attempted a more comprehensive analysis of gene islands in *Bacteria* involved in H₄MPT-linked C₁ transfers via analysis of available genomic sequences,

via expanding gene databases by cosmid walking, and via mutagenesis and phylogenetic analysis. The following major objectives were pursued: (i) to determine whether gene clustering patterns are conserved within specific groups of *Bacteria*, (ii) to determine whether such patterns reflect phylogenies of the respective genes, (iii) to determine whether these patterns reflect organismal phylogenies, and (iv) to attempt identification of novel C₁ transfer genes based on the clustering patterns.

MATERIALS AND METHODS

Sequence analysis. The genome of *Methylobacterium extorquens* AM1 was analyzed as described in reference 3. The genome of *Methylobacillus flagellatus* KT was analyzed as described in reference 14. The genomes of *Methylococcus capsulatus* Bath, *Rhodopirellula baltica*, *Gemmata obscuriglobus*, and *Burkholderia xenovorans* LB400 were analyzed as described in reference 5). The sequences of interest were retrieved from the genome of *Methylobium petroleophilum* PM1 (http://genome.jgi-psf.org/draft_microbes/metpe/metpe.home.html) via BLAST analyses using the sequences of *B. xenovorans* as queries as described previously (5). Partial sequences of gene islands encoding H₄MPT-linked C₁ transfer reactions from *Xanthobacter autotrophicus*, *Hyphomicrobium zavarzinii*, *Methylosinus* sp. strain LW2, *Methylomonas* sp. strain LW13, and *Methylomicrobium* sp. strain AMO were obtained via cosmid walking as described in reference 5. Environmental sequences of interest were detected as described in reference 14.

Phylogenetic analysis. For phylogenetic analyses, the PHYLIP package (9) was used. Distance and parsimony methods were used, and 100 bootstrap analyses were performed. Concatenated polypeptide sequences were generated as follows. The respective polypeptide sequences were translated from the respective cosmid sequences or retrieved from the respective genomic databases as described in reference 5. Separate polypeptide sequences (or truncated sequences) were aligned by using the CLUSTAL W program (23), and all of the sequences were truncated to be of the same length. The truncated polypeptide sequences of each organism were then fused together. To clarify the position of *M. flagellatus* sequences, the following polypeptides were concatenated in the following order (ranging in length from 1,410 to 1,544 amino acid residues): OrfY-Mch-Orf5-Orf7-Fae-Orf17. To clarify the position of *X. autotrophicus* se-

* Corresponding author. Mailing address: 231 Wilcox Hall, Box 352125, University of Washington, Seattle, WA 98195. Phone: (206) 543-6683. Fax: (206) 616-5721. E-mail: milachis@u.washington.edu.

† Present address: Department of Microbiology, University of Washington, Seattle, WA 98195.

‡ Present address: Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138.

TABLE 1. Genes mutated in this study

Gene ^a	Known or predicted function	C ₁ growth defect ^b
<i>mxoE1</i> *	Unknown function in methanol oxidation	No*
<i>mxoD1</i> *	Methanol oxidation, interaction between methanol dehydrogenase and cytochrome c _L	No*
<i>mxoD2</i> *	Methanol oxidation, interaction between methanol dehydrogenase and cytochrome c _L	No*
<i>pabB</i> *	Component 1 of PABA synthase, biosynthesis of H ₄ F and H ₄ MPT	Yes*
<i>pcbD</i> *	Pterin-4a-carbinolamine dehydratase, regeneration of BH ₄	No*
<i>pts</i> *	6-Pyruvoyl tetrahydropterin synthase, biosynthesis of BH ₄	No*
<i>fae2</i> *	Formaldehyde-activating enzyme	No*
<i>fae3</i> *	Formaldehyde-activating enzyme	No*
<i>fae2 fae3</i>		No*
<i>orf1</i> *	ATP-utilizing protein, ATP grasp family	Yes [†]
<i>afpA</i> *	Archaeal flavoprotein, FMN-binding electron transfer protein	Yes [†]
<i>orf5</i> [†]	Unknown function in biosynthesis of H ₄ MPT	Yes [†]
<i>mtdB</i> [†]	Methylene-H ₄ MPT dehydrogenase	Yes [†]
<i>mptG</i> [†]	Ribofuranosylaminobenzene 5-phosphate synthase, biosynthesis of H ₄ MPT	Yes [†]
<i>orfY</i> *	Predicted ATP-dependent carboligase	Yes [†]

^a *, Function is predicted based on sequence similarity; †, function has been characterized in *M. extorquens*. See text for references.

^b *, Mutated in *M. extorquens*; †, mutated in *M. flagellatus*.

quences, the following polypeptides were concatenated in the following order (ranging in length from 2,333 to 2,401 amino acid residues): Mch-Orf5-Orf9-FhcC-FhcD-FhcA-FhcB. In the case of *G. obscuriglobus*, two homologs of *fhcD* are found in the genome (53 and 39% identity with the *fhcD* from *M. extorquens*, respectively, at the amino acid level). The one with the higher identity to the proteobacterial FhcDs was used in the analysis. The concatenated sequences were realigned by using the CLUSTAL W program, and the alignments were manually curated.

Mutant generation and mutant complementation. Mutations in the following genes of *M. extorquens* were generated essentially as described in reference 18: *mxoE1*, *mxoD2*, *mxoD3*, *pabB*, *pcbD*, *pts*, *fae2*, and *fae3* (Table 1). A double *fae2 fae3* mutant was generated essentially as described in reference 18. Mutants in the following genes of *M. flagellatus* were generated by using the pCM184 suicide vector (18): *orf5*, *orfY*, *mtdB*, *mptG*, *orf1*, and *afpA* (Table 1). For mutant complementation, the previously described broad-host-range expression vector pCM80 (17) was used.

Mutant characterization. Growth characteristics of the *M. extorquens* mutants were tested on methanol-supplemented or succinate-supplemented media or on succinate-supplemented media in the presence of methanol vapors, as described earlier (6). To complement the *pabB* mutation, a range of concentrations (1 nM to 1 mM) of *para*-aminobenzoic acid (PABA) was used as a supplement. To characterize the phenotypes of the *M. flagellatus* mutants, a standard medium with added formaldehyde was used as described previously (4).

RESULTS

Gene islands encoding H₄MPT-linked C₁ transfer reactions in Bacteria. A cluster of genes with homologs in methanogenic archaea (the “archaeal-like” gene island) has been characterized in *M. extorquens*, and most of the “archaeal-like” genes have been shown to be involved in oxidation of formaldehyde to formate via H₄MPT-linked derivatives (6, 7, 27). Analysis of the yet-incomplete genome of *M. extorquens* revealed no other “archaeal” gene islands or gene homologs in the chromosome, with the exceptions of two distant homologs of *fae*, *fae2* and *fae3*, of unknown function (5). In the present study, we expanded analysis of genes involved in H₄MPT-linked reactions to include other proteobacterial species, as well as representatives of *Planctomycetes*. Genomic sequences are now available for three methylotrophic proteobacteria other than *M. extorquens*: *Methylococcus capsulatus* Bath (28), *Methylobium petroleophilum* PM1 (http://genome.jgi-psf.org/draft_microbes/metpe/metpe.home.html), and *Methylobacillus flagellatus* KT (http://genome.jgi-psf.org/draft_microbes/metfl/metfl.home.html). In addition, genomic sequences of some nonmethylotro-

phic bacteria were shown to contain genes involved in H₄MPT-linked C₁ transfers, i.e., the genome of *Burkholderia xenovorans* LB400 and the genomes of the three planctomycetes: *Gemmata obscuriglobus*, *Gemmata* Wa1-1, and *Rhodopirellula baltica* (5, 10, 20). We also expanded “archaeal” gene analysis in *Proteobacteria* by partially sequencing these chromosomal regions from *Methylomicrobium* sp. strain AMO, *Methylomonas* sp. strain LW13, *Hyphomicrobium zavarzinii*, *Xanthobacter autotrophicus*, and *Methylosinus* sp. strain LW2 (5). In addition, clusters of genes homologous to the “archaeal” genes in *M. extorquens* have been identified in the environmental shotgun sequence database generated from the Sargasso Sea DNA (14). The newly identified gene clusters were aligned with the gene cluster in *M. extorquens* and analyzed in terms of gene order conservation (Fig. 1).

In general, we observed a high degree of gene order conservation in methylotrophic proteobacteria. However, various signs of chromosomal rearrangements particular to specific lineages were observed, such as insertions, deletions, and inversions. Analysis of the genomic sequence of *M. capsulatus*, a γ -proteobacterial methanotroph, revealed that the genes in question were not clustered together on the chromosome but formed three different gene islands separated by large distances on the chromosome. An additional homolog of *fhcA* was also present, one not linked to other C₁ genes (28). Gene order in the main island was highly conserved relative to the order of genes in *M. extorquens* between *fhcC* and *orf9*. However, *mtdB* was not found in the main island in *M. capsulatus*. Instead, two copies of *mtdB* were found as a separate island on the chromosome. There is an insertion of an open reading frame (*orf1*) between *orf17* and *orf9* of *M. capsulatus* not found in the *M. extorquens* island or anywhere in the almost complete genome of *M. extorquens*. This gene has homologs in archaeal genomes, and its translated product possesses kinase motifs, based on COG (for clusters of orthologous genes) analysis (<http://www.ncbi.nlm.nih.gov/COG/>), but nothing is known about its function. Although in *M. extorquens* three genes with no archaeal homologs (homologous to *mxoE* and *mxoD* genes) were found as part of a methylotrophy gene island, separating *orf9* from *orf19* to -22 in the *M. capsulatus* island, a homolog of *pabB*, a gene predicted to encode component I of PABA syn-




FIG. 1. Clusters of genes encoding reactions of the H₄MPT-linked C₁ transfer pathway in *Bacteria*. Genes for C₁ transfer reactions are shown in red; previously characterized genes for cofactor biosynthesis or not-yet-identified functions are shown in yellow. Genes with representatives mutated in the present study are shown in blue (C₁-negative phenotype) or purple (C₁-positive phenotype). Genes not involved in this study are left blank. Genes not contiguous on chromosomes are separated by double lines.

these, was found downstream of *orf9*, together with *orf21*. *orf22*, *orf19*, and *orf20* formed a separate gene island in the *M. capsulatus* chromosome. The partially sequenced gene islands from two other γ -proteobacterial methanotrophs, *Methylomicrobium* sp. strain AMO and *Methylomonas* sp. strain LW13 revealed a gene order similar to the one in the main island in *M. capsulatus* (Fig. 1); however, in *Methylomicrobium* sp. strain AMO, insertions of genes not conserved in other clusters were found (Fig. 1).

Analysis of the recently generated genomic sequence of a β -proteobacterial methylothermophile, *M. flagellatus* revealed the presence of two “archaeal-like” gene islands. The main island aligned very well with the main island of *M. capsulatus*; however, three insertions are present in the *M. flagellatus* island. The gene for *mtdB* is located between *mptG* and *orfY*, as in *M. extorquens*. Three genes are inserted between *orf7* and *fae*, encoding, respectively, transaldolase (*tal*), hexulosephosphate synthase (*hps*), and hexulosephosphate isomerase (*hpi*), enzymes of the ribulose monophosphate cycle for formaldehyde assimilation (1). Another novel gene, tentatively designated *pcbD*, is found downstream of *orf21* in *M. flagellatus*, which is

predicted to encode pterin-4a-carbinolamine dehydratase (an enzyme involved in regeneration of the cofactor tetrahydrobiopterin, BH₄ [24]), based on COG analysis. However, no nucleotide or amino acid conservation exists between characterized eukaryotic (24) and putative bacterial *pcbD* genes (National Center for Biotechnology Information). A *pcbD* homolog is present in a larger C₁ island of *M. extorquens*, separated by 16.5 kb from *fhcC* and by ca. 7 kb from *fch*, which encodes methenyl tetrahydrofolate (H₄F) cyclohydrolase, an enzyme involved in methylothermophily (3). A *pcbD* homolog in *M. capsulatus* is found ca. 7 kb apart from *fch*-*mtdA* pair, but no other C₁ genes are present in the vicinity. *MtdA* encodes methylene tetrahydrofolate dehydrogenase, which is also involved in methylothermophily in *M. extorquens* AM1 (3).

The second gene island of *M. flagellatus*, containing *orf19*, -20, and -22, aligns with the respective island in *M. capsulatus*. However, one additional gene is present upstream of *orf20*, having homologs in *Archaea*, encoding a conserved archaeal flavoprotein (*afpA*) (8). In addition to the two gene islands, three other archaeal gene homologs were found in the genome. One of them was highly similar to *fae* in the main gene

island, whereas two others were similar to the *fae* homologs, *fae2* and *fae3*, previously identified in *M. extorquens*.

We have recently identified gene islands involved in H₄MPT-linked C₁ transfers in the shotgun sequence library of environmental DNA from the Sargasso Sea that belong to as-yet-identified bacteria (14). One of these clusters is shown in Fig. 1. The gene order in this cluster shares many similarities with that for *M. capsulatus* and *M. flagellatus*.

Three partial cluster sequences from α -proteobacteria representing *Hyphomicrobium*, *Methylosinus*, and *Xanthobacter* were aligned with the cluster in *M. extorquens* (Fig. 1). A partial cluster of nine genes has been sequenced from another *Methylobacterium* species closely related to *M. extorquens* (16), showing identical gene order and a high degree of sequence conservation at the DNA level, and thus is not shown in Fig. 1. The partial gene islands from *H. zavarzinii* and *Methylosinus* sp. strain LW2 revealed a high degree of gene order conservation with the *Methylobacterium* species, with some exceptions. For example, the cluster from *H. zavarzinii* contained no insert of *mxg* gene homologs found in *M. extorquens*, whereas the cluster in *Methylosinus* sp. strain LW2 contained two *fae* homologs. In contrast, the gene order in the partial cluster from *X. autotrophicus* was strikingly different from all other known clusters: no *mptG* was present upstream of *mtdB* (a region of 5 kb has been sequenced [data not shown]), no *orfY* was present between *mtdB* and *mch*, and no genes were conserved in their order downstream of *orf5*. Instead, *orf5* was followed by *orf1* and *orf9*, downstream of which an inverted cluster of the *fhc* genes was found. Additional sequence of 15 kb downstream of *fhcB* did not reveal the presence of any other genes of interest.

The genomes of two other β -proteobacteria were involved in our analyses, those of closely related organisms belonging to *Burkholderiaceae*, *M. petroleophilum* PM1 and *B. xenovorans* LB400. Although *M. petroleophilum* can grow on some C₁ compounds (K. M. Scow and K. R. Hristova, unpublished data), *B. xenovorans* is not known to be a methylotroph (20). Analysis of the genome of *M. petroleophilum* revealed that most of the genes of interest were located in a single gene island. The gene order between *mptG* and *fae* was conserved with relation to the order in most of the islands discussed above, whereas the rest of the island was represented by an inversion of the cluster of genes represented by *afpA-orf17* in the environmental clone, in which *pabB* was missing. The *fhc* gene cluster was located downstream of *orf17* and was inverted relative to its position in other methylotrophs belonging to α -, β -, and γ -proteobacterial groups. A homolog of *fae2* from *M. extorquens* and *M. flagellatus* was found downstream of *fhcC*, followed by *pabB*. The C₁ transfer gene island in *B. xenovorans* had a very similar structure with the following exceptions. A cluster of *cbh* genes (encoding genes of the Calvin-Benson-Bassham cycle) was inserted between *orf20* and *orf19* (20; data not shown in Fig. 1), no *fae2* homolog was present in the cluster or anywhere in the genome, a *pabB* homolog was present elsewhere in the genome, and a *pcbD* homolog was present upstream of *mptG*.

As is seen in Fig. 1, C₁ transfer gene clustering in *Planctomycete* genomes is much looser compared to proteobacterial genomes. However, the *mch-orf5* order was conserved in the genome of *R. baltica*, and the order of *orfY-mch-orf5-orf7* was conserved in *G. obscuriglobus*. Besides this, clusters of three

(*fhcADC*) and two (*fhcDC*) genes are found in the genomes of *R. baltica* and *G. obscuriglobus*, respectively (Fig. 1). However, no homolog of *fhcB* was detected in *R. baltica* (10), whereas in *G. obscuriglobus* two homologs of *fhcD* were identified that greatly diverged in their sequences (46% identity between the translated amino acid sequences). The low coverage of the genome of *Gemmata* sp. strain Wa1-1 (5) allowed detection of only a few gene clusters, and these were the same as in *G. obscuriglobus* (data not shown). In all *Planctomycete* genomes, *orf19* was linked to *mptG* and *orf9* was linked to *orf21*, showing novel clustering types. In *R. baltica*, a *pabB* homolog was found linked to the *mch-orf5* cluster, whereas in *G. obscuriglobus*, a homolog of *pts*, a gene potentially encoding an enzyme involved in BH₄ biosynthesis, 6-pyruvoyl-tetrahydropterin synthase, was found downstream of *orf7*. *fae* genes in *Planctomycetes*, unlike those in *Proteobacteria*, were found linked to genes revealing homology to genes encoding either MtdA or MtdB enzymes (44 to 49% and 28 to 30% identity at the amino acid level, respectively). MtdA and MtdB have related sequences (ca. 30% identity), MtdA being a NADP-specific dehydrogenase that utilizes both methylene-H₄F and methylene-H₄MPT, whereas MtdB uses both NAD and NADP but is specific to H₄MPT (3, 12).

Phylogenetic analysis. Analysis of these gene clusters has revealed a great deal of divergence in gene order within the groups of the α -proteobacterial and β -proteobacterial species. The gene order in the *M. flagellatus* cluster was more similar to that in γ -proteobacterial clusters than to that in clusters of other β -proteobacteria, whereas the gene order in *X. autotrophicus* was unique within the bacterial set sampled. These distinct organizations may indicate that the C₁ clusters in these strains have a separate evolutionary history relative to other methylotrophs within their own respective classes of proteobacteria. To test this hypothesis, we performed phylogenetic analyses with the subsets of sequences available. In accordance with our previous experience, due to the divergence of the genes in question, analysis of single polypeptide sequences often results in unresolved branching patterns, reflected by low bootstrap values. However, using concatenated polypeptide sequences results in increased phylogenetic signal and thus a better resolution in phylogenetic analyses (14). To assess the phylogenetic position of the “archaeal” genes in *M. flagellatus* relative to other bacterial species, we used concatenates of six polypeptide sequences as described in Materials and Methods. Both distance and parsimony analyses resulted in identical branching patterns, placing *M. flagellatus* sequences with the sequences of γ -proteobacterial methylotrophs, with high bootstrap values, while the sequences of other β -proteobacteria, *B. xenovorans* and *M. petroleophilum*, branched together (Fig. 2A). Such a branching pattern suggests separate evolutionary histories for the two types of C₁ clusters found in β -proteobacteria. Similar analyses were performed with a subset of sequences to obtain insights into the position of the *X. autotrophicus* sequences within the group tested. Concatenates of seven polypeptide sequences were used as described in Materials and Methods. Distance and parsimony analyses produced identical branching patterns in phylogenetic trees (Fig. 2B). In these trees, the sequences of *X. autotrophicus* clearly separated from the sequences of *M. extorquens* and from all




FIG. 2. Phylogenetic analysis of concatenated polypeptides representing subsets of H₄MPT-linked C₁ transfer genes (see Materials and Methods). Bootstrap values are shown for distance/parsimony analyses. (A) Tree resolving the position of the *M. flagellatus* sequences using six sequences in the concatenates. (B) Tree resolving the position of the *X. autotrophicus* sequences using seven sequences in the concatenates. A, P, α, β, and γ indicate *Archaea*, *Planctomycetes*, and α-, β-, and γ-*Proteobacteria*, respectively. AACY010825 is the sequence recovered from the Sargasso Sea metagenome.

other sequences, suggesting that the genes in question have distinct evolutionary histories within α-proteobacteria as well.

Functional analysis of novel genes conserved in C₁ transfer clusters. (i) Mutagenesis in *M. extorquens*. An insert of three nonarchaeal genes is present in the gene cluster of *M. extorquens* (Fig. 1). One of these genes is homologous to *mxoE*, a gene of unknown function in the gene cluster (*mox* cluster) involved in methanol oxidation (3), while two others are both homologous to *mxoD*, another gene in the *mox* cluster that has been shown to be involved in interaction between methanol dehydrogenase and cytochrome *c_L* (25). We mutated all three *mox* gene homologs to test their involvement in C₁ metabolism. The resulting mutants did not reveal any visible defects in growth on methanol (Table 1); thus, the roles of these *mxo* gene homologs remain unknown.

Another gene persistently present in methylotroph genomes and in some cases in the C₁ transfer clusters is the gene tentatively designated *pcbD*, based on COG analysis. Known *pcbD* genes encode pterin-4a-carbinolamine dehydratase (24), an enzyme involved in regeneration of tetrahydrobiopterin (BH₄). This enzyme also possesses a second function, as a regulatory protein (24). A gene encoding a homolog of an enzyme involved in the BH₄ biosynthesis pathway, 6-pyruvoyl tetrahydropterin synthase (*pts* [24]), is clustered with the C₁ transfer genes in *G. obscuriglobus*, and this gene is recognizable in many of the sequenced genomes. We used *M. extorquens* as a model to test for possible involvement of these genes in C₁ metabolism but did not observe any deficiency for growth on methanol for mutants lacking either *pcbD* or *pts* (Table 1).

We have previously identified two distant homologs of *fae* in

the genome of *M. extorquens*, tentatively designated *fae2* and *fae3*, which were not part of any C₁ clusters (5). Homologs of both *fae2* and *fae3* were identified in the genome of *M. flagellatus*, a close homolog of *fae3* was identified in *R. baltica* (10), and a homolog of *fae2* was identified in the genome of *M. petroleophilum*. In the latter case, *fae2* was part of the C₁ transfer cluster (Fig. 1). We mutated *orf2* and *orf3* in *M. extorquens*, and we also obtained a double *fae2 fae3* mutant. All three mutants grew with wild-type characteristics on both succinate and methanol (Table 1). Overexpression of either *fae2* or *fae3* in the background of a *fae* mutant of *M. extorquens* (27) did not lead to complementation of the phenotype of the mutant (14, the present study).

Analysis of H₄MPT-linked C₁ transfer gene islands also revealed the presence of a nonarchaeal gene, a homolog of *pabB*. This gene encodes component 1 of PABA synthase, the enzyme involved in biosynthesis of PABA, the precursor in biosyntheses of both H₄F and H₄MPT (11). To test if mutants in *pabB* would reveal a specific phenotype, we mutated a *pabB* homolog in *M. extorquens*. This gene is not a part of the C₁ island shown in Fig. 1, and it was identified in the genomic database as the highest hit with *pabB* homologs present in other C₁ islands. Null mutants in *pabB* were selected on succinate-supplemented medium with or without added PABA (1 mM). *pabB* mutants grew on succinate or methanol plates supplemented with 1 mM PABA but not on either substrate in the absence of PABA supplementation. The presence of wild-type on the same plates as *pabB* mutants permitted slow growth of the mutants in a manner consistent with cross-feeding. This phenomenon likely explains our success in isolating

pabB mutants in the absence of PABA: release of PABA from lysed cells and/or single-crossover exconjugants may have obviated the need for PABA supplementation. We determined the minimal PABA supplement concentration to allow growth of the mutants on succinate to be ca. 10 nM. We then tested the influence of added methanol on growth of *pabB* mutants in the presence of the minimal supplementary concentration of PABA. The addition of ca. 1 mM methanol caused significant growth inhibition of the *pabB* mutant (Table 1). Methanol sensitivity has been shown to be a trait of mutants defective in H₄MPT-linked formaldehyde oxidation (6, 12, 19, 27). Increasing the concentration of PABA supplement gradually alleviated methanol sensitivity, and the mutants could grow on methanol with wild-type rate in the presence of 1 mM PABA.

(ii) **Mutagenesis in *M. flagellatus*.** Two novel genes were found conserved in a number of proteobacterial gene clusters that were not identifiable in the genome of *M. extorquens* (3) but have homologs in *Archaea*: *orf1* and *afpA*. No function has been proposed for homologs of *orf1* in *Archaea*; however, AfpA in *Archaeoglobus fulgidus* has been proposed to be involved in electron transfer between ferredoxin and the CO dehydrogenase/acetyl coenzyme A synthase (CODH/ACS) complex (8). The possible involvement of *orf1* and *afpA* in the C₁ transfer pathway was tested via mutagenesis in *M. flagellatus*. Previously, *mch* from this organism has been mutated, and the phenotype of this mutant suggested that the pathway was not essential for methylotrophy in this organism but played an auxiliary function in formaldehyde detoxification (4). However, in the main model organism for studying H₄MPT-linked C₁ transfer reactions, *M. extorquens*, we demonstrated that mutations in different genes involved in the pathway resulted in phenotypes that differed in the degree of formaldehyde sensitivity (6, 19). To provide a reference database for characterization of the mutants in genes of unknown functions in *M. flagellatus*, we also mutated genes previously characterized in *M. extorquens*, i.e., *mptG*, *mtdB*, and *orf5*. We also tested the function of *orfY* by mutation. The function of this gene remained unknown in *M. extorquens*. We had originally reported obtaining knockout mutants in this gene in *M. extorquens* (7), but later tests have shown these mutants were single-crossover recombinants (unpublished data). Our recent attempts to generate *orfY*-null mutants in *M. extorquens* were unsuccessful, suggesting that this gene is essential in this organism.

Mutants of *M. flagellatus* were generated via allelic exchange as described in Materials and Methods. Double-crossover mutations were generated in all of the genes tested, as judged by diagnostic PCR (data not shown). Their sensitivity to formaldehyde was investigated as previously described (4) and compared to the phenotypes of the wild-type strain and the previously characterized *mch* mutant, respectively. All of the mutants investigated showed increased formaldehyde sensitivity similar to the previously described mutant in *mch*; however, the new mutants showed slightly lower formaldehyde sensitivity (between 2.5 and 3 mM versus 2 mM for the *mch* mutant) (Table 1). The functions of *mtdB* and *mptG* have been determined (12, 22), and a function has been proposed for *orf5* in H₄MPT biosynthesis (6). Based on the phenotypes of the new mutants and also on gene location within the formaldehyde oxidation islands, *orfY*, *orf1*, and *afpA* are likely to be involved in the H₄MPT-linked pathway as well. To test whether *orfY*

from *M. extorquens* could complement *orfY* mutants of *M. flagellatus*, we expressed *orfY* from *M. extorquens* in the *orfY* mutant of *M. flagellatus* and tested the recombinants for formaldehyde sensitivity. The recombinants carrying the copy of the *M. extorquens* gene in *trans* showed wild-type sensitivity toward formaldehyde.

Expression of *orf1* and *afpA* in the *dmrA* mutant of *M. extorquens*. Although *orf1* and *afpA* genes are present in many of the “archaeal” gene islands characterized here, neither of these genes is identifiable in the almost-complete genome of *M. extorquens*. Conversely, a gene has been identified in *M. extorquens* that is involved in the last step of the H₄MPT biosynthetic pathway, a gene that encodes dihydromethanopterin reductase (*dmrA* [2, 21, 22]), that is not recognizable in the genomes of *M. capsulatus*, *M. flagellatus*, *B. xenovorans*, and *M. petroleophilum* or in the planctomycete genomes. Mutagenesis of *orf1* and *afpA* in *M. flagellatus* suggests their involvement in some step of the H₄MPT-linked C₁ transfer pathway (see above). To test whether *orf1* or *afpA* could fulfill the function of *dmrA*, we expressed the *orf1* genes from *M. capsulatus* and *Methylomonas* sp. strain LW13, and the *afpA* genes from *M. flagellatus* and *B. xenovorans* in the *dmrA* mutant of *M. extorquens* and tested the transconjugants for growth on methanol and methanol sensitivity. Although expression of *orf1* genes had no effect on the phenotype of the *dmrA* mutant, expression of *afpA* genes resulted in reversal of the *dmrA* mutant phenotype to the wild-type phenotype, i.e., AfpA could fulfill the function of DmrA. Since the substrate for the DmrA reaction, dihydromethanopterin, is not commercially available, we were not able to confirm the function of AfpA via enzyme activity measurements.

DISCUSSION

In this study we investigated the structure of clusters of genes encoding reactions of H₄MPT-linked C₁ transfer and correlated these with specific bacterial groups. We demonstrated that, whereas tight clustering of the genes in question is characteristic of *Proteobacteria* and a great deal of gene order conservation occurs, the cluster structure and the degree of clustering vary greatly both within the specific groups and between the groups. We have previously reported that gene sequences of *Planctomycetes* involved in H₄MPT-linked reactions diverge significantly from their proteobacterial counterparts and occupy an intermediate phylogenetic position between *Archaea* and *Proteobacteria* (5). Here we demonstrate that the degree of gene clustering in *Planctomycetes* is also intermediate between *Archaea* and *Proteobacteria*: whereas almost no clustering for the genes in question occurs in *Archaea*, a certain degree of clustering takes place in *Planctomycetes*. However, only two instances of a cluster structure common to both *Planctomycetes* and *Proteobacteria* were found (*mch-orf5* and *fhcDC*). The results of this analysis point toward long and separate histories of the genes in question in *Archaea*, *Proteobacteria*, and *Planctomycetes*.

In a previous study, we argued against a single recent event of lateral transfer of genes for H₄MPT-linked reactions between *Archaea* and *Bacteria* (5). However, the data presented here suggest that lateral transfers within groups—specifically, within *Proteobacteria*—have likely occurred. The difference in

gene cluster structure and in the phylogenetic positions of β -proteobacterial genes represented by *M. flagellatus* on one hand and by *B. xenovorans* and *M. petroleophilum* on the other may have two alternative explanations, both involving lateral transfers. First, an ancestor of *M. flagellatus* may have lost the gene cluster typical of β -proteobacteria and subsequently acquired a new cluster via a lateral transfer from a γ -proteobacterial methanotroph. Second, the genes found in modern representatives of *Burkholderiaceae* are a result of lateral transfer into the ancestor of both *Burkholderia* and *Methylobium* from a source not known at this time. Similarly, the history of the genes for H₄MPT-linked reactions in *X. autotrophicus* may involve a lateral transfer from an unknown source since they diverge significantly in both sequence and clustering patterns from genes in other α -proteobacteria. It is interesting that, whereas the cluster structure found in environmental clones from the Sargasso Sea is similar to that from γ -proteobacteria and *M. flagellatus*, gene sequences diverge significantly from both and form a separate branch on phylogenetic trees (14). It is possible that the clusters found in the Sargasso Sea group and γ - and β -proteobacterial methyloproteobacteria have descended from a single ancestral cluster but were exposed to different types of selective pressures, resulting in great sequence divergence.

Clustering of H₄MPT-linked C₁ transfer genes on the chromosome of *M. extorquens* has been advantageous for gaining insights into details of C₁ metabolic pathways, including gene discovery and identification. Involving more species in gene cluster analyses as described here led to the discovery of additional genes potentially involved in the H₄MPT-linked C₁ transfer pathway (Table 1), at least in some of the organisms producing methanopterin or its derivatives. *orfY* is predicted to encode an ATP-grasp family protein, possibly a carboxylase, in accordance with COG analysis. Its role in H₄MPT-linked C₁ transfer remains unknown. Since all of the genes for the catalytic reactions in the pathway are known, *orfY* may be involved in either cofactor biosynthesis or regulation of the pathway. The presence of *orfI* in a number of C₁ transfer gene clusters (Fig. 1) and the phenotype of the *orfI* mutant in *M. flagellatus* suggest this gene is also involved in the pathway. However, *orfI* is not present in the almost complete genome of *M. extorquens*. This may imply either that *orfI* is involved in a reaction not essential in *M. extorquens* (for example, a cofactor modification) or that a nonhomologous substitution for this gene is present in *M. extorquens*. Our data point toward *pabB* being involved in the biosynthesis of PABA, a precursor of both H₄F and H₄MPT. Based on mutant analysis in *M. extorquens*, higher PabB activity is required during growth on C₁ compounds compared to multicarbon compounds, a finding in agreement with previous data on the major role of the H₄MPT-linked formaldehyde oxidation pathway in C₁ oxidation. *afpA* appears to encode a novel dihydromethanopterin reductase. Ding and Ferry have recently purified an AfpA homolog from *A. fulgidus* and demonstrated that it is an FMN-binding electron carrier protein (8). These authors also analyzed the distribution of AfpAs in prokaryotes based on the genomic sequences available at the time and concluded that these were restricted to methanogenic archaea. Ding and Ferry proposed a function for AfpA in an electron transport chain in *A. fulgidus* in which electrons originating from the CODH/ACS complex are trans-

ferred to AfpA mediated by ferredoxin, thus assigning a role for AfpA in energy generation. We demonstrated here that *afpA* homologs are present in the C₁ gene islands of most of the bacteria synthesizing H₄MPT or its derivatives. Our data on complementation of the *dmrA* mutant, as well as the absence of CODH/ASC in the bacteria in question, suggest that *afpA* homologs perform an alternative physiological function in the biosynthesis of H₄MPT. In methanogenic *Archaea* not relying on CODH/ACS complex for energy generation (such as *M. thermoautotrophicus* or *M. jannaschii*), *afpA* homologs may also encode DMR enzymes. *M. extorquens* possesses a DMR that is a homolog of dihydrofolate reductase, encoded by *dmrA* (2, 21, 22). However, thus far, *dmrA* is unique to *M. extorquens* and not found in the genomes of other *Bacteria* and *Archaea* synthesizing H₄MPT. Interestingly, the genome of *M. capsulatus* lacks both *dmrA* and *afpA*, suggesting that yet another nonhomologous DMR may exist.

ACKNOWLEDGMENTS

This study was supported by the NSF Microbial Observatories program (MCB-0131957) and in part by a grant from the NIH (GM36296).

The Joint Genome Institute is acknowledged for sequencing the genomes of *M. flagellatus* and *M. petroleophilum*, and The Institute for Genomic Research is acknowledged for early release of the genome sequence of *G. obscuriglobus* (all three projects are funded by the DOE).

REFERENCES

1. Anthony, C. 1982. Biochemistry of methyloproteobacteria. Academic Press, London, England.
2. Caccamo, M. A., C. M. Malone, and M. E. Rasche. 2004. Biochemical characterization of a dihydromethanopterin reductase involved in tetrahydromethanopterin biosynthesis in *Methylobacterium extorquens* AM1. *J. Bacteriol.* **186**:2068–2073.
3. Chistoserdova, L., S.-W. Chen, A. Lapidus, and M. E. Lidstrom. 2003. Methyloproteobacteria in *Methylobacterium extorquens* AM1 from a genomic point of view. *J. Bacteriol.* **185**:2980–2987.
4. Chistoserdova, L., L. Gomelsky, J. A. Vorholt, M. Gomelsky, Y. D. Tsygankov, and M. E. Lidstrom. 2000. Analysis of two formaldehyde oxidation pathways in *Methylobacillus flagellatus* KT, a ribulose monophosphate cycle methyloproteobacter. *Microbiology* **146**:233–238.
5. Chistoserdova, L., C. Jenkins, M. G. Kalyuzhnaya, C. J. Marx, A. Lapidus, J. A. Vorholt, J. T. Staley, and M. E. Lidstrom. 2004. The enigmatic planktonocytes may hold a key to the origins of methanogenesis and methyloproteobacteria. *Mol. Biol. Evol.* **21**:1234–1241.
6. Chistoserdova, L., M. E. Rasche, and M. E. Lidstrom. 2005. Novel dephosphotetrahydromethanopterin biosynthesis genes discovered via mutagenesis in *Methylobacterium extorquens* AM1. *J. Bacteriol.* **187**:2508–2512.
7. Chistoserdova, L., J. A. Vorholt, R. K. Thauer, and M. E. Lidstrom. 1998. C₁ transfer enzymes and coenzymes linking methyloproteobacteria and methanogenic archaea. *Science* **281**:99–102.
8. Ding, Y. H., and J. G. Ferry. 2004. Flavin mononucleotide-binding flavoprotein family in the domain *Archaea*. *J. Bacteriol.* **186**:90–97.
9. Felsenstein, J. 2003. Inferring phylogenies. Sinauer Associates, Inc., Sunderland, Mass.
10. Glöckner, F. O., M. Kube, M. Bauer, H. Teeling, T. Lombardot, W. Ludwig, D. Gade, A. Beck, K. Borzym, K. Heitmann, R. Rabus, H. Schlesner, R. Amann, and R. Reinhardt. 2003. Complete genome sequence of the marine planktonic *Pirellula* sp. strain 1. *Proc. Natl. Acad. Sci. USA* **100**:8298–8303.
11. Graham, D. E., and R. H. White. 2002. Elucidation of methanogenic coenzyme biosyntheses: from spectroscopy to genomics. *Nat. Prod. Rep.* **19**:133–147.
12. Hagemeier, C. H., L. Chistoserdova, M. E. Lidstrom, R. K. Thauer, and J. A. Vorholt. 2000. Characterization of a second methylene tetrahydromethanopterin dehydrogenase from *Methylobacterium extorquens* AM1. *Eur. J. Biochem.* **267**:3762–3769.
13. Kalyuzhnaya, M. G., M. E. Lidstrom, and L. Chistoserdova. 2004. Utility of environmental probes targeting ancient enzymes: methyloproteobacteria detection in Lake Washington. *Microb. Ecol.* **48**:463–472.
14. Kalyuzhnaya, M. G., O. Nercessian, A. Lapidus, and L. Chistoserdova. 8 April 2005, posting date. Fishing for biodiversity: novel methanopterin-linked C₁ transfer genes deduced from the Sargasso Sea metagenome. *Environ. Microbiol.* doi:10.1111/j.1462-2920.2005.00798.x.

15. Kalyuzhnaya, M. G., O. Nercessian, M. E. Lidstrom, and L. Chistoserdova. Development and application of polymerase chain reaction primers based on *fhcD* for environmental detection of methanopterin-linked C₁-metabolism in bacteria. *Environ. Microbiol.*, in press.
16. Kayser, M. F., Z. Ucurum, and S. Vuilleumier. 2002. Dichloromethane metabolism and C1 utilization genes in *Methylobacterium* strains. *Microbiology* **148**:1915–1922.
17. Marx, C. J., and M. E. Lidstrom. 2001. Development of improved versatile broad-host-range vectors for use in methylotrophs and other gram-negative bacteria. *Microbiology* **147**:2065–2075.
18. Marx, C. J., and M. E. Lidstrom. 2002. A broad-host-range *cre-lox* system for antibiotic marker recycling in gram-negative bacteria. *BioTechniques* **33**:1062–1067.
19. Marx, C. J., L. Chistoserdova, and M. E. Lidstrom. 2003. Formaldehyde-detoxifying role of the tetrahydromethanopterin-linked pathway in *Methylobacterium extorquens* AM1. *J. Bacteriol.* **185**:7160–7168.
20. Marx, C. J., J. A. Miller, L. Chistoserdova, and M. E. Lidstrom. 2004. Multiple formaldehyde oxidation/detoxification pathways in *Burkholderia fungorum* LB400. *J. Bacteriol.* **186**:2173–2178.
21. Marx, C. J., B. N. O'Brien, J. Breezee, and M. E. Lidstrom. 2003. Novel methylotrophy genes of *Methylobacterium extorquens* AM1 identified by using transposon mutagenesis including a putative dihydromethanopterin reductase. *J. Bacteriol.* **185**:669–673.
22. Rasche, M. E., S. A. Havemann, and M. Rosenzvaig. 2004. Characterization of two methanopterin biosynthesis mutants of *Methylobacterium extorquens* AM1 by use of a tetrahydromethanopterin bioassay. *J. Bacteriol.* **186**:1565–1570.
23. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight Matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
24. Thöny, B., G. Auerbach, and N. Blau. 2000. Tetrahydrobiopterin biosynthesis, regulation and functions. *Biochem. J.* **347**:1–16.
25. Toyama, H., H. Inagaki, K. Matsushita, C. Anthony, and O. Adachi. 2003. The role of the MxaD protein in the respiratory chain of *Methylobacterium extorquens* during growth on methanol. *Biochim. Biophys. Acta* **1647**:372–375.
26. Vorholt, J. A., L. Chistoserdova, S. M. Stolyar, R. K. Thauer, and M. E. Lidstrom. 1999. Distribution of tetrahydromethanopterin-dependent enzymes in methylotrophic bacteria and phylogeny of methenyl tetrahydromethanopterin cyclohydrolases. *J. Bacteriol.* **181**:5750–5757.
27. Vorholt, J. A., C. J. Marx, M. E. Lidstrom, and R. K. Thauer. 2000. Novel formaldehyde-activating enzyme in *Methylobacterium extorquens* AM1 required for growth on methanol. *J. Bacteriol.* **182**:6645–6650.
28. Ward, N., (swsl)O. Larsen, J. Sakwa, L. Bruseth, H. Khouri, A. S. Durkin, G. Dimitrov, L. Jiang, D. Scanlan, K. H. Kang, M. Lewis, K. E. Nelson, B. Methe, M. Wu, J. F. Heidelberg, I. T. Paulsen, D. Fouts, J. Ravel, H. Tettelin, Q. Ren, T. Read, R. T. DeBoy, R. Seshadri, S. L. Salzberg, H. B. Jensen, N. K. Birkeland, W. C. Nelson, R. J. Dodson, S. H. Grindhaug, I. Holt, I. Eidhammer, I. Jonassen, S. Vanaken, T. Utterback, T. V. Feldblyum, C. M. Fraser, J. R. Lillehaug, and J. A. Eisen. 2004. Genomic insights into methanotrophy: the complete genome sequence of *Methylococcus capsulatus* (Bath). *PLoS Biol.* **2**:e303.