

## Expression of *Pseudomonas aeruginosa aer-2*, One of Two Aerotaxis Transducer Genes, Is Controlled by RpoS

Chang Soo Hong,<sup>1</sup> Akio Kuroda,<sup>1</sup> Noboru Takiguchi,<sup>1</sup> Hisao Ohtake,<sup>2</sup> and Junichi Kato<sup>1\*</sup>

Department of Molecular Biotechnology, Graduate School of Advanced Sciences of Matter, Hiroshima University, Higashi-Hiroshima, Hiroshima,<sup>1</sup> and Department of Biotechnology, Graduate School of Engineering, Osaka University, Suita, Osaka,<sup>2</sup> Japan

Received 8 October 2004/Accepted 12 November 2004

***Pseudomonas aeruginosa* possesses two aerotaxis transducers, Aer and Aer-2. A deletion-insertion mutation of alternative sigma factor RpoS eliminated Aer-2-mediated aerotaxis but not Aer-mediated aerotaxis. Transcriptional analysis revealed that *cheY2*, *cheA2*, *cheW2*, and *aer-2* were expressed in an RpoS-dependent manner as a single transcript.**

An aerotactic response in *Pseudomonas aeruginosa* PAO1 (10) has previously been characterized as the movement of a cell toward oxygen (18). In *P. aeruginosa* PAO1, chemotaxis proteins such as CheA, CheB, CheR, CheW, and CheY are required and two methyl-accepting chemotaxis proteins, Aer and Aer-2, function as independent sensor-transducers for aerotaxis (11). In *Escherichia coli*, two methyl-accepting chemotaxis proteins, Aer and Tsr, independently sense and transduce aerotactic signals (2, 17). Recently, it was demonstrated that *E. coli* Aer is a methylation-independent transducer (3) and the PAS domain of *E. coli* Aer requires the C-terminal HAMP domain for native-fold formation and flavin adenine dinucleotide binding (9, 21).

The aerotactic responses of *P. aeruginosa* cells were induced during the transition from exponential to stationary growth phase (12). In the previous study, we demonstrated that *aer* is transcriptionally regulated by the anaerobic regulator ANR (12). ANR is involved in the anaerobic induction of various enzymatic systems, including those required for arginine fermentation, cyanogenesis, and denitrification (1, 7, 8, 23). ANR activates target promoters by binding to ANR boxes, the consensus sequences shared by the ANR-dependent promoters (1, 8). The *aer* promoter contains two ANR boxes at –42.5 and –93.5 bp upstream of the transcriptional start site of *aer*, and both of them are essential for expression of the *aer* gene (12). The *anr* mutation eliminated Aer-mediated aerotaxis, but not Aer-2-mediated aerotaxis, suggesting that *aer-2* expression is regulated by a factor other than ANR. In the present study, we report that the alternative sigma factor RpoS is required for Aer-2-mediated aerotaxis and the transcription of *aer-2* is dependent on RpoS.

The sigma factor RpoS is known to have a role in regulating the expression of stationary-phase genes in a wide range of bacteria, including *P. aeruginosa* (13, 19). To assess the possibility that RpoS is involved in the stationary induction of Aer-2-mediated aerotaxis, the *rpoS* gene was disrupted by inserting

a *tet* (conferring tetracycline resistance) cassette (22) into the wild-type gene in the *P. aeruginosa* PAO1 genome as described previously (14). The resulting *rpoS* mutant, designated PAO-CH1, was fully motile and grew as well as the parent strain, PAO1. PAO-CH1 was examined for the ability to exhibit aerotaxis.

Aerotactic responses of *P. aeruginosa* were assessed by the chemotaxis well chamber method (18). Stationary-phase cells of the *rpoS* mutant PAO-CH1 harboring the green fluorescent protein expression vector pMRP9-1 (16) exhibited decreased but significant aerotaxis. The entire *rpoS* gene (19) was cloned into pMRP9-1 to construct pCSH9-11. Plasmid pCSH9-11 complemented the mutation of PAO-CH1 (Fig. 1), showing that the mutation phenotype was not due to polar effects. The intensity of aerotaxis by PAO-CH1 was as strong as those of the *aer* and *aer-2* single mutants (11) (Fig. 1). These results suggest the possibility that RpoS regulates only one of the aerotaxis transducer genes. To confirm this possibility, we constructed the *aer rpoS* and *aer-2 rpoS* double mutants by inserting a *kan* (conferring kanamycin resistance) cassette into the wild-type *aer* and *aer-2* genes in the PAO-CH1 genome, respectively. The *aer rpoS* and *aer-2 rpoS* double mutants were designated PAO-CH2 and PAO-CH3, respectively. Aerotaxis assays revealed that PAO-CH2 failed to exhibit aerotaxis, whereas PAO-CH3 showed the same level of aerotactic responses as the *aer-2* single mutant (Fig. 1). These results demonstrated that Aer-2-mediated aerotaxis, but not Aer-mediated aerotaxis, requires RpoS.

The *aer-2* gene is located near an *mcp*-like gene (*tlpF*) and a complete set of chemotaxis-like genes (*cheY2*, *cheA2*, *cheW2*, *cheR2*, and *cheB2*) encoding homologues of CheY, CheA, CheW, CheR, and CheB (11) (Fig. 2). These genes are juxtaposed and have the same transcriptional polarity. *tlpF*, *cheY2*, *cheA2*, and *cheW2* are located upstream of *aer-2*. Their coding regions overlap or are separated by short intergenic regions (27 to 197 bp), suggesting that these genes are expressed as a single transcript. To locate the promoter of the transcript containing *aer-2*, we constructed promoter fusions in the broad-host-range transcriptional fusion vector pQF50 (5). Regions upstream of *aer-2* were isolated and inserted individually upstream from the promoterless *lacZ* gene in pQF50 (Fig. 2). Each of five constructs was transformed into PAO1 and PAO-CH1.  $\beta$ -Galac-

\* Corresponding author. Mailing address: Department of Molecular Biotechnology, Graduate School of Advanced Sciences of Matter, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8530, Japan. Phone: 81-82-424-7757. Fax: 81-82-424-7047. E-mail: jun@hiroshima-u.ac.jp.




FIG. 1. Aerotactic responses by wild-type and mutant strains of *P. aeruginosa*. Changes in the green fluorescent protein (GFP) fluorescence intensity of the upper well were measured by a fluorescence spectrophotometer. Symbols: ○, PAO1(pMRP9-1) (wild type); △, TLPG01(pMRP9-1) (*aer* mutant) (11); □, TLPG01(pMRP9-1) (*aer-2* mutant) (11); ◇, PAO-CH1(pMRP9-1) (*rpoS* mutant); ▲, PAO-CH3(pMRP9-1) (*aer-2 rpoS* mutant); ■, PAO-CH2(pMRP9-1) (*aer rpoS* mutant); ◆, PAO-CH1(pCSH9-11).

tosidase activities were then measured in stationary-phase cells of transformants at 37°C (15). High levels of β-galactosidase activity were detected with PAO1(pCSH10.5 [carrying *tlpF cheY2A2W2 aer-2::lacZ*]) and PAO1(pCSH10.4 [carrying *cheY2A2W2 aer-2::lacZ*]) (Fig. 2). In PAO1 harboring pCSH10.3 (carrying *cheA2W2 aer-2::lacZ*), pCSH10.2 (carrying *cheW2 aer-2::lacZ*), and pCSH10.1 (carrying *aer-2::lacZ*), β-galactosidase levels were about 150 U, which was similar to the basal levels seen in PAO1 harboring the control plasmid pQF50. These results suggest that *cheY2*, *cheA2*, *cheW2*, and *aer-2* are expressed as a single transcript and the transcript starts from the region upstream of *cheY2*. The *lacZ* fusion pCSH10.4 gave a basal level of β-galactosidase activity in PAO-CH1, demonstrating that RpoS is required for transcription of the *cheY2A2W2* and *aer-2* genes.

To identify the 5' end of the mRNA containing *cheY2A2W2* and *aer-2*, we carried out rapid amplification of cDNA ends (RACE) as described previously (6, 12). *cheY2* sequence-specific primer Y2SPI (5'-GCGTCACTCGAGCAGTTTC-3') was used for reverse transcription of the total RNA from PAO1. A nested PCR using forward primers T17ADP (5'-GAGTCGACTCGAGAATTC-3') and ADP (5'-GAGTCGACTCGAGAATTC-3') and *cheY2* sequence-specific primers Y2SPII (5'-TTCATCGCGTCGCTC GATTC-3') and Y2SPIII (5'-ATTCGGTGGTCAGCATGAG G-3') enabled amplification of a RACE product (data not shown) that was subsequently isolated and subcloned into pUC118 (20). DNA sequencing indicated that the 5' ends were located at an A nucleotide at position -101 and a G nucleotide at position -100 (relative to the *cheY2* start codon) (Fig. 3). A potential RpoS -10 region (CTTTACT) was located at positions -13 to -7 upstream of the transcription start point (the

### Che cluster 4




FIG. 2. Physical map of Che cluster 4 of *P. aeruginosa* PAO1 (11) and regions cloned in the promoter probe vector pQF50. Specific restriction sites used to isolate each fragment are shown on the map. Restriction sites: B, BamHI; Nr, NruI; Sa, Sau3AI; Sp, SphI; Xm, XmnI. The locations and orientations of *tlpF*, *cheY2*, *cheA2*, *cheW2*, *aer-2*, *cheR2*, PAO174, and *cheB2* are indicated by horizontal arrows. Gene identification numbers used in the *P. aeruginosa* genome sequencing project (<http://www.pseudomonas.com/>) are indicated below the gene names. Open bars are *P. aeruginosa* chromosomal DNA fragments subcloned into pQF50. The *lacZ* gene is shown by the black arrow. β-Galactosidase activities were determined in *P. aeruginosa* wild-type PAO1 and its *rpoS* mutant, PAO-CH1, containing the *aer-2::lacZ* transcriptional fusion plasmids shown. β-Galactosidase activity is shown along with the standard deviation (mean of four independent experiments).




FIG. 3. Promoter region of *P. aeruginosa cheY2*. The transcriptional start sites, which were determined by 5' RACE and DNA sequencing, are indicated by the angled arrow. Numbering is relative to the transcriptional start site (the A nucleotide). The putative translation start site, ribosome-binding site (RBS), and -35 promoter sequence are underlined. Double underlining indicates a putative RpoS -10 region.

A nucleotide (Fig. 3), with six of the seven bases being identical to the consensus sequence (CTATACT) (4).

In summary, *P. aeruginosa* possesses two aerotaxis transducers, Aer and Aer-2. *aer* expression is dependent on the anaerobic regulator ANR, which is converted to its active form under low oxygen supply. *aer-2* is transcribed together with *cheY2A2W2* (and probably with *cheR2B2*). The stationary-phase sigma factor RpoS is required for transcription of the operon.

#### REFERENCES

1. Arai, H., M. Mizutani, and Y. Igarashi. 2003. Transcriptional regulation of the *nos* genes for nitrous oxide reductase in *Pseudomonas aeruginosa*. *Microbiology* **149**:29–36.
2. Bibikov, S. I., R. Biran, K. E. Rudd, and J. S. Parkinson. 1997. A signal transducer for aerotaxis in *Escherichia coli*. *J. Bacteriol.* **179**:4075–4079.
3. Bibikov, S. I., A. C. Miller, K. K. Gosink, and J. S. Parkinson. 2004. Methylation-independent aerotaxis mediated by the *Escherichia coli* Aer protein. *J. Bacteriol.* **186**:3730–3737.
4. Espinosa-Urgel, M., C. Chamizo, and A. Tormo. 1996. A consensus sequence structure for  $\sigma^S$ -dependent promoters. *Mol. Microbiol.* **21**:657–659.
5. Farinha, M. A., and A. M. Kropinski. 1990. Construction of broad-host-range plasmid vectors for easy visible selection and analysis of promoters. *J. Bacteriol.* **172**:3496–3499.
6. Frohman, M. A., M. Dush, and G. R. Martin. 1988. Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc. Natl. Acad. Sci. USA* **85**:8998–9002.
7. Galimand, M., M. Gamper, A. Zimmermann, and D. Haas. 1991. Positive FNR-like control of anaerobic arginine degradation and nitrate respiration in *Pseudomonas aeruginosa*. *J. Bacteriol.* **173**:1598–1606.
8. Haas, D., M. Gamper, and A. Zimmermann. 1992. Anaerobic control in *Pseudomonas aeruginosa*, p. 177–187. In E. Galli, S. Silver, and B. Witholt (ed.), *Pseudomonas: molecular biology and biotechnology*. American Society for Microbiology, Washington, D.C.
9. Herrmann, S., Q. Ma, M. S. Johnson, A. V. Repik, and B. L. Taylor. 2004. PAS domain of the Aer redox sensor requires C-terminal residues for native-fold formation and flavin adenine dinucleotide binding. *J. Bacteriol.* **186**:6782–6791.
10. Holloway, B. W., V. Krishnapillai, and A. F. Morgan. 1979. Chromosomal genetics of *Pseudomonas*. *Microbiol. Rev.* **43**:73–102.
11. Hong, C. S., M. Shitashiro, A. Kuroda, T. Ikeda, N. Takiguchi, H. Ohtake, and J. Kato. 2004. Chemotaxis proteins and transducers for aerotaxis in *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* **231**:247–252.
12. Hong, C. S., A. Kuroda, T. Ikeda, N. Takiguchi, H. Ohtake, and J. Kato. 2004. The aerotaxis transducer gene *aer*, but not *aer-2*, is transcriptionally regulated by the anaerobic regulator ANR in *Pseudomonas aeruginosa*. *J. Biosci. Bioeng.* **97**:184–190.
13. Jorgensen, F., M. Bally, V. Chapomherve, G. Michel, A. Lazdunski, P. Williams, and G. S. Stewart. 1999. RpoS-dependent stress tolerance in *Pseudomonas aeruginosa*. *Microbiology* **145**:835–844.
14. Kato, J., T. Nakamura, A. Kuroda, and H. Ohtake. 1999. Cloning and characterization of chemotaxis genes in *Pseudomonas aeruginosa*. *Biosci. Biotechnol. Biochem.* **63**:151–161.
15. Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
16. Parsek, M. R., and E. P. Greenberg. 1999. Quorum sensing signals in development of *Pseudomonas aeruginosa* biofilms. *Methods Enzymol.* **31**:43–55.
17. Rebbapragata, A., M. S. Johnson, G. P. Harding, A. J. Zuccarelli, H. M. Fletcher, I. B. Zhulin, and B. L. Taylor. 1997. The Aer protein and the serine chemoreceptor Tsr independently sense intracellular energy levels and transduce oxygen, redox, and energy signals for *Escherichia coli* behavior. *Proc. Natl. Acad. Sci. USA* **94**:10541–10546.
18. Shitashiro, M., T. Fukumura, J. Kato, A. Kuroda, T. Ikeda, N. Takiguchi, and H. Ohtake. 2003. Evaluation of bacterial aerotaxis for its potential use in detecting the toxicity of chemicals to microorganisms. *J. Biotechnol.* **101**:11–18.
19. Tanaka, K., and H. Takahashi. 1994. Cloning, analysis and expression of an *rpoS* homolog gene from *Pseudomonas aeruginosa* PAO1. *Gene* **150**:81–85.
20. Vieira, J., and J. Messing. 1987. Production of single-stranded plasmid DNA. *Methods Enzymol.* **153**:3–11.
21. Watts, K. J., Q. Ma, M. S. Johnson, and B. L. Taylor. 2004. Interactions between the PAS and HAMP domains of the *Escherichia coli* aerotaxis receptor Aer. *J. Bacteriol.* **186**:7440–7449.
22. Wu, H., J. Kato, A. Kuroda, T. Ikeda, N. Takiguchi, and H. Ohtake. 2000. Identification and characterization of two chemotactic transducers for inorganic phosphate in *Pseudomonas aeruginosa*. *J. Bacteriol.* **182**:3400–3404.
23. Zimmermann, A., C. Reimmann, M. Galimand, and D. Haas. 1991. Anaerobic growth and cyanide synthesis of *Pseudomonas aeruginosa* depend on *anr*, a regulator gene homologous with *fnr* of *Escherichia coli*. *Mol. Microbiol.* **5**:1483–1490.