

Structural Classification of Bacterial Response Regulators: Diversity of Output Domains and Domain Combinations†

Michael Y. Galperin*

National Center for Biotechnology Information, National Library of Medicine,
 National Institutes of Health, Bethesda, Maryland 20894

Received 12 December 2005/Accepted 28 March 2006

CheY-like phosphoacceptor (or receiver [REC]) domain is a common module in a variety of response regulators of the bacterial signal transduction systems. In this work, 4,610 response regulators, encoded in complete genomes of 200 bacterial and archaeal species, were identified and classified by their domain architectures. Previously uncharacterized output domains were analyzed and, in some cases, assigned to known domain families. Transcriptional regulators of the OmpR, NarL, and NtrC families were found to comprise almost 60% of all response regulators; transcriptional regulators with other DNA-binding domains (LytTR, AraC, Spo0A, Fis, YcbB, RpoE, and MerR) account for an additional 6%. The remaining one-third is represented by the stand-alone REC domain (~14%) and its combinations with a variety of enzymatic (GGDEF, EAL, HD-GYP, CheB, CheC, PP2C, and HisK), RNA-binding (ANTAR and CsrA), protein- or ligand-binding (PAS, GAF, TPR, CAP_ED, and HPT) domains, or newly described domains of unknown function. The diversity of domain architectures and the abundance of alternative domain combinations suggest that fusions between the REC domain and various output domains is a widespread evolutionary mechanism that allows bacterial cells to regulate transcription, enzyme activity, and/or protein-protein interactions in response to environmental challenges. The complete list of response regulators encoded in each of the 200 analyzed genomes is available online at http://www.ncbi.nlm.nih.gov/Complete_Genomes/RRcensus.html.

It has been 20 years since the first bacterial response regulator genes were sequenced (35, 39, 89) and the CheY-like receiver (phosphoacceptor) domain was identified as their common regulatory module (95, 124, 138; see references 55, 57, 127, and 150 for reviews and reference 43 for a historical perspective). Structural characterization of the CheY-like receiver domain (hereafter, the REC domain) from *Escherichia coli* and other organisms confirmed that it is an autonomously folding, evolutionarily stable, compact structural unit (125, 126, 147) whose conformation undergoes a distinctive change upon phosphorylation (65, 75). These conformational changes, which increase the propensity of the REC domain to form dimers, are used by bacterial and archaeal cells to transmit and propagate a wide variety of environmental and intracellular signals (29, 127, 150).

The typical scheme of bacterial two-component signal transduction involves signal sensing (ligand binding or other conformational change) by a sensory histidine kinase that leads to its autophosphorylation, followed by phosphoryl transfer to the Asp residue in the N-terminal REC domain of the cognate response regulator, which affects the properties of the C-terminal DNA-binding domain (127, 150). Structural characterization of the DNA-binding domains of various response regulators revealed several variations on the common helix-turn-helix (HTH) theme, exemplified by the NarL-type, OmpR-type “winged helix,” Spo0A-type, and Fis-like structures (12, 78, 79,

87, 88, 100). Sequence comparisons revealed additional types of DNA-binding domains in certain response regulators, some of which have been experimentally verified (94, 113).

In addition to forming associations with various DNA-binding domains, the REC domain can function as a stand-alone module. Its propensity to protein-protein interactions plays a key role in the chemotaxis machinery, as well as in some other regulatory pathways (10, 130). It also forms other types of response regulators by associating with certain enzymatic or protein-binding domains, e.g., with the methyltransferase (CheB) and chemotaxis modulator (CheW) domains, both of which participate in the adaptation to attractants during chemotaxis (61). In the CheB-type response regulator, the N-terminal REC domain packs against the active site of the C-terminal methyltransferase domain and inhibits methyltransferase activity by restricting access to the active site (34). The conformational change occurring upon phosphorylation of the REC domain apparently disrupts this interaction and relieves the methyltransferase domain from inhibition by REC (34). In the *Bacillus subtilis* CheV protein, the presence of the N-terminal CheW-like domain was shown to stabilize the phosphorylated state of the C-terminal REC domain (40, 61).

The REC domain can also form combinations with other signaling domains, e.g., in the *Caulobacter crescentus* PleD response regulator, which controls flagellum ejection and stalk formation during the *C. crescentus* life cycle (1, 2, 52). In PleD, the N-terminal REC domain is fused to an inactivated REC domain and a C-terminal GGDEF domain, which has diguanylate cyclase activity that produces bis-(3'→5')-cyclic diguanosine monophosphate (c-di-GMP), a secondary messenger in bacteria (99, 112; see references 58 and 109 for reviews). In *Vibrio cholerae* response regulator VieA, which controls biofilm for-

* Mailing address: National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD 20894. Phone: (301) 435-5910. Fax: (301) 435-7793. E-mail: galperin@ncbi.nlm.nih.gov.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

mation by this organism (137), the REC domain is associated with the EAL domain, a c-di-GMP-specific phosphodiesterase (22, 30, 114, 131). A combination of the REC domain with another c-di-GMP-specific phosphodiesterase, the HD-GYP domain, controls biosynthesis of extracellular polysaccharide in *Xanthomonas campestris* (45, 111, 121).

The response regulator sets encoded in *E. coli*, *B. subtilis*, *Synechococcus* sp., *Streptococcus pneumoniae*, and several other bacteria have been tabulated and classified (48, 51, 74, 90, 91, 97, 146, 148). There have been several attempts to compile comprehensive lists of response regulators in all sequenced microbial genomes (11, 66, 70, 141), as well as to list response regulators in microbial genome databases, such as KEGG, SENTRA, and COG (60, 84, 134). However, the extreme diversity of REC domain combinations encoded in recently sequenced microbial genomes continues to generate problems during genome annotation (43, 157). We have already discussed the most common errors in annotating signaling proteins and suggested that, unless exact function of the newly described protein is known, its annotation should be based on its domain composition, rather than on sequence similarity of any given domain (38, 109). To assist in this process, I present here a comprehensive domain-based classification of the response regulators encoded in completely sequenced genomes of 200 bacterial and archaeal species and describe domain architectures that involve the REC domain and known DNA-binding, enzymatic, and ligand-binding domains, as well as several new domains, not described previously.

MATERIALS AND METHODS

Data sources. The analyzed set included complete genome sequences of 200 bacterial and archaeal species, sequenced by 20 September 2005, as listed on the NCBI's Entrez Genomes website (151). Only one representative genome per species was chosen, typically the first publicly released one, according to the NCBI Genomes database listing. As an exception, two strains of *E. coli*, K12 and O157:H7, and three serovars of *Salmonella enterica*, Paratyphi, Typhi, and Typhimurium, were included in the list, as described earlier (42). The genome sequences were downloaded from the NCBI's Genomes database or searched directly on the NCBI website (<http://www.ncbi.nlm.nih.gov/BLAST/>) using the PSI-BLAST (3) tool.

Identification of REC-containing proteins. The complete list of REC-containing proteins encoded in each of the 200 analyzed genomes was compiled separately for each bacterial phylum based on the results of iterative PSI-BLAST searches (3) of the NCBI's Reference Sequence (RefSeq) database (103). In each case, the search was initiated with the 122-amino-acid sequence of the *Thermotoga maritima* CheY protein TM0468 (GenBank accession no. AAD35552; UniProt entry Q9WYT9) and a precomputed position-specific scoring matrix. The query sequence and the position-specific scoring matrix are both available in the supplemental material. The database search space was limited to the given phylum (e.g., Firmicutes[orgn]); sequences of incomplete genomes were excluded using the `srdc_refseq_provisional[prop]` modifier. The PSI-BLAST searches used strict inclusion threshold E values of 10^{-4} to 10^{-5} , adjusting as necessary, and were repeated until convergence. The phylogenetic representation and the total numbers of REC-containing proteins encoded in each given genome were estimated using the Taxonomy Report option in the BLAST output. Potential false-positive hits were checked at every step of PSI-BLAST using the CDD Domain viewer (86) and manually removed (unselected) from the hit list for each iteration of PSI-BLAST. Low-scoring hits (potential false negatives) were inspected on a case-by-case basis by comparing them against the CDD and Pfam domain databases (14, 85) using the CD search tool (86) with relaxed cutoff values.

Classification of REC-containing proteins. The lists of REC-containing proteins from each complete genome were checked against the CDD, COG, and Pfam domain databases (14, 85, 134), using the CDD Domain viewer (86), and grouped according to their domain composition. The sequences were also

matched against UniProt (13), using the UniProt batch retrieval tool, and linked to the corresponding entries in the Pfam database (14). The families of response regulators were named after their experimentally characterized representatives, where available (e.g., OmpR-like and Spo0A-like). In the absence of experimentally characterized representatives, families of response regulators were named based on their domain architectures (e.g., REC-AraC and RpoE-REC). To account for the cases where different species (e.g., three species of *Brucella*) had merged RefSeq entries, PSI-BLAST searches were also run against selected genomes using the NCBI's Genomic BLAST tool (31), followed by manual analysis of the outputs. Whenever possible, the protein counts were compared to the published data, and discrepancies were resolved on a case-by-case basis. In most instances, discrepancies could be attributed to different treatments of highly diverged and truncated sequences. In this work, such sequences were defined as ones showing up in the PSI-BLAST searches but not recognized as containing the REC domain by RPS-BLAST comparison against CDD, COG, or Pfam databases with default or relaxed (E value of 1.0) parameters. These were typically excluded from the total count but still listed (marked with asterisks) in Table S1 in the supplemental material. All N-terminally truncated sequences were manually inspected to see if the apparent truncation could be due to the incorrect selection of the start codon during automated annotation. Corrected open reading frames for nine genes, for which this proved to be the case (*Bacillus cereus* BC5352; *Bacteroides thetaiotaomicron* BT3258; *Bradyrhizobium japonicum* bsl1713, bsl2179, and bsr2863; *Haloarcula marismortui* rMAC1630 and rMAC3385; and *Mesorhizobium loti* msl3517 and mlr3700) (see the file RRseqUpdate in the supplemental material), have been submitted to the NCBI RefSeq database. The domain architecture of the Mlr3700 protein is discussed in the Results.

RESULTS

Response regulators encoded in prokaryotic genomes show a great variety of output domains and domain combinations. Unfortunately, owing to the relatively high sequence conservation of the REC domains, sequence database searches for relatives of certain response regulators often retrieve nonhomologous protein sequences that share only the REC domain (43). Hence, classification of response regulators has to be based primarily on their domain architectures and structures of the constituent domains. To achieve proper granularity, HTH domains of the response regulators have been grouped here by sequence similarity, following the approach utilized previously in the COG database (135). Response regulators that contained RNA-binding, protein-binding, or enzymatic output domains have been put into separate categories (Table 1). Any sequence fragments longer than 90 amino acid residues, not assigned to any known protein domain, were analyzed separately and, if appropriate, identified as members of existing or novel domain families. The results of this analysis are shown in Tables 1 and 2.

Response regulators with DNA-binding output domains. The great majority of all response regulators encoded in complete prokaryotic genomes combines the REC domain with a DNA binding HTH domain, typically either of NarL-type or of OmpR-type winged HTH family (Fig. 1). These domains are well studied and have known three-dimensional structures (12, 107). A factor of inversion stimulation (Fis)-type DNA binding domain (71, 73) is also commonly found in response regulators, either directly fused to the REC domain or containing an additional AAA+-type ATPase domain (Fig. 1). Response regulators of the latter type are referred to here as the NtrC family, after its best-studied representative (65, 76, 95, 145). The former type of domain architecture was first described in the RegA and PrrA response regulators from *Rhodobacter sphaeroides* (36, 73, 101, 115) and is referred to here as the PrrA family. Other widespread DNA-binding domains in re-

TABLE 1. Widespread types of bacterial response regulators^a

Response regulator				Output domain		
Type ^b	COG no.	Size (aa)	No. in the set (% of total)	Name	Pfam entry	Reference(s)
Stand-alone REC	0784	120	659 (14.3%)	REC	PF00072	125, 146
DNA-binding						
OmpR-like	0745	240	1526 (33.1%)	wHTH	PF00486	88
NarL-like	2197	240	862 (18.7%)	HTH	PF00196	12
NtrC-like	2204	450	393 (8.5%)	AAA-FIS	PF00158 PF02954	76, 95
LytR-like	3279 ^c	250	139 (3.0%)	LytTR	PF04397	94
PrrA-like	4567	170	48 (1.0%)	FIS	PF02954	73
REC-AraC (YesN-like)	4753	260	48 (1.0%)	HTH_AraC	PF00165	106
RNA-binding, AmiR/NasR-like	3707	180	40 (0.9%)	ANTAR	PF03861	119
Enzymatic						
Methylesterase (CheB-like)	2201	300	107 (2.3%)	CheB	PF01339	34
Diguanylate cyclase (PleD-like)	3706	310	154 (3.4%) ^d	GGDEF	PF00990	28
c-di-GMP phosphodiesterase (VieA-like)	2200 ^c	370	83 (1.8%) ^d	EAL	PF00563	46
c-di-GMP phosphodiesterase (RpfG-like)	3437	360	72 (1.6%)	HD-GYP		45, 111
Protein phosphatase (RsbU-like)	2208	350	46 (1.0%)	PP2C (SpoIIE)	PF07228	116
Histidine kinase	0642	350	88 (1.9%)	HisKA-HATPase	PF00512 PF02518	19
Protein-binding						
CheW-REC (CheV-like)	0835 ^c	300	72 (1.6%)	CheW	PF01584	110

^a The table lists response regulators with at least 40 representatives in the analyzed set, found in at least two bacterial phyla.

^b Description of the domain structure, if available, or sequence alignment.

^c This COG contains only the output domain of this response regulator.

^d Includes 50 response regulators that contain both GGDEF and EAL domains.

sponse regulators include the AraC-type HTH domain (106) and the LytTR domain (94), whose structure is still unknown (Table 1).

This work revealed a number of additional combinations of the REC domain with previously described DNA-binding domains, including cases where the HTH domain occupies the N-terminal part of the protein (Table 2). In some instances, newly identified output domains could be linked to known HTH domains by PSI-BLAST or CD searches. Thus, a previously uncharacterized N-terminal output domain in response regulators of the CV0537 family, encoded in *Chromobacterium violaceum* and *Burkholderia* spp. (*beta*-proteobacterial transcriptional regulator, BetR, deposited in the Pfam database as domain PF08667), was shown to be related to the XRE-type HTH domain (PF01381), found in Cro and cI repressors (see Fig. S1 in the supplemental material). In some cases, however, response regulators that were originally assumed to be DNA-binding showed no statistically significant similarity to any known DNA-binding domain. For example, nine haloarchaeal response regulators annotated as Hlx1 through Hlx6 (Table 2) contain a new output domain (renamed here HalX) (see Fig. S2 in the supplemental material), whose similarity, if any, to the homeodomain-containing animal Hlx proteins is totally spurious. All such response regulators have been classified as “uncharacterized.” Most of these newly described domain combinations are relatively rare and have a narrow phylogenetic distribution (Table 2).

Response regulators with RNA-binding output domains. Of all the numerous RNA-binding domains described in the past several years (7), only one, ANTAR (119), has been commonly found in response regulators (Table 1). In transcriptional reg-

ulators of the AmiR and NasT type, this domain stimulates transcription by preventing transcription termination at rho-independent terminators (96, 119). A single instance of a response regulator formed by another RNA-binding domain, a fusion of the REC domain with an N-terminal CsrA domain (50, 108), was found in *Rhodospirellula baltica* protein RB8820 (Table 2).

Response regulators with enzymatic output domains. Recognition of the REC domain as a universal regulatory domain was prompted by sequencing of the chemotaxis protein CheB, which combines an N-terminal REC domain with a C-terminal methylesterase domain. The latter domain retained catalytic activity in the absence of REC or when the REC domain was phosphorylated (81, 120, 124). Subsequent studies revealed response regulators that contain the REC domain in association with other enzymatic domains. PleD-like response regulators (1, 2, 52) have a GGDEF-type output domain, which has diguanylate cyclase activity (28, 112), whereas VieA-type response regulators combine the REC domain with the EAL domain, which has c-di-GMP-specific phosphodiesterase activity, and a DNA-binding domain (131, 137). Counting the response regulators containing any of these domains shows that each of them comprises more than 1% of the total set and can be found in various phylogenetic lineages (Fig. 1 and Table 1).

Two more common enzymatic output domains in response regulators are the protein phosphatase of PP2C type (RsbU-like) and the phosphodiesterase HD-GYP, a variant of the widespread HD-type phosphohydrolase (PF01966). The former domain dephosphorylates Ser~P and Thr~P residues in a variety of proteins (64, 116) and is often linked to the regulation of RNA polymerase sigma factors (143, 153). The HD-

TABLE 2. Response regulators with unusual domain organizations

Family name, domain organization	Total no.	Example		Output domain		
		Organism name	Gene name (accession no.)	Pfam entry	COG no.	Reference(s)
DNA-binding, REC-HTH						
Spo0A-like	15	<i>Bacillus subtilis</i>	BSU24220 (CAB14353)	PF005154		78
YcbB (GlnL)-like	14	<i>Bacillus subtilis</i>	BSU02450 (CAB12039)	PF08664		113
REC-SARP (REC-HTH-BTAD)	5	<i>Deinococcus radiodurans</i>	DR2556 (AAF12095)	PF00486 PF03704	3947	152, 154
REC-CRP	2	<i>Gloeobacter violaceus</i>	glr1768 (BAC89709)	PF00325	0664	49
DNA-binding, HTH-REC						
RpoE-REC	15	<i>Caulobacter crescentus</i>	CC3477 (AAK25439)	PF04542 PF04545	1595	26
MerR-REC	6	<i>Treponema denticola</i>	TDE0855 (AAS11346)	PF00376	2452	25
BetR-REC	5	<i>Chromobacterium violaceum</i>	CV0537 (AAQ58214)	PF08667		This work
AlpA-REC	2	<i>Dehalococcoides ethenogenes</i>	DET1294 (AAW39407)	PF05930	3311	139
HlxR-REC	1	<i>Haloarcula marismortui</i>	rrnB0301 (AAV48466)	PF01638		1733
LysR-REC	1	<i>Silicibacter pomeroyi</i>	SPO3037 (AAV96273)	PF00126	0583	93
RNA-binding, CsrA-REC	1	<i>Rhodopirellula baltica</i>	RB8820 (CAD75938)	PF02599		50
Enzymatic						
Ser/Thr protein kinase REC-RsbW or REC-RsbU-RsbW	14	<i>Leptospira interrogans</i>	LA2951 (AAN50149)		2172	153
Phosphoaspartate phosphatase REC-CheC	13	<i>Vibrio cholerae</i>	VCA0189 (AAF96102)	PF04509	1776	98, 129
Adenylate cyclase REC-ACyc	6	<i>Bradyrhizobium japonicum</i>	Blr2288 (BAC47553)	PF00211	2114	156
ParA family ATPase REC-Soj	5	<i>Ralstonia solanacearum</i>	RSc0653 (CAD14183)	PF00991	0455	77
NAD(P)H:disulfide oxidoreductase REC-TrxB	3	<i>Nocardia farcinica</i>	Nfa52850 (BAD60137)	PF00070	2345	72
Poly-N-acetylglucosamine deacetylase REC-LmbE	1	<i>Leifsonia xyli</i>	Lxx09320 (AAT88820)	PF02585	2120	132
Glycosyltransferase REC-WcaA	1	<i>Methanobacterium thermoautotrophicum</i>	MTH548 (AAB85054)	PF00535	1215	133
PilB-like ATPase REC-PuIE	2	<i>Dechloromonas aromatica</i>	Daro_0966 (AAZ45722)	PF00437	2804	102
Predicted phosphatase REC-PglZ	1	<i>Porphyromonas gingivalis</i>	PG0928 (AAQ66063)	PF08665		This work
Predicted phosphodiesterase REC-HDOD	4	<i>Vibrio cholerae</i>	VC1081 (AAF94240)	PF08668	1639	This work
Ser/Thr protein kinase REC-STYK	1	<i>Rhodopirellula baltica</i>	RB10329 (CAD78826)	PF00069	0515	117
Protein-protein interaction						
REC-TPR (VieB-like)	20	<i>Desulfovibrio vulgaris</i>	DVU2937 (AAS97409)	PF00515	0457	21
Hnr-like	18	<i>Escherichia coli</i>	RssB (AAC74317)			15
REC-HPt	9	<i>Xylella fastidiosa</i>	XF2359 (AAF85158)	PF01627	2198	63
REC-PAS	4	<i>Haloarcula marismortui</i>	RrnAC0674 (AAV45676)	PF01627	2202	136
REC-GAF	4	<i>Geobacter sulfurreducens</i>	GSU1316 (AAR34692)	PF01590	2203	54
Ligand-binding						
REC-PilZ	2	<i>Geobacter sulfurreducens</i>	GSU0877 (AAR34207)	PF07238	3215	4
REC-cNMPbinding	2	<i>Bdellovibrio bacteriovorus</i>	Bd3065 (CAE80821)	PF00027	0664	33
Uncharacterized						
PatA-type	21	<i>Nostoc</i> sp. PCC7120	all0521 (BAB72479)			80, 83
Ycf55-type	7	<i>Nostoc</i> sp. PCC7120	alr0960 (BAB72917)			This work
REC-HalX	9	<i>Halobacterium</i> sp. NRC-1	VNG2036G (AAG20197)	PF08663		This work
OsaB-type	2	<i>Streptomyces coelicolor</i>	SCO5749 (CAA19850)			20

GYP domain has been proposed to function as a c-di-GMP hydrolase, based on its peculiar phyletic pattern, complementing that of the EAL domain (45, 46). Although this suggestion was consistent with participation of the HD-GYP-containing response regulator RpfG in regulation of extracellular polysaccharide production in *Xanthomonas campestris* (121) and with the wide distribution of the RpfG-type response regulators (Table 1), it has long remained unverified. Recently, the c-di-GMP-specific phosphodiesterase activity of the HD-GYP domain in RpfG has been demonstrated in a direct biochemical assay (111).

Response regulators whose output domain is a protein phosphatase of CheC type, which dephosphorylates the Asp~P residue of REC (98, 129), are far less common and found mostly in gamma-proteobacteria. The PglZ domain, found in response regulators from *Porphyromonas gingivalis* and *Cyto-*

phaga hutchinsonii, is another likely phosphatase. The *pglZ* (phage growth limitation) gene of *Streptomyces coelicolor* A3 (2) is part of a four-gene operon that somehow controls phage propagation in this organism (128). Its homologs, which include *Salmonella enterica* serovar Typhimurium protein STM4492, are often encoded on integrative plasmids (23). While its exact function is unknown, the PglZ protein and its close homologs have all the hallmarks of the alkaline phosphatase protein superfamily (see Fig. S3 in the supplemental material). Given that members of this superfamily typically function as phosphatases, phosphomutases, or phosphodiesterases (44), PglZ is likely to have a similar enzymatic activity.

Hybrid histidine kinases and REC-HisK response regulators. In addition to response regulators, the REC domain is often found at the C terminus of sensory histidine kinases, which are then referred to as “hybrid” kinases. The search




FIG. 1. Domain architectures of the most common types of bacterial response regulators. The sectors in the chart indicate the respective share of each domain combination and are colored similarly to the corresponding output domains. The sector marked as REC-Fis REC-AraC represents all rare DNA-binding response regulators listed in Table 2.

method used in this work retrieved hybrid histidine kinases as well as response regulators; these were subsequently separated based on their domain architectures. In *E. coli*, five histidine kinases (ArcB, BarA, EvgS, RcsC, and TorS) out of 30 are hybrid; in *B. subtilis*, there are no hybrid kinases with the exception of the YwpD protein that combines a truncated REC domain with a HisK domain (for simplicity, two separate sub-domains of histidine kinases, the dimerization/phosphoacceptor one, HisKA, represented by Pfam entries PF00512, PF07568, PF07730, and PF02518, and the catalytic one, HATPase_c or PF02518, are treated here as a single HisK domain). However, the YwpD protein differs from typical hybrid kinases in that its REC domain is located at the N terminus, and there is no (other) sensory domain (Fig. 2). This means that the only signal that could be sensed by the YwpD protein is the phosphorylation (or other modification) of its N-terminal REC domain. Therefore, YwpD-like histidine kinases can be considered yet another form of response regulators with enzymatic output domains whose activity is modulated by appropriate sensory histidine kinases. Alternatively, the REC and HisK domains of such proteins might not be regulated at all and function solely as sinks for the phosphoryl residues. In any case, similarly to the response regulators of the PleD, VieA, and RpfG families, YwpD-like histidine kinases are members of complex signal transduction cascades operating inside the bacterial cell (41).

Unfortunately, the separation between YwpD-like response regulators and hybrid histidine kinases is not clear-cut. Genome analysis reveals numerous intermediate forms with additional signaling domains, typically PAS and/or GAF, which could affect the enzymatic activity of the output domain (Fig. 2). For the purposes of this work, only YwpD-like histidine kinases without any discernible signaling domains intervening between REC and HisK were counted as bona fide response regulators.

The search for REC-containing proteins also revealed hy-

brid histidine kinases which, in addition to the REC domains, contain DNA-binding domains at their C-termini. These “one-component” signal transduction systems (141), exemplified by *B. cereus* protein BC3207, look like fusions of classical histidine kinases with full-length response regulators. Such proteins are encoded in a variety of bacterial genomes, sometimes in significant numbers. In *Bacteroides thetaiotaomicron*, for example, 36 out of 44 hybrid histidine kinases have tandem AraC-like DNA-binding domains at their C-termini (species-specific listings of hybrid histidine kinases are available in the files that are linked to Table S1 in the supplemental material). In *Bacteroides fragilis* strain YCH46, this domain organization is seen in 11 out of 17 hybrid histidine kinases. Some of such fusion proteins contain periplasmic, extracellular, or integral membrane sensory domains, so that they can only act as transcriptional regulators while their binding sites on the chromosomal DNA are positioned in the vicinity of the membrane.

Response regulators with protein-binding domains. In some response regulators, output domains are devoid of (known) enzymatic activity and exert their effects through protein-protein interactions. These include chemotaxis response regulators of the CheV type, combining REC with the protein-binding CheW-like domain (Table 1), and response regulators of the Hnr-type (see below), as well as response regulators that contain PAS, GAF, TPR, or HPt domains as the only output domains (Table 2). While PAS and GAF domains are known to bind ligands, signal transduction likely involves their interactions with other protein domains, not ligand binding per se. The same could be true for other ligand binding domains, such as CAP_ED, cNMP-binding, and PilZ domains (Table 2).

Clade-specific response regulators. While response regulators of OmpR, NarL, and many other families are widespread in *Bacteria*, some response regulators are found only in representatives of certain phylogenetic lineages. Some of them can be considered true synapomorphies—shared derived characteristics of the representatives of a particular group to the




FIG. 2. Domain architectures of response regulators with the histidine kinase (HisK) output domain. Domain symbols are drawn approximately to scale. Some of the domains are highly diverged and could be recognized only by searching the CDD database using relaxed stringency parameters. Source organisms, gene names, and NCBI protein database accession numbers for the domain architectures shown above are as follows (domain architecture number and accession number) *B. subtilis* *ywpD* (1; CAB05945) or *Nostoc* sp. PCC 7120 *all3764* (1; BAB75463); *Nostoc* sp. PCC 7120 *all1279* (2; BAB73236), *alr1968* (16; BAB73667), and *all4097* (17; BAB75796); *Bradyrhizobium japonicum* *bll1030* (3; BAC46295); *Haloarcula marismortui* *rmAC0574* (4; AAV45588), *rmAC3379* (5; AAV48063), *rmAC2533* (6; AAV47335), *rmAC0075* (7; AAV45155), *rmAC0848* (8; AAV45829), *rmAC1626* (9; AAV46539), *rmAC3361* (10; AAV48050), *rmAC1495* (12; AAV46418), and *pNG7156* (13; AAV44864); *Synechocystis* sp. PCC 6803 *slr0222* (11; BAA10222); *Anabaena variabilis* *Ava_B0191* (14; ABA24903); *Gloeobacter violaceus* *gll0634* (15; BAC88575); *Desulfovibrio vulgaris* *DVU2677* (18; AAS97149); *Dechloromonas aromatica* *Daro_2847* (19; AAZ47577). Proteins with domain architecture numbers 1, 14, and 18 were considered bona fide response regulators.

exclusion of other groups—whereas others are encoded only in organisms with larger genomes and could have been lost in streamlined genomes of parasites and other small-genome organisms.

Spo0A. The master regulator of sporulation, Spo0A, is shared by all spore-forming representatives of *Firmicutes*, namely, the bacteria belonging to the orders *Bacillales*, *Clostridiales*, and *Thermoanaerobacteriales* (122, 123). Remarkably, in the genome sequence of *Clostridium perfringens* strain 13,

the *spo0A* gene is inactivated by what authors claim is a verified frameshift (118). If so, the sequenced strain should be unable to sporulate and produce toxins, which would make it a convenient laboratory strain but hardly a representative of the naturally occurring clostridia. In any case, Spo0A combines the REC domain with an unusual HTH domain that has additional α -helices (78). Outside *Firmicutes*, this HTH domain has been seen so far only in the spore-forming high-GC gram-positive bacterium *Symbiobacterium thermo-*

philum, which is currently assigned to the *Actinobacteria* but is apparently a member of the *Firmicutes* (140). Thus, Spo0A is a typical synapomorphy, common to the spore-forming bacteria of the *Firmicute* lineage.

YcbB. *B. subtilis* response regulator YcbB has been originally described as one of the genes whose disruption increased cell resistance to protonophores (105). This effect remained enigmatic until a recent study showed that YcbB controls the glutamine utilization operon (113). YcbB has been renamed GlnL, which is bound to cause confusion, as it is unrelated to the *E. coli* GlnL (NtrC) response regulator. In fact, the C-terminal domain of YcbB does not show obvious relation to any previously described HTH domain (see Fig. S4 in the supplemental material). Since DNA binding by YcbB has now been experimentally demonstrated (113), its C-terminal domain can be considered a new type of the DNA-binding HTH domain.

PatA. Response regulators of the PatA type, which control heterocyst development, phototaxis, and chromatic adaptation in cyanobacteria (18, 80), combine the C-terminal REC domain with N-terminal PATAN (PatA N terminus) domain and an HTH domain in the middle (83). Although the PATAN domain itself is found in a variety of bacteria, the PATAN-REC combination appears to be limited to cyanobacteria and δ -proteobacteria (83).

RpoE-REC. Rhizobia, brucellae, bartonellae, and other members of the alpha-subdivision of proteobacteria encode an unusual response regulator that combines a C-terminal REC domain with an N-terminal sigma-24 subunit of RNA-polymerase (σ^E or RpoE) (Table 2). With the sole exception of the marine bacterium *Silicibacter pomeroyi*, this response regulator is encoded in every alpha-proteobacterial genome that is larger than 1.6 Mb, often adjacent to the gene coding for a histidine kinase of the recently described HWE family (62). Sequence comparisons identified the RpoE-REC domain combination in the genome of *Mesorhizobium loti* (the *mlr3700* gene product), where the original genome annotation included only its C-terminal REC domain. Response regulators with the RpoE-REC domain combination have not been described in the available literature, and the only indication of their biological function(s) so far is the observation of increased expression of the *C. crescentus* response regulator CC3477 in the minimal medium compared to the rich medium (56). Given the DNA-binding properties of σ^E , these proteins can be predicted to serve as environmentally modulated transcriptional regulators. It remains to be seen whether their RpoE-like components could function as genuine sigma subunits of the RNA polymerase.

RssB. The Hnr-type response regulator, also referred to as RssB or SprE, regulates turnover of the stress sigma factor RpoS in *E. coli*, *Salmonella*, and several other enterobacteria (53). Due to the high degree of sequence conservation and narrow phyletic distribution, its output domain was previously considered to be unique (92). However, sequence analysis showed that this output domain is a degraded version of the PP2C-type Ser/Thr protein phosphatase that is shorter than the typical PP2C domain, has lost some of its Mn^{2+} -binding Asp residues (Fig. 3), and is apparently devoid of the protein phosphatase activity. These observations are in line with the earlier suggestion (15) that RssB is a former anti-sigma factor


that during evolution was recruited to serve as a recognition factor for proteolysis. However, RssB has apparently evolved from a protein phosphatase, not a protein kinase (an anti-sigma). As noted previously (69), phylogenetic distribution of this response regulator is limited to the gamma-proteobacterial families *Enterobacteriaceae* and *Vibrionaceae*, indicating that Hnr-dependent regulation of RpoS turnover is a peculiar feature of *E. coli* and its closest relatives. In pseudomonads, the degradation of the PP2C domain is much less pronounced, the key metal-binding residues are conserved, and the Hnr-related response regulators are likely to have the protein phosphatase activity (Fig. 3).

REC-HDOD. Several proteobacteria, including *Vibrio cholerae* and *Pseudomonas aeruginosa*, encode a response regulator with an unusual output domain, assigned to the COG1639 (41). In addition, this output domain is often encoded in the genomes of the same bacteria in a stand-alone form (see Fig. S5 in the supplemental material). The structure of one such protein, *Campylobacter jejuni* CJ0248, has been solved by the Joint Center for Structural Genomics and deposited in the Protein Data Bank under accession code 1VQR. Sequence and structural analysis of this output domain (deposited in the Pfam database as domain PF08668) showed its distant relationship to the widespread HD phosphohydrolase (PF01966) domain (9). This domain has been named the HD-related output domain, or HDOD, but the lack of conservation of the key metal-binding residues of the typical HD domain did not allow us to predict whether it still has a phosphatase or phosphodiesterase activity and, accordingly, what its role is, if any, in the signal transduction phosphorelay.

Other response regulators. Comparative analysis revealed several other clade-specific response regulators that are found exclusively in cyanobacteria (for example, the Slr2024 family), halobacteria (REC-HalX family), alpha-proteobacteria (CC0440 family), or epsilon-proteobacteria (CJ1608 family). A cyanobacteria-specific response regulator, Sll1879, combines the REC-domain with the Ycf55 domain, found in a stand-alone form in chloroplasts of green plants and red algae. No mention of experimental studies of these response regulators could be found in the available literature, and their functions remain unknown.

Avoidance of certain response regulator types. The availability of a complete genome sequence allows one to determine not only which proteins are encoded there but also which proteins (protein families) are missing. Absence of certain types of response regulators in all representatives of a given phylum could be an important feature of that phylum that might help in understanding mechanisms of signal transduction in its members.

Actinobacteria, for example, encode numerous transcriptional regulators of the SARP family (152), which combine an HTH domain with the BTAD domain, whose function is unknown (154). However, the current set of 390 actinobacterial response regulators does not include a single instance of a REC-SARP domain combination. The reasons for that remain unclear; other organisms, such as *Deinococcus radiodurans* and *Clostridium acetobutylicum*, do encode response regulators with this domain combination (Table 2). None of the available actinobacterial genomes encodes response regulators of the NtrC type, which is consistent with the absence of σ^{54} in these

**Enterobacteriaceae**

HNR_ECOLI 170-337 (337) HCRVNYRQLVAADKPG**L**VLD**I**AAALSEN-DLAFYCLD**V**TR-AGHNGVLAALLLRALFNGLLQE-29-
 MviA_Salty 170-337 (337) HCRINRQLVSADQPG**L**VLD**I**APLSDN-ELAFYCLD**V**TR-AGDNGVLAALLLRALFNGLLQD-29-
 Hnr_Yerpe 170-337 (338) HCRINRQLTMVEKPG**L**VLD**I**AALSDK-DLAFYCLD**V**TR-AGDNGVLAALLLRVLFNGILQE-29-
 RssB_Erwch 170-338 (338) RCRVNYRQLMAADRPG**L**VLD**I**AAALSGD-DLAFYCLD**V**TAAANNHGVLAAALLLRALFNSSLQE-29-
 ExpM_Pecto 170-338 (338) RCRVNYRQLTTAEQPG**L**VLD**I**AAALSET-ELAFYCLD**V**TQGVNNGT**L**AAALLLRALFNGLLQE-29-
 RssB_Panto 170-337 (338) NCRVNYRQLTMAEQPG**L**VLD**I**AAALSDK-DLAFYCLD**V**TR-AGDNGVLAALLLRALFNGLLQE-29-
 Plu2502 171-338 (341) HCKVNYRKLNAADKPG**L**VLD**I**AAALSEN-DLAFYCLD**V**TR-AGHNGVLAALLLRALFNGLLQE-29-

Vibrionaceae

VC1050 200-362 (368) SWKCSYRLLQSAELMP**L**V**F**D**Y**AWLMNG-QFV**F**Y**V**V**D**AAS-HRECGA**A**ST**L**L**V**RALFHDYLRN-25-
 VP2183 199-361 (368) DWRCYRLLQ**S**ADLMP**L**V**F**D**Y**RWLMNG-Q**F**AFY**L**V**D**SAS-QENGGVAT**L**L**V**RALFDDYLRN-25-
 VV2417 199-361 (368) DWKFSYKLLQ**S**TDMMP**L**V**F**D**Y**RWLMNG-Q**F**AFY**L**V**D**SAS-VEKQGVAT**L**L**R**ALFDDYLRN-25-
 VF1689 203-365 (368) NWKLSYRLLQ**S**TDGMP**L**I**F**D**Y**EWLLDG-RL**F**F**F**Y**L**V**D**SHS-GGENGV**I**SC**L**L**I**RA**F**F**N**D**F**IRE-25-

Pseudomonadaceae

PA2798 176-345 (394) GLEFSHRI**I**PSLYLSG**D**F**V**D**Y**FRVDER-RVAFY**L**A**D**V**S**G-HGASSAF**V**T**V**L**L**K**F**M**T**R**L**L**L**YE-29-
 PP2165 176-344 (393) EF**A**FEHQ**I**I**P**S**L**Y**L**S**G**D**F**V**D**Y**F**R**V**D**E**R-R**I**A**F**Y**L**A**D**V**S**G-HGASSAF**V**T**V**L**L**K**F**M**T**R**L**M**F**E-28-
 Pfl_3849 195-364 (412) E**F**R**F**AHQ**I**I**P**S**L**Y**L**S**G**D**F**V**D**Y**F**R**V**D**E**R-R**V**A**F**Y**L**A**D**V**S**G-HGASSAF**V**T**V**L**L**K**F**M**T**R**L**L**F**E-29-

Alteromonadaceae

SO3196 174-344 (367) AAKVDYSL**F**KSSD**I**SS**Y**F**I**D**S**TMVGS**D**-Y**L**I**M**Y**M**A**H**L**P**-EDNRA**A**F**A**C**V**L**L**R**S**F**V**N**Q**L**K**S-30-
 IL1096 166-325 (329) N**G**Y**Q**F**N**C**S**I**N**D**S**G**K**A**P**I**W**I**D**Y**R**P**L**T**E**G-R**V**L**V**I**M**A**A**A**Q**N-**A**T**E**Q**N**L**L**P**L**L**V**L**K**T**L**I**D**P**L**I**R**Q-29-

Ser/Thr phosphatases

PDB: 1A6Q 20-248 (382) NGLRYGLSSMQW-R**V**E**M**E**D**A**H**T**A**V**I**-9-S**F**F**A**V**Y**D**G**H**A**-GSQ**V**A**K**Y**C**E**H**L**L**D**H**I**T**N**N**Q**D**F-39-
 PDB: 1TXO 3-197 (237) L**V**L**R**Y**A**A**R**S**D**R**G**L**V**R**A**N**N**E**D**S**V**Y**A**G**A**---R**L**L**A**L**A**D**G**M**G****4**G**E**V**A**S**Q**L**V**I**A**A**L**A**H**L**D**D**D**E**P**G**G**-29-
 RSBX_BACSU 8-165 (199) E**H**I**Q**T**L**V**Y**Q**L**N**K**E**G**K**S**I**C**G**D**S**F**F**M**K**A**-4-L**I**C**A**V**A**D**G**L**G**-S**G**S**L**A**N**E**S**S**A**I**K**D**L**V**E**N**Y**A**S**E-16-
 RSBU_BACSU 118-283 (335) Q**E**E**A**L**D**I**G**A**I**S**V**P**A**K**Q**M**S**G**D**Y**H**F**V**K-4-I**N**I**A**I**A**D**V**I**G**-K**G**I**P**A**A**L**C**M**S**M**I**K**Y**A**M**D**S**L**P**E**T**-22-
 RSBP_BACSU 188-357 (403) V**N**D**Q**V**Q**I**D**S**Y**N**A**S**S**E**L**S**G**D**L**Y**G**Y**Y**Q-5-Y**G**I**I**L**D**V**M**G-H**G**I**S**S**A**L**I**T**M**S**L**H**P**L**F**Q**R**Q**I**T**Q**-26-
 SP2E_BACSU 591-755 (827) Y**R**V**S**T**G**A**A**A**A**K**G**G**L**V**S**G**D**S**Y**S**M**M**E**-5-Y**A**A**A**I**S**D**G**M**G**-N**G**A**R**A**H**F**E**S**N**E**T**I**K**L**L**E**K**I**L**E**S**-22-

**Enterobacteriaceae**

HNR_ECOLI GQ**F**P**L**L**V**G**Y**Y**H**R**E**L**K**N**L**I**L**V**S**A**G**L**N**A**T**L**N**T**G**E**H**Q**V**Q**I**S**N**G**V**P**L**T**L**G**N**A**Y**L**N**Q**L**S**Q**R**C**D-A**W**Q**C**Q**I**W**T**G**G**R**L**R**L**M**L**S**A**E
 MviA_Salty GQ**F**P**L**F**V**G**Y**Y**H**S**E**L**K**N**L**I**L**V**S**A**G**L**N**A**T**L**N**T**G**A**H**Q**V**Q**I**S**S**G**V**P**L**T**L**G**N**A**Y**L**N**Q**L**S**Q**R**C**D-S**W**Q**C**Q**I**W**G**A**G**R**L**R**L**M**L**S**A**E
 Hnr_Yerpe GQ**F**P**L**L**V**G**Y**Y**H**V**E**Y**Q**N**L**I**L**V**S**A**G**L**H**A**N**I**S**A**G**K**N**Q**I**Q**L**N**N**G**V**P**L**T**L**K**T**A**H**L**N**Q**V**S**Q**R**C**E-A**W**Q**C**Q**V**W**G**G**S**R**L**R**L**M**L**A**E**
 RssB_Erwch GQ**F**P**L**L**V**G**Y**Y**H**R**E**L**K**R**L**I**L**I**S**A**G**L**H**A**T**L**S**V**E**E**Q**H**I**G**L**N**N**G**V**P**L**T**L**E**A**A**Y**L**N**Q**I**S**Y**Q**C**E-S**W**Q**C**Q**M**W**S**G**G**R**L**R**L**M**L**S**T**E
 ExpM_Pecto G**R**F**P**L**L**V**G**Y**Y**H**R**P**L**K**Q**L**I**L**I**S**A**G**L**N**A**T**L**N**V**N**E**Q**Q**V**L**N**S**G**V**P**L**T**L**E**G**A**Y**L**N**Q**L**N**Y**Q**C**E-T**W**Q**C**Q**I**W**G**G**G**R**L**R**L**M**L**T**T**E
 RssB_Panto GQ**F**P**L**L**V**G**Y**Y**H**R**G**L**K**N**L**I**L**V**S**A**G**L**N**A**T**L**H**V**H**A**H**Q**I**Q**L**S**N**G**V**P**L**T**M**G**S**T**H**L**N**Q**I**S**Q**R**A**E-N**W**Q**C**H**V**W**G**A**G**R**I**R**L**M**L**S**T**E
 Plu2502 GQ**F**P**L**L**V**G**Y**Y**H**R**E**L**K**N**L**I**L**V**S**A**G**L**N**A**T**L**N**T**G**E**H**Q**V**Q**I**S**N**G**V**P**L**T**L**G**N**A**Y**L**N**Q**L**S**Q**R**C**D-A**W**Q**C**Q**I**W**N**T**N**S**Q**I**N**L**M**L**S**S**M**

Vibrionaceae

VC1050 G**K**V**S**A**L**F**G**V**A**D**L**A**D**G**T**L**S**L**P**A**G**L**D**C**Q**W**S**N**G**Y**V**H**N**R**I**T**S**G**V**Q**L**G---D**N**S**L**R**N**F**I**T**K**D**L**P**I**R**A**A**C**Q**I**S**L**S**K**P**S**A**Y**S**F**T**L**D**I**
 VP2183 G**A**V**K**A**V**F**G**V**A**D**L**A**S**S**T**V**S**L**P**A**G**L**D**S**Q**W**S**D**G**R**E**N**Q**H**I**A**A**G**N**R**L**G---D**G**S**M**K**N**F**I**T**R**D**L**P**L**K**T**T**G**Q**L**S**I**G**L**C**L**G**A**S**S**F**S**L**N**I
 VV2417 C**P**V**N**A**L**F**G**I**A**D**L**V**D**G**S**V**E**I**L**P**A**G**L**D**S**Q**W**S**T**G**F**I**N**Q**H**I**A**A**G**N**R**L**G**---H**N**S**L**K**N**F**I**T**R**D**L**N**I**G**S**G**S**Q**L**T**L**G**C**L**G**S**A**S**F**S**L**N**I**
 VF1689 Q**A**M**S**G**V**F**G**Y**V**D**L**S**E**Q**A**V**F**I**L**S**A**G**L**D**A**I**W**E**S**N**N**Q**R**N**V**L**R**A**N**Q**L**T---T**H**A**S**T**N**Q**V**H**R**F**S**V**K**E**G**M**N**S**L**S**L**N**N**V**C**S**S**F**K**L**E**I

Pseudomonadaceae

PA2798 K**H**V**T**M**L**G**V**I**D**L**E**-K**N**S**L**T**Y**S**I**G**G**H**L**P**L**P**V**L**F**V**E**G**Q**A**G**Y**L**E**G**R**G**---6-D**D**A**T**Y**D**D**R**V**M**E**L**P**P**S**F**S**L**S**L**F**S**G**I**L**D**V**L**P**G**A
 PP2165 K**H**V**T**M**V**G**V**I**D**E**E**-S**G**L**L**T**Y**A**V**G**G**H**L**P**L**P**V**L**T**P**E**H**C**R**Y**L**E**G**R**G---6-D**E**A**T**Y**Q**D**L**V**V**E**L**P**P**Q**F**S**L**S**L**M**S**G**I**L**D**L**L**P**G**C
 Pfl_3849 K**H**V**T**M**V**G**V**I**D**E**E**-T**G**L**L**T**Y**S**I**G**G**H**L**P**L**P**V**L**T**P**D**S**V**R**Y**L**E**G**R**G---6-N**E**A**T**Y**E**D**H**V**L**E**L**P**P**T**F**S**L**T**L**M**S**G**I**L**D**L**L**P**E**F

Alteromonadaceae

SO3196 I**T**V**D**I**V**Y**V**A**I**E**L**T**R**Y**R**A**A**I**A**Q**A**G**K**L**R**C**Y**L**R**N**D**E**G**L**S**P**L**A**L**S**D**S---6-D**W**G**K**P**S**V**Q**F**R**T**I**L**P**Q**E**Q**L**C**I**A**T**E**P**E**H**K**Q**L**L**I
 IL1096 T**A**F**D**A**L**V**I**V**O**T**T**K**H**S**L**W**A**Q**A**G**D**K**F**E**P**E**P**G**A**K**P**D**L**A**L**G**I**W**Q**H**A**-----N**F**R**E**H**R**I**E**H**L**K**L**H**C**G**Y**E**N**T**E**L**I**V**T**A**V**

Ser/Thr phosphatases

PDB: 1A6Q S**G**S**T**A**V**G**V**L**I**S**P**---Q**H**T**Y**F**I**N**C**D**S**R**G**L**L**C**R**N**R**K**V**H**F**T**Q**D**H**K**P**-50-P**E**P**E**V**H**D**I**E**R**S**E**D**D**Q**F**I**I**L**A**C**D**G**I**W**D**V**M**G**N**E
 PDB: 1TXO M**G**T**T**L**T**A**I**L**F**A**G**---N**R**L**G**L**V**H**I**C**D**S**R**G**Y**L**L**R**D**G**E**L**T**Q**I**T**K**D**D**T**F**-32-V**E**P**T**L**T**M**R**E---A**R**A**G**D**R**Y**L**L**C**S**G**L**S**D**P**V**S**D**E**
 RSBX_BACSU R**G**A**T**A**S**I**L**K**I**N**F**E**Q**R**Q**F**T**Y**C**S**V**G**N**V**R**F**I**L**H**S**P**S**G**E**S**F**Y**L**P**I**S**E---4-K**P**Q**K**Y**K**T**H**T**A**T**Y**E**K**G**S**K**F**I**I**H**T**O**G**L**N**V**P**D**I**R**S**
 RSBU_BACSU M**F**I**T**M**F**Y**A**N**Y**N**M**D**K**H**Q**F**T**Y**A**S**A**C**H**E**P**G**F**Y**S**Q**K**D**N**T**F**Y**D**L**E**A**K**---6-Q**D**Y**D**Y**K**Q**F**D**Q**H**L**E**K**G**D**M**I**V**L**F**S**D**G**V**T**E**C**R**T**E**N**
 RSBP_BACSU H**Y**C**T**A**I**Y**L**E**I**D**I**A**R**Q**R**I**D**Y**V**N**A**C**H**P**A**L**W**-Q**D**D**S**G**T**Q**H**L**L**H**A**T**S**---6-E**D**L**E**F**Q**S**S**S**L**S**Y**T**E**D**G**R**L**L**L**Y**T**O**G**V**M**D**P**T**A**S**C**
 SP2E_BACSU I**Y**S**T**L**D**L**S**I**I**D**L**Q**D**A**S**C**K**F**L**V**G**S**T**P**S**F**I**K**R**G**D**Q**V**M**K**V**Q**A**S**N**L**P---4-N**E**F**D**V**E**V**V**S**E**Q**L**K**A**G**D**L**L**I**M**M**S**G**I**F**E**G**P**K**H**V

FIG. 3. Multiple sequence alignment of the Hnr-type output domain and its comparison with the protein phosphatase domains of the PP2C family. The top sequence is that of *E. coli* protein RssB (GenBank accession no. AAC74317). Conserved hydrophobic amino acid residues are shaded yellow; other conserved residues are shown in bold typeface. Metal-binding residues of PP2C-type protein phosphatases are shown in white

genomes. The only exception is *Symbiobacterium thermophilum*, which is not really an actinobacterium (140) (see above). As noted earlier (42), not even the largest actinobacterial genomes encode any chemotaxis proteins; this includes the absence of the CheB, CheC, and CheV family proteins, as well as a relatively low number of stand-alone REC domains. Although actinobacteria encode signaling proteins with HisK, GGDEF, EAL, HD-GYP, and ACyc output domains, none of them is found in actinobacterial response regulators. Finally, only a small fraction of actinobacterial histidine kinases are hybrids. These features illuminate the peculiarity of the signal transduction in actinobacteria and suggest that representatives of this phylum utilize a rather straightforward mechanism of signal transduction that includes tight pairing of each HisK with a cognate DNA-binding (or RNA-binding) response regulator.

Archaeal transcriptional regulators and signal transducers are generally similar to those in bacteria (16, 47). However, archaeal response regulators differ dramatically from bacterial ones: no sequenced archaeal genome encodes OmpR, NarL, NtrC, LytR, or PrrA family response regulators (see Table S1 in the supplemental material; see also reference 141). The only response regulators that are common for bacteria and archaea are the stand-alone REC domain and CheB-like and REC-HisK combinations. Other archaeal response regulators include REC-PAS and REC-GAF domain combinations, as well as combinations of the REC domain with uncharacterized lineage-specific domains, found, for example, only in halobacteria or only in metanosarcinas.

Statistics of response regulators. The number of response regulators encoded in each complete microbial genome shows strong positive correlation with the genome size. In the logarithmic coordinates, this dependence shows a slope of approximately 1.8, meaning that it is close to the quadratic function (Fig. 4) (see also reference 141). There are certain deviations from the general trend: some representatives of beta-, delta- and epsilon-proteobacteria, such as *Geobacter sulfurreducens*, *Wolinella succinogenes*, and *Thiobacillus denitrificans*, encode substantially more response regulators than other organisms with the same genome sizes. At the same time, other organisms encode relatively few response regulators or none at all (e.g., *Sulfolobus* spp.) (Fig. 4).

DISCUSSION

The sequence analysis of REC domain-containing response regulators presented in this work shows that functions of these proteins are not limited to transcriptional regulation and chemotaxis. While transcriptional regulators with DNA-binding output domains comprise about two-thirds of all response regulators, the remaining third consists of stand-alone REC domains, response regulators with enzymatic or protein-binding output domains, and response regulators whose output domains have not been characterized and whose functions re-

main unknown. In a somewhat similar analysis by Ulrich and colleagues (141), stand-alone REC domains and response regulators with unknown output domains were not considered, resulting in a much higher fraction of DNA-binding response regulators in their set.

Mechanisms of REC-mediated signaling. Stand-alone REC domains, such as chemotaxis regulator CheY and sporulation regulator Spo0F, have two principal modes of transmitting regulatory signals. Both depend on protein-protein interactions of the phosphorylated form REC~P with its various targets but differ in the fate of the phosphoryl group, which can stay on the REC domain or be transferred to another protein with REC~P functioning as an intermediate in a multicomponent phosphorelay (37). In some cases, REC~P might function just as a sink for phosphoryl groups (10, 130). In most bacteria, chemotactic signaling by CheY involves direct binding of its phosphorylated form, CheY~P, to the flagellar switch protein, FliM (24, 149). In archaea, which have an entirely different flagellar machinery that does not include FliM, CheY-like proteins interact with a different, still unidentified, protein (130). In cyanobacteria, whose motility apparatus is related to the type IV pili, CheY~P interacts with yet another type of switch complex (18). This is also the case for type IV pili-mediated social motility of *Myxococcus xanthus* (144). In addition to their role in chemotaxis, stand-alone REC domains participate in developmental processes, including regulation of sporulation in *B. subtilis* (Spo0F), heterocyst formation in *Nostoc* sp. (DevR), and cyst cell development in *Rhodospirillum centenum* (17, 27, 37). Furthermore, separately expressed REC and methylesterase domains of the CheB protein have been shown to form a complex even in the absence of a linker (6). These data suggest that REC domains could participate in a variety of protein-protein interactions with yet unknown target proteins.

Another big group of response regulators combines the REC domain with various enzymatic domains that are involved in signal transduction. Besides chemotactic methylesterase CheB and protein phosphatase CheC, these include the diguanylate cyclase (GGDEF), c-di-GMP phosphodiesterase (EAL and HD-GYP), adenylate cyclase, Ser/Thr-specific protein phosphatase, and histidine kinase output domains. In some cases, phosphorylation of the REC domain has been shown to activate the attached enzymatic domains (34, 99). While this activation could be due to the relief of enzyme inhibition by nonphosphorylated REC (34), it appears that in the case of CheB, both relief of inhibition and genuine activation take place (5, 6).

Another mechanism of REC-mediated signaling is the well-documented propensity of phosphorylated REC domains to form dimers. Phosphorylation-induced dimerization plays a key role in the regulation of DNA-binding response regulators, dramatically improving their binding to the palindromic or direct-repeat chromosomal binding sites. This mechanism has




FIG. 4. Correlation between the genome size and total number of encoded response regulators. Each symbol indicates the logarithm of the number of response regulators encoded in a single bacterial genome with the following color scheme: actinobacteria, black circles; cyanobacteria, cyan circles; alpha-proteobacteria, brown diamonds; beta-, delta-, and epsilon-proteobacteria, yellow diamonds; gamma-proteobacteria, blue squares; firmicutes, open squares; members of other bacterial phyla (*Aquificales*, *Bacteroidetes*, *Chlamydiae*, *Deinococcus-Thermus*, *Planctomycetes*, *Spirochetes*, and *Thermotogales*), red triangles. Archaeal response regulators are indicated by yellow triangles. Organisms that encode no response regulators are indicated by symbols on the x axis ($\log N = 0$); these, however, were ignored for the calculation of the slope and the correlation coefficient of the regression line. The numbers in the circles indicate the data points corresponding to the following organisms: 1, *Dechloromonas aromatica*; 2, *Geobacter sulfurreducens*; 3, *Wolinella succinogenes*; 4, *Dehalococcoides* sp. strain CBDB1; 5, *Sulfolobus solfataricus*.

been demonstrated for the response regulators of the OmpR, NarL, LytR, and PrrA families (12, 73, 88, 94). In the response regulators of the NtrC family, all three domains appear to participate in oligomerization, which regulates the activity of its AAA+ ATPase domain (67, 76).

The interplay of dimerization and relief-of-inhibition mechanisms could be particularly important for response regulators involved in c-di-GMP turnover. In response regulators of the REC-GGDEF type, phosphorylation of the REC domain activates the diguanylate cyclase by enabling formation of a GGDEF dimer, the active form of the enzyme (99, 112). In contrast, the c-di-GMP-specific phosphodiesterase (the EAL domain) is active as a monomer (114). Therefore, activation of EAL phosphodiesterase by REC phosphorylation in response regulators of the REC-EAL type (131, 137) appears to be due to the relief-of-inhibition mechanism. This distinction is particularly important for the response regulators of REC-GGDEF-EAL domain architecture, which comprise 1.1% of all response regulators in the analyzed set. Although in many such proteins either the GGDEF or EAL domain appears to be inactivated (114), there are some without obvious deviations from the consensus (109). Such response regulators should have the only the

phosphodiesterase activity in their monomeric form but might acquire diguanylate cyclase activity upon dimerization. It is entirely possible that upon phosphorylation of their REC domains, the overall activity of such response regulators changes from c-di-GMP-hydrolase to c-di-GMP synthetase and vice versa.

In addition to dimerization, relief-of-inhibition, direct activation, and protein-protein interactions with or without phosphorylation, REC-mediated signaling might involve other mechanisms, such as, for example, allosteric regulation of the output domains. This versatility of the REC domain is consistent with the diversity of signal output domains that interact with it.

New types of response regulators. The same regulatory mechanisms mentioned above—dimerization, protein-protein interaction, and activation/relief-of-inhibition—can be expected to work in the newly described response regulators. Among the new types of DNA-binding response regulators (Table 2), DNA-binding domains of LysR, MerR, and HxlR types are known to function as dimers (25, 93, 155), as is the RNA-binding CsrA domain (50). DNA-binding domains of the XRE-family, which includes cI and Cro regulators, also function as dimers, suggesting that the BetR domain, a newly described distant member of this family, would function the same way. However, the RpoE-like domain of the

CC3477-type (RpoE-REC) response regulators is not known to form dimers, so they most likely act through relief of inhibition and/or protein-protein interactions of their C-terminal REC domains.

Among enzymatic output domains of response regulators (Tables 1 and 2), adenylate and diguanylate cyclases, histidine kinases, and RsbW-type Ser/Thr protein kinases (anti-sigma factors) are known to function as dimers (19, 28, 82, 156). In these response regulators, both phosphorylation-induced dimerization of the REC domains and relief-of-inhibition could contribute to the activity of the output domains. In all other cases, a nonphosphorylated REC domain can be expected to block enzyme activity the same way it does that in the CheB structure (34). However, just as in CheB, a direct enzyme activation by REC~P, as well as an allosteric regulation, also remains a possibility.

Phylogenetic distribution of response regulators. The census of response regulators confirmed several trends seen previously in the census of signal transduction proteins (42). In accordance with previous observations (46, 68, 141, 142), the total number of response regulators encoded in completely sequenced prokaryotic genomes positively correlates with the genome size (Fig. 4) and the total number of encoded proteins. The largest deviation from the general trend is seen in the same representatives of beta-, delta-, and epsilon-proteobacteria that have been previously found to encode more signaling proteins than others with the same genome size (42). Given only a minimal overlap between the two data sets, these observations show that certain environmental bacteria encode a disproportionately higher number of sensor, signal transduction, and response regulator proteins than other prokaryotes with the same genome size. On the other end of the spectrum are bacteria and archaea that encode fewer response regulators than the average bacterium of the same genome size. These are typically obligate parasites, such as *Yersinia pestis*, or organisms such as *Sulfolobus* spp. or *Thermoplasma* spp. that inhabit relatively stable and unique environments where few parameters, if any, ever change. These data validate the use of the term “bacterial IQ” (42) to describe the fraction of the bacterial proteome dedicated to environmental sensing and signal transduction.

Origin and function of diverse domain architectures. The diversity of output domains in response regulators (Tables 1 and 2) suggests that virtually any DNA-binding or enzymatic domain could be fused to the REC domain to form a response regulator. For DNA-binding domains, this is probably correct, as representatives of all major classes of the HTH domain (8) could be found in response regulators (Table 2). As discussed above, the propensity of the REC domain to dimerize upon phosphorylation offers an effective mechanism of transcriptional regulation. Remarkably, HTH domains such as LysR or MerR that are typically localized at the N-termini of the corresponding transcriptional regulators have the same position in response regulators (Table 2), a clear deviation from the standard REC-HTH architecture. This diversity makes it even more difficult to explain the predominance of OmpR-type and NarL-type response regulators, far fewer numbers of the LytR-type, PrrA-type, and YesN-type response regulators, and only sporadic appearance of the response regulators with other types of the HTH domain (Fig. 1 and Tables 1 and 2).

Most enzymatic output domains found in response regulators (Tables 1 and 2) are signal transduction domains themselves, often found in transmembrane receptors (42). Such fusions add yet another layer of regulation in the complex intracellular signal transduction network, allowing modulation of the cellular levels of cAMP and c-di-GMP, as well as the activity of Ser/Thr protein kinases and phosphatases, by sensory histidine kinases in response to specific environmental cues (41). Fusions with other enzymes typically have a narrow phylogenetic distribution, and it is not known if they are ever expressed. Such combinations must have arisen from random gene translocation and fusion events and were later fixed in evolution because they allowed tight regulation of the enzyme activity. In one such case, a fusion of the REC and thioredoxin reductase domains could play an important role in environmental regulation of the cellular dithiol-disulfide ratio. This domain combination has been found in several actinobacteria and cyanobacteria and in *Solibacter usitatus*, a recently sequenced representative of the phylum *Acidobacteria*. Plant pathogen *Leifsonia xyli* encodes a unique response regulator with the LmbE output domain, recently shown to have *N*-acetylglucosamine deacetylase activity (Table 2). Such domain combination could allow this bacterium to regulate degradation of *N*-acetylglucosamine-containing plant polymers, such as lignin and chitosan, depending on the environmental conditions.

In general, it appears that fusing an existing metabolic enzyme to the REC domain is an evolutionarily harmless procedure: the resulting fused protein is inactive unless its REC domain is phosphorylated by a cognate histidine kinase. The same logic could be applied to the fusions between REC domains and other signaling, ligand-binding, or protein-binding domains. Indeed, the large number of rare domain fusions (Table 2; see Table S1 in the supplemental material) supports the view that signaling domain fusions follow the Lego principle: almost any fusion can occur but only some of them remain fixed in the evolution (43). If this logic is correct, many more types of response regulators can be expected to be found in microbial genomes. Several relatively common, previously uncharacterized rare or unique domains identified in this study are listed in the supplemental material and will soon be available in the public domain databases. An experimental study of these novel response regulators and analysis of their potential functions will remain the task for future work, which will provide further understanding of the organization and evolution of the prokaryotic signal transduction systems.

ACKNOWLEDGMENTS

I thank Andrei Gabrielian, Bill Klimke, Darren Natale, Tao Tao, and Yuri Wolf for helpful advice; Mark Gomelsky, Igor Zhulin, and two reviewers for critical comments; and many other colleagues for suggestions.

This study was supported by the Intramural Research Program of the National Library of Medicine at the U.S. National Institutes of Health.

REFERENCES

1. Aldridge, P., and U. Jenal. 1999. Cell cycle-dependent degradation of a flagellar motor component requires a novel-type response regulator. *Mol. Microbiol.* **32**:379–391.
2. Aldridge, P., R. Paul, P. Goymer, P. Rainey, and U. Jenal. 2003. Role of the GGDEF regulator PleD in polar development of *Caulobacter crescentus*. *Mol. Microbiol.* **47**:1695–1708.

3. Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zheng, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST—a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
4. Amikam, D., and M. Y. Galperin. 2006. PilZ domain is part of the bacterial c-di-GMP binding protein. *Bioinformatics* **22**:3–6.
5. Anand, G. S., P. N. Goudreau, and A. M. Stock. 1998. Activation of methyl-esterase CheB: evidence of a dual role for the regulatory domain. *Biochemistry* **37**:14038–14047.
6. Anand, G. S., and A. M. Stock. 2002. Kinetic basis for the stimulatory effect of phosphorylation on the methyl-esterase activity of CheB. *Biochemistry* **41**:6752–6760.
7. Anantharaman, V., E. V. Koonin, and L. Aravind. 2002. Comparative genomics and evolution of proteins involved in RNA metabolism. *Nucleic Acids Res.* **30**:1427–1464.
8. Aravind, L., V. Anantharaman, S. Balaji, M. M. Babu, and L. M. Iyer. 2005. The many faces of the helix-turn-helix domain: transcription regulation and beyond. *FEMS Microbiol. Rev.* **29**:231–262.
9. Aravind, L., and E. V. Koonin. 1998. The HD domain defines a new superfamily of metal-dependent phosphohydrolases. *Trends Biochem. Sci.* **23**:469–472.
10. Armitage, J. P. 1999. Bacterial tactic responses. *Adv. Microb. Physiol.* **41**:229–289.
11. Ashby, M. K. 2004. Survey of the number of two-component response regulator genes in the complete and annotated genome sequences of prokaryotes. *FEMS Microbiol. Lett.* **231**:277–281.
12. Baikalov, I., I. Schroder, M. Kaczor-Grzeskowiak, K. Grzeskowiak, R. P. Gunsalus, and R. E. Dickerson. 1996. Structure of the *Escherichia coli* response regulator NarL. *Biochemistry* **35**:11053–11056.
13. Bairoch, A., R. Apweiler, C. H. Wu, W. C. Barker, B. Boeckmann, S. Ferro, E. Gasteiger, H. Huang, R. Lopez, M. Magrane, M. J. Martin, D. A. Natale, C. O'Donovan, N. Redaschi, and L. S. Yeh. 2005. The Universal Protein Resource (UniProt). *Nucleic Acids Res.* **33**:D154–D159.
14. Bateman, A., L. Coin, R. Durbin, R. D. Finn, V. Hollich, S. Griffiths-Jones, A. Khanna, M. Marshall, S. Moxon, E. L. Sonnhammer, D. J. Studholme, C. Yeats, and S. R. Eddy. 2004. The Pfam protein families database. *Nucleic Acids Res.* **32**:D138–D141.
15. Becker, G., E. Klauk, and R. Hengge-Aronis. 2000. The response regulator RssB, a recognition factor for σ^S proteolysis in *Escherichia coli*, can act like an anti- σ^S factor. *Mol. Microbiol.* **35**:657–666.
16. Bell, S. D. 2005. Archaeal transcriptional regulation—variation on a bacterial theme? *Trends Microbiol.* **13**:262–265.
17. Berleman, J. E., and C. E. Bauer. 2005. Involvement of a Che-like signal transduction cascade in regulating cyst cell development in *Rhodospirillum rubrum*. *Mol. Microbiol.* **56**:1457–1466.
18. Bhaya, D., A. Takahashi, and A. R. Grossman. 2001. Light regulation of type IV pilus-dependent motility by chemosensor-like elements in *Synechocystis PCC6803*. *Proc. Natl. Acad. Sci. USA* **98**:7540–7545.
19. Bilwes, A. M., L. A. Alex, B. R. Crane, and M. I. Simon. 1999. Structure of CheA, a signal-transducing histidine kinase. *Cell* **96**:131–141.
20. Bishop, A., S. Fielding, P. Dyson, and P. Herron. 2004. Systematic insertional mutagenesis of a streptomycete genome: a link between osmoadaptation and antibiotic production. *Genome Res.* **14**:893–900.
21. Blatch, G. L., and M. Lasse. 1999. The tetratricopeptide repeat: a structural motif mediating protein-protein interactions. *Bioessays* **21**:932–939.
22. Bobrov, A. G., O. Kirillina, and R. D. Perry. 2005. The phosphodiesterase activity of the HmsP EAL domain is required for negative regulation of biofilm formation in *Yersinia pestis*. *FEMS Microbiol. Lett.* **247**:123–130.
23. Boltner, D., C. MacMahon, J. T. Pembroke, P. Strike, and A. M. Osborn. 2002. R391: a conjugative integrating mosaic comprised of phage, plasmid, and transposon elements. *J. Bacteriol.* **184**:5158–5169.
24. Bren, A., and M. Eisenbach. 1998. The N terminus of the flagellar switch protein, FlhM, is the binding domain for the chemotactic response regulator, CheY. *J. Mol. Biol.* **278**:507–514.
25. Brown, N. L., J. V. Stoyanov, S. P. Kidd, and J. L. Hobman. 2003. The MerR family of transcriptional regulators. *FEMS Microbiol. Rev.* **27**:145–163.
26. Campbell, E. A., J. L. Tupy, T. M. Gruber, S. Wang, M. M. Sharp, C. A. Gross, and S. A. Darst. 2003. Crystal structure of *Escherichia coli* σ^E with the cytoplasmic domain of its anti- σ RseA. *Mol. Cell* **11**:1067–1078.
27. Campbell, E. L., K. D. Hagen, M. F. Cohen, M. L. Summers, and J. C. Meeks. 1996. The *devR* gene product is characteristic of receivers of two-component regulatory systems and is essential for heterocyst development in the filamentous cyanobacterium *Nostoc* sp. strain ATCC 29133. *J. Bacteriol.* **178**:2037–2043.
28. Chan, C., R. Paul, D. Samoray, N. C. Amiot, B. Giese, U. Jenal, and T. Schirmer. 2004. Structural basis of activity and allosteric control of diguanylate cyclase. *Proc. Natl. Acad. Sci. USA* **101**:17084–17089.
29. Cho, H. S., J. G. Peltton, D. Yan, S. Kustu, and D. E. Wemmer. 2001. Phosphoaspartates in bacterial signal transduction. *Curr. Opin. Struct. Biol.* **11**:679–684.
30. Christen, M., B. Christen, M. Folcher, A. Schauerer, and U. Jenal. 2005. Identification and characterization of a cyclic di-GMP-specific phosphodiesterase and its allosteric control by GTP. *J. Biol. Chem.* **280**:30829–30837.
31. Cummings, L., L. Riley, L. Black, A. Souvorov, S. Resenchuk, I. Dondoshansky, and T. Tatusova. 2002. Genomic BLAST: custom-defined virtual databases for complete and unfinished genomes. *FEMS Microbiol. Lett.* **216**:133–138.
32. Das, A. K., N. R. Helps, P. T. Cohen, and D. Barford. 1996. Crystal structure of the protein serine/threonine phosphatase 2C at 2.0 Å resolution. *EMBO J.* **15**:6798–6809.
33. Diller, T. C., Madhusudan, N. H. Xuong, and S. S. Taylor. 2001. Molecular basis for regulatory subunit diversity in cAMP-dependent protein kinase: crystal structure of the type II beta regulatory subunit. *Structure* **9**:73–82.
34. Djordjevic, S., P. N. Goudreau, Q. Xu, A. M. Stock, and A. H. West. 1998. Structural basis for methyl-esterase CheB regulation by a phosphorylation-activated domain. *Proc. Natl. Acad. Sci. USA* **95**:1381–1386.
35. Drury, L. S., and R. S. Buxton. 1985. DNA sequence analysis of the dye gene of *Escherichia coli* reveals amino acid homology between the dye and OmpR proteins. *J. Biol. Chem.* **260**:4236–4242.
36. Eraso, J. M., and S. Kaplan. 1994. PrrA, a putative response regulator involved in oxygen regulation of photosynthesis gene expression in *Rhodospirillum rubrum*. *J. Bacteriol.* **176**:32–43.
37. Fabret, C., V. A. Feher, and J. A. Hoch. 1999. Two-component signal transduction in *Bacillus subtilis*: how one organism sees its world. *J. Bacteriol.* **181**:1975–1983.
38. Fedorova, N. D., A. N. Nikolskaya, and M. Y. Galperin. 2003. Using genomic context in the analysis of multi-component systems, p. 167–194. *In* M. Y. Galperin and E. V. Koonin (ed.), *Frontiers in computational genomics*, vol. 3. Caister Academic Press, Wymondham, United Kingdom.
39. Ferrari, F. A., K. Trach, D. LeCoq, J. Spence, E. Ferrari, and J. A. Hoch. 1985. Characterization of the *spoA* locus and its deduced product. *Proc. Natl. Acad. Sci. USA* **82**:2647–2651.
40. Fredrick, K. L., and J. D. Helmann. 1994. Dual chemotaxis signaling pathways in *Bacillus subtilis*: a σ^D -dependent gene encodes a novel protein with both CheW and CheY homologous domains. *J. Bacteriol.* **176**:2727–2735.
41. Galperin, M. Y. 2004. Bacterial signal transduction network in a genomic perspective. *Environ. Microbiol.* **6**:552–567.
42. Galperin, M. Y. 2005. A census of membrane-bound and intracellular signal transduction proteins in bacteria: bacterial IQ, extroverts and introverts. *BMC Microbiol.* **5**:35. [Online.] <http://www.biomedcentral.com/1471-2180/5/35>.
43. Galperin, M. Y., and M. Gomelsky. 2005. Bacterial signal transduction modules: from genomics to biology. *ASM News* **71**:326–333.
44. Galperin, M. Y., and M. J. Jedrzejas. 2001. Conserved core structure and active site residues in alkaline phosphatase superfamily enzymes. *Proteins* **45**:318–324.
45. Galperin, M. Y., D. A. Natale, L. Aravind, and E. V. Koonin. 1999. A specialized version of the HD hydrolase domain implicated in signal transduction. *J. Mol. Microbiol. Biotechnol.* **1**:303–305.
46. Galperin, M. Y., A. N. Nikolskaya, and E. V. Koonin. 2001. Novel domains of the prokaryotic two-component signal transduction systems. *FEMS Microbiol. Lett.* **203**:11–21.
47. Geiduschek, E. P., and M. Ouhammouch. 2005. Archaeal transcription and its regulators. *Mol. Microbiol.* **56**:1397–1407.
48. Grebe, T. W., and J. B. Stock. 1999. The histidine protein kinase superfamily. *Adv. Microb. Physiol.* **41**:139–227.
49. Green, J., C. Scott, and J. R. Guest. 2001. Functional versatility in the CRP-FNR superfamily of transcription factors: FNR and FLP. *Adv. Microb. Physiol.* **44**:1–34.
50. Gutierrez, P., Y. Li, M. J. Osborne, E. Pomerantseva, Q. Liu, and K. Gehring. 2005. Solution structure of the carbon storage regulator protein CsrA from *Escherichia coli*. *J. Bacteriol.* **187**:3496–3501.
51. Hagiwara, D., T. Yamashino, and T. Mizuno. 2004. Genome-wide comparison of the His-to-Asp phosphorelay signaling components of three symbiotic genera of *Rhizobia*. *DNA Res.* **11**:57–65.
52. Hecht, G. B., and A. Newton. 1995. Identification of a novel response regulator required for the swarmer-to-stalked-cell transition in *Caulobacter crescentus*. *J. Bacteriol.* **177**:6223–6229.
53. Hengge-Aronis, R. 2002. Signal transduction and regulatory mechanisms involved in control of the σ^S (RpoS) subunit of RNA polymerase. *Microbiol. Mol. Biol. Rev.* **66**:373–395.
54. Ho, Y. S., L. M. Burden, and J. H. Hurley. 2000. Structure of the GAF domain, a ubiquitous signaling motif and a new class of cyclic GMP receptor. *EMBO J.* **19**:5288–5299.
55. Hoch, J. A., and T. J. Silhavy (ed.). 1995. Two-component signal transduction. ASM Press, Washington, D.C.
56. Hottes, A. K., M. Meewan, D. Yang, N. Arana, P. Romero, H. H. McAdams, and C. Stephens. 2004. Transcriptional profiling of *Caulobacter crescentus* during growth on complex and minimal media. *J. Bacteriol.* **186**:1448–1461.
57. Inouye, M., and R. Dutta (ed.). 2003. Histidine kinases in signal transduction. Academic Press, San Diego, Calif.
58. Jenal, U. 2004. Cyclic di-guanosine-monophosphate comes of age: a novel

- secondary messenger involved in modulating cell surface structures in bacteria? *Curr. Opin. Microbiol.* **7**:185–191.
59. Jones, D. T. 1999. Protein secondary structure prediction based on position-specific scoring matrices. *J. Mol. Biol.* **292**:195–202.
 60. Kanehisa, M., S. Goto, S. Kawashima, Y. Okuno, and M. Hattori. 2004. The KEGG resource for deciphering the genome. *Nucleic Acids Res.* **32**:D277–D280.
 61. Karatan, E., M. M. Saulmon, M. W. Bunn, and G. W. Ordal. 2001. Phosphorylation of the response regulator CheV is required for adaptation to attractants during *Bacillus subtilis* chemotaxis. *J. Biol. Chem.* **276**:43618–43626.
 62. Karniol, B., and R. D. Vierstra. 2004. The HWE histidine kinases, a new family of bacterial two-component sensor kinases with potentially diverse roles in environmental signaling. *J. Bacteriol.* **186**:445–453.
 63. Kato, M., T. Mizuno, T. Shimizu, and T. Hakoshima. 1997. Insights into multistep phosphorelay from the crystal structure of the C-terminal HPT domain of ArcB. *Cell* **88**:717–723.
 64. Kennelly, P. J. 2002. Protein kinases and protein phosphatases in prokaryotes: a genomic perspective. *FEMS Microbiol. Lett.* **206**:1–8.
 65. Kern, D., B. F. Volkman, P. Luginbuhl, M. J. Nohaile, S. Kustu, and D. E. Wemmer. 1999. Structure of a transiently phosphorylated switch in bacterial signal transduction. *Nature* **402**:894–898.
 66. Kim, D., and S. Forst. 2001. Genomic analysis of the histidine kinase family in bacteria and archaea. *Microbiology* **147**:1197–1212.
 67. Klose, K. E., A. K. North, K. M. Stedman, and S. Kustu. 1994. The major dimerization determinants of the nitrogen regulatory protein NtrC from enteric bacteria lie in its carboxy-terminal domain. *J. Mol. Biol.* **241**:233–245.
 68. Konstantinidis, K. T., and J. M. Tiedje. 2004. Trends between gene content and genome size in prokaryotic species with larger genomes. *Proc. Natl. Acad. Sci. USA* **101**:3160–3165.
 69. Koonin, E. V., L. Aravind, and M. Y. Galperin. 2000. A comparative-genomic view of the microbial stress response, p. 417–444. *In* G. Storz and R. Hengge-Aronis (ed.), *Bacterial stress responses*. ASM Press, Washington, D.C.
 70. Koretke, K. K., A. N. Lupas, P. V. Warren, M. Rosenberg, and J. R. Brown. 2000. Evolution of two-component signal transduction. *Mol. Biol. Evol.* **17**:1956–1970.
 71. Kostrewa, D., J. Granzin, C. Koch, H. W. Choe, S. Raghunathan, W. Wolf, J. Labahn, R. Kahmann, and W. Saenger. 1991. Three-dimensional structure of the *E. coli* DNA-binding protein FIS. *Nature* **349**:178–180.
 72. Kuriyan, J., T. S. Krishna, L. Wong, B. Guenther, A. Pahler, C. H. Williams, Jr., and P. Model. 1991. Convergent evolution of similar function in two structurally divergent enzymes. *Nature* **352**:172–174.
 73. Laguri, C., M. K. Phillips-Jones, and M. P. Williamson. 2003. Solution structure and DNA binding of the effector domain from the global regulator PrrA (RegA) from *Rhodobacter sphaeroides*: insights into DNA binding specificity. *Nucleic Acids Res.* **31**:6778–6787.
 74. Lange, R., C. Wagner, A. de Saizieu, N. Flint, J. Molnos, M. Stieger, P. Caspers, M. Kamber, W. Keck, and K. E. Amrein. 1999. Domain organization and molecular characterization of 13 two-component systems identified by genome sequencing of *Streptococcus pneumoniae*. *Gene* **237**:223–234.
 75. Lee, S. Y., H. S. Cho, J. G. Pelton, D. Yan, E. A. Berry, and D. E. Wemmer. 2001. Crystal structure of activated CheY. Comparison with other activated receiver domains. *J. Biol. Chem.* **276**:16425–16431.
 76. Lee, S. Y., A. De La Torre, D. Yan, S. Kustu, B. T. Nixon, and D. E. Wemmer. 2003. Regulation of the transcriptional activator NtrC1: structural studies of the regulatory and AAA+ ATPase domains. *Genes Dev.* **17**:2552–2563.
 77. Leonard, T. A., P. J. Butler, and J. Lowe. 2005. Bacterial chromosome segregation: structure and DNA binding of the Soj dimer—a conserved biological switch. *EMBO J.* **24**:270–282.
 78. Lewis, R. J., S. Krzywda, J. A. Brannigan, J. P. Turkenburg, K. Muchova, E. J. Dodson, I. Barak, and A. J. Wilkinson. 2000. The trans-activation domain of the sporulation response regulator Spo0A revealed by X-ray crystallography. *Mol. Microbiol.* **38**:198–212.
 79. Lewis, R. J., D. J. Scott, J. A. Brannigan, J. C. Ladds, M. A. Cervin, G. B. Spiegelman, J. G. Hoggett, I. Barak, and A. J. Wilkinson. 2002. Dimer formation and transcription activation in the sporulation response regulator Spo0A. *J. Mol. Biol.* **316**:235–245.
 80. Liang, J., L. Scappino, and R. Haselkorn. 1992. The *patA* gene product, which contains a region similar to CheY of *Escherichia coli*, controls heterocyst pattern formation in the cyanobacterium *Anabaena* 7120. *Proc. Natl. Acad. Sci. USA* **89**:5655–5659.
 81. Lupas, A., and J. Stock. 1989. Phosphorylation of an N-terminal regulatory domain activates the CheB methyltransferase in bacterial chemotaxis. *J. Biol. Chem.* **264**:17337–17342.
 82. Madec, E., A. Laszkiewicz, A. Iwanicki, M. Obuchowski, and S. Seror. 2002. Characterization of a membrane-linked Ser/Thr protein kinase in *Bacillus subtilis*, implicated in developmental processes. *Mol. Microbiol.* **46**:571–586.
 83. Makarova, K. S., E. V. Koonin, R. Haselkorn, and M. Y. Galperin. 16 March 2006, posting date. Cyanobacterial response regulator PatA contains a conserved N-terminal domain (PATAN) with an alpha-helical insertion. *Bioinformatics* [Online.] doi:10.1093/bioinformatics/btl096.
 84. Maltsev, N., E. Marland, G. X. Yu, S. Bhatnagar, and R. Lusk. 2002. Sentra, a database of signal transduction proteins. *Nucleic Acids Res.* **30**:349–350.
 85. Marchler-Bauer, A., J. B. Anderson, P. F. Cherukuri, C. DeWeese-Scott, L. Y. Geer, M. Gwadz, S. He, D. I. Hurwitz, J. D. Jackson, Z. Ke, C. J. Lanczycki, C. A. Liebert, C. Liu, F. Lu, G. H. Marchler, M. Mullokandov, B. A. Shoemaker, V. Simonyan, J. S. Song, P. A. Thiessen, R. A. Yamashita, J. J. Yin, D. Zhang, and S. H. Bryant. 2005. CDD: a Conserved Domain Database for protein classification. *Nucleic Acids Res.* **33**:D192–D196.
 86. Marchler-Bauer, A., and S. H. Bryant. 2004. CD-Search: protein domain annotations on the fly. *Nucleic Acids Res.* **32**:W327–W331.
 87. Martinez-Hackert, E., and A. M. Stock. 1997. The DNA-binding domain of OmpR: crystal structures of a winged helix transcription factor. *Structure* **5**:109–124.
 88. Martinez-Hackert, E., and A. M. Stock. 1997. Structural relationships in the OmpR family of winged-helix transcription factors. *J. Mol. Biol.* **269**:301–312.
 89. Matsumura, P., J. J. Rydel, R. Linzmeier, and D. Vacante. 1984. Overexpression and sequence of the *Escherichia coli* *cheY* gene and biochemical activities of the CheY protein. *J. Bacteriol.* **160**:36–41.
 90. Mizuno, T. 1997. Compilation of all genes encoding two-component phosphotransfer signal transducers in the genome of *Escherichia coli*. *DNA Res.* **4**:161–168.
 91. Mizuno, T., T. Kaneko, and S. Tabata. 1996. Compilation of all genes encoding bacterial two-component signal transducers in the genome of the cyanobacterium, *Synechocystis* sp. strain PCC 6803. *DNA Res.* **3**:407–414.
 92. Muffler, A., D. Fischer, S. Altuvia, G. Storz, and R. Hengge-Aronis. 1996. The response regulator RssB controls stability of the σ^S subunit of RNA polymerase in *Escherichia coli*. *EMBO J.* **15**:1333–1339.
 93. Muraoka, S., R. Okumura, N. Ogawa, T. Nonaka, K. Miyashita, and T. Senda. 2003. Crystal structure of a full-length LysR-type transcriptional regulator, CbnR: unusual combination of two subunit forms and molecular bases for causing and changing DNA bend. *J. Mol. Biol.* **328**:555–566.
 94. Nikolskaya, A. N., and M. Y. Galperin. 2002. A novel type of conserved DNA-binding domain in the transcriptional regulators of the AlgR/AgrA/LytR family. *Nucleic Acids Res.* **30**:2453–2459.
 95. Nixon, B. T., C. W. Ronson, and F. M. Ausubel. 1986. Two-component regulatory systems responsive to environmental stimuli share strongly conserved domains with the nitrogen assimilation regulatory genes *ntrB* and *ntrC*. *Proc. Natl. Acad. Sci. USA* **83**:7850–7854.
 96. O'Hara, B. P., R. A. Norman, P. T. Wan, S. M. Roe, T. E. Barrett, R. E. Drew, and L. H. Pearl. 1999. Crystal structure and induction mechanism of AmiC-AmiR: a ligand-regulated transcription antitermination complex. *EMBO J.* **18**:5175–5186.
 97. Ohmori, M., M. Ikeuchi, N. Sato, P. Wolk, T. Kaneko, T. Ogawa, M. Kanehisa, S. Goto, S. Kawashima, S. Okamoto, H. Yoshimura, H. Katoh, T. Fujisawa, S. Ehira, A. Kamei, S. Yoshihara, R. Narikawa, and S. Tabata. 2001. Characterization of genes encoding multi-domain proteins in the genome of the filamentous nitrogen-fixing cyanobacterium *Anabaena* sp. strain PCC 7120. *DNA Res.* **8**:271–284.
 98. Park, S. Y., X. Chao, G. Gonzalez-Bonet, B. D. Beel, A. M. Bilwes, and B. R. Crane. 2004. Structure and function of an unusual family of protein phosphatases: the bacterial chemotaxis proteins CheC and CheX. *Mol. Cell* **16**:563–574.
 99. Paul, R., S. Weiser, N. C. Amiot, C. Chan, T. Schirmer, B. Giese, and U. Jenal. 2004. Cell cycle-dependent dynamic localization of a bacterial response regulator with a novel di-guanylate cyclase output domain. *Genes Dev.* **18**:715–727.
 100. Pelton, J. G., S. Kustu, and D. E. Wemmer. 1999. Solution structure of the DNA-binding domain of NtrC with three alanine substitutions. *J. Mol. Biol.* **292**:1095–1110.
 101. Phillips-Jones, M. K., and C. N. Hunter. 1994. Cloning and nucleotide sequence of *regA*, a putative response regulator gene of *Rhodobacter sphaeroides*. *FEMS Microbiol. Lett.* **116**:269–275.
 102. Planet, P. J., S. C. Kachlany, R. DeSalle, and D. H. Figurski. 2001. Phylogeny of genes for secretion NTPases: identification of the widespread *tadA* subfamily and development of a diagnostic key for gene classification. *Proc. Natl. Acad. Sci. USA* **98**:2503–2508.
 103. Pruitt, K. D., T. Tatusova, and D. R. Maglott. 2005. NCBI Reference Sequence (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res.* **33**:D501–D504.
 104. Pullen, K. E., H. L. Ng, P. Y. Sung, M. C. Good, S. M. Smith, and T. Alber. 2004. An alternate conformation and a third metal in PstP/Ppp, the *M. tuberculosis* PP2C-Family Ser/Thr protein phosphatase. *Structure* **12**:1947–1954.
 105. Quirk, P. G., A. A. Guffanti, S. Clejan, J. Cheng, and T. A. Krulwich. 1994. Isolation of Tn917 insertional mutants of *Bacillus subtilis* that are resistant to the protonophore carbonyl cyanide m-chlorophenylhydrazone. *Biochim. Biophys. Acta* **1186**:27–34.

106. Rhee, S., R. G. Martin, J. L. Rosner, and D. R. Davies. 1998. A novel DNA-binding motif in MarA: the first structure for an AraC family transcriptional activator. *Proc. Natl. Acad. Sci. USA* **95**:10413–10418.
107. Robinson, V. L., T. Wu, and A. M. Stock. 2003. Structural analysis of the domain interface in DrrB, a response regulator of the OmpR/PhoB subfamily. *J. Bacteriol.* **185**:4186–4194.
108. Romeo, T. 1998. Global regulation by the small RNA-binding protein CsrA and the non-coding RNA molecule CsrB. *Mol. Microbiol.* **29**:1321–1330.
109. Römling, U., M. Gomelsky, and M. Y. Galperin. 2005. C-di-GMP: The dawning of a novel bacterial signalling system. *Mol. Microbiol.* **57**:629–639.
110. Rosario, M. M., K. L. Fredrick, G. W. Ordal, and J. D. Helmann. 1994. Chemotaxis in *Bacillus subtilis* requires either of two functionally redundant CheW homologs. *J. Bacteriol.* **176**:2736–2739.
111. Ryan, R. P., Y. Fouhy, J. F. Lucey, L. C. Crossman, S. Spiro, Y.-W. He, L.-H. Zhang, S. Heeb, M. Cámara, P. Williams, and J. M. Dow. 2006. Cell-cell signaling in *Xanthomonas campestris* involves an HD-GYP domain protein that functions in cyclic di-GMP turnover. *Proc. Natl. Acad. Sci. USA*, **103**:6712–6717.
112. Ryjenkov, D. A., M. Tarutina, O. M. Moskvina, and M. Gomelsky. 2005. Cyclic diguanylate is a ubiquitous signaling molecule in bacteria: insights into biochemistry of the GGDEF protein domain. *J. Bacteriol.* **187**:1792–1798.
113. Satomura, T., D. Shimura, K. Asai, Y. Sadaie, K. Hirooka, and Y. Fujita. 2005. Enhancement of glutamine utilization in *Bacillus subtilis* through the GlnK-GlnL two-component regulatory system. *J. Bacteriol.* **187**:4813–4821.
114. Schmidt, A. J., D. A. Ryjenkov, and M. Gomelsky. 2005. Ubiquitous protein domain EAL encodes cyclic diguanylate-specific phosphodiesterase: enzymatically active and inactive EAL domains. *J. Bacteriol.* **187**:4774–4781.
115. Sganga, M. W., and C. E. Bauer. 1992. Regulatory factors controlling photosynthetic reaction center and light-harvesting gene expression in *Rhodobacter capsulatus*. *Cell* **68**:945–954.
116. Shi, L. 2004. Manganese-dependent protein O-phosphatases in prokaryotes and their biological functions. *Front. Biosci.* **9**:1382–1397.
117. Shi, L., M. Potts, and P. J. Kennelly. 1998. The serine, threonine, and/or tyrosine-specific protein kinases and protein phosphatases of prokaryotic organisms: a family portrait. *FEMS Microbiol. Rev.* **22**:229–253.
118. Shimizu, T., K. Ohtani, H. Hirakawa, K. Ohshima, A. Yamashita, T. Shiba, N. Ogasawara, M. Hattori, S. Kuhara, and H. Hayashi. 2002. Complete genome sequence of *Clostridium perfringens*, an anaerobic flesh-eater. *Proc. Natl. Acad. Sci. USA* **99**:996–1001.
119. Shu, C. J., and I. B. Zhulin. 2002. ANTAG: an RNA-binding domain in transcription antitermination regulatory proteins. *Trends Biochem. Sci.* **27**:3–5.
120. Simms, S. A., M. G. Keane, and J. Stock. 1985. Multiple forms of the CheB methyltransferase in bacterial chemotaxis. *J. Biol. Chem.* **260**:10161–10168.
121. Slater, H., A. Alvarez-Morales, C. E. Barber, M. J. Daniels, and J. M. Dow. 2000. A two-component system involving an HD-GYP domain protein links cell-cell signalling to pathogenicity gene expression in *Xanthomonas campestris*. *Mol. Microbiol.* **38**:986–1003.
122. Sonenshein, A. L. 2000. Control of sporulation initiation in *Bacillus subtilis*. *Curr. Opin. Microbiol.* **3**:561–566.
123. Stephenson, K., and R. J. Lewis. 2005. Molecular insights into the initiation of sporulation in gram-positive bacteria: new technologies for an old phenomenon. *FEMS Microbiol. Rev.* **29**:281–301.
124. Stock, A., D. E. Koshland, Jr., and J. Stock. 1985. Homologies between the *Salmonella typhimurium* CheY protein and proteins involved in the regulation of chemotaxis, membrane protein synthesis, and sporulation. *Proc. Natl. Acad. Sci. USA* **82**:7989–7993.
125. Stock, A. M., E. Martinez-Hackert, B. F. Rasmussen, A. H. West, J. B. Stock, D. Ringe, and G. A. Petsko. 1993. Structure of the Mg²⁺-bound form of CheY and mechanism of phosphoryl transfer in bacterial chemotaxis. *Biochemistry* **32**:13375–13380.
126. Stock, A. M., J. M. Mottonen, J. B. Stock, and C. E. Schutt. 1989. Three-dimensional structure of CheY, the response regulator of bacterial chemotaxis. *Nature* **337**:745–749.
127. Stock, A. M., V. L. Robinson, and P. N. Goudreau. 2000. Two-component signal transduction. *Annu. Rev. Biochem.* **69**:183–215.
128. Sumby, P., and M. C. Smith. 2002. Genetics of the phage growth limitation (Pgl) system of *Streptomyces coelicolor* A3(2). *Mol. Microbiol.* **44**:489–500.
129. Szurmant, H., T. J. Muff, and G. W. Ordal. 2004. *Bacillus subtilis* CheC and FliY are members of a novel class of CheY-P-hydrolyzing proteins in the chemotactic signal transduction cascade. *J. Biol. Chem.* **279**:21787–21792.
130. Szurmant, H., and G. W. Ordal. 2004. Diversity in chemotaxis mechanisms among the bacteria and archaea. *Microbiol. Mol. Biol. Rev.* **68**:301–319.
131. Tamayo, R., A. D. Tischler, and A. Camilli. 2005. The EAL domain protein VieA is a cyclic diguanylate phosphodiesterase. *J. Biol. Chem.* **280**:33324–33330.
132. Tanaka, T., T. Fukui, S. Fujiwara, H. Atomi, and T. Imanaka. 2004. Concentrated action of diacetylchitobiose deacetylase and exo- β -D-glucosaminidase in a novel chitinolytic pathway in the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1. *J. Biol. Chem.* **279**:30021–30027.
133. Tarbouriech, N., S. J. Charnock, and G. J. Davies. 2001. Three-dimensional structures of the Mn and Mg dTDP complexes of the family GT-2 glycosyltransferase SpsA: a comparison with related NDP-sugar glycosyltransferases. *J. Mol. Biol.* **314**:655–661.
134. Tatusov, R. L., N. D. Fedorova, J. D. Jackson, A. R. Jacobs, B. Kiryutin, E. V. Koonin, D. M. Krylov, R. Mazumder, S. L. Mekhedov, A. N. Nikolskaya, B. S. Rao, S. Smirnov, A. V. Sverdlov, S. Vasudevan, Y. I. Wolf, J. J. Yin, and D. A. Natale. 11 September 2003, posting date. The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**:41. [Online.] doi:10.1186/1471-2105-4-41.
135. Tatusov, R. L., M. Y. Galperin, D. A. Natale, and E. V. Koonin. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* **28**:33–36.
136. Taylor, B. L., and I. B. Zhulin. 1999. PAS domains: internal sensors of oxygen, redox potential, and light. *Microbiol. Mol. Biol. Rev.* **63**:479–506.
137. Tischler, A. D., and A. Camilli. 2004. Cyclic diguanylate (c-di-GMP) regulates *Vibrio cholerae* biofilm formation. *Mol. Microbiol.* **53**:857–869.
138. Trach, K. A., J. W. Chapman, P. J. Piggot, and J. A. Hoch. 1985. Deduced product of the stage 0 sporulation gene *spoOF* shares homology with the Spo0A, OmpR, and SfrA proteins. *Proc. Natl. Acad. Sci. USA* **82**:7260–7264.
139. Trempy, J. E., J. E. Kirby, and S. Gottesman. 1994. Alp suppression of Lon: dependence on the *slpA* gene. *J. Bacteriol.* **176**:2061–2067.
140. Ueda, K., A. Yamashita, J. Ishikawa, M. Shimada, T. O. Watsuji, K. Morimura, H. Ikeda, M. Hattori, and T. Beppu. 2004. Genome sequence of *Symbiobacterium thermophilum*, an uncultivable bacterium that depends on microbial commensalism. *Nucleic Acids Res.* **32**:4937–4944.
141. Ulrich, L. E., E. V. Koonin, and I. B. Zhulin. 2005. One-component systems dominate signal transduction in prokaryotes. *Trends Microbiol.* **13**:52–56.
142. van Nimwegen, E. 2003. Scaling laws in the functional content of genomes. *Trends Genet.* **19**:479–484.
143. Vijay, K., M. S. Brody, E. Fredlund, and C. W. Price. 2000. A PP2C phosphatase containing a PAS domain is required to convey signals of energy stress to the σ^B transcription factor of *Bacillus subtilis*. *Mol. Microbiol.* **35**:180–188.
144. Vlamakis, H. C., J. R. Kirby, and D. R. Zusman. 2004. The Che4 pathway of *Myxococcus xanthus* regulates type IV pilus-mediated motility. *Mol. Microbiol.* **52**:1799–1811.
145. Volkman, B. F., M. J. Nohaile, N. K. Amy, S. Kustu, and D. E. Wemmer. 1995. Three-dimensional solution structure of the N-terminal receiver domain of NtrC. *Biochemistry* **34**:1413–1424.
146. Volz, K. 1993. Structural conservation in the CheY superfamily. *Biochemistry* **32**:11741–11753.
147. Volz, K., and P. Matsumura. 1991. Crystal structure of *Escherichia coli* CheY refined at 1.7-Å resolution. *J. Biol. Chem.* **266**:15511–15519.
148. Wang, L., Y. P. Sun, W. L. Chen, J. H. Li, and C. C. Zhang. 2002. Genomic analysis of protein kinases, protein phosphatases and two-component regulatory systems of the cyanobacterium *Anabaena* sp. strain PCC 7120. *FEMS Microbiol. Lett.* **217**:155–165.
149. Welch, M., K. Oosawa, S. Aizawa, and M. Eisenbach. 1993. Phosphorylation-dependent binding of a signal molecule to the flagellar switch of bacteria. *Proc. Natl. Acad. Sci. USA* **90**:8787–8791.
150. West, A. H., and A. M. Stock. 2001. Histidine kinases and response regulator proteins in two-component signaling systems. *Trends Biochem. Sci.* **26**:369–376.
151. Wheeler, D. L., T. Barrett, D. A. Benson, S. H. Bryant, K. Canese, D. M. Church, M. DiCuccio, R. Edgar, S. Federhen, W. Helmberg, D. L. Kenton, O. Khovayko, D. J. Lipman, T. L. Madden, D. R. Maglott, J. Ostell, J. U. Pontius, K. D. Pruitt, G. D. Schuler, L. M. Schriml, E. Sequeira, S. T. Sherry, K. Sirotkin, G. Starchenko, T. O. Suzek, R. Tatusov, T. A. Tatusova, L. Wagner, and E. Yaschenko. 2005. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res.* **33**:D39–D45.
152. Wietzorrek, A., and M. Bibb. 1997. A novel family of proteins that regulates antibiotic production in streptomycetes appears to contain an OmpR-like DNA-binding fold. *Mol. Microbiol.* **25**:1181–1184.
153. Yang, X., C. M. Kang, M. S. Brody, and C. W. Price. 1996. Opposing pairs of serine protein kinases and phosphatases transmit signals of environmental stress to activate a bacterial transcription factor. *Genes Dev.* **10**:2265–2275.
154. Yeats, C., S. Bentley, and A. Bateman. 6 February 2003, posting date. New knowledge from old: in silico discovery of novel protein domains in *Streptomyces coelicolor*. *BMC Microbiol.* **3**:3. [Online.] doi:10.1186/1471-2180-3-3.
155. Yurimoto, H., R. Hirai, N. Matsuno, H. Yasueda, N. Kato, and Y. Sakai. 2005. HxIR, a member of the DUF24 protein family, is a DNA-binding protein that acts as a positive regulator of the formaldehyde-inducible *hxlAB* operon in *Bacillus subtilis*. *Mol. Microbiol.* **57**:511–519.
156. Zhang, G., Y. Liu, A. E. Ruoho, and J. H. Hurley. 1997. Structure of the adenyllyl cyclase catalytic core. *Nature* **386**:247–253.
157. Zhulin, I. B., A. N. Nikolskaya, and M. Y. Galperin. 2003. Common extracellular sensory domains in transmembrane receptors for diverse signal transduction pathways in bacteria and archaea. *J. Bacteriol.* **185**:285–294.