

Anaerobic Survival of *Pseudomonas aeruginosa* by Pyruvate Fermentation Requires an Usp-Type Stress Protein

Kerstin Schreiber,¹ Nelli Boes,¹ Martin Eschbach,¹ Lothar Jaensch,² Juergen Wehland,²
Thomas Bjarnsholt,³ Michael Givskov,³ Morten Hentzer,³† and Max Schobert^{1*}

Institute of Microbiology, Technical University Braunschweig, Spielmannstr. 7, D-38106 Braunschweig, Germany¹; Department of Cell Biology, German Research Centre for Biotechnology, Mascheroder Weg 1, D-38124 Braunschweig, Germany²; and Center for Biomedical Microbiology, BioCentrum-DTU, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark³

Received 1 July 2005/Accepted 13 October 2005

Recently, we identified a pyruvate fermentation pathway in *Pseudomonas aeruginosa* sustaining anaerobic survival in the absence of alternative anaerobic respiratory and fermentative energy generation systems (M. Eschbach, K. Schreiber, K. Trunk, J. Buer, D. Jahn, and M. Schobert, *J. Bacteriol.* 186:4596–4604, 2004). Anaerobic long-term survival of *P. aeruginosa* might be essential for survival in deeper layers of a biofilm and the persistent infection of anaerobic mucus plaques in the cystic fibrosis lung. Proteome analysis of *P. aeruginosa* cells during a 7-day period of pyruvate fermentation revealed the induced synthesis of three enzymes involved in arginine fermentation, ArcA, ArcB, and ArcC, and the outer membrane protein OprL. Moreover, formation of two proteins of unknown function, PA3309 and PA4352, increased by factors of 72- and 22-fold, respectively. Both belong to the group of universal stress proteins (Usp). Long-term survival of a PA3309 knockout mutant by pyruvate fermentation was found drastically reduced. The oxygen-sensing regulator Anr controls expression of the *P*_{PA3309}-*lacZ* reporter gene fusion after a shift to anaerobic conditions and further pyruvate fermentation. PA3309 expression was also found induced during the anaerobic and aerobic stationary phases. This aerobic stationary-phase induction is independent of the regulatory proteins Anr, RpoS, RelA, GacA, RhlR, and LasR, indicating a currently unknown mechanism of stationary-phase-dependent gene activation. PA3309 promoter activity was detected in the deeper layers of a *P. aeruginosa* biofilm using a *P*_{PA3309}-*gfp* (green fluorescent protein gene) fusion and confocal laser-scanning microscopy. This is the first description of an Anr-dependent, anaerobically induced, and functional Usp-like protein in bacteria.

Pseudomonas aeruginosa is a highly adaptable bacterium that colonizes various environmental niches. It is also a leading opportunistic pathogen in human infections. *P. aeruginosa* is the dominant pathogen causing chronic respiratory infections of cystic fibrosis (CF) patients. This results in progressive lung damage and is the major cause of morbidity and mortality in CF patients (43). Recent data indicate that anaerobic conditions play an important role during persistent infection of the CF lung. *P. aeruginosa* forms biofilm-like microcolonies in the CF lung mucus embedded in an anaerobic environment (28, 47). Under these conditions, nitrate serves as an alternative electron acceptor sustaining growth under anaerobic conditions. Moreover, nitrate seems to favor the formation of more robust anaerobic biofilms (20, 50). Biofilm formation protects *P. aeruginosa* in hostile environments like the CF lung from the immune response of the host and leads to high antibiotic tolerance of the cells (7, 25, 41). A recent publication shows that oxygen-restricted cells in deeper layers of a colony biofilm are highly resistant to antibiotics due to stationary-phase conditions caused by oxygen depletion (4).

P. aeruginosa has a limited potential to survive in an anaerobic environment. Growth is supported by denitrification with

nitrate or nitrite (5, 8, 51). Moreover, arginine sustains moderate anaerobic growth via fermentation (44). Recently, we have shown that pyruvate—one of the most abundant metabolites in all cells—allows anaerobic long-term survival of *P. aeruginosa* (13). However, in contrast to denitrification and arginine fermentation, pyruvate fermentation does not sustain anaerobic growth. We identified three enzymes essential for pyruvate fermentation: phosphotransacetylase (Pta), acetate kinase (AckA), and a lactate dehydrogenase (LdhA). The operon encoding Pta and AckA is induced in response to oxygen limitation in dependence of the anaerobic regulatory protein Anr (13). The *ackA-pta* locus in *Escherichia coli* is induced in response to starvation during aerobic conditions (31). Pyruvate fermentation in *P. aeruginosa* might also play an important role as a general endogenous survival metabolism in response to energy starvation.

Here, we report the investigation of the physiological basis of pyruvate fermentation in *P. aeruginosa* starting with a proteome approach via two-dimensional (2D) gel electrophoresis. We identified two anaerobically induced Usp-type stress proteins (PA3309 and PA4352) and investigated the role of PA3309 during pyruvate fermentation and biofilm growth in more detail. In contrast to *E. coli* or Usp-type proteins investigated in *Mycobacteria* (33), our data indicate the presence of Usp-type proteins in *P. aeruginosa*, which are produced in response to oxygen limitation under the control of the oxygen-sensing regulator Anr. We found evidence for induction of the PA3309 gene in the stationary phase by an Anr-independent unknown regulator.

* Corresponding author: Mailing address: Institute of Microbiology, Technical University Braunschweig, Spielmannstr. 7, D-38106 Braunschweig, Germany. Phone: 49 531 3915857. Fax: 49 531 3915854. E-mail: m.schobert@tu-bs.de.

† Present address: Carlsberg Research Center, Biosector, Gamle Carlsberg Vej 10, DK-2500 Valby, Denmark.

TABLE 1. Strains of *P. aeruginosa* and *E. coli* and plasmids used in this study

Bacterial strain or plasmid	Genotype or phenotype	Source or reference
Strains		
<i>P. aeruginosa</i>		
PAO1	Wild type	11
PAO6261	PAO1 Δanr	49
PAO-MW20	PAO1 <i>rpoS::aacC1</i> Gm ^r	46
PAO6251	PAO1 $\Delta arcDABC$	17
PAO6281	PAO1 <i>gacA::Ω-Sp/Sm</i>	36
KS06	PAO1 <i>attB::(P_{PA3309}-lacZ)</i>	This study
KS08	PAO6261 <i>attB::(P_{PA3309}-lacZ)</i>	This study
KS10	PAO-MW20 <i>attB::(P_{PA3309}-lacZ)</i>	This study
KS11	PAO1 <i>attB::(mini-CTX-lacZ)</i>	This study
KS15	PAO1 <i>attB::(P_{PA3309}-gfp)</i>	This study
KS16	PAO1 $\Delta PA3309::aacC1$ Gm ^r	This study
KS17	PAO1 $\Delta PA3309$	This study
KS27	PAO1 <i>attB::(mini-CTX2)</i>	This study
KS28	KS17 <i>attB::(mini-CTX2)</i>	This study
KS29	KS17 <i>attB::(pKS13)</i>	This study
KS31	PAO1 <i>rhlR attB::(P_{PA3309}-lacZ)</i>	This study
KS32	PAO1 <i>lasR attB::(P_{PA3309}-lacZ)</i>	This study
KS33	PAO6281 <i>attB::(P_{PA3309}-lacZ)</i>	This study
KS35	PAO1 $\Delta relA$	This study
KS37	PAO1 $\Delta relA attB::(PPA3309-lacZ)$	This study
NB015	PAO1 $\Delta PA4352$	This study
<i>E. coli</i>		
DH10B	F ⁻ <i>mcrA</i> $\Delta(mrr-hsdRMS-mcrBC)$ $\phi 80dlacZ\Delta M15$ $\Delta lacX74$ <i>deoR recA1 endA1 araD139 $\Delta(ara leu)$ 7697 <i>galU galk</i> λ^{-} <i>rpsL nupG</i></i>	Gibco BRL (Invitrogen)
S17 λ pir	<i>pro thi hsdR⁺ Tc^r Sm^r</i> ; chromosome::RP4-2 Tc::Mu-Km::Tn7/ λ pir	9
SM10	<i>thi-1 thr leu tonA lacY supE recA::RP4-2-Tc::Mu</i> (Km ^r)	9
Plasmids		
pEX18Ap	Ap ^r <i>oriT⁺ sacB⁺</i> ; gene replacement vector with MCS from pUC18	23
Mini-CTX- <i>lacZ</i>	Tc ^r ; promoterless <i>lacZ</i> gene	2
Mini-CTX2	Tc ^r ; integration vector	24
pSB219.9A	Gm ^r ; <i>lasR::Gm</i> suicide vector	1
pSB224.10A	Tc ^r ; <i>rhlR::Tc</i> suicide vector	1
pPS858	Ap ^r Gm ^r ; source of gentamicin cassette	23
pFLP2	Ap ^r ; source of FLP recombinase	23
pMH305	Ap ^r Gm ^r ; source of <i>gfp</i> cassette	Morten Hentzer
pKS08	Ap ^r Gm ^r ; pMH305 with 526 bp of PA3309 promoter between EcoRI and BamHI	This study
pKS09	Tc ^r ; mini-CTX2 with NotI fragment liberated from pKS08 containing PA3309 promoter fused to <i>gfp</i>	This study
pKS10	Ap ^r Gm ^r ; pEX18Ap with 723-bp promoter of PA3309, Gm ^r - <i>gfp</i> fragment from pPS858 and 625 bp downstream of coding region of PA3309 between EcoRI and SalI	This study
pKS13	Tc ^r ; mini-CTX2 with 1,200-bp PCR fragment covering entire PA3309 gene and 455 bp of putative promoter region and 289 bp of downstream region between EcoRI and HindIII	This study
pKS15	Tc ^r ; mini-CTX- <i>lacZ</i> containing 492-bp fragment of putative promoter region of PA3309 gene between EcoRI and BamHI	This study
pKS18	Ap ^r Gm ^r ; pEX18Ap with 796-bp promoter of <i>relA</i> (PA0934), Gm ^r - <i>gfp</i> fragment from pPS858, and 824 bp of the 3' coding region of <i>relA</i> between EcoRI and HindIII	This study
pNB007	Ap ^r Gm ^r ; pEX18Ap with 596-bp promoter PA4352, Gm ^r - <i>gfp</i> fragment from pPS858, and 617 bp downstream of coding region of PA4352 between SacI and HindIII	This study

MATERIALS AND METHODS

Bacterial strains and growth conditions. The bacterial strains and plasmids used in this study are shown in Table 1. For standard molecular biology protocols, *E. coli* and *P. aeruginosa* strains were grown in LB medium as described before (13). Antibiotics were used at the following concentrations for *P. aeruginosa*: carbenicillin at 500 μ g/ml, gentamicin at 200 μ g/ml, and tetracycline at 100 μ g/ml.

Pyruvate fermentation and anaerobic energy starvation experiments. Pyruvate fermentation experiments with *P. aeruginosa* were performed with a slightly modified protocol. Instead of OS medium, we used a potassium phosphate (100 mM)-buffered LB medium at a pH value of 7.4. *P. aeruginosa* was grown aerobically up to an optical density at 578 nm (OD₅₇₈) of 0.3. Cells were shifted to anaerobic conditions by transferring the culture to rubber-stoppered bottles and immediate addition of 40 mM pyruvate. Incubation was carried out at 37°C

without shaking. Viable cell counts were determined as described before (13). Formation of fermentation products and consumption of pyruvate were analyzed using high-performance liquid chromatography analysis as outlined previously (13). Control experiments revealed no consumption of pyruvate during aerobic growth in LB medium up to an OD₅₇₈ of 0.3.

For anaerobic long-term energy starvation experiments, 40 ml LB medium was inoculated with 10⁸ *P. aeruginosa* cells/ml and incubated under anaerobic conditions for 20 days. Cell numbers increased up to 4 \times 10⁸ cells/ml within 24 h. After 24 h, cells faced severe energy starvation, resulting in a dramatic reduction in the number of cells/ml.

Construction and testing of the promoter-*lacZ* reporter gene fusions. Chromosomal promoter-*lacZ* reporter gene fusions were constructed using the mini-CTX-*lacZ* vector. A 492-bp PCR product, covering the region from 450 bp

upstream and 26 bp downstream of the translational start of the PA3309 gene, was generated using primer Pa-3309-for (5'-CGGAATTCGCCATGGACGAGGAATG-3') and Pa-3309-rev (5'-CGGGATCTCCACGGCTACCA GAATG-3'). Pa-3309-for contained an EcoRI restriction site (underlined) at the 5' end, and Pa-3309-rev contained a restriction site for BamHI (underlined) also at its 5' end. The EcoRI- and BamHI-digested PCR product was cloned into the EcoRI and BamHI sites of mini-CTX-lacZ to generate pKS15. Transfer of pKS15 in *P. aeruginosa* was carried out by a diparental mating using *E. coli* S17 λ -pir as donor. The CTX integrase of pKS15 promoted integration of the vector into the *attB* site of the *P. aeruginosa* genome. The vector was transferred into PAO1, and the *anr*, *rpoS*, *rhlR*, *lasR*, *gacA*, and *relA* mutant strains to generate the *P. aeruginosa* strains KS06, KS08, KS10, KS31, KS32, KS33, and KS37, respectively (see also Table 1). In these mutant strains, parts of the mini-CTX-lacZ vector containing the tetracycline resistance cassette were deleted using a flip-pase (FLP) recombinase encoded on the pFLP2 plasmid. Reporter gene fusion assays were performed as outlined before in detail (13, 37). The obtained activities were given in Miller units (29).

Construction of a P_{PA3309} -gfp fusion. We used a transcriptional P_{PA3309} -gfp (green fluorescent protein [GFP] gene) reporter gene fusion to monitor the promoter activity of PA3309 in biofilms. A 526-bp fragment of the PA3309 promoter region without the translational start codon was amplified by PCR with primers oKS16 (5'-GGAATTCGCCATGGACGAGGAATG-3'), containing an EcoRI restriction site at the 5' end, and oKS17 (5'-CGGGATCCAAGGTGT CCCTCCAGAGTG-3'), with a BamHI site. The PCR product was cloned into pMH305 upstream of *gfp* to generate pKS08. The P_{PA3309} -gfp reporter gene fusion was liberated via NotI digestion and ligated into mini-CTX2 to generate pKS09. This vector was transferred and integrated into the *P. aeruginosa* genome using the procedure described above. The resulting strain was named KS15.

Construction of *P. aeruginosa* Δ PA3309, Δ PA4352, Δ relA, *rhlR*:Tc^r, and *lasR*::Gm^r mutants. Unmarked gene deletion mutants were obtained using the well-established strategies based on *sacB* counter selection and FLP recombinase excision (23). First a suicide vector, pKS10, was constructed to replace the PA3309 gene with a gentamicin cassette. After transfer of this vector into *P. aeruginosa*, a double-crossover mutant was obtained by *sacB*-based counterselection. The resulting mutant, KS16, was verified by Southern blot analysis. Finally, FLP recombinase encoded on the pFLP2 plasmid removed the FRT-flanked gentamicin cassette to generate KS17. To construct the suicide vector pKS10, the BamHI-digested gentamicin resistance cassette of pPS858 was cloned between two PCR fragments of the PA3309 gene in the multiple cloning site of pEX18Ap. The two PCR fragments contained DNA homologous to the upstream and downstream areas of the PA3309 gene. A 723-bp fragment containing the upstream promoter region of the PA3309 gene was amplified using primers oKS10 (5'-CGGAATTCGAACAAGGCGCTGAAG-3'), with an EcoRI restriction site at the 5' end, and oKS11 (5'-CGCGGATCCAACTCAAGGACACTGTA-3'), with a BamHI site. The primers oKS12 (5'-CGGGATCTCCGGTCTGCTGGT-3'), with a BamHI restriction site, and oKS13 (5'-ACGCGTTCGACACGCCATCATCGTCCT-3'), with a SalI restriction site, were used for the amplification of 625 bp of the PA3309 downstream region.

The unmarked Δ PA4352 mutant (NB015) was generated by the same strategy described above using the suicide vector pNB007. In this suicide vector, the following PCR fragments flanked the gentamicin resistance cassette: The 596-bp fragment of the upstream region of the PA4352 gene was amplified using primers oNB01 (5'-CGAGCTCTACGGCGACTTCTGTAAGG-3'), with a SacI restriction site, and oNB02 (5'-CGGGATCCAAGCGGATGCTTCGGACT-3'), with a BamHI site. The primers oNB03 (5'-CGCGGATCCCTTCGCGCGCGCTGA-3'), with a BamHI site, and oNB04 (5'-CCCAAGCTTCCCTGGCGCCGTGACC-3'), with a HindIII site, amplified 617 bp of the corresponding downstream region of PA4352.

The Δ relA mutant KS35 was constructed by the same strategy described above. For construction of the suicide vector pKS18, the primers oKS40 (5'-GGAATTCGGCCAGTGCATTGCTGTG-3'), with an EcoRI restriction site at the 5' end, and oKS41 (5'-CGGGATCTTACCACGGTGCAGCTAG-3'), with a BamHI site, amplified 796 bp of the putative promoter region of the *relA* gene. The primers oKS42 (5'-CGCGGATCCCGAGCAGCTGAGATCA-3'), with a BamHI site, and oKS43 (5'-CCCAAGCTTGGGCGAGTTCGAGAGC-3'), with a HindIII site, were used to amplify 824 bp of the 3' end of *relA*.

The two quorum-sensing mutants KS31 (*rhlR*:Tc^r) and KS32 (*lasR*::Gm^r) were constructed as described previously (1) using KS06 as the parent strain.

Plasmid construction for complementation of the PA3309 knockout mutant. For construction of the complementation plasmid for the chromosomal PA3309 knockout mutant, a 1,200-bp PCR product covering 455 bp of the PA3309 promoter region, the PA3309 gene, and 289 bp downstream of PA3309 was amplified using primers oKS14 (5'-CGGAATTCGCCATGGACGAGGAAC

TTATCCACGTGCGGATGGTC-3'), with a HindIII site. The product was digested with EcoRI and HindIII and ligated into mini-CTX2 to generate pKS13. The vector was transferred into *P. aeruginosa* KS17 to generate KS29 by a diparental mating with *E. coli* S17 λ pir and integrated at the *attB* site in the genome as described above. As a control, the empty mini-CTX2 vector was integrated into the genome of PAO1 and KS17 to generate KS27 and KS28, respectively.

Proteomic analysis. To prevent changes of the cellular protein pattern during cell harvesting and cell extract preparation, the culture was mixed with a double volume of ice-cold potassium phosphate-buffer (0.1 M; pH 7.4) and allowed to cool for 20 min. Cells were centrifuged at 8,000 \times g for 30 min at 4°C and washed twice with potassium phosphate buffer. Cells were resuspended in a small volume of potassium phosphate buffer. The protein concentration in whole-cell suspensions was determined using the bicinchoninic acid protein assay (Sigma, Taufkirchen, Germany). Cells were disrupted by incubation of a 360- μ l culture aliquot with 150 μ l NaOH for 1 h at 70°C. For protein isolation, we modified a protocol described previously (19) and extracted proteins directly from whole cells with phenol and a subsequent acetone precipitation. The precipitated proteins were solubilized in sample buffer consisting of 7 M urea, 2 M thiourea, 4% 3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate (CHAPS), 50 mM dithiothreitol, and 2% ampholytes (Bio-Lyte; Bio-Rad, Munich, Germany). The protein concentration was determined in the sample buffer using the PlusOne 2D Quant kit (Amersham Biosciences, Freiburg, Germany). 2D gel electrophoresis was performed using immobilized pH gradient (IPG) strips 11 or 17 cm in length covering two different pH ranges (pH 4.7 to 5.9 or 5 to 8) (IPG Ready Strips; Bio-Rad, Munich, Germany). For the narrow pH range, the IPG strips (11 cm) were rehydrated overnight in rehydration buffer containing 50 μ g of protein. Isoelectric focusing (IEF) was carried out at 20°C under mineral oil in the PROTEAN IEF cell (Bio-Rad, Munich, Germany) for a total of 35,000 V \cdot h. The focused IPG strips were reduced for 15 min in a sodium dodecyl sulfate (SDS) equilibration solution containing 15 mM dithiothreitol and afterwards alkylated twice for 15 min in the same buffer containing 150 mM iodoacetamide prior to SDS-polyacrylamide gel electrophoresis (SDS-PAGE). The IPG strips were transferred to 10.5 to 14% SDS-PAGE (Criterion Tris-HCl gel; Bio-Rad, Munich, Germany) gels, and electrophoresis was performed at a constant 200 V for 55 min. Large IPG strips (17 cm, pH 5 to 8) were loaded with 700 μ g of protein, and IEF was conducted for a total of 110,000 V \cdot h. SDS-PAGE was performed at a constant temperature of 20°C with 1 W per gel for approximately 20 h using 10% polyacrylamide gels (25.5 by 20.5 cm). All gels were stained with ruthenium(II)-tris-(bathophenanthroline disulfonate) (RuBPS) as described before (35). Gels were documented with an FX-Scanner (Bio-Rad, Munich, Germany). Analysis and quantification of differential protein spot patterns were performed by using the software Z3 (Compugen, Tel Aviv, Israel). Gel spots were excised and treated using a method described before introducing minor modifications (40). Briefly, the gel pieces were washed with water, dehydrated with acetonitrile (ACN), and digested with trypsin (sequencing grade; Promega). Peptides were extracted and collected in four elution steps (each 15 min, 37°C) using 25 mM NH₄HCO₃, ACN, 5% formic acid, and again ACN. Extracted peptides were purified using ZipTip C₁₈ microcolumns (Millipore), following the manufacturer's instructions. Proteins were identified by peptide-mass fingerprint (PMF) as well as post-source decay fragmentation data recorded on a Bruker Ultraflex matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometer. PMF data were analyzed using an internal MASCOT server at the GBF (version 1.9; Matrix Science) (34) and the NCBI database (restricted to the taxon *Pseudomonas aeruginosa*). Only peptides with a MASCOT rank of 1 were considered significant and used for the combined peptide score. The criteria used to accept protein identifications based on PMF data included the extent of sequence coverage (minimum of 30%), the number of peptides matched (minimum of 5), and the score of probability (minimum of 70 for the Mowse score). Lower-scoring proteins were either verified manually or rejected.

Biofilm experiments. Biofilm studies were performed in three-channel flow cells with individual channel dimensions of 1 by 4 by 40 mm supplied with a flow of 3 ml h⁻¹ of AB medium (22) supplemented with 300 μ M glucose and where indicated with 50 mM KNO₃ (6). The system was assembled and cultures for inoculation were prepared as described before (6, 21). Biofilms were stained with Syto62 (Molecular Probes) to visualize the biofilm matrix. Dead cells in the biofilm were visualized with propidium iodide (Sigma). Microscopy and image acquisition were done as outlined in reference 22 with a Zeiss LSM 510 confocal laser-scanning microscope (CLSM) (Carl Zeiss, Jena, Germany). Quantitative image analysis was done using the COMSTAT software (22). Images were processed using the Imaris software (Bitplane AG, Zurich, Switzerland). For protein analysis, biofilms were grown in silicone tubes (30 cm; inner




FIG. 1. (A) Two-dimensional image of crude extracts of *P. aeruginosa* PAO1 incubated under pyruvate fermentation conditions for 7 days. Outlined areas in the gel indicate zones of the 2D gel that are represented in Fig. 1B. (B) Enlarged 2D gel images showing protein extracts of the *P. aeruginosa* PAO1 aerobic culture immediately before the shift to anaerobic pyruvate fermentation conditions and the pyruvate fermentation culture shown in Fig. 1A. Numbers of the boxed spots indicate identified proteins that are synthesized at higher levels during pyruvate fermentation; the numbers correlate with the numbers given in Table 2.

square, 0.5 cm), in AB medium as described above and supplied with a flow rate of 20 ml h⁻¹.

RESULTS

Proteome analysis of *P. aeruginosa* under pyruvate fermentation conditions. Recently, we showed that *P. aeruginosa* can survive anaerobically for several weeks by pyruvate fermentation in the absence of nitrate, nitrite, or arginine (13). Pyruvate fermentation allows survival but not growth of *P. aeruginosa* cells during severe anaerobic energy starvation conditions. We used a proteomics approach to identify proteins which are preferentially synthesized during pyruvate fermentation. Long-term survival pyruvate fermentation experiments using cultures grown in LB medium were set up as described previously (13). For this purpose, cells were grown aerobically to an OD₅₇₈ of 0.3, immediately shifted to anaerobic flasks containing pyruvate at a final concentration of 40 mM, and further incubated for 7 days at 37°C. Viable cell counts of three independent cultures were determined and remained almost constant during the whole experiment ($2.8 \times 10^8 \pm 1.3 \times 10^8$ cells per ml of medium at day 0 and $1.6 \times 10^8 \pm 1.0 \times 10^7$ at day 7).

We compared the protein pattern of the aerobically grown culture in phosphate buffered LB medium (OD₅₇₈ = 0.3) with that of the culture incubated for 7 days under pyruvate fermentation conditions (phosphate-buffered LB medium plus 40 mM pyruvate, Fig. 1A and B). The concentration of 11 proteins was found highly increased at least ninefold under anaerobic pyruvate fermentation conditions (Table 2). MALDI-TOF

analysis identified the 11 separated proteins as representing 6 different proteins. Three proteins involved in arginine fermentation (ArcA, ArcB, and ArcC) were found highly upregulated as well as the outer membrane protein OprL and two proteins, PA3309 and PA4352, which are annotated as conserved hypothetical proteins but which each have universal stress protein motifs (Table 2 and Fig. 1B). None of these proteins was expected to be involved in or contribute to pyruvate fermentation. The results of the proteome analysis raised the question of whether proteins involved in arginine fermentation or the two hypothetical proteins contribute to survival during pyruvate fermentation.

Contribution of *arcDABC* and PA3309 to anaerobic long-term survival of *P. aeruginosa*. Since we used a complex medium containing arginine, we investigated if arginine fermentation contributes to survival under anaerobic pyruvate fermentation conditions. The $\Delta arcDABC$ mutant PAO6251 cannot grow anaerobically by arginine fermentation. Viable cell counts of the $\Delta arcDABC$ mutant decreased by a factor of 30 compared to the wild type during 20 days of pyruvate fermentation. In contrast, the viable cell counts of the wild-type control, which was incubated under anaerobic conditions without pyruvate, decreased by a factor of 500 (Fig. 2). These results demonstrated that arginine fermentation contributes to anaerobic long-term survival: however, to a lesser extent than the PA3309 protein.

Next, we investigated if the hypothetical protein PA3309 promotes survival during pyruvate fermentation. First, a PA3309 chromosomal knockout mutant strain was constructed

TABLE 2. Identified proteins under pyruvate fermentation in LB medium^a

Spot no.	PA no.	Protein	Description	Regulation (fold) ^b
1	PA5171	ArcA	Arginine deiminase	132
2	PA5171	ArcA	Arginine deiminase	9
3	PA5171	ArcA	Arginine deiminase	Unmatched
4	PA5172	ArcB	Catabolic ornithine carbamoyltransferase	29
5	PA5172	ArcB	Catabolic ornithine carbamoyltransferase	14
6	PA5173	ArcC	Carbamate kinase	Unmatched
7	PA5173	ArcC	Carbamate kinase	60
8	PA4352		Conserved hypothetical protein	22
9	PA0973	OprL	Outer membrane protein	99
10	PA0973	OprL	Outer membrane protein	Unmatched
11	PA3309		Conserved hypothetical protein	72

^a PA number, protein short name, and description are shown according to the *Pseudomonas* database (www.pseudomonas.com).

^b Shown are the ratios of the amount of proteins from a *P. aeruginosa* PAO1 culture incubated for 7 days under pyruvate fermentation conditions at 37°C in phosphate-buffered LB medium to the amount of protein in a culture grown under aerobic conditions (see Materials and Methods). Proteins were separated by 2D gel electrophoresis and stained with RuBPS (see Materials and Methods for details). Protein ratios of 2D gel images were determined by the Z3 software (Compugen, Tel Aviv, Israel). "Unmatched" describes a protein spot which is detected only in the protein pattern of the pyruvate fermentation culture.

as described in Materials and Methods. The PA3309 mutant strain exhibited a clear 400-fold reduction of anaerobic long-term survival over a 20-day period (Fig. 2). We were able to complement the PA3309 mutant by chromosomal integration of pKS13, which contains the cloned PA3309 gene and the putative promoter region (Table 1). Survival rates of the complemented mutant strain (KS29) do not completely reach the wild-type level, but were found significantly increased (100-fold) compared to those of the PA3309 mutant (KS17) and the PA3309 mutant containing the empty mini-CTX2 vector integrated in the *attB* locus (KS28; data for this mutant not shown). A PA4352 knockout mutant (NB015) without the second Usp-type stress protein showed no reduced survival during pyruvate fermentation (Fig. 2).

Our studies revealed an essential role of PA3309 for anaerobic long-term survival during pyruvate fermentation.

The conserved hypothetical protein PA3309 belongs to the group of anaerobically induced universal stress proteins. The conserved hypothetical protein PA3309 consists of 151 amino acids with a calculated mass of 16,496 Da and a theoretical pI of 5.31. The Pfam database indicates the presence of a single Usp domain (Pfam accession number PF00582) which originates from the universal stress protein A (UspA) of *E. coli*. PA3309 shares the single conserved Usp domain, a similar molecular mass, and a moderate amino acid sequence identity of 37% with *E. coli* UspA. Six Usp-type stress protein paralogues have been identified in *E. coli* which are produced in response to a variety of different stress conditions, most of them leading to growth arrest (27). We tested if the PA3309 knockout mutant showed similar phenotypes to the *E. coli uspA* mutant in response to UV stress and stationary-phase survival. However, no similar behavior was observed (data not shown). Despite the observed phenotype of the PA3309 knockout mutant during pyruvate fermentation, we did not observe a defect in anaerobic denitrifying growth using different media

supplemented with 50 mM nitrate. However, PA3309 contributes to survival of severe anaerobic energy stress conditions. We incubated wild-type *P. aeruginosa* and the PA3309 mutant anaerobically in LB medium without nitrate, nitrite, or pyruvate. Within 20 days, cell numbers of the wild-type strain decreased by a factor of 9.0×10^2 , while cell numbers of the mutant strain decreased by a factor of 9.7×10^4 .

PA3309 is induced upon a shift to anaerobic conditions by the oxygen regulator Anr. We studied the regulatory behavior of the PA3309 promoter towards various environmental stimuli and corresponding metabolic conditions with a chromosomal transcriptional promoter-*lacZ* reporter gene fusion. We monitored β -galactosidase activities of the P_{PA3309} -*lacZ* reporter gene fusion (KS06) during pyruvate fermentation for the first 4 days. The β -galactosidase activities increased linearly up to $1,594 \pm 20$ Miller units during the first 4 days (Fig. 3A), indicating strong anaerobic induction. Anr is a global transcriptional regulatory protein of the Crp-Fnr family which activates gene expression in *P. aeruginosa* in response to oxygen limitation (16, 39). We detected a putative Anr box 86 bp upstream of the translational start codon in the presumed promoter region of PA3309 using tools of the PRODORIC database (30). We also checked dependence of the PA3309 promoter on Anr. No significant increase was detected when the P_{PA3309} -*lacZ* reporter gene fusion was monitored in the *anr* mutant strain PAO6261 (Fig. 3A). We also measured a strong increase in β -galactosidase activity of the P_{PA3309} -*lacZ* reporter gene fusion in wild-type *P. aeruginosa* upon a shift to anaerobic conditions in the absence of pyruvate. Again, under these conditions the *anr* mutant failed to induce the PA3309




FIG. 2. Anaerobic survival of *P. aeruginosa* wild type (■), the PAO6251 ($\Delta arcDABC$) mutant (×), the $\Delta PA4352$ mutant (▲), the $\Delta PA3309$ mutant (●), and the complemented $\Delta PA3309$ mutant (○) in the presence of 40 mM pyruvate. CFU of the *P. aeruginosa* wild type without pyruvate served as a control (□). The *P. aeruginosa* wild type and mutants were grown aerobically in phosphate-buffered LB medium. At an OD_{578} of 0.3, cultures were transferred to rubber-stoppered bottles and 40 mM pyruvate was added. Survival under anaerobic conditions without alternative electron acceptors was determined with viable cell counts on agar plates. Graphs represent the results of at least three independent experiments. Standard deviations were 44% until day 5 and below 5% between days 10 and 20 and are omitted for the sake of clarity.




FIG. 3. (A) Diagram showing the expression pattern of the *P. aeruginosa* wild type containing the P_{PA3309} -*lacZ* reporter gene fusion (strain KS06; gray bars) and the *anr* mutant (PAO6261) containing the P_{PA3309} -*lacZ* reporter gene fusion (KS08; white bars) during the first 4 days of a pyruvate fermentation experiment. Cells were grown in phosphate buffered LB medium at 37°C (for details see Materials and Methods). Time point 0 is the aerobic culture shortly before the transfer to an anaerobic flask. (B) Diagram showing the expression pattern of the *P. aeruginosa* wild type containing the P_{PA3309} -*lacZ* reporter gene fusion (strain KS06; gray bars) and the *anr* mutant (PAO6261) containing the P_{PA3309} -*lacZ* reporter gene fusion (KS08; white bars) in a shift experiment. Cells were grown at 37°C in LB medium plus 50 mM KNO_3 up to an OD_{578} of 0.4 (time point 0) and transferred to a rubber-stoppered flask for a subsequent 4-h incubation (time point 4). β -Galactosidase activities were determined at the indicated time points. Experiments were repeated three times.

promoter activity (Fig. 3B). Clearly, Anr induces anaerobic expression of PA3309 independent of the presence of pyruvate.

PA3309 is induced in stationary phase. Further examination of the P_{PA3309} -*lacZ* behavior during various anaerobic growth phases revealed a 5.3-fold induction during anaerobic exponential growth compared to aerobic exponential growth. Interestingly, the PA3309 promoter was further induced 1.8-fold in

the anaerobic stationary phase (Table 3). Next, we investigated the behavior of the PA3309 promoter during the aerobic stationary phase. To our surprise, a 5.4-fold induction was observed. Gene expression of PA3309 in the aerobic stationary phase was further confirmed by 2D gel electrophoresis (Fig. 4B and C). In contrast to pyruvate fermentation, where we identified only one protein spot representing PA3309 (Fig. 4D), two spots representing PA3309 were identified in the protein pattern of aerobic stationary-phase cells (Fig. 4C). This might indicate a phosphorylation of PA3309 as reported for *E. coli* UspA (15). The Anr regulator was the candidate responsible for the aerobic induction of the P_{PA3309} -*lacZ* reporter gene fusion upon entry into the stationary phase, since the strong respiration of a high-cell-density culture leads to oxygen limitation. However, expression of the P_{PA3309} -*lacZ* reporter gene fusion in the *anr* mutant still increased 4.2-fold upon entry in the aerobic stationary phase (Table 3). Therefore, promoter activity in the aerobic stationary phase is independent of Anr. Expression of the *uspA*, *uspC*, *uspD*, and *uspE* genes in *E. coli* upon entry in the stationary phase requires stringent control via the nucleotide guanosine 3', 5'-bisdiphosphate (ppGpp) and the RelA protein (14, 18, 26). In *P. aeruginosa*, the RelA protein (PA0934) synthesizes ppGpp in response to amino acid starvation conditions or in the stationary phase (12, 42). Deletion of the *P. aeruginosa* *relA* gene abolishes the production of ppGpp and also decreases production of RpoS (12, 42). Expression of the *P. aeruginosa* PA3309 promoter in a *P. aeruginosa* *relA* mutant remained unchanged compared to that under wild-type conditions (Table 3). Furthermore, we did not detect increased PA3309 promoter activity when wild-type cells faced carbon starvation during growth in 1/20 LB medium (data not shown). Therefore, the stringent response system does not contribute to PA3309 regulation under the tested conditions.

The stationary-phase sigma factor RpoS activates gene expression upon entry into the stationary phase. Anr-dependent genes like *azu* encoding the blue copper protein azurin have been reported to be RpoS dependent (45). However, an *rpoS* mutant showed no influence on the PA3309 promoter activity under aerobic or anaerobic stationary-phase conditions (Table 3).

In *Mycobacterium smegmatis*, a two-component regulator similar to the *Mycobacterium tuberculosis* DevR protein was shown to induce expression of three genes encoding Usp-type proteins (3, 32). DevR shares domain organization and 37% identity on the amino acid sequence level to the *P. aeruginosa* GacA regulator of the global GacA/GacS system (global antibiotics and cyanide control). This system regulates the expression of multiple phenotypes in pseudomonads. Again, no change in PA3309 promoter activity was found in a *gacA* mutant strain (Table 3). Similar observations were made using *lasR* and *rhlR* mutants carrying defects in both quorum-sensing systems of *P. aeruginosa* (Table 3).

So far, we have not checked if PA3309 expression in the stationary phase is RecA/FtsK dependent, as shown for *E. coli* *uspA* (10). However, since the PA3309 deletion mutant is not sensitive to UV exposure (data not shown), a RecA/FtsK-dependent regulation seems unlikely.

Expression of PA3309 in biofilms. We checked the spatial distribution of PA3309 promoter induction within biofilms, its dependence on the Anr regulatory protein, and the effect of

TABLE 3. Expression of the P_{PA3309} -*lacZ* reporter gene fusion in *P. aeruginosa* wild type and mutants

<i>P. aeruginosa</i> strain	β -Galactosidase activity (Miller units) or fold induction ^a					
	Aerobic			Anaerobic		
	Exponential phase ^b	Stationary phase ^c	Fold induction ^d	Exponential phase ^b	Stationary phase ^c	Fold induction ^e
Wild type	214.4 \pm 9.6	1,149.2 \pm 70.4	5.4	1,339.1 \pm 171.7	2,377.5 \pm 270.8	1.8
<i>anr</i> mutant	218.1 \pm 8.0	914.1 \pm 23.5	4.2	ND ^f	ND	ND
<i>rpoS</i> mutant	284.3 \pm 164.1	1,160.5 \pm 500.1	4.1	956.2 \pm 117.6	2,262.4 \pm 0	2.4
<i>gacA</i> mutant	229.7 \pm 11.3	985.4 \pm 47.0	4.3	1,161.2 \pm 118.8	2,524.9 \pm 154.4	2.2
<i>rhIR</i> mutant	248.5 \pm 4.0	1,342.2 \pm 87.5	5.4	1,314.8 \pm 56.3	2,747.9 \pm 818.2	2.1
<i>lasR</i> mutant	217.1 \pm 2.9	1,139.8 \pm 79.4	5.3	1,172.6 \pm 44.8	2,106.6 \pm 65.5	1.8
<i>relA</i> mutant	455.5 \pm 20.8	1,577.5 \pm 70.0	3.5	ND	ND	ND

^a Shown are the β -galactosidase activities of the P_{PA3309} -*lacZ* reporter gene fusion in wild-type PAO1 and various mutants under aerobic and anaerobic conditions as described in Materials and Methods.

^b Exponential growth phase at an OD₅₇₈ of 0.5 or 0.3 (aerobic and anaerobic culture, respectively).

^c Stationary phase represents 16 h after entry into the stationary phase.

^d Fold induction from aerobic exponential phase to aerobic stationary phase.

^e Fold induction from anaerobic exponential phase to anaerobic stationary phase.

^f ND, not determined.

nitrate. To visualize the promoter activity of PA3309 in biofilms, a P_{PA3309} -*gfp* reporter gene fusion was constructed and transferred to the *P. aeruginosa* chromosome (see Materials and Methods). Biofilms were grown for 6 days in flow cells and inspected by CLSM. As depicted in Fig. 5A, PA3309 promoter activity visualized by GFP fluorescence, represented as yellow areas in the red-colored biofilm matrix, was only detectable in the deeper layers of the biofilm. Quantitative analysis of the biofilms using COMSTAT revealed a twofold increase of GFP

fluorescence when biofilms grew in the presence of nitrate (Fig. 5B). Since GFP requires small amounts of oxygen for activity, the deeper layers of the inspected biofilms were not strictly anaerobic but oxygen limited. In control experiments, biofilms were stained with propidium iodide to visualize dead cells. Only a small portion of the biofilm cells (1 to 2%) were not alive (data not shown).

To examine if the expression of PA3309 in biofilms is Anr dependent, a proteome approach was applied. Biofilms of *P.*




FIG. 4. (A) 2D gel image of crude extracts of *P. aeruginosa* PAO1 incubated under pyruvate fermentation conditions for 7 days. The boxed area contains the one or two protein spots marked by an ellipse, which were identified to be PA3309 and represented in detail in panels B to F. Enlarged 2D gel images showing the area outlined in panel A of protein extracts of *P. aeruginosa* PAO1 aerobic culture (B) growing exponentially at an OD₅₇₈ of 0.4; (C) 16 h after entering stationary phase; (D) after 7 days under pyruvate fermentation; (E) of a 6-day-old biofilm (*P. aeruginosa* PAO1), and (F) a protein extract of the *anr* mutant PAO6261 grown as a biofilm for 6 days. Growth conditions of the biofilm experiments are outlined in Materials and Methods.




FIG. 5. Vertical sections showing spatial structures of 6-day-old *P. aeruginosa* KS15 expressing a P_{PA3309} -*gfp* transcriptional fusion. Biofilms were grown in AB minimal medium containing 300 μ M glucose without (A) and supplemented with (B) 50 mM KNO_3 . Scale bars represent 50 μ m. The biofilms were stained with Syto62 (Molecular Probes) to visualize the biofilm matrix. Promoter activity is visualized by GFP fluorescence, represented as yellow areas in the red-colored biofilm matrix. Using the COMSTAT software (22), GFP was calculated to be expressed in 13.9% of the total biomass in the absence of nitrate and 29.8% when medium contained 50 mM nitrate.

aeruginosa PAO1 and the *anr* mutant strain PAO6261 were grown for 6 days under aerobic conditions, since the *anr* mutant is unable to grow and, consequently, cannot form biofilms under strict anaerobic conditions. Figure 4E and F show the partial enlargements of a 2D gel image containing two PA3309 protein spots. Both strains contain similar amounts of PA3309. Therefore, we conclude that under these microaerobic conditions induction of the PA3309 promoter is independent of Anr, comparable to the Anr-independent induction in the aerobic stationary phase.

DISCUSSION

During pyruvate fermentation, the survival of *P. aeruginosa* depends on PA3309, a protein with the signature domain of universal stress proteins (Usp). The PA3309 protein contains a single Usp domain and shares 37% amino acid sequence identity with UspA of *E. coli*. Universal stress proteins have been identified in bacteria, archaea, plants, and fungi. Six different Usp proteins (UspA, UspC, UspD, UspE, UspF, and UspG) are present in *E. coli* (27). Although much is known about the regulation, the function of Usp-type proteins still remains unknown (27). The *E. coli* proteins are produced in response to a large number of different stresses, including DNA damage, stationary phase, and growth arrest. In some cases, Usp-type proteins have been linked to the resistance to DNA-damaging agents and respiratory uncouplers (18). However, a *P. aeruginosa* PA3309 deletion mutant shows no comparable phenotype to an *E. coli* *uspA* mutant strain, regarding UV resistance and survival in the stationary phase (data not shown). The phenotype of the PA3309 mutant seems to be restricted to anaerobic conditions. Besides the essential role of PA3309 during pyruvate fermentation, PA3309 also contributes to survival during

long-term anaerobic energy starvation in the absence of pyruvate. During these stress conditions, cell numbers of the wild type decrease dramatically by a factor of 9.0×10^2 while cell numbers of the PA3309 mutant decrease by a factor of 9.7×10^4 . Consistent with the anaerobic phenotypes, regulation of the PA3309 gene was found to be dependent on the oxygen-sensing regulatory protein Anr. The β -galactosidase activities of a P_{PA3309} -*lacZ* reporter gene fusion remained unchanged in an *anr* mutant strain upon a shift to anaerobic conditions, while they increased four- to eightfold (Fig. 3) in wild-type *P. aeruginosa*. As shown in Fig. 3, this anaerobic induction of the PA3309 promoter is independent of the presence of pyruvate. The regulatory behavior is consistent with the observed anaerobic phenotype of the PA3309 mutant. However, we also observed production of the PA3309 protein in the aerobic stationary phase and monitored a fivefold-increased β -galactosidase activity of the P_{PA3309} -*lacZ* reporter gene fusion. We showed that this induction in the stationary phase is independent of Anr. It is also independent of other regulators which induce promoter activity in the stationary phase, during starvation, or at high cell densities, such as RpoS, RelA, and GacA and the quorum-sensing regulators RhlR and LasR. We also did not find a phenotype of the PA3309 mutant during the aerobic stationary phase in LB medium with or without supplemented pyruvate as reported for the *E. coli* *uspA* mutant. These experiments clearly indicate a different role and regulation of the Usp-type protein PA3309 in *P. aeruginosa* compared to the Usp proteins in *E. coli*. Stationary-phase expression of the *P. aeruginosa* PA3309 promoter is independent of RelA, which induces expression of *usp* genes in *E. coli*, and GacA, which shows similarities to DevR of *M. smegmatis*.

To identify the role of PA3309 during the aerobic stationary phase, we are currently investigating the PA3309 promoter in more detail and aim to identify the regulator involved.

We also investigated the spatial distribution of PA3309 promoter induction within biofilms. Biofilms are dense bacterial communities attached to a surface and surrounded by an exopolysaccharide matrix. Differences in density and architecture determine the access to nutrients and oxygen within the biofilm. Oxygen limitation can start within the first 30 μ m below the surface of an aerobically grown *P. aeruginosa* biofilm (48). Proteome studies of *P. aeruginosa* biofilms indicate that a large portion of a mature biofilm population is under oxygen limitation (38). Moreover, *P. aeruginosa* forms even more robust biofilms when grown anaerobically in the presence of nitrate (50). Since the CF airway mucus is anaerobic (47), anaerobic biofilms might mirror a persistent infection situation in the CF lung. Oxygen limitation contributes to antibiotic resistance of the oxygen-restricted layers of aerobically grown biofilms (4). Previously, increased expression of PA3309 in aerobic and anaerobic biofilms was described (50). Our biofilm experiments clearly showed an induction of the PA3309 promoter in the deeper layers of biofilms and an increased promoter activity when biofilms were grown in the presence of nitrate. However, 2D gel electrophoresis revealed that PA3309 production in biofilms is independent of the oxygen-sensing regulator Anr. Currently, we are investigating if PA3309 also contributes to survival of cells in deeper layers of a biofilm.

2D gel analysis of *P. aeruginosa* cells during pyruvate fermentation revealed production of a second Usp-type stress

protein, PA4352, as well as proteins involved in arginine fermentation. While a PA4352 mutant had no phenotype during pyruvate fermentation, a mutant with a defect in arginine fermentation, Δ *arcDABC*, showed decreased survival. Since we used a complex medium and shifted the culture to anaerobic conditions during the early exponential phase, minor amounts of arginine could support survival. We determined arginine levels in LB medium to be approximately 1.6 mM (data not shown), but these amounts do not allow survival in the absence of pyruvate (see the wild-type control in Fig. 2). However, the mutant with a defect in arginine fermentation also grows poorly, even in aerobiosis (17). This suggests that the arginine deiminase operon in general supports growth and survival of *P. aeruginosa*, including pyruvate fermentation.

This is the first report of an Usp-type stress protein in *P. aeruginosa* which contributes to anaerobic survival and pyruvate fermentation and which is produced in response to anaerobiosis in an Anr-dependent manner. Usp-type stress proteins recently gained attention from investigations using *Mycobacteria*. Data from proteome and transcriptome analysis showed that Usp-type proteins of *Mycobacteria* were produced upon oxygen limitation and phagocytosis (33). The stationary-phase regulator DevR, which is required for oxygen starvation, was identified to control *usp* gene expression in *Mycobacterium smegmatis* (3, 32). Oxygen limitation has been shown to induce a transition from active growth to a nonreplicative persistent stage important for *Mycobacterium tuberculosis* latency and infection (33). Our results indicate that an Usp-type protein in *P. aeruginosa* also contributes to anaerobic survival and may play a role in survival in anaerobic mucus plaques in CF lungs.

ACKNOWLEDGMENTS

We thank Dieter Jahn for critically reading the manuscript. We are indebted to Dieter Haas (Universite de Lausanne, Switzerland) for providing the *P. aeruginosa gacA*, *arcDABC*, and *anr* mutants. We thank E. P. Greenberg (University of Iowa, Ames) for supplying the *P. aeruginosa rpoS* mutant and Miguel Camara (University of Nottingham, Nottingham, Great Britain) for providing the plasmids pSB219.9A and pSB224.10A for construction of the *lasR* and *rhlR* knockout mutants in *P. aeruginosa*. We thank H. P. Schweizer (University of Colorado, Boulder) for providing the pEX18Ap, mini-CTX-*lacZ*, pFLP2, and mini-CTX2 plasmids.

This investigation was funded by funds from the Deutsche Forschungsgemeinschaft, the German Research Centre for Biotechnology, and the Fonds der Chemischen Industrie. K.S. was supported by the DFG-European Graduate College 653.

REFERENCES

1. Beatson, S. A., C. B. Whitchurch, A. B. T. Semmler, and J. S. Mattick. 2002. Quorum sensing is not required for twitching motility in *Pseudomonas aeruginosa*. *J. Bacteriol.* **184**:3598–3604.
2. Becher, A., and H. P. Schweizer. 2000. Integration-proficient *Pseudomonas aeruginosa* vectors for isolation of single-copy chromosomal *lacZ* and *lux* gene fusions. *BioTechniques* **29**:948–950, 952.
3. Boon, C., and T. Dick. 2002. *Mycobacterium bovis* BCG response regulator essential for hypoxic dormancy. *J. Bacteriol.* **184**:6760–6767.
4. Borriello, G., E. Werner, F. Roe, A. M. Kim, G. D. Ehrlich, and P. S. Stewart. 2004. Oxygen limitation contributes to antibiotic tolerance of *Pseudomonas aeruginosa* in biofilms. *Antimicrob. Agents Chemother.* **48**:2659–2664.
5. Carlson, C. A., and J. L. Ingraham. 1983. Comparison of denitrification by *Pseudomonas stutzeri*, *Pseudomonas aeruginosa*, and *Paracoccus denitrificans*. *Appl. Environ. Microbiol.* **45**:1247–1253.
6. Christensen, B. B., C. Sternberg, J. B. Andersen, R. J. Palmer, Jr., A. T. Nielsen, M. Givskov, and S. Molin. 1999. Molecular tools for study of biofilm physiology. *Methods Enzymol.* **310**:20–42.
7. Costerton, J. W., P. S. Stewart, and E. P. Greenberg. 1999. Bacterial biofilms: a common cause of persistent infections. *Science* **284**:1318–1322.
8. Davies, K. J., D. Lloyd, and L. Boddy. 1989. The effect of oxygen on denitrification in *Paracoccus denitrificans* and *Pseudomonas aeruginosa*. *J. Gen. Microbiol.* **135**:2445–2451.
9. de Lorenzo, V., and K. N. Timmis. 1994. Analysis and construction of stable phenotypes in gram-negative bacteria with Tn5- and Tn10-derived minitransposons. *Methods Enzymol.* **235**:386–405.
10. Diez, A., N. Gustavsson, and T. Nystrom. 2000. The universal stress protein A of *Escherichia coli* is required for resistance to DNA damaging agents and is regulated by a RecA/FtsK-dependent regulatory pathway. *Mol. Microbiol.* **36**:1494–1503.
11. Dunn, N. W., and B. W. Holloway. 1971. Pleiotrophy of p-fluorophenylalanine-resistant and antibiotic hypersensitive mutants of *Pseudomonas aeruginosa*. *Genet. Res.* **18**:185–197.
12. Erickson, D. L., J. L. Lines, E. C. Pesci, V. Venturi, and D. G. Storey. 2004. *Pseudomonas aeruginosa relA* contributes to virulence in *Drosophila melanogaster*. *Infect. Immun.* **72**:5638–5645.
13. Eschbach, M., K. Schreiber, K. Trunk, J. Buer, D. Jahn, and M. Schobert. 2004. Long-term anaerobic survival of the opportunistic pathogen *Pseudomonas aeruginosa* via pyruvate fermentation. *J. Bacteriol.* **186**:4596–4604.
14. Farewell, A., A. A. Diez, C. C. DiRusso, and T. Nyström. 1996. Role of the *Escherichia coli* FadR regulator in stasis survival and growth phase-dependent expression of the *uspA*, *fad*, and *fab* genes. *J. Bacteriol.* **178**:6443–6450.
15. Freestone, P., T. Nystrom, M. Trinei, and V. Norris. 1997. The universal stress protein, UspA, of *Escherichia coli* is phosphorylated in response to stasis. *J. Mol. Biol.* **274**:318–324.
16. Galimand, M., M. Gamper, A. Zimmermann, and D. Haas. 1991. Positive FNR-like control of anaerobic arginine degradation and nitrate respiration in *Pseudomonas aeruginosa*. *J. Bacteriol.* **173**:1598–1606.
17. Gamper, M., A. Zimmermann, and D. Haas. 1991. Anaerobic regulation of transcription initiation in the *arcDABC* operon of *Pseudomonas aeruginosa*. *J. Bacteriol.* **173**:4742–4750.
18. Gustavsson, N., A. Diez, and T. Nystrom. 2002. The universal stress protein paralogs of *Escherichia coli* are co-ordinately regulated and co-operate in the defence against DNA damage. *Mol. Microbiol.* **43**:107–117.
19. Hanna, S. L., N. E. Sherman, M. T. Kinter, and J. B. Goldberg. 2000. Comparison of proteins expressed by *Pseudomonas aeruginosa* strains representing initial and chronic isolates from a cystic fibrosis patient: an analysis by 2-D gel electrophoresis and capillary column liquid chromatography-tandem mass spectrometry. *Microbiology* **146**:2495–2508.
20. Hassett, D. J., J. Cuppoletti, B. Trapnell, S. V. Lyman, J. J. Rowe, S. Sun Yoon, G. M. Hilliard, K. Parvatiyar, M. C. Kamani, D. J. Wozniak, S. H. Hwang, T. R. McDermott, and U. A. Ochsner. 2002. Anaerobic metabolism and quorum sensing by *Pseudomonas aeruginosa* biofilms in chronically infected cystic fibrosis airways: rethinking antibiotic treatment strategies and drug targets. *Adv. Drug Deliv. Rev.* **54**:1425–1443.
21. Heydorn, A., B. K. Ersboll, M. Hentzer, M. R. Parsek, M. Givskov, and S. Molin. 2000. Experimental reproducibility in flow-chamber biofilms. *Microbiology* **146**:2409–2415.
22. Heydorn, A., A. T. Nielsen, M. Hentzer, C. Sternberg, M. Givskov, B. K. Ersboll, and S. Molin. 2000. Quantification of biofilm structures by the novel computer program COMSTAT. *Microbiology* **146**:2395–2407.
23. Hoang, T. T., R. R. Karkhoff-Schweizer, A. J. Kutchma, and H. P. Schweizer. 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**:77–86.
24. Hoang, T. T., A. J. Kutchma, A. Becher, and H. P. Schweizer. 2000. Integration-proficient plasmids for *Pseudomonas aeruginosa*: site-specific integration and use for engineering of reporter and expression strains. *Plasmid* **43**:59–72.
25. Hoiby, N., H. Krogh Johansen, C. Moser, Z. Song, O. Ciofu, and A. Kharazmi. 2001. *Pseudomonas aeruginosa* and the *in vitro* and *in vivo* biofilm mode of growth. *Microbes Infect.* **3**:23–35.
26. Kvint, K., C. Husbond, A. Farewell, O. Nybroe, and T. Nystrom. 2000. Emergency derepression: stringency allows RNA polymerase to override negative control by an active repressor. *Mol. Microbiol.* **35**:435–443.
27. Kvint, K., L. Nachin, A. Diez, and T. Nystrom. 2003. The bacterial universal stress protein: function and regulation. *Curr. Opin. Microbiol.* **6**:140–145.
28. Lam, J., R. Chan, K. Lam, and J. W. Costerton. 1980. Production of mucoid microcolonies by *Pseudomonas aeruginosa* within infected lungs in cystic fibrosis. *Infect. Immun.* **28**:546–556.
29. Miller, J. M. 1992. A short course in bacterial genetics. A laboratory manual and handbook for *Escherichia coli* and related bacteria. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
30. Munch, R., K. Hiller, H. Barg, D. Heldt, S. Linz, E. Wingender, and D. Jahn. 2003. PRODORIC: prokaryotic database of gene regulation. *Nucleic Acids Res.* **31**:266–269.
31. Nystrom, T., C. Larsson, and L. Gustafsson. 1996. Bacterial defense against aging: role of the *Escherichia coli* ArcA regulator in gene expression, readjusted energy flux and survival during stasis. *EMBO J.* **15**:3219–3228.
32. O'Toole, R., M. J. Smeulders, M. C. Blokpoel, E. J. Kay, K. Lougheed, and H. D. Williams. 2003. A two-component regulator of universal stress protein expression and adaptation to oxygen starvation in *Mycobacterium smegmatis*. *J. Bacteriol.* **185**:1543–1554.

33. O'Toole, R., and H. D. Williams. 2003. Universal stress proteins and *Mycobacterium tuberculosis*. Res. Microbiol. **154**:387–392.
34. Perkins, D. N., D. J. Pappin, D. M. Creasy, and J. S. Cottrell. 1999. Probability-based protein identification by searching sequence databases using mass spectrometry data. Electrophoresis **20**:3551–3567.
35. Rabilloud, T., J. M. Strub, S. Luche, A. van Dorselaer, and J. Lunardi. 2001. A comparison between Sypro Ruby and ruthenium II tris (bathophenanthroline disulfonate) as fluorescent stains for protein detection in gels. Proteomics **1**:699–704.
36. Reimmann, C., M. Beyeler, A. Latifi, H. Winteler, M. Foglino, A. Lazdunski, and D. Haas. 1997. The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer N-butyryl-homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. Mol. Microbiol. **24**:309–319.
37. Rompf, A., C. Hungerer, T. Hoffmann, M. Lindenmeyer, U. Romling, U. Gross, M. O. Doss, H. Arai, Y. Igarashi, and D. Jahn. 1998. Regulation of *Pseudomonas aeruginosa* hemF and hemN by the dual action of the redox response regulators Anr and Dnr. Mol. Microbiol. **29**:985–997.
38. Sauer, K., A. K. Camper, G. D. Ehrlich, J. W. Costerton, and D. G. Davies. 2002. *Pseudomonas aeruginosa* displays multiple phenotypes during development as a biofilm. J. Bacteriol. **184**:1140–1154.
39. Sowers, R. G. 1991. Identification and molecular characterization of a transcriptional regulator from *Pseudomonas aeruginosa* PAO1 exhibiting structural and functional similarity to the FNR protein of *Escherichia coli*. Mol. Microbiol. **5**:1469–1481.
40. Shevchenko, A., I. Chernushevich, M. Wilm, and M. Mann. 2000. De novo peptide sequencing by nanoelectrospray tandem mass spectrometry using triple quadrupole and quadrupole/time-of-flight instruments. Methods Mol. Biol. **146**:1–16.
41. Stoodley, P., K. Sauer, D. G. Davies, and J. W. Costerton. 2002. Biofilms as complex differentiated communities. Annu. Rev. Microbiol. **56**:187–209.
42. van Delden, C., R. Comte, and M. Bally. 2001. Stringent response activates quorum sensing and modulates cell density-dependent gene expression in *Pseudomonas aeruginosa*. J. Bacteriol. **183**:5376–5384.
43. Van Delden, C., and B. H. Iglewski. 1998. Cell-to-cell signaling and *Pseudomonas aeruginosa* infections. Emerg. Infect. Dis. **4**:551–560.
44. Vander Wauven, C., A. Piérard, M. Kley-Raymann, and D. Haas. 1984. *Pseudomonas aeruginosa* mutants affected in anaerobic growth on arginine: evidence for a four-gene cluster encoding the arginine deiminase pathway. J. Bacteriol. **160**:928–934.
45. Vijgenboom, E., J. E. Busch, and G. W. Canters. 1997. *In vivo* studies disprove an obligatory role of azurin in denitrification in *Pseudomonas aeruginosa* and show that *azu* expression is under control of *rpoS* and ANR. Microbiology **143**:2853–2863.
46. Whiteley, M., M. R. Parsek, and E. P. Greenberg. 2000. Regulation of quorum sensing by RpoS in *Pseudomonas aeruginosa*. J. Bacteriol. **182**:4356–4360.
47. Worlitzsch, D., R. Tarran, M. Ulrich, U. Schwab, A. Cekici, K. C. Meyer, P. Birrer, G. Bellon, J. Berger, T. Weiss, K. Botzenhart, J. R. Yankaskas, S. Randell, R. C. Boucher, and G. Doring. 2002. Effects of reduced mucus oxygen concentration in airway *Pseudomonas* infections of cystic fibrosis patients. J. Clin. Investig. **109**:317–325.
48. Xu, K. D., P. S. Stewart, F. Xia, C.-T. Huang, and G. A. McFeters. 1998. Spatial physiological heterogeneity in *Pseudomonas aeruginosa* biofilm is determined by oxygen availability. Appl. Environ. Microbiol. **64**:4035–4039.
49. Ye, R. W., D. Haas, J.-O. Ka, V. Krishnapillai, A. Zimmermann, C. Baird, and J. M. Tiedje. 1995. Anaerobic activation of the entire denitrification pathway in *Pseudomonas aeruginosa* requires Anr, an analog of Fnr. J. Bacteriol. **177**:3606–3609.
50. Yoon, S. S., R. F. Hennigan, G. M. Hilliard, U. A. Ochsner, K. Parvatiyar, M. C. Kamani, H. L. Allen, T. R. DeKievit, P. R. Gardner, U. Schwab, J. J. Rowe, B. H. Iglewski, T. R. McDermott, R. P. Mason, D. J. Wozniak, R. E. Hancock, M. R. Parsek, T. L. Noah, R. C. Boucher, and D. J. Hassett. 2002. *Pseudomonas aeruginosa* anaerobic respiration in biofilms: relationships to cystic fibrosis pathogenesis. Dev. Cell **3**:593–603.
51. Zumft, W. G. 1997. Cell biology and molecular basis of denitrification. Microbiol. Mol. Biol. Rev. **61**:533–616.