

Positive Selection of the Hrp Pilin HrpE of the Plant Pathogen *Xanthomonas*

Ernst Weber and Ralf Koebnik*

Institute of Genetics, Martin-Luther-University, 06099 Halle, Germany

Received 27 September 2005/Accepted 21 November 2005

The plant-pathogenic bacterium *Xanthomonas campestris* pv. *vesicatoria* possesses a type III secretion (TTS) system which is encoded by the 23-kb *hrp* (hypersensitive response and pathogenicity) gene cluster. The TTS system is necessary for pathogenicity in susceptible hosts and induction of the hypersensitive response in resistant plants. At the cell surface, the TTS system is associated with an extracellular filamentous structure, the Hrp pilus, which serves as a conduit for the transfer of bacterial proteins into the plant cell cytosol. The major pilus component, the HrpE pilin, is unique to xanthomonads. Previous work showed that HrpE contains two regions: a hypervariable surface-exposed domain, including the N-terminal secretion signal, and a C-terminal polymerization domain. In this study, the evolutionary rate of the *hrpE* gene was analyzed. Twenty-one alleles were cloned, sequenced, and compared with five known *hrpE* alleles. The ratio of synonymous (K_s) and nonsynonymous (K_a) substitution rates shows that parts of the HrpE N terminus are subjected to positive selection and the C terminus is subjected to purifying selection. The trade-off between positive and purifying selection at the very-N terminus allowed us to ascertain the amphipathic α -helical nature of the TTS signal. This is the first report of a surface structure from a plant-pathogenic bacterium that evolved under the constraint of positive selection and hints to the evolutionary adaptation of this extracellular appendage to avoid recognition by the plant defense surveillance system.

The type III secretion (TTS) system is a hallmark of many gram-negative bacterial pathogens. This specialized secretion system is responsible for the transport of proteins across the two bacterial membranes, across the host cell plasma membrane, and in some cases across the plant cell wall into the host cell interior. A defect in this system leads to a complete loss of bacterial pathogenicity, as demonstrated for the animal pathogens *Yersinia* spp., *Shigella* spp., *Salmonella* spp., and enteropathogenic and enterohemorrhagic *Escherichia coli* and for the plant pathogens *Ralstonia solanacearum*, *Pseudomonas syringae*, *Erwinia* spp., and *Xanthomonas* spp. (10). TTS systems are encoded by approximately 20 genes, 11 of which are conserved between different bacterial species, thus suggesting that they constitute the core of the secretion machinery. While the TTS apparatus is conserved, the secreted substrates differ considerably. The recognition of these proteins by the TTS system is still not well understood. The TTS signal is located in the first 15 to 20 amino acids of the protein and is not conserved on the amino acid level (17, 23). Additionally, the 5' region of the corresponding mRNA may contribute to recognition by the secretion machinery (1).

Our laboratory studies the plant pathogen *Xanthomonas campestris* pathovar *vesicatoria*, which is the causal agent of bacterial spot disease in pepper and tomato plants. The TTS system of *X. campestris* pv. *vesicatoria* is encoded by the 23-kb *hrp* (hypersensitive response and pathogenicity) gene cluster, which is essential to cause disease in susceptible plants as well as to induce the hypersensitive response in resistant plants (9). A key component of the *X. campestris* pv. *vesicatoria* pathoge-

nicity machinery is the Hrp pilus, which is thought to serve as a conduit for protein translocation between the bacterium and the plant host cell (30). The Hrp pilus is a filamentous structure that extends up to several micrometers from the bacterial cell surface and is mainly composed of a large number of identical pilin subunits, the 10-kDa HrpE protein, which is unique for the genus *Xanthomonas* (30). HrpE itself is a TTS substrate and harbors a secretion signal in the first 17 N-terminal amino acids (29). Sequence alignment and secondary structure predictions of five *hrpE* sequences showed that the very-C terminus is entirely conserved and predominantly α -helical, whereas the N terminus is hypervariable and mostly devoid of regular secondary structure. Reporter protein fusions and pentapeptide insertions demonstrated that HrpE contains two functional domains, a surface-exposed domain, including the TTS signal, and the polymerization domain (29). The sequence hypervariability of the surface-exposed domain prompted us to analyze the evolutionary rate of 26 *hrpE* sequences, representing *Xanthomonas* strains belonging to 14 different pathovars. Our results reveal different modes of selection acting on different portions of the protein. The N terminus of HrpE undergoes positive selection, whereas the C terminus is subjected to purifying selection. Since the positively selected parts are exposed at the surface of the assembled Hrp pilus, the plant pathogen *Xanthomonas* thus generates surface diversity, most probably to evade recognition by the plant perception and defense systems. Furthermore, we propose an amphipathic α -helix as the TTS signal of HrpE which is on one side highly conserved and polar and on the opposite side positively selected and hydrophobic.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The following pathovars/strains of *X. campestris* were used: alfalfae KS, amoraciae XA, begoniae 077-3382, citri 50E, dieffenbachiae 729, glycines 202, malvacearum G-34, phaeseoli 85-6, vesicatoria

* Corresponding author. Mailing address: Martin-Luther-Universität, Institut für Genetik, Weinbergweg 10, 06120 Halle (Saale), Germany. Phone: 49-345-5526293. Fax: 49-345-5527151. E-mail: koebnik@gmx.de.

56, vesicatoria 85-10, vignicola 81-30, and vitians 164 (4); amoraciae 3838, amoraciae 5 (48A), raphani 23, and raphani 25 (R. E. Stall, University of Florida, Gainesville); and amoraciae LMG535, amoraciae LMG7383, raphani LMG860, raphani LMG7505, and raphani LMG8134 (Belgium coordinated collections of microorganisms, Brussels, Belgium). *Escherichia coli* cells were cultivated at 37°C in LB medium, and *Xanthomonas* strains were grown at 30°C in NYG broth (6) or on NYG 1.5% agar. Antibiotics were added to the medium at the following final concentrations: ampicillin, 100 µg/ml; kanamycin, 25 µg/ml; rifampin, 100 µg/ml.

Sequence data. The coding regions of 21 *hrpE* alleles were amplified using PCR primers derived from conserved regions of *hrpD6* (*hrpE*up, 5'-CAACATCGTCGCGCACATCGC) and *hpaB* (*hrpE*down, 5'-TCGAATCGCGCGCTGCTCAT) located upstream and downstream of the *hrpE* coding sequence, respectively. For colony PCR, a single colony of each strain was used as the source of template DNA. PCR conditions were an initial 95°C step for 300 s, followed by 35 cycles of 95°C for 30 s, 58°C for 30 s, and 72°C for 40 s, and a final step at 72°C for 10 min. PCR products were cloned into pCR 2.1-TOPO and transformed into DH10b (Invitrogen, Carlsbad, CA). Five clones of each *hrpE* allele were pooled and sequenced using vector-specific primers (TOPO TA cloning kit; Invitrogen).

Additionally, the *hrpE* nucleotide sequences of *X. campestris* pv. vesicatoria strain 75-3 (GenBank accession number [AC] AF056246), *X. oryzae* pv. oryzae strain KACC 10331 (AC AE013598), *X. axonopodis* pv. glycines strain 8ra (AC AF499777), *X. axonopodis* pv. citri strain 306 (AC AE011665), and *X. campestris* pv. campestris strain ATCC 33913 (AC AE012221) were used for bioinformatic analyses.

Sequence analysis. The nucleotide sequences were translated and aligned using CLUSTAL X (26). The multiple alignment was then manually edited using the BIOEDIT program (11). The amino acid alignments were transposed back to the nucleotide sequence level to gain a codon-based alignment. Mean diversity estimates within sequences were performed with the MEGA 2 software, version 2.1 (15), using the modified Nei-Gojobori method with the Jukes-Cantor corrected distance and the complete-deletion-of-gaps option. Unweighted pair group method with averages (UPGMA) trees were constructed using PAUP 3.1.1 (25). Distances were determined by using the Kimura two-parameter model. Support for nodes was assessed with a bootstrap confidence level using 1,000 replicates.

Calculation of synonymous and nonsynonymous mean pairwise diversities. The method of Comeron, as implemented in the K-Estimator program, version 6.1 (5), was applied to test whether positive or purifying selection has taken place on *hrpE*. K-Estimator takes into account the divergence values, number of nucleotides or codons, the transition/transversion substitution ratio (α/β), the amino acid composition, and the G+C content at the third codon position. The confidence intervals for K_a and K_s were obtained by Monte Carlo simulations (5). Synonymous mean pairwise diversity per synonymous site (K_s) and nonsynonymous mean pairwise diversity per nonsynonymous site (K_a) were calculated with the complete-deletion-of-gaps option and a sliding window of 15 nucleotides with a step size of 9 nucleotides.

Nucleotide sequence accession numbers. The nucleotide sequences for the 21 *hrpE* alleles were submitted to GenBank (accession numbers DQ286679 to DQ286699).

RESULTS

Rate of synonymous and nonsynonymous substitutions (K_s and K_a). To analyze the general extent of sequence variation in the genus *Xanthomonas*, we analyzed five genes from five *Xanthomonas* strains (Table 1). Generally, a neutral rate test compares the number of synonymous (silent) substitutions per synonymous site (K_s) with the number of nonsynonymous (amino acid-changing) substitutions per nonsynonymous site (K_a) in protein coding sequences. Thus, the ratio K_a/K_s measures the difference between the two rates, in which K_s is often regarded as a value of the underlying mutation rate (19). If an amino acid change is neutral, it will be fixed at the same rate as a synonymous mutation, with K_a/K_s of 1. If the amino acid change is deleterious, purifying selection will reduce its fixation rate, and thus K_a/K_s is <1. However, only if the amino acid change offers a selective advantage will it be fixed at a higher rate than a synonymous mutation, with K_a/K_s of >1 (positive

TABLE 1. Sequence variations of five genes among five *Xanthomonas* strains^a

Gene	Size (bp)	Mean pairwise value (10^2)		K_a/K_s ratio
		K_s	K_a	
<i>hrpE</i>	279	49.4 ± 6.4	27.8 ± 4.3	0.560
<i>hrcS</i>	261	25.8 ± 4.8	0.80 ± 0.1	0.031
<i>hrcT</i>	828	22.2 ± 2.3	3.90 ± 0.6	0.170
<i>gapA</i> ^b	999	17.2 ± 1.8	0.70 ± 0.2	0.040
<i>mdh</i> ^b	987	18.8 ± 2.0	2.50 ± 0.5	0.130

^a *X. campestris* pv. vesicatoria strain 75-3, *X. oryzae* pv. oryzae strain KACC 10331, *X. axonopodis* pv. glycines strain 8ra, *X. axonopodis* pv. citri strain 306, and *X. campestris* pv. campestris strain ATCC 33913.

^b *X. axonopodis* pv. glycines strain 8ra was not included in the analysis.

selection) (31). Three of the analyzed genes (*hrcS*, *hrcT*, and *hrpE*) are located in the *hrp* gene cluster and encode crucial components of the TTS system. For reference, two housekeeping genes, malate dehydrogenase (*mdh*) and glyceraldehyde-3-phosphate dehydrogenase (*gapA*), were used. The most notable feature is an unusually high level of amino acid substitutions in the *hrpE* gene product, which is reflected by the K_a value of 0.278 (Table 1). In contrast, the K_a values for *hrcT*, *hrcS*, *mdh*, and *gapA* are uniformly low and fall within the narrow range of 0.007 to 0.039. In addition, the K_s value for *hrpE* is strongly elevated (0.494) in comparison with those of *hrcT*, *hrcS*, *mdh*, and *gapA* (0.172 to 0.258) (Table 1). The K_a/K_s ratio, which describes the evolutionary rate of a protein coding sequence, is much higher for *hrpE* than for the other four genes. Although the K_a value does not exceed the K_s value for the full-length coding sequence of *hrpE*, this high value indicates that parts of *hrpE* may be subjected to positive selection.

Comparison of *hrpE* sequences. For a more detailed analysis, the *hrpE* regions of 21 *Xanthomonas* strains belonging to 12 pathovars were cloned and sequenced. The phylogenetic tree of all 26 *hrpE* sequences was constructed using the UPGMA method (Fig. 1). Interestingly, the genealogy revealed two major branches (groups I and II).

Five of 13 *hrpE* sequences in group I and 3 of 13 *hrpE* sequences in group II were identical. These siblings were excluded from further analyses. Figure 2 shows an amino acid sequence alignment of the nonredundant HrpE proteins. The last 24 amino acid residues of HrpE are highly conserved in all xanthomonads, suggesting that this region is crucial for HrpE function. Another conserved sequence motif (PxxxSxxxRxxQxxD) is located at the very-N terminus, whereas the region between the two conserved parts is hypervariable. Intriguingly, the conserved N-terminal amino acid pattern which corresponds to the TTS signal is predicted to form an α -helix (30). A helical wheel plot revealed that the conserved polar residues are located on one side of the helix, whereas the opposite side consists of hydrophobic amino acids (Fig. 3). These findings strongly suggest that the HrpE secretion signal is formed by an amphipathic α -helix.

Whether or not recombination in the *hrpE* sequence has occurred was analyzed by assessing the degree of phylogenetic congruence. Based on the amino acid sequence alignment (Fig. 2), the *hrpE* alleles were split into two halves: a variable region consisting of codons 4 to 50 and the relatively conserved region




FIG. 1. UPGMA tree of *hrpE* sequences from 26 *Xanthomonas* strains. Numbers on branches indicate the number of times (percent) that the node was supported by 1,000 replicates of the bootstrap analysis (only values greater than 80% are shown).

encompassing codons 51 to 93. Phylogenetic trees were then estimated for each half. No significant differences in topologies of the trees were observed (data not shown), thus indicating that no recombination between *hrpE* alleles belonging to groups I and II had occurred.




FIG. 3. Type III secretion signal of the *X. campestris* pv. *vesicatoria* 85-10 HrpE in a helical wheel representation. Conserved amino acids and hydrophobic residues are shaded in gray and black, respectively. The N-terminal proline residue is oriented to the bottom.

Modes of selection operating on *hrpE*. Using the enlarged set of *hrpE* sequences, we investigated whether certain parts of HrpE show different modes of selection. A sliding window analysis of K_a and K_s values was performed. Since the first amino acid of HrpE from group I and the first four amino acids of HrpE from group II give no substantial information, they were not considered. Windows in which K_a exceeded K_s (one tailed Z-test, $P < 0.05$) (Fig. 4A) were found in the N-terminal and central regions of the protein. This finding shows that diversifying selection had occurred at these regions. In contrast, K_s exceeded K_a in windows 98 and 215 to 266 (one tailed Z-test, $P < 0.05$ [Fig. 4A]; the number indicates the central nucleotide of the window), indicating that these regions evolved under purifying selection.

The observation that the *hrpE* sequences belong to two clearly distinct phylogenetic groups prompted us to analyze whether different modes of selection operated after separation




FIG. 2. Nonredundant amino acid sequence alignment of HrpE. HrpE sequences of group I and II xanthomonads were aligned using CLUSTAL X (26). Residues which are identical in at least 90% of the sequences are shown in white on a black background. Proposed functional domains are indicated on top of the alignment (29).




FIG. 4. Mean pairwise diversity plots of K_s and K_a values. The plots were constructed for windows of 15 nucleotides sliding along the *hrpE* sequence alignment. The first codon of group I and the first four codons of group II sequences are not included. (A) Comparison of all *hrpE* sequences. (B) Group I sequence comparison. (C) Group II sequence comparison. Filled boxes indicate the windows (step size of 9 nucleotides) in which nonsynonymous differences significantly exceed synonymous differences (one-tailed Z test, $P < 0.05$). White boxes indicate positions in the sequence where the reverse condition applies ($K_a > K_s$; $P < 0.05$) or sequences which are identical. Solid line, mean K_a ; dashed line, mean K_s .

of groups I and II. Individual analyses of group I and II sequences showed that positive selection occurred at the very-N terminus in both groups (group I, 8 to 26; group II, 17 to 44), whereas the C terminus was subjected to purifying selection (group I, 224 to 266; group II, 242 to 266) (Fig. 4B and C). Intriguingly, the intermediate parts of *hrpE* evolved differently in groups I and II. Most notably, the region between windows 71 and 125 shows strong signatures of positive selection in group I, while in group II this region evolved mainly under purifying constraints.

The positively selected positions are surface exposed. Certain modes of selection act often highly localized within genes. For many plant resistance proteins, plant chitinases, and the outer membrane porin PorB of *Neisseria meningitidis*, positive selection has been detected primarily for solvent-exposed




FIG. 5. Positively selected parts of HrpE are surface exposed. Mean hydrophobicity plots of group I (dashed line) and group II (solid line) HrpE proteins were calculated over a sliding window of five amino acid residues, using the Kyte and Doolittle hydrophobicity scale (16). Values below 0.4 are considered hydrophilic. Positively selected parts in groups I and II are indicated by open and filled boxes, respectively.

amino acids (3, 7, 28). Since solvent-exposed parts of a protein are in general rather hydrophilic and core regions of proteins are more hydrophobic, we performed hydrophobicity plot analyses of groups I and II and compared the localizations of positively selected and hydrophilic regions. As shown in Fig. 5, the HrpE regions which are positively selected in group I or II correspond to hydrophilic regions. Of special interest is the region between amino acids 25 and 35, which is hydrophilic in group I and hydrophobic in group II. This different physicochemical character is perfectly reflected by different modes of selection, indicating that this region might be surface exposed and therefore positively selected in group I. Notably, for the C-terminal hydrophilic region an indication of positive selection is missing. This finding may be related to the proposed function of this domain in the polymerization process (29).

DISCUSSION

Many gram-negative plant pathogens possess a surface-exposed Hrp pilus that mediates the long-distance transport of TTS substrates across the plant cell wall. Although the major pilus subunits of different genera do not share sequence homology, they share several features with each other, such as small size (6 to 11 kDa), predominantly α -helical secondary structure, and sequence hypervariability (14). Interestingly, sequence hypervariability is not only observed between different species but also between pathovars of the same species (12). In this study, we analyzed the sequence variability of the *Xanthomonas* Hrp pilus subunit HrpE, showed that the observed sequence hypervariability is caused by positive selection, and speculated about the biological significance of hypervariability.

A sliding window analysis of 18 *hrpE* alleles revealed several regions in the N-terminal two-thirds in which K_a exceeds K_s , thus indicating that these regions are positively selected. In contrast, the far C terminus is almost identical or subjected to purifying selection in all *Xanthomonas* sequences. Positive selection has been reported as a beacon for proteins involved in genetic conflict and has been documented in genes that encode

surface proteins of pathogen (2, 28) and host defense systems, such as the human major histocompatibility complex (13), plant chitinases (3), and plant resistance genes (7). The correlation between positive selection and host-pathogen interactions is particularly strong. For example, a GenBank survey uncovered remarkably few sequences (0.45%) evolving under positive selection, but more than half of these sequences were involved in host-pathogen interactions (8). Only six examples of bacterial surface proteins which have undergone positive selection have been described, but all of them originated from animal pathogens (27). This is the first report of a surface protein from a plant pathogen that is subjected to positive selection. This points to an unexpected analogy between animal and plant pathogenic bacteria in their strategy to avoid recognition by the host.

Despite the lack of an adaptive immune system, plants possess recognition capacities for several microbial compounds often indispensable for the microbial lifestyle. This set of general elicitors are referred to as pathogen-associated molecular patterns (PAMPs) and include the eubacterial flagellin, lipopolysaccharide of gram-negative bacteria, fungal cell wall-derived glucans, chitins, mannans, and surface proteins of microbial pathogens (20). An Hrp pilus of 500 nm in length would consist of approximately 1,000 HrpE subunits. As an abundant key component of the TTS system which is crucial for pathogenicity, HrpE would be a prime candidate to act as a PAMP. However, there is no report of a plant species or cultivar which has evolved a mechanism to detect HrpE or Hrp pilins from other plant pathogens as a PAMP.

Diversification of the Hrp pilus surface by positive selection acting on surface-exposed parts of the protein would make perfect sense in order to conceal the attack by *Xanthomonas* spp. Such a surface-limited diversification is a well-known strategy of human pathogenic bacteria (2, 28). Indeed, in both phylogenetic groups of *Xanthomonas* a remarkable correlation between hydrophilic regions and positive selection was found. Notably, only the first four hydrophilic stretches of the HrpE sequence were positively selected, whereas the C-terminal hydrophilic stretch was not. These findings are in perfect agreement with the proposed domain structure of HrpE (29). In this previous work it was found that the N-terminal two-thirds of the protein are tolerant to pentapeptide insertions, whereas the C-terminal third is not. Moreover, we isolated four dominant-negative insertions in the C-terminal third, thus supporting our hypothesis that this domain is conserved due to its function in polymerization. In line with this model, the C-terminal hydrophilic stretch would be hidden within the assembled Hrp pilus, and varying this sequence would give no selective advantage. In conclusion, the N-terminal surface-exposed domain functions as a camouflage suit for the highly conserved assembly domain at the C terminus.

An alternative mechanism to achieve sequence variation is based on a high frequency of recombination, as has been reported for components of the outer membrane or cell surface appendages of mammalian pathogens, such as the outer membrane protein PorB and the pilus protein Pile of *N. meningitidis* and the flagellin FliC of pathogenic *E. coli* (2, 22, 28). However, we did not observe discrepancies in the phylogenies between the full-length sequence and the separately analyzed N-terminal and C-terminal halves, indicating that recombina-

tion is not responsible for the diversification of *hrpE*. This finding is supported by the absence of a second copy of *hrpE*-related sequences in the completely sequenced genomes of five *Xanthomonas* strains. Interestingly however, the plant pathogens *P. syringae* and *R. solanacearum* possess a second copy of their Hrp pilin genes. It has been speculated that these second copies might act as capping or scaffolding proteins (12). Alternatively, one might speculate about a possible role of these extra genes in surface variation.

The very-N terminus of HrpE was shown to serve as a TTS signal (29). Despite 15 years of intense research, and even with the three-dimensional structures of the N termini of TTS substrates at hand, the nature of the TTS signal is still enigmatic (24). The fact that the N termini of these substrates do not share any obvious consensus sequence at the primary sequence level makes it difficult to understand how TTS substrates are recognized and secreted even by heterologous TTS machineries (24). Comparisons of *Yersinia* TTS substrates indicated that a more general property serves as secretion signal. Yersinial secretion signals consist of an amphipathic series of randomly alternating polar and hydrophobic residues, and even synthetic amphipathic signals consisting of serine and isoleucine were able to mediate secretion of a reporter protein (18). Similar biophysical features were identified in TTS substrates from the plant pathogen *P. syringae* (21). However, amphipathic N-terminal sequences derived from cytosolic components of the yersinial Yop regulon did not promote secretion of the reporter protein YopE (17). These findings imply that, in addition to the amphipathic character, there must be a sequence dependence on N-terminal secretion signals that is currently not understood.

While most studies on TTS signals have compared the N termini of distantly related or unrelated TTS substrates or have introduced point mutations into the N terminus of TTS substrates (21, 24), we analyzed a set of naturally occurring HrpE variants. From previous work it is known that the HrpE TTS signal is located in the first 17 amino acids (29). Since all HrpE N termini were obtained from pathogens, it is reasonable to assume that they are all functional TTS signals. The trade-off between sequence diversification (to avoid host recognition) and functional conservation (to allow secretion) operating at the HrpE N terminus offered a unique advantage to identify residues and principles that are of crucial importance for function as a TTS signal. The α -helix that would be formed by the amphipathic N-terminal sequence displays a polar and a hydrophobic side. In addition to the conserved amphipathic character of the α -helix, the five amino acid residues at the polar side seem to be critical for function, as indicated by their strong conservation. Probably, these two features are crucial for function of the HrpE TTS signal. To what extent this rule may apply to other TTS substrates remains to be clarified in future studies.

ACKNOWLEDGMENTS

We thank Ulla Bonas for generous support of this study, Hannelore Espenhahn for excellent technical assistance, Jens Boch, Daniela Büttner, Sabine Kay, Diana Kühn, and Ute Wahrmund for critical reading of the manuscript, and Martin Beye and Martin Hasselmann for fruitful discussions.

E.W. was supported by a GradFG fellowship from Sachsen-Anhalt.

REFERENCES

1. **Anderson, D. M., and O. Schneewind.** 1997. A mRNA signal for the type III secretion of Yop proteins by *Yersinia enterocolitica*. *Science* **278**:1140–1143.
2. **Andrews, T. D., and T. Gojobori.** 2004. Strong positive selection and recombination drive the antigenic variation of the PilE protein of the human pathogen *Neisseria meningitidis*. *Genetics* **166**:25–32.
3. **Bishop, J. G., A. M. Dean, and T. Mitchell-Olds.** 2000. Rapid evolution in plant chitinases: molecular targets of selection in plant-pathogen coevolution. *Proc. Natl. Acad. Sci. USA* **97**:5322–5327.
4. **Bonas, U., R. Schulte, S. Fenselau, G. V. Minsavage, B. J. Staskawicz, and R. E. Stall.** 1991. Isolation of a gene cluster from *Xanthomonas campestris* pv. vesicatoria that determines pathogenicity and the hypersensitive response on pepper and tomato. *Mol. Plant-Microbe Interact.* **4**:81–88.
5. **Comeron, J. M.** 1999. K-Estimator: calculation of the number of nucleotide substitutions per site and the confidence intervals. *Bioinformatics* **15**:763–764.
6. **Daniels, M. J., C. E. Barber, P. C. Turner, M. K. Sawczyc, R. J. W. Byrde, and A. H. Fielding.** 1984. Cloning of genes involved in pathogenicity of *Xanthomonas campestris* pv. *campestris* using the broad host range cosmid pLAFR1. *EMBO J.* **3**:3323–3328.
7. **Ellis, J., P. Dodds, and T. Pryor.** 2000. The generation of plant disease resistance gene specificities. *Trends Plant Sci.* **5**:373–379.
8. **Endo, T., K. Ikeo, and T. Gojobori.** 1996. Large-scale search for genes on which positive selection may operate. *Mol. Biol. Evol.* **13**:685–690.
9. **Fenselau, S., I. Balbo, and U. Bonas.** 1992. Determinants of pathogenicity in *Xanthomonas campestris* pv. *vesicatoria* are related to proteins involved in secretion in bacterial pathogens of animals. *Mol. Plant-Microbe Interact.* **5**:390–396.
10. **Ghosh, P.** 2004. Process of protein transport by the type III secretion system. *Microbiol. Mol. Biol. Rev.* **68**:771–795.
11. **Hall, T. A.** 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **1999**:95–98.
12. **He, S. Y., K. Nomura, and T. S. Whittam.** 2004. Type III protein secretion mechanism in mammalian and plant pathogens. *Biochim. Biophys. Acta* **1694**:181–206.
13. **Hughes, A. L., and M. Nei.** 1988. Pattern of nucleotide substitution at major histocompatibility complex class I loci reveals overdominant selection. *Nature* **335**:167–170.
14. **Koebnik, R.** 2001. The role of bacterial pili in protein and DNA translocation. *Trends Microbiol.* **9**:586–590.
15. **Kumar, S., K. Tamura, I. B. Jakobsen, and M. Nei.** 2001. MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics* **17**:1244–1245.
16. **Kyte, J., and R. F. Doolittle.** 1982. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**:105–132.
17. **Lloyd, S. A., A. Forsberg, H. Wolf-Watz, and M. S. Francis.** 2001. Targeting exported substrates to the Yersinia TTSS: different functions for different signals? *Trends Microbiol.* **9**:367–371.
18. **Lloyd, S. A., M. Norman, R. Rosqvist, and H. Wolf-Watz.** 2001. Yersinia YopE is targeted for type III secretion by N-terminal, not mRNA, signals. *Mol. Microbiol.* **39**:520–531.
19. **Miyata, T., T. Yasunaga, and T. Nishida.** 1980. Nucleotide sequence divergence and functional constraint in mRNA evolution. *Proc. Natl. Acad. Sci. USA* **77**:7328–7332.
20. **Nürnberg, T., F. Brunner, B. Kemmerling, and L. Piater.** 2004. Innate immunity in plants and animals: striking similarities and obvious differences. *Immunol. Rev.* **198**:249–266.
21. **Petnicki-Ocwieja, T., D. J. Schneider, V. C. Tam, S. T. Chancey, L. Shan, Y. Jamir, L. M. Schechter, M. D. Janes, C. R. Buell, X. Tang, A. Collmer, and J. R. Alfano.** 2002. Genomewide identification of proteins secreted by the Hrp type III protein secretion system of *Pseudomonas syringae* pv. *tomato* DC3000. *Proc. Natl. Acad. Sci. USA* **99**:7652–7657.
22. **Reid, S. D., R. K. Selander, and T. S. Whittam.** 1999. Sequence diversity of flagellin (*flhC*) alleles in pathogenic *Escherichia coli*. *J. Bacteriol.* **181**:153–160.
23. **Schechter, L. M., K. A. Roberts, Y. Jamir, J. R. Alfano, and A. Collmer.** 2004. *Pseudomonas syringae* type III secretion system targeting signals and novel effectors studied with a Cya translocation reporter. *J. Bacteriol.* **186**:543–555.
24. **Sorg, J. A., N. C. Miller, and O. Schneewind.** 2005. Substrate recognition of type III secretion machines -testing the RNA signal hypothesis. *Cell. Microbiol.* **7**:1217–1225.
25. **Swofford, D. L.** 1993. PAUP: phylogeny analysis using parsimony, version 3.1.1. Illinois Natural History Survey, Champaign, Ill.
26. **Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins.** 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**:4876–4882.
27. **Tümmler, B., and P. Cornelis.** 2005. Pyoverdine receptor: a case of positive Darwinian selection in *Pseudomonas aeruginosa*. *J. Bacteriol.* **187**:3289–3292.
28. **Urwin, R., E. C. Holmes, A. J. Fox, J. P. Derrick, and M. C. Maiden.** 2002. Phylogenetic evidence for frequent positive selection and recombination in the meningococcal surface antigen PorB. *Mol. Biol. Evol.* **19**:1686–1694.
29. **Weber, E., and R. Koebnik.** 2005. Domain structure of HrpE, the Hrp pilus subunit of *Xanthomonas campestris* pv. *vesicatoria*. *J. Bacteriol.* **187**:6175–6186.
30. **Weber, E., T. Ojanen-Reuhs, E. Huguet, G. Hause, M. Romantschuk, T. K. Korhonen, U. Bonas, and R. Koebnik.** 2005. The type III-dependent Hrp Pilus is required for productive interaction of *Xanthomonas campestris* pv. *vesicatoria* with pepper host plants. *J. Bacteriol.* **187**:2458–2468.
31. **Yang, Z., and J. P. Bielawski.** 2000. Statistical methods for detecting molecular adaptation. *Trends Ecol. Evol.* **15**:496–503.