

σ^E Regulates and Is Regulated by a Small RNA in *Escherichia coli*^{∇†}

Karl M. Thompson,^{1‡} Virgil A. Rhodius,² and Susan Gottesman^{1*}

Laboratory of Molecular Biology, National Cancer Institute, National Institutes of Health, Bethesda, Maryland 20892,¹ and
Department of Microbiology and Immunology, University of California, San Francisco,
San Francisco, California 94143²

Received 4 January 2007/Accepted 26 March 2007

RybB is a small, Hfq-binding noncoding RNA originally identified in a screen of conserved intergenic regions in *Escherichia coli*. Fusions of the *rybB* promoter to *lacZ* were used to screen plasmid genomic libraries and genomic transposon mutants for regulators of *rybB* expression. A number of plasmids, including some carrying *rybB*, negatively regulated the fusion. An insertion in the *rep* helicase and one upstream of *dnaK* decreased expression of the fusion. Multicopy suppressors of these insertions led to identification of two plasmids that stimulated the fusion. One contained the gene for the response regulator OmpR; the second contained *mipA*, encoding a murein hydrolase. The involvement of MipA and OmpR in cell surface synthesis suggested that the *rybB* promoter might be dependent on σ^E . The sequence upstream of the +1 of *rybB* contains a consensus σ^E promoter. The activity of *rybB-lacZ* was increased in cells lacking the RseA anti-sigma factor and when σ^E was overproduced from a heterologous promoter. The activity of *rybB-lacZ* and the detection of RybB were totally abolished in an *rpoE*-null strain. In vitro, σ^E efficiently transcribes from this promoter. Both a *rybB* mutation and an *hfq* mutation significantly increased expression of both *rybB-lacZ* and *rpoE-lacZ* fusions, consistent with negative regulation of the σ^E response by RybB and other small RNAs. Based on the plasmid screens, NsrR, a repressor sensitive to nitric oxide, was also found to negatively regulate σ^E -dependent promoters in an RseA-independent fashion.

Prokaryotic small regulatory RNA molecules have long been known to regulate plasmid replication and phage development (60). Recent studies have found that, in addition, regulatory RNAs have important roles for bacterial cell growth and physiology. A number of genome-wide searches performed by various groups using computational and biochemical methods have uncovered close to 100 small RNAs (sRNAs) in *Escherichia coli* (6, 8, 27, 47, 57, 62). More than a dozen of these RNAs have been characterized thus far and have been found to affect various physiological pathways. Characterization of additional sRNAs continues to yield surprising insights into cell physiology.

In *Escherichia coli*, the largest class of small regulatory RNAs binds to the RNA chaperone Hfq and regulates the stability and translation of specific mRNAs. The Hfq-binding sRNAs act by pairing with regions of complementarity in their target mRNAs. The vast majority of the Hfq-binding sRNAs act negatively, to either inhibit translation or stimulate the degradation of the target mRNA, also resulting in degradation of the sRNA itself (26, 33, 56). In some cases, sRNAs of this class can act positively. For instance, DsrA and RprA stimulate translation of the stationary-phase/general stress sigma factor RpoS (sigma S or σ^S) (32, 51).

RybB, an 81-nucleotide sRNA, was identified in a computational genome-wide search for novel sRNAs, as well as in an

expression-based search (57, 62). RybB binds the RNA chaperone protein Hfq, suggesting that the RNA acts by pairing with mRNAs. Steady-state levels of RybB could be detected only in stationary phase in rich media (57, 62). However, the conditions leading to the synthesis of RybB and its effects on the cell remained largely uncharacterized until recently.

In characterizing the regulation of RybB synthesis, we found that RpoE (sigma E or σ^E) was absolutely necessary for its synthesis and was a direct regulator of its transcription. σ^E is a member of the extracytoplasmic family of alternative sigma factors (31). σ^E guides the core RNA polymerase in binding to the promoter region and initiating transcription of specific genes in response to extracytoplasmic stress (excess heat shock or misfolded outer membrane proteins [OMPs]) (36, 52). RybB also represses σ^E synthesis, creating an autoregulatory loop.

(A portion of this work was used in partial fulfillment of the Ph.D. degree to K.M.T. from Howard University [Department of Microbiology].)

MATERIALS AND METHODS

Strains, plasmids, and media. The strains used are listed in Table S1 in the supplemental material. Most are derivatives of *E. coli* K-12 MG1655. Strains were grown in Luria-Bertani (LB) medium (KD Medical, Columbia, MD) in an Innova 3100 water bath shaker (New Brunswick Scientific, Edison, NJ) at approximately 200 rpm at 37°C unless otherwise indicated. Unless otherwise indicated, marked mutations were moved into the desired strain background using bacteriophage P1 transduction (49). Transductants were selected on LB plates supplemented with 25 μ g/ml kanamycin, 10 μ g/ml chloramphenicol, or 25 μ g/ml tetracycline (KD Medical, Columbia, MD), depending on the marker used. Tetrazolium (2,3,5-triphenyltetrazolium chloride [TTC]) (Sigma)-lactose (TTC-Lac) plates were prepared as previously described (38). Insertion mutants were screened on MacConkey-lactose plates supplemented with 25 μ g/ml kanamycin (KD Medical) or TTC-Lac plates supplemented with 25 μ g/ml kanamycin (TTC-Lac-Kan). All plasmids used in this study are listed in Table S2 in the supplemental material. Plasmids were generally introduced into strains by TSS (trans-

* Corresponding author. Mailing address: Bldg. 37, Rm. 5132, National Cancer Institute, Bethesda, MD 20892. Phone: (301) 496-3524. Fax: (301) 496-3875. E-mail: susang@helix.nih.gov.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

‡ Present address: Dept. of Microbiology and Immunology, Virginia Commonwealth University, Richmond, VA 23298-0678.

[∇] Published ahead of print on 6 April 2007.

formation and storage solution) transformation (10). Transformants were selected on LB agar plates supplemented with 100 µg/ml ampicillin (KD Medical). Genomic DNA transformants were screened on MacConkey-lactose plates supplemented with 50 µg/ml of ampicillin (Mac-Lac-Amp) (KD Medical).

Construction of *rybB-lacZ* transcriptional fusions. Single-copy chromosomal *rybB-lacZ* transcriptional fusions used in this study were first constructed in plasmid pRS1553 (<http://www.mimg.ucla.edu/bobs/vectors/Alpha-lac/alphaLac.htm>), a derivative of the vectors described in reference 50. To construct the multicopy *rybB-lacZ* long transcriptional fusion, the region from -77 to +22 of *rybB* was PCR amplified from MG1655 genomic DNA using primers KT12 and KT14 (see Table S3 in the supplemental material), digested with EcoRI and BamHI, and ligated to a pRS1553 EcoRI/BamHI digest to yield pKMT30. Similarly, in the *rybB-lacZ* short transcriptional fusion, the region from -36 to +22 of *rybB* was PCR amplified using primers KT13 and KT14 (Table S3) and ligated to pRS1553 to yield pKMT31. Strains containing these plasmids were then crossed with λRS468; Lac⁺ recombinants yielded transducing phages λKMT30 and λKMT31. λKMT30 and λKMT31 were used to lysogenize DJ480, yielding strains KMT12000 and KMT11000, respectively.

Construction of an *rpoE-lacZ* translational fusion. A region of *rpoE* from 160 nucleotides (nt) upstream to 270 nt downstream of the ATG was amplified from MG1655 genomic DNA using primers KT132 and KT143 (see Table S3 in the supplemental material), digested with EcoRI and BamHI, and ligated to EcoRI/BamHI-treated pRS1551 to yield pKMT35. Plasmid pKMT35 was then crossed with λRS468 (50) to yield λKMT35. Strain NM6010 was lysogenized with λKMT35, resulting in strain KMT14000.

Construction of pBAD-*rybB* and pBAD-*nsrR* plasmids. To clone the *rybB* gene under the control of an arabinose-inducible promoter, plasmid pNM12, a derivative of pBAD24 used for expression of sRNAs, was used as the vector (32). Oligonucleotides KT119 and KT120 (see Table S3 in the supplemental material) were annealed, and the resulting product was ligated to an MscI/PstI digest of pNM12, resulting in pKMT5. To clone *nsrR* under the control of an arabinose-inducible promoter, *nsrR* was amplified from MG1655 genomic DNA using primers KT191 and KT192, digested with EcoRI and PstI, and ligated to pBAD24, resulting in pKMT8. The constructs were confirmed by DNA sequencing.

Transposon mutant screen. For screen II, a pool of random insertion mutant derivatives of DJ480 was created by infecting DJ480 with ANK1316 (29) and selecting kanamycin-resistant colonies; a P1 lysate made on this pool was used to deliver the random insertions into KMT12000 by transduction. A total of 4,200 kanamycin-resistant transductants were screened on TTC-Lac-Kan; 22 Lac⁻ colonies were picked and assayed for immunity to phage *limm21*. Only two of the Lac⁻ transductants were still immune to the phage and thus retained the chromosomal *rybB-lacZ* fusion. The two insertions, *rb2::kan* and *rb4::kan*, were retransduced into KMT12000 to confirm the linkage of the *kan* insertion to the phenotype.

Insertions were mapped by an inverse PCR method as previously described (9, 43). The first PCR uses a random primer (RandomPrim1) and a transposon-specific primer (TnKm1) (see Table S3 in the supplemental material). In the second PCR, a nested set of primers (Revprim2 and TnKm2) (Table S3) was used. TnKm2 was used to sequence the PCR product to obtain the location of the *kan* gene. The *rb2::kan* insertion mapped to the middle of the *rep* gene, at nucleotide 3958864, and was renamed *rep::kan*; the strain carrying this insertion is KMT12002. The *rb4::kan* insertion mapped to an intergenic region directly upstream of the *dnaK* gene, at nucleotide 12109, and was renamed *zaa-1000::kan*; the strain carrying this insertion is KMT12003.

Genomic DNA library screens. For screen I, a genomic DNA library was introduced into KMT12000 (long *rybB-lacZ* fusion) and KMT11000 (short *rybB-lacZ* fusion) by electroporation (54). Following electroporation, 1 ml of LB medium was added to the electroporation mixtures, the culture was incubated at 37°C for 1 h, and 50-µl aliquots were spread on Mac-Lac-Amp. For KMT12000, approximately 10,500 colonies were screened; for KMT11000, approximately 3,500 colonies were screened.

In addition to screening KMT12000 and KMT11000, the genomic library was also introduced into KMT12002 and KMT12003 (*rep::kan* and *zaa-1000::kan* mutant derivatives of KMT12000, Lac⁻ phenotype, from screen II) to screen for multicopy suppressors (screen III). Transformants were screened for a Lac⁺ phenotype on Mac-Lac-Amp.

Construction of chromosomal *rybB* and *rseA* mutants. Kanamycin-resistant deletion insertions in *rseA* and *rybB* were constructed using the mini-λ-based λ Red recombineering system (15). A *kan* cassette with 50 bp of homology to regions flanking *rybB* was amplified using primers KT18 and KT19 (Table S3). A *kan* cassette with 50 bp of homology to *rseA* was amplified using oligonucleotides KT76 and KT77 (see Table S3 in the supplemental material). A culture of DJ480

containing mini-λ::*tet* (NM300) was grown at 32°C to mid-log phase and switched to 42°C to induce expression of the λ Red proteins. Aliquots of the uninduced and induced cultures were made electrocompetent, and approximately 100 ng of the respective PCR products was electroporated into a 50-µl aliquot of the electrocompetent cells. Recombinants were selected on LB agar plates containing 50 µg/ml of kanamycin.

RNA isolation and Northern blots. Total RNA was extracted in mid-exponential or stationary phase using the hot-phenol method (2). RNA samples were run on 8% acrylamide gels, and nonisotopic Northern blotting was performed as previously described (34). The 5'-biotinylated oligonucleotide KT98 was used as a probe for the detection of RybB RNA (see Table S3 in the supplemental material).

In vitro transcription assays. Single-round in vitro transcription experiments were performed to test the *rybB* promoter. The *rybB* promoter DNA template was prepared by PCR from MG1655 genomic DNA and contained sequences from -126 to +112 with respect to the transcription start and also the *rybB* terminator, using oligonucleotide primers *rybB* prom and *rybB* term (see Table S3 in the supplemental material). The control *PdegP* DNA template was prepared from the vector pUA66, carrying the native *PdegP* σ^E promoter sequences from -65 to +20 (46). PCR was used to amplify a *degP* promoter fragment from -200 to +175 with the primers pMS201-FW200 and pMS201-RV2 terminator (Table S3). Note that the pMS201-RV2 terminator primer contains the *rpoC* terminator sequence (35) to provide efficient transcription termination at the underlined C in Table S3.

RNA polymerase core enzyme was purified as described in reference 65, and His₆-tagged σ^E was purified using a QIAGEN Ni²⁺ affinity column per the manufacturer's instructions. The transcription assays were performed as described in reference 45 with the following modifications. Binding reaction mixtures (6 µl) contained 5 nM template DNA, 250 nM core RNA polymerase, 1,250 nM σ^E, and 1× binding buffer (5% glycerol, 20 mM Tris [pH 8.0], 300 mM potassium acetate, 5 mM magnesium acetate, 0.1 mM EDTA, 1 mM dithiothreitol, 50 µg/ml bovine serum albumin, 0.05% Tween). Single-round transcriptions were initiated with 2 µl of a 2× XTP-heparin mix (to give a final concentration of 200 µM ATP/UTP/GTP, 10 µM CTP, 2.5 µCi [α-³²P]CTP [3,000 Ci/mmol], 100 µg/ml heparin in 1× binding buffer), mixtures were incubated for 5 min at 37°C, and then reactions were terminated with 4 µl stop solution (20 mM EDTA, 80% deionized formamide, 0.05% [wt/vol] bromophenol blue, 0.05% [wt/vol] xylene cyanol FF). The RNA transcripts were resolved by electrophoresis after heating at 90°C for 2 min and loading 3 µl on a 6% denaturing polyacrylamide sequencing gel together with DNA sequencing reaction products that functioned as size markers. Transcripts were visualized using a Molecular Dynamics Storm 560 PhosphorImager scanning system.

β-Galactosidase assays. β-Galactosidase assays were carried out in 96-well plates and read in a SpectraMax 250 (Molecular Devices, Sunnyvale, CA) as previously described (66). Briefly, 50-ml LB cultures (KD Medical, Columbia, MD) were grown at 32°C or 37°C. At various time points, 100-µl aliquots of the cell culture were added to a well containing permeabilization buffer. Specific activity units are 25-fold lower than Miller units.

RESULTS

Genetic screens for regulators of RybB synthesis. In the study that initially identified RybB, its expression could be detected only in stationary phase (62). This pattern of expression was later confirmed, and the start site of *rybB* transcription was determined by 5' rapid amplification of cDNA ends (57), but the regulators of RybB expression were not determined.

Two *rybB-lacZ* transcriptional fusion strains were initially used to screen for RybB regulators. One fusion, the *rybB-lacZ* long fusion, contained -79 to +22 of *rybB*, with respect to the mapped transcription start site; the other, the *rybB-lacZ* short fusion, contained -36 to +22 of *rybB*. The *rybB-lacZ* long-fusion strain was Lac⁺ on Mac-Lac plates; the *rybB-lacZ* short-fusion strain was less strongly Lac⁺. The activity of both fusions increased upon entry of cells into stationary phase, consistent with previous reports where RybB RNA was detected only in stationary phase (57, 62). Expression of the *rybB-lacZ* fusions was not altered in an *rpoS*-null mutant, suggesting that station-

TABLE 1. Screen I: genomic library clones that modulate *rybB-lacZ* fusion expression

Library screen host ^a	Plasmid isolate (fragment boundaries)	Lac phenotype	Gene		Comments
			Name ^b	Description ^c	
<i>rybB-lacZ</i> (long)	pK-09 (886844–890902); pK-235 (884802–887483)	Lac ⁻	>'ybjK	Putative DeoR-type transcriptional regulator	RybB negative regulation of σ^E ; see text
			<'rybB	sRNA	
	pK-215 (4402773–4406851); pK-221 (4402773–4404689) ^a	Lac ⁻	>'purA	Adenylosuccinate synthetase	NsrR effect; this work
			>nsrR	Transcriptional repressor	
	pK-222 (2869026–2871368)	Lac ⁻	>ybjM	Unknown function	Cell division and cell surface effects?
			<grxA'	Glutaredoxin I	
			<truD	tRNA pseudouridine 13 synthase	
<ispF			2-C-Methyl-D-erythritol-2,4-cyclodiphosphate synthetase, isoprenoid biosynthesis		
pK-121 (3025059–3026683)	Lac ⁻	<ispD	4-Diphosphocytidyl-2C-methyl-D-erythritol synthetase	Unknown mechanism	
		<ftsB	Essential cell division protein		
		<ygbE'	Putative cytochrome oxidase subunit		
pK-126 (835278–777897)	Lac ⁻	>'guaD	Guanine deaminase	Unknown mechanism	
		>ygfQ	Unknown function (transporter?)		
pK-220 (3242589–3244529)	Lac ⁻	<ygfS'	Unknown function (oxidoreductase?)	Unknown mechanism	
		>'ybiB	Unknown function (transferase/phosphorylase?)		
		>ybiC	Unknown (dehydrogenase?)		
		<ybiJ	Unknown function		
		<ybiI	Unknown function		
pK-212 (982071–986106)	Lac ⁻	<ybiX'	Unknown function	σ^E regulated (45, 46)	
		>'uxaA	Altronate dehydratase		
		>exuT	Hexuronate transporter		
<i>rybB-lacZ</i> (short)	pK-304 (4102714–4104832)	Lac ⁻	>exuR'	Negative transcriptional regulator of the Exu operon	Regulator of periplasmic functions, may down-regulate σ^E stress
			<'cpxA	Membrane sensor kinase in CpxAR two-component signal transduction system	
			<cpxR	Response regulator in CpxAR two-component signal transduction system	
pK-407 (3341506–3345714)	Lac ⁻	>cpxP	Periplasmic protein activated by CpxAR, induced in alkaline pH, suppresses toxic envelope protein effects	σ^E -regulated operon σ^E competition? (45, 46)	
		>'yhbN	ABC transporter subunit		
		>yhbG	ABC transporter subunit		
		>rpoN	σ^{54} , nitrogen starvation sigma factor		
		>yhbH	Probable σ^{54} modulation protein		
>ptsN	Pts, regulates N metabolism				
>yhbJ'	ATPase				

Continued on following page

TABLE 1—Continued

Library screen host ^a	Plasmid isolate (fragment boundaries)	Lac phenotype	Gene		Comments
			Name ^b	Description ^c	
	pK-402 (1646935–1649118)	Lac ⁻	' <i>rzpQ</i>	Unknown function (cryptic prophage)	Cell division defect?
			> <i>dicF</i>	sRNA, inhibits <i>ftsZ</i>	
			> <i>dicB</i>	Control of cell division, activates MinC	
			> <i>ydfD</i>	Unknown function (cryptic prophage)	
			> <i>ydfE</i>	Unknown function (cryptic prophage)	
			> <i>insD'</i>	Cryptic prophage, transposase?	
<i>rybB-lacZ</i> (long)	pK-601 (2037829–2040601)	Lac ⁺⁺	> <i>yedY</i>	Unknown function	Unknown mechanism; increase membrane stress?
			> <i>yedZ</i>	Unknown function (inner membrane protein)	
			> <i>yodA</i>	Cadmium-induced metal binding protein; SoxS, Fur regulated	
			> <i>yodB</i>	Unknown function (cytochrome?)	

^a *rybB-lacZ* long fusion, KMT12000; *rybB-lacZ* short fusion, KMT11000.

^b >, clockwise gene; <, counterclockwise gene; ', incomplete gene.

^c Gene descriptions are from EcoCyc (27, 28).

ary-phase induction of *rybB* expression is RpoS independent (data not shown). The short fusion was used only in the first of the screens described below; tests with this fusion suggest that it is qualitatively similar in its regulation to the long fusion but may be missing some promoter sequences necessary for optimal expression.

The initial screen for regulators of *rybB* (screen I) was of an *E. coli* genomic DNA plasmid library in the *rybB-lacZ* fusion strains (54). A total of 11,000 transformants of the *rybB-lacZ* long fusion and 3,000 transformants of the *rybB-lacZ* short fusion were screened for Lac⁻ or Lac⁺⁺ phenotypes on Mac-Lac-Amp or TTC-Lac-Amp agar plates; candidate plasmids were confirmed by isolation and retransformation. Twelve independent plasmids, representing 10 different genomic regions, that reduced fusion expression were identified (Lac⁻ phenotype) (Table 1). Two independent plasmids carried the *rybB* gene itself. These plasmids also contained the upstream flanking gene, *ybjK*, which is a predicted DeoR-type transcriptional regulator. However, a null mutation of *ybjK* in the chromosome did not affect the expression of *rybB-lacZ* activity, suggesting that YbjK is not required for expression of *rybB* (data not shown). One clone gave a Lac⁺⁺ phenotype for the *rybB-lacZ* long fusion; it did not carry any known transcriptional regulators (Table 1). No explanation for the phenotypes resulting from these plasmids was immediately obvious; we return to some of them below.

Because the *rybB-lacZ* long fusion resulted in strains that were relatively Lac⁺, it proved challenging to identify plasmids leading to higher levels of expression (positive regulators); only one was identified in the previous screen. In order to identify possible positive regulators of *rybB* synthesis, a second screen for random insertion mutations that modulated expression of the *rybB-lacZ* long fusion was carried out. Approximately 25,000 transposon insertions into the *rybB-lacZ* strain were screened for changes in the Lac phenotype (screen II). Only

two insertions that changed expression in a reproducible fashion were isolated; both reduced *rybB* expression. Unexpectedly, one was in *rep*, encoding an ATP-dependent DNA helicase; the second was upstream of *dnaK* (*zaa-1000::kan*). The strain carrying the *rep::kan* insertion grew slowly and was somewhat cold sensitive. The *zaa-1000::kan* insertion is presumed to be polar on *dnaK* expression. However, the effect of *zaa-1000::kan* mutation on DnaK levels was not determined.

While it seemed unlikely that either Rep or DnaK was a direct regulator of *rybB* transcription, the Lac⁻ phenotype of the strains carrying these insertions allowed an easier screen for multicopy plasmids that increased expression from the *rybB-lacZ* long fusion. Using strains carrying the *rybB-lacZ* long fusion containing either the *rep::kan* (KMT12002) or *zaa-1000::kan* (KMT12003) insertion, we once again introduced the plasmid genomic library and screened for suppression of the Lac⁻ phenotype on Mac-Lac-Amp plates (screen III). A total of 14,000 transformants were screened for each strain. Four plasmids that increased expression of the fusions were identified; one plasmid was found twice, once in each mutant strain (Table 2). None of the plasmids carried either the *rep* or *dnaK* genes. Of the plasmids identified in this screen, only two carried known regulators, *emrR* and *ompR*.

EmrR is a negative regulator of genes involved in exporting drugs from the cell (30, 63). While an *emrR* plasmid stimulated expression of both the long and short *rybB-lacZ* fusions modestly, a mutation in *emrR* also had a modest stimulatory effect, suggesting both positive and negative effects of EmrR (data not shown). Epistasis experiments (see below and Fig. 1B) suggested that the EmrR effect was likely to be indirect, and it was not further pursued.

OmpR is the response regulator component of the EnvZ-OmpR two-component signal transduction system. OmpR acts as a transcriptional regulator of OMPs, most notably OmpC and OmpF, in response to changes in osmolarity of the envi-

TABLE 2. Genomic library clones isolated as multicopy suppressors; increased expression of *rybB-lacZ* (screen III)

Plasmid isolate (fragment boundaries)	Multicopy suppression		Gene	
	Original mutation	Modulatory effect in <i>ompR</i> ⁻ ?	Name	Description ^a
pK4-14 (1862417–1865705)	<i>rrb2</i> (<i>rep::kan</i>) and <i>rrb4</i> (<i>zaa-1000::kan</i>)	Yes; stimulation	> <i>yeaD</i> < <i>yeaE</i> < <i>mipA</i> > <i>yeaG</i>	Unknown function, conserved Methylglyoxal reductase Scaffolding protein for murein polymerase and murein hydrolase Unknown function, conserved
pK4-29 (1403515–1405770)	<i>rrb4</i> (<i>zaa-1000::kan</i>)	No	> <i>abgR</i> < <i>iszA</i> > <i>ydaL</i> < <i>ydaM</i>	Putative LysR-type transcriptional regulators sRNA Unknown function Unknown function, conserved
pK4-55 (3531018–3535617)	<i>rrb4</i> (<i>zaa-1000::kan</i>)	Yes; stimulation (complementation)	> <i>pckA</i> < <i>envZ</i> < <i>ompR</i> > <i>greB</i> > <i>yhgF</i>	Phosphoenolpyruvate carboxykinase Inner membrane osmosensor protein, regulates activity of OmpR Transcriptional regulator involved in osmoregulation of OmpC and OmpF Transcriptional elongation factor Unknown function
pK2-8 (2805651–2809436)	<i>rrb2</i> (<i>rep::kan</i>)	No	> <i>proX</i> > <i>ygaY</i> > <i>ygaZ</i> > <i>ygaH</i> > <i>emrR</i>	High-affinity transport for glycine, betaine, and proline Uncharacterized member of the major facilitator superfamily of transporters Predicted transporter Predicted inner membrane protein Negatively regulates transcription of the <i>emrAB</i> operon encoding subunits of a multidrug efflux pump; acts here via OmpR

^a Description of genes from EcoCyc (27, 28) and results from this paper.

ronment (41, 64). OmpR is also a transcriptional activator of two sRNAs, OmrA and OmrB, involved in the posttranscriptional regulation of several OMPs (22). To assess the role of OmpR in the transcription of *rybB*, a Δ *ompR* derivative of the *rybB-lacZ* strain was examined. Consistent with multicopy OmpR increasing expression of RybB (Fig. 1B and Table 2), the specific activity of the long *rybB-lacZ* was reduced approximately fivefold in the Δ *ompR* mutant in stationary phase (Fig. 1A), suggesting that OmpR is necessary for the full activity of the *rybB-lacZ* fusion.

If OmpR is the sole direct regulator of *rybB* transcription, we would expect the *ompR* mutant to be epistatic to all of the stimulatory plasmids from screen III, except for those that carry *ompR* itself. The four plasmids listed in Table 2 were each transformed into the Δ *ompR* derivative of the long *rybB-lacZ* fusion, and the Lac phenotype was assessed (Fig. 1B; Table 2). As expected, the plasmid carrying the *ompR* gene produced a strong Lac⁺ phenotype (Fig. 1B) in the otherwise Lac⁻ Δ *ompR* background. Two other plasmids, including the one carrying *emrR*, failed to increase expression in the *ompR* mutant (Fig. 1B), suggesting that the stimulatory activity of these plasmids was upstream of OmpR and depends upon OmpR. However, the plasmid carrying the *mipA* (*yeaF*) and *yeaE* genes was able to stimulate the *rybB-lacZ* fusion in both wild-type and Δ *ompR* strains, in both exponential phase (Fig.

1C) and stationary phase (data not shown). Parallel results were found with the short fusion: the *mipA* plasmid and the *ompR* plasmid increased activity in an *ompR* mutant derivative (data not shown).

The finding that *mipA-yeaE* was epistatic to *ompR* suggested that OmpR was not the sole primary regulator of *rybB*. While the function of YeaE is unknown, MipA has been shown to interact with murein synthase and murein hydrolase in vitro (59). MipA is therefore thought to be a scaffolding protein for murein synthesis (59).

These results suggested that possibly neither OmpR nor MipA-YeaE was a direct regulator of *rybB*. If so, and if they act in a similar way, the stimulation of the *rybB* promoter upon overproduction of either of these proteins should reflect changes in the activity or synthesis of a transcription factor that acts directly at the *rybB* promoter. A clue to the identity of this factor came from the realization that Δ *ompR* strains have about a fivefold reduction in both *rybB* expression and the activity of σ^E (Fig. 1A) (36). Given the role of MipA in murein synthesis, it seemed possible that these plasmids both act to activate σ^E , the alternative sigma factor that is activated by periplasmic protein misfolding (4, 36, 37, 48), and that σ^E promotes transcription of *rybB*.

An examination of the promoter region of *rybB* suggested that this was likely to be the case (Fig. 2). The -35 and -10




FIG. 1. OmpR and MipA-YeaE regulation of *rybB*. (A) β -Galactosidase activity of wild-type (KMT12000) and $\Delta ompR$ (KMT12005) *rybB-lacZ* fusion strains. Cells were grown in LB medium at 37°C and assayed as described in Materials and Methods. Numbers on the y axis are machine units as described in Materials and Methods. (B) A $\Delta ompR$ *rybB-lacZ* fusion strain (KMT12005) was transformed with plasmid pHDB3 (vector) or library clone pK2-8, pK4-14, pK4-29, or pK4-55 (Table 2), plated on Mac-Lac-Amp plates, and grown overnight at 37°C. (C) Wild-type *rybB-lacZ* fusion strains carrying either vector control or pK4-14 and $\Delta ompR$ *rybB-lacZ* fusion strains carrying either vector control or pK4-14 were grown to an optical density at 600 nm of approximately 0.5 and assayed for β -galactosidase activity. The values are means for three independent experiments. (D) A *degP-lacZ* (PND254) transcriptional fusion strain carrying either pHDB3 (a vector control) or pK4-14 (*mipA-yeaE* plasmid) was grown to an optical density at 600 nm of approximately 2.5 to 3.0 and assayed for β -galactosidase activity as described above. Note that PND254 is mutant for *cpxA* (see Table S1 in the supplemental material), and therefore activation likely reflects only the σ^E -dependent promoter.

	-35	spacer	-10
<i>E. coli rybB</i>	ACCGC GAACTT TTCCGCAGGGCATCAGCT T TAATTAGTG		
<i>Salmonella rybB</i>	TAGGTC GAACTT TTTCGTAAAGCATCAGCTACCTATTG		
<i>Klebsiella rybB</i>	CAGGC GAACTT TTTCTCAGGGCATCAGCTAAGTAATG		
<i>Yersinia rybB</i>	TCCCC GAACTT TCTTATCAGGCGGGGCTGAATTAATG		
<i>Photobacterium rybB</i>	CACGA GAACTT TTTTCGCAACCCGAGCTGAATTAATG		
<i>Erwinia rybB</i>	AACTC GAACTT TTTCTTTGGTCCACAGCTGAATAAGTG		
<i>E. coli degP</i>	GGA ACTTCAGGCTATAAAACGAATCTGA		
<i>E. coli rpoH P3</i>	TGA ACTTGTGGATAAAATCACGGTCTGA		
<i>E. coli rpoE P2</i>	GGA ACTTTACAAAACGAGACACTCTAA		
<i>E. coli</i> σ^E consensus	GGA ACTT	16 nt	TCAA A

FIG. 2. Alignment of the *rybB* promoter. The *E. coli rybB* region from -1 to -39 was aligned with the corresponding sequences from *Salmonella*, *Klebsiella*, *Yersinia*, *Photobacterium*, and *Erwinia*. Below the *rybB* sequences are the promoter sequences for other well-known *E. coli* σ^E -dependent promoters (46). Below the alignment is the consensus sequence previously defined for σ^E -dependent promoters (46).

consensus sequences for *E. coli* σ^E -dependent promoters are GGAAGCTT and TCAA, respectively, separated by a 16-nt spacer region (46). The *rybB* upstream region contains highly conserved sequences characteristic of the σ^E -10 and -35 sequences beginning at -11 and -33 nucleotides upstream from the RybB +1, respectively (Fig. 2). The putative *rybB* -35 sequence is identical to the -35 sequence for other well-characterized strong σ^E -dependent promoters in *E. coli*, including the *degP*, *rpoHP3*, and *rpoEP2* promoters (Fig. 2). The putative *rybB* -10 sequence has a high degree of similarity to these same promoters, with only one nucleotide change.

Overexpression of OmpR, by increasing the expression of the major OMPs, would be expected to increase σ^E activity (3) and therefore *rybB* expression. Does MipA overexpression also




FIG. 3. σ^E regulation of *rybB*. (A) β -Galactosidase activity of wild-type (KMT12000) and *rseA* (KMT12041) *rybB-lacZ* strains was assayed in both exponential (optical density at 600 nm [OD₆₀₀], 0.5) and stationary (OD₆₀₀, approximately 2.5 to 3.0) phases as described above. (B) β -Galactosidase activity of the wild-type *rybB-lacZ* strain (KMT12000) containing the vector control pTrc99A (dashed lines) or pLC245 (solid lines) was measured in the presence (solid symbols) or absence (open symbols) of IPTG. Cultures were grown to an OD₆₀₀ of 0.3, IPTG was added to a final concentration of 1 mM, and aliquots of the culture were taken every 5 min for 30 min. (C) β -Galactosidase activity of wild-type (KMT12000) and $\Delta\rho E$ (KMT12047) *rybB-lacZ* strains were assayed in stationary-phase growth as described above. (D) β -Galactosidase activity of wild-type (KMT12000) and *ppGpp*⁰ (KMT12031) *rybB-lacZ* strains throughout growth.

increase σ^E activity? We tested this by determining whether multicopy *mipA-yeaE* induced transcription of a known σ^E -dependent promoter, *degP*. In the presence of the *mipA-yeaE* plasmid, there is a clear induction of a *degP-lacZ* transcriptional fusion, compared to the vector control (Fig. 1D), consistent with the idea that *rybB* transcription may be regulated by σ^E .

σ^E is necessary for transcriptional induction of *rybB*. The results described above were consistent with the idea that *rybB* expression was dependent upon σ^E . We further examined this possibility by testing expression of the *rybB-lacZ* fusion and RybB RNA levels in strains where σ^E activity or synthesis was either increased or abolished.

In the absence of RseA, the σ^E -specific anti-sigma factor, the activity of σ^E -dependent promoters is significantly increased (16); *rybB* should be constitutively activated if it is transcribed from a σ^E -dependent promoter. Indeed, in the absence of RseA, the specific activity of the *rybB-lacZ* fusion

was ~23-fold higher in exponential phase and 7-fold higher in stationary phase (Fig. 3A). The smaller effect in stationary phase is likely to be due to the fact that σ^E is already partially activated in stationary-phase cells (14). In addition, overexpression of σ^E from a plasmid carrying *rhoE* under the control of an IPTG (isopropyl- β -D-thiogalactopyranoside)-inducible promoter led to an immediate and significant increase in *rybB* expression, consistent with σ^E acting as a direct regulator of *rybB* transcription (Fig. 3B).

Conversely, eliminating σ^E prevented expression of *rybB*. Although *rhoE* is essential, it can be deleted in the presence of uncharacterized suppressor mutations (4), allowing us to compare the specific activity of the *rybB-lacZ* fusion and RybB RNA levels in the presence and absence of *rhoE*. On MacConkey-lactose plates the $\Delta\rho E$ *rybB-lacZ* fusion strain was totally white (Lac⁻) (data not shown). The stationary-phase activity of *rybB-lacZ* seen in the wild type was completely abolished in the $\Delta\rho E$ background (Fig. 3C). Northern blots




FIG. 4. σ^E -dependent RybB RNA expression in vivo and in vitro. (A) Northern blot of RybB RNA isolated from wild-type (KMT12000), $\Delta rpoE$ (KMT12047), and $\Delta rybB$ (KMT12054) strains. Total RNA was isolated from exponential (optical density at 600 nm [OD_{600}], 0.5 to 0.7) and stationary (OD_{600} , 2.5 to 3.0) phases of growth and processed as described in Materials and Methods. (B) Northern blot for RybB RNA isolated in exponential phase (OD_{600} , 0.5 to 0.7). The wild-type *rybB-lacZ* strain (KMT12000) containing the vector control pTrc99A or pLC245 was grown to an OD_{600} of 0.3 and treated with 100 μ M IPTG for 60 min, and total RNA was isolated from each sample and corresponding no-IPTG control and processed as described in Materials and Methods. (C) In vitro transcription assay with purified core RNA polymerase, purified σ^E , and the *rybB* DNA template. Lanes 1 to 3, purified core RNA polymerase only; lanes 4 to 6, purified core RNA polymerase plus purified σ^E with the respective templates: P_{degP} , a σ^E -dependent promoter (lanes 1 and 4); P_{rybB} (lanes 2 and 5); and P_{yudF} , a σ^{70} -dependent promoter (lanes 3 and 6).

confirmed that RybB RNA was not detected in a $\Delta rpoE$ background in stationary phase (Fig. 4A). In cells carrying the plasmid with an IPTG-inducible *rpoE* gene, expression of the *rybB* fusion was restored (data not shown) and RybB RNA could be detected even in exponential phase (Fig. 4B).

It was recently shown that the activity of σ^E -dependent promoters increases upon entry into stationary phase (14), which is consistent with our observations of *rybB-lacZ* activity (Fig. 1A). This stationary-phase induction of σ^E -dependent promoters was shown to be dependent on the alarmone guanosine 3',5'-bispyrophosphate (ppGpp) (14). If the stationary-phase induction of the *rybB* promoter is ppGpp dependent, then the stationary-phase induction of *rybB-lacZ* should be abolished in a ppGpp⁰ ($\Delta relA \Delta spoT$) genetic background. In order to determine if the *rybB* promoter was ppGpp dependent, a $\Delta relA \Delta spoT$ double mutant of the *rybB-lacZ* transcriptional fusion was constructed and assayed for β -galactosidase activity throughout growth. Consistent with the observations of Costanzo and Ades (14), the stationary-phase induction of *rybB-lacZ* is defective in a ppGpp⁰ genetic background ($\Delta relA \Delta spoT$) (Fig. 3D). Altogether, these results strongly support the idea that transcriptional induction of *rybB* is completely σ^E dependent. The short *rybB-lacZ* fusion gave parallel results, showing full dependence on *rpoE* on plates and increased synthesis in an *rseA* mutant (data not shown).

As final evidence that σ^E directly regulates *rybB*, in vitro transcription reactions were carried out to determine if there was a biochemical interaction between σ^E and the *rybB* promoter. In these experiments, the *rybB* transcript was expressed 10-fold more than the *degP* transcript (Fig. 4C), generally considered a good σ^E -dependent promoter (20). In a control ex-

periment using the vegetative sigma factor, σ^{70} , no transcript was detected for the *rybB* gene (data not shown). The transcript start determined here is consistent with that predicted from the promoter and previously determined in vivo (Fig. 2) (57). Taken together, the in vivo and in vitro data show that *rybB* is completely σ^E dependent and that in the presence of σ^E , the *rybB* promoter is highly transcribed.

RybB modulation of σ^E activity. Plasmids containing *rybB* were isolated in screen I as negative regulators of the *rybB-lacZ* fusion (Table 1), suggesting that RybB may negatively regulate itself. This was tested further by expressing *rybB* under the control of the pBAD promoter. The activity of *rybB-lacZ* was decreased 2-fold in exponential phase and 10-fold in stationary phase when RybB was overexpressed by induction with arabinose (Fig. 5A). Thus, overexpression of RybB is sufficient to down-regulate the σ^E -dependent *rybB* promoter. The short *rybB-lacZ* fusion gave similar results; RybB overexpression decreased expression of the fusion 10-fold (data not shown).

As discussed above, σ^E is the direct regulator of *rybB* transcription. Therefore, it is possible that RybB autorepression is due to a RybB regulatory effect on σ^E synthesis or activity. This was tested by assaying an *rpoE-lacZ* translational fusion in the presence of *rybB* expressed from the pBAD promoter. This fusion contains the σ^E -dependent promoter for *rpoE* but apparently not a functional σ^{70} promoter; it is fully σ^E dependent (data not shown). It should reflect both transcriptional and translational regulation of *rpoE*. The activity of *rpoE-lacZ* decreased approximately 3-fold in exponential and approximately 11-fold in stationary phase when RybB was overexpressed (Fig. 5B). These results parallel those of the *rybB-lacZ* fusion, indi-




FIG. 5. Regulation of σ^E by sRNAs. Aliquots of the cultures were assayed for β -galactosidase activity as described in Materials and Methods. The values are means from three independent experiments. Error bars represent standard deviations. Numbers on the y axis are machine units as described in Materials and Methods. (A and B) β -Galactosidase activities of a Δ *ara* *rybB::kan rybB-lacZ* long-transcriptional-fusion strain (KMT12120) containing the vector control pNM12 or pKMT5 (pBAD-*rybB*⁺) (A) and the wild-type *rpoE-lacZ* fusion strain (KMT14000) containing the vector control pNM12 or pKMT5 (B) in the presence of arabinose (0.02%) were assayed in both exponential and stationary phases of growth. (C and D) β -Galactosidase activities of wild-type (KMT12000), Δ *rybB* (KMT12054), and Δ *hfq* (KMT12055) *rybB-lacZ* strains (C) and wild-type (KMT14000), Δ *rybB* (KMT14002), and Δ *hfq* (KMT14003) *rpoE-lacZ* translational fusion strains (D) were assayed during growth in LB medium.

cating that RybB overexpression is sufficient to inhibit σ^E synthesis and/or activity.

Introduction of a *rybB*-null mutation into the *rybB-lacZ* transcriptional fusion strain led to a modest increase (less than twofold) in fusion expression (Fig. 5C). Not surprisingly, this effect was detectable only in stationary phase, when RybB is normally expressed. Since RybB was reported to bind to the RNA chaperone Hfq, and most Hfq-binding sRNAs require Hfq for their activity (55), mutating *hfq* should also increase the activity of the *rybB-lacZ* fusion. In fact, a null mutation in *hfq* increased expression of *rybB-lacZ* dramatically, in both exponential and stationary phase (Fig. 5C). Parallel effects were seen with the *rpoE-lacZ* fusion (Fig. 5D). Deletion of the *rybB* gene led to a twofold increase in the activity of the *rpoE-lacZ* fusion in stationary phase, and the absence of *hfq* increased the activity of the *rpoE-lacZ* fusion throughout growth

about 5- to 10-fold (Fig. 5D). Thus, it seems likely that, in addition to RybB, other Hfq-dependent RNAs also negatively regulate σ^E synthesis or activity. Furthermore, the high level of expression of these σ^E reporters in an *hfq* mutant under otherwise unperturbed growth conditions suggests that the σ^E stress response is significantly activated under these conditions. It is interesting that the activity of both *rybB-lacZ* and *rpoE-lacZ* fusions still increases in stationary phase, even in the absence of *hfq*. Other recent reports of Hfq-dependent negative regulation of σ^E in *E. coli* (19, 23, 42) (see Discussion), in *Salmonella* (21), and in *Vibrio cholerae* (18) are consistent with our findings.

Characterization of sRNA regulation of σ^E . One model for the RybB negative regulation of σ^E activity by sRNAs is by negative regulation of OMP synthesis, leading to a decrease in periplasmic stress. OMPs, including the major outer mem-




FIG. 6. RybB autorepression is RseA independent. β -Galactosidase activities were assayed as for Fig. 5. (A) β -Galactosidase activity of the Δara *rybB::kan rybB-lacZ* transcriptional fusion strain (KMT12120) containing the vector control pNM12 or pKMT5 and the $\Delta rseA$ *rybB::kan rybB-lacZ* transcriptional fusion strain (KMT12119) containing the vector control pNM12 or pKMT5 (pBAD-*rybB*⁺) grown with arabinose (0.02%); results for exponential growth are shown. (B) β -Galactosidase activity of the $\Delta rseA$ *rpoE-lacZ* fusion strain (KMT14001) containing the vector control pNM12 or pKMT5 grown with arabinose and assayed in both exponential and stationary phases of growth.

brane porin OmpC, contain a C-terminal Y-X-F tripeptide motif. This region has been shown to interact with the PDZ domain of the periplasmic protease DegS, leading to cleavage of the RseA anti-sigma factor (1, 61). As noted above, a mutation in *ompR* that reduces the synthesis of the major outer membrane porins OmpC and OmpF leads to decreased expression of σ^E -dependent reporters (Fig. 1A). This suggests that decreasing the flux of proteins to the cell surface is sufficient to decrease periplasmic stress. A number of sRNAs that negatively regulate OMPs with this C-terminal motif have been identified (reviewed in references 22 and 58). This set of sRNAs that target OMPs may provide an explanation for the strong effect of *hfq* mutants on σ^E activity.

If RybB autorepression and RpoE repression are indirect, via down-regulating OMP synthesis, we would expect the effect on σ^E -dependent promoters to depend upon RseA. This was tested with three different σ^E -dependent reporters, *rybB-lacZ*, *rpoE-lacZ* (Fig. 6), and *rpoHP3-lacZ* (data not shown). A wild type and a $\Delta rseA$ derivative of each reporter were tested for the effect of overexpressing RybB from an arabinose-inducible promoter. The *rseA* mutation significantly increased the basal level of expression of the *rybB-lacZ*, *rpoEP2-lacZ* (Fig. 6; compare Fig. 6B to Fig. 5B) and *rpoHP3-lacZ* (data not shown) fusions, as expected. RybB was unable to down-regulate the expression of any of these fusions in the $\Delta rseA$ strain background (Fig. 6 and data not shown).

Reexamination of plasmids from screens. The identification of σ^E as an essential regulator of *rybB-lacZ* suggested that plasmids and mutations that modulated *rybB-lacZ* expression in our screens should also act through effects on σ^E synthesis or activity, as RybB itself seems to. We reexamined the plasmids isolated in these screens to see whether they provided any further insights into σ^E signaling. Regulation of expression of the σ^E -dependent reporter might occur in various ways: (i) by

increasing or decreasing periplasmic stress by changes in synthesis of OMPs mediated through RseA, the anti-sigma factor for σ^E (1, 4, 5, 61); (ii) by titration of σ^E by the presence of σ^E promoter on a multicopy plasmid; or (iii) by affecting σ^E translation or activity. In addition, it is possible that a given σ^E reporter might be sensitive to an additional regulator that is not active on other σ^E promoters.

The stimulatory plasmids described in Table 2 all either directly or indirectly affect the synthesis of OMPs that signal to the RseA protease. Overexpression of OmpR presumably leads to the overexpression of OmpC or OmpF, both known to induce σ^E activity. Similarly, MipA, an OMP involved in murein synthesis (39, 59), carries the DegS-activating sequence in its C terminus (61), and activates not only the *rybB-lacZ* fusion but also another σ^E -dependent fusion, *degP-lacZ* (Fig. 1D). The other two plasmids isolated in screen III (Table 2) acted upstream of OmpR and therefore, we believe, act by signaling to OmpR, indirectly increasing Omp synthesis. It seems likely that overexpression of many genes perturbs the cell envelope and/or osmolarity control sufficiently to cause induction of the EnvZ/OmpR system and therefore stimulate σ^E expression.

In a reciprocal manner, we expect that the plasmids in screen I (Table 1) that decreased expression of the *rybB-lacZ* fusion decrease the synthesis and/or activity of σ^E . Three of the plasmids contain the regions previously identified either as carrying σ^E promoters or showing regulation by σ^E in genome-wide expression arrays (Table 1) (46). Titration of σ^E is one possible explanation for the down-regulation of the *rybB* promoter by these plasmids, but it is also likely that the products are made in a σ^E -dependent fashion because they contribute to decreasing periplasmic stress. One plasmid carries *cpxR* and part of *cpxA*; *cpxR* is the response regulator for a set of genes also involved in periplasmic stress relief (44). Presumably, overex-




FIG. 7. NsrR regulation of *rybB* and *rpoE* expression (A) β -Galactosidase activities of the $\Delta ara714$ *leu::Tn10* (*Tet*^r) *rybB-lacZ* transcriptional fusion strain (KMT12069) containing the vector control pNM12 or pKMT8 (pBAD-*nsrR*⁺) and the $\Delta ara714$ *leu::Tn10* (*Tet*^r) $\Delta rseA$ *rybB-lacZ* transcriptional fusion strain (KMT12078) containing the vector control pNM12 or pKMT8 assayed in the presence of arabinose (0.02%) in late exponential phase. (B) β -Galactosidase activities of the wild type (KMT14000) and the $\Delta rseA$ *rpoE-lacZ* fusion strain (KMT14001) containing the vector control pNM12 or pKMT8 assayed in late exponential phase.

pressing CpxR turns on genes that reduce the basal level signaling to σ^E . Consistent with this idea, overexpression of CpxR is known to functionally compensate for lack of σ^E at high temperatures (13).

One plasmid identified in screen I carried *nsrR*, a transcriptional repressor that responds to nitric oxide and helps the cell respond to the stress of nitric oxide exposure (7). Overexpression of *nsrR* from the pBAD promoter decreased expression of the *rybB-lacZ* fusion, as well as expression of *rpoE-lacZ* (Fig. 7A and B, respectively). Therefore, the activity of the original plasmid appears to be due to NsrR. The overexpression of *nsrR* inhibited these two σ^E reporters even in a $\Delta rseA$ strain (Fig. 7). This identifies NsrR as a novel, RseA-independent regulator of σ^E -dependent genes, although whether regulation is direct or indirect remains to be determined. This suggests that the nitrosative stress response may somehow communicate with the envelope stress response. None of the other plasmids from screen I were tested in this way.

DISCUSSION

The σ^E regulon of *E. coli* controls a set of periplasmic proteases and chaperones, as well as proteins that act to properly direct proteins to the outer membrane (46). The work described here, along with recent publications and reports from other groups, extends the regulon to noncoding regulatory RNAs and demonstrates that these regulatory RNAs have an important role in reducing the basal level of induction of the σ^E system, presumably by reducing the flux of OMPs to the cell surface (24, 42).

RybB, the sRNA studied here, is a conserved 81-nt RNA that binds Hfq and is expressed during stationary-phase growth (57, 62). In the present study, we examined the regulation and function of RybB. We found that *rybB* was fully dependent

upon σ^E for expression in vivo, was transcribed efficiently by σ^E in vitro, and has a promoter that fits the previously described consensus for σ^E promoters (46). These findings indicate that *rybB* is an sRNA member of the σ^E regulon. We further found that RybB function is to negatively regulate the σ^E regulon: RybB overexpression decreases expression of several σ^E promoters, including *rybB* itself; deletion of *rybB* has the converse effect.

While the manuscript was in preparation, two groups working with *E. coli* and *Salmonella* reported that RybB and another regulatory RNA, MicA, require σ^E for their transcription, down-regulate σ^E activity, and have as their direct targets several OMPs (24, 42, 53). The only known target of MicA is the major OMP OmpA; RybB down-regulates transcripts for at least two porins in *E. coli* and 14 transcripts for OMPs in *Salmonella*. Some targets down-regulated after overexpression of σ^E , for instance, *ompX*, have no known sRNA regulators, suggesting that there may be other σ^E -regulated sRNAs as well (23, 46).

The primary role of σ^E has been thought to be responding to envelope stress that results from environmental change. The emerging evidence that the σ^E regulon encodes multiple sRNAs that negatively regulate porins and other envelope proteins changes this view in a very important way. It has long been known that cells have mechanisms to balance production of OMPs. For example, upon overexpression of some OMPs, the expression of other OMPs is lowered (11, 12, 17). Therefore, there must be a mechanism for coordination of OMP expression that functions above the level of individual transcription factors such as OmpR. We propose that σ^E , through its sRNA arm, performs this coordination function. It appears that during normal cell growth, some, if not most, OMP genes are transcribed and would be translated in excess, beyond the level needed for function. The relative levels of many of these pro-

teins are determined by sRNAs of the regulon utilizing two mechanisms: (i) sRNAs directly target certain mRNAs for destruction, and (ii) relative levels of these targets in turn alter σ^E activity, either directly by activating DegS to initiate RseA cleavage (OmpC, OmpW) or indirectly by altering the envelope folding environment (OmpA). Together, these mechanisms allow the cell to set the relative levels of many envelope constituents. Moreover, additional features enhance the reach of this integrative function of σ^E . Many sRNAs not encoded in the σ^E regulon have targets that affect σ^E activity (e.g., MicF and MicC), thereby allowing their effects to be coordinated with those of σ^E . Taken together, these results suggest that a primary function of σ^E is to coordinate expression of envelope constituents, especially porins, under all growth conditions.

The importance of this coordination can be best appreciated when it is disrupted. Overexpression of the σ^E regulon causes cell growth problems and lysis in *E. coli* (25, 40). In *Salmonella*, a strain with an *rseA* deletion rapidly acquired secondary mutations that decreased σ^E activity, and an *rseA hfq* double mutation proved to be lethal (21). Presumably under these conditions, the σ^E regulon is fully induced and there is abundant machinery for folding OMPs and inserting them into the outer membrane. Thus, the tendency for these cells to lyse may instead reflect too high a protein load in the outer membrane, rather than the accumulation of unfolded OMPs in the periplasm. The limitation of OMP synthesis in a σ^E -dependent fashion thus appears to reflect a critical role in regulating both the flux of OMPs (to allow proper folding and insertion) and the capacity of the outer membrane itself.

Other levels of regulation of the RybB promoter. In this picture of the σ^E network, RybB and MicA are made in response to σ^E stress and down-regulate that stress, either directly or indirectly by down-regulating the synthesis of OMPs that are the substrates for the σ^E functions. The strong phenotype of an *hfq* mutant in expressing σ^E genes even in exponential growth suggests that the down-regulation of other OMPs by sRNAs, probably not all under σ^E control, is operating under all growth conditions. RybB overexpression down-regulates three different σ^E -dependent fusions tested, *rybB-lacZ*, *rpoHP3-lacZ*, and *rpoEP2-lacZ*, all in an RseA-dependent fashion. We expect all OMP signaling to be via RseA cleavage of this anti-sigma factor. The observation that the three fusions are not down-regulated by RybB in an *rseA* mutant strongly suggests that the RybB effect on σ^E activity and/or synthesis is RseA dependent (i.e., via OMPs). This is consistent with the observations of others that RybB down-regulates OMPs (24, 42).

The *rybB-lacZ* fusion proved to be a sensitive reporter for examining the σ^E signaling pathway. As noted above, one of the plasmids isolated in screen I which down-regulated the *rybB-lacZ* fusion (Table 1) contained *nsrR*, which proved to be an RseA-independent regulator of two σ^E -dependent fusions, *rybB-lacZ* and *rpoE-lacZ* (Fig. 7). The identification of NsrR as a regulator of σ^E -dependent genes suggests a connection between periplasmic stress and nitric oxide stress that warrants further investigation.

Upstream signals that affect σ^E . The approach used in this study to investigate *rybB* regulation identified both multicopy plasmids and mutations that affected the *rybB-lacZ* fusion, now known to reflect σ^E function. Among the plasmids that re-

duced function were some that may titrate σ^E or reduce stress that signals to σ^E ; some stimulatory plasmids identified in screen III act by activating OmpR, presumably increasing the synthesis of outer membrane proteins. Previous plasmid screens for stimulators of a σ^E -dependent promoter gave a nonoverlapping list of candidates (36), suggesting that there are significant numbers of genes and pathways that can lead to induction.

Most puzzling of the genetic loci identified in our screens was the insertion mutation in the *rep* gene, encoding a DNA helicase. This insertion significantly decreases expression of the fusion, and therefore presumably σ^E activity. The Rep helicase is involved in the restart of damaged replication forks, and in cells devoid of *rep*, there may be accumulation of DNA for early regions of the chromosome (in front of possible replication errors) compared to later regions. The genes for the major outer membrane porins, *ompC*, *ompF*, and *ompA*, are all relatively far from *oriC*. Their location may cause a lower level of OMP synthesis in the *rep* mutant. If this is the explanation for the lower activity in the *rep* mutant, it may be serendipitous, or it may reflect a mechanism for ensuring that the dosage of these genes does not increase until near the end of the replication cycle, when cells, and therefore the available cell surface, are larger.

Our investigation of *rybB* regulation, combined with work by others, points to a critical role for Hfq-regulated sRNAs in the control of the cell surface and the transport of OMPs to the cell surface. At the core of this response are MicA and RybB, made in large amounts only when the export machinery is stressed, and capable of down-regulating major outer membrane porins. The combination of multiple sRNAs, regulated in a variety of ways and targeting a variety of different OMP transcripts, provides the flexibility and capability of responding to many signals that the cell is likely to need.

ACKNOWLEDGMENTS

We thank members of the Gottesman lab for their help at various stages of this project and Carol Gross for her support and comments on the work and the manuscript. In particular, we thank Nadim Majdalani for advice and assistance with various strain constructions. Also, we thank Carin Vanderpool and Mitsuoki Kawano for help with sequencing of the transposon insertions and for providing the pool of random kanamycin insertions. We thank Gisela Storz, Nadim Majdalani, Carin Vanderpool, Colleen McCullen, Maude Guillier, Alexander Boudour, Pierre Mandin, and Kyung Moon for their comments and critical reading of the manuscript.

This research was supported in part by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research.

REFERENCES

- Ades, S. E., L. E. Connolly, B. M. Alba, and C. A. Gross. 1999. The *Escherichia coli* σ^E -dependent extracytoplasmic stress response is controlled by the regulated proteolysis of an anti- σ factor. *Genes Dev.* **13**:2449–2461.
- Aiba, H., S. Adhya, and B. de Crombrughe. 1981. Evidence for two functional *gal* promoters in intact *Escherichia coli* cells. *J. Biol. Chem.* **256**:11905–11910.
- Alba, B. M., and C. A. Gross. 2004. Regulation of the *Escherichia coli* sigma-dependent envelope stress response. *Mol. Microbiol.* **52**:613–619.
- Alba, B. M., H. J. Ji Zhong, J. C. Pelayo, and C. A. Gross. 2001. *degS* (*hhoB*) is an essential *Escherichia coli* gene whose indispensable function is to provide σ^E activity. *Mol. Microbiol.* **40**:1323–1333.
- Alba, B. M., J. A. Leeds, C. Onufryk, C. Z. Lu, and C. A. Gross. 2002. DegS and YaeL participate sequentially in the cleavage of RseA to activate the σ^E dependent extracytoplasmic stress response. *Genes Dev.* **16**:2156–2168.
- Argaman, L., R. Hershsberg, J. Vogel, G. Bejerano, E. G. Wagner, H.

- Margalit, and S. Altuvia. 2001. Novel small RNA-encoding genes in the intergenic regions of *Escherichia coli*. *Curr. Biol.* **11**:941–950.
7. Bodenmiller, D. M., and S. Spiro. 2006. The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-sensitive transcriptional regulator. *J. Bacteriol.* **188**:874–881.
 8. Chen, S., E. A. Lesnik, T. A. Hall, R. Sampath, R. H. Griffey, D. J. Ecker, and L. B. Blyn. 2002. A bioinformatics based approach to discover small RNA genes in the *Escherichia coli* genome. *Biosystems* **65**:157–177.
 9. Chun, K. T., H. J. Edenberg, M. R. Kelley, and M. G. Goebel. 1997. Rapid amplification of uncharacterized transposon-tagged DNA sequences from genomic DNA. *Yeast* **13**:233–240.
 10. Chung, C. T., S. L. Niemela, and R. H. Miller. 1989. One-step preparation of competent *Escherichia coli*: transformation and storage of bacterial cells in the same solution. *Proc. Natl. Acad. Sci. USA* **86**:2172–2175.
 11. Click, E. M., G. A. McDonald, and C. A. Schnaitman. 1988. Translational control of exported proteins that results from OmpC porin overexpression. *J. Bacteriol.* **170**:2005–2011.
 12. Click, E. M., and C. A. Schnaitman. 1989. Export-defective LamB protein is a target for translational control caused by OmpC porin overexpression. *J. Bacteriol.* **171**:616–619.
 13. Connolly, L., A. De Las Penas, B. M. Alba, and C. A. Gross. 1997. The response to extracytoplasmic stress in *Escherichia coli* is controlled by partially overlapping pathways. *Genes Dev.* **11**:2012–2021.
 14. Costanzo, A., and S. E. Ades. 2006. Growth phase-dependent regulation of the extracytoplasmic stress factor, sigma E, by guanosine 3',5'-bispyrophosphate (ppGpp). *J. Bacteriol.* **188**:4627–4634.
 15. Court, D. L., S. Swaminathan, D. Yu, H. Wilson, T. Baker, M. Bubunenko, J. Sawitzke, and S. K. Sharan. 2003. Mini-lambda: a tractable system for chromosome and BAC engineering. *Gene* **315**:63–69.
 16. De Las Peñas, A., L. Connolly, and C. A. Gross. 1997. The σ^E -mediated response to extracytoplasmic stress in *Escherichia coli* is transduced by RseA and RseB, two negative regulators of σ^E . *Mol. Microbiol.* **24**:373–385.
 17. Diedrich, D. L., and J. A. Fralick. 1982. Relationship between the OmpC and LamB proteins of *Escherichia coli* and its influence on the protein mass of the outer membrane. *J. Bacteriol.* **149**:156–160.
 18. Ding, Y., B. M. Davis, and M. K. Waldor. 2004. Hfq is essential for *Vibrio cholerae* virulence and downregulates sigma expression. *Mol. Microbiol.* **53**:345–354.
 19. Douchin, V., C. Bohn, and P. Bouloc. 2006. Down-regulation by porins by a small RNA bypasses the essentiality of the regulated intramembrane proteolysis protease RseP in *Escherichia coli*. *J. Biol. Chem.* **281**:12253–12259.
 20. Erickson, J. W., V. Vaughn, W. A. Walter, F. C. Neidhardt, and C. A. Gross. 1987. Regulation of the promoters and transcripts of *rpoH*, the *Escherichia coli* heat shock regulatory gene. *Genes Dev.* **1**:419–432.
 21. Figueroa-Bossi, N., S. Lemire, D. Maloriol, R. Balbontin, J. Casadesus, and L. Bossi. 2006. Loss of Hfq activates the sigma E-dependent envelope stress response in *Salmonella enterica*. *Mol. Microbiol.* **62**:838–852.
 22. Guillier, M., and S. Gottesman. 2006. Remodelling of the *Escherichia coli* outer membrane by two small regulatory RNAs. *Mol. Microbiol.* **59**:231–247.
 23. Guisbert, E., V. A. Rhodius, N. Ahuja, E. Witkin, and C. A. Gross. 2007. Hfq modulates the σ^E -mediated envelope stress response and the σ^{32} -mediated cytoplasmic stress response in *Escherichia coli*. *J. Bacteriol.* **189**:1963–1973.
 24. Johansen, J., A. A. Rasmussen, M. Overgaard, and P. Valentin-Hansen. 2006. Conserved small non-coding RNAs that belong to the σ^E regulon: role in down-regulation of outer membrane proteins. *J. Mol. Biol.* **364**:1–8.
 25. Kabir, M. S., D. Yamashita, S. Koyama, T. Oshima, K. Kurokawa, M. Maeda, R. Tsunedomi, M. Murata, C. Wada, H. Mori, and M. Yamada. 2005. Cell lysis directed by σ^E in early stationary phase and effect of induction of the *rpoE* gene on global gene expression in *Escherichia coli*. *Microbiology* **151**:2721–2735.
 26. Kawamoto, H., T. Morita, A. Shimizu, T. Inada, and H. Aiba. 2005. Implication of membrane localization of target mRNA in the action of a small RNA: mechanism of post-transcriptional regulation of glucose transporter in *Escherichia coli*. *Genes Dev.* **19**:328–338.
 27. Kawano, M., A. A. Reynolds, J. Miranda-Rios, and G. Storz. 2005. Detection of 5'- and 3'-UTR-derived small RNAs and cis-encoded antisense RNAs in *Escherichia coli*. *Nucleic Acids Res.* **33**:1040–1050.
 28. Keseler, I. M., J. Collado-Vides, S. Gama-Castro, J. Ingraham, S. Paley, I. T. Paulsen, M. Peralta-Gil, and P. D. Karp. 2005. EcoCyc: a comprehensive database resource for *Escherichia coli*. *Nucleic Acids Res.* **33**:D334–D337.
 29. Kleckner, N., J. Bender, and S. Gottesman. 1991. Uses of transposons with emphasis on Tn10. *Methods Enzymol.* **204**:139–180.
 30. Lomovskaya, O., K. Lewis, and A. Matin. 1995. EmrR is a negative regulator of the *Escherichia coli* multidrug resistance pump EmrAB. *J. Bacteriol.* **177**:2328–2334.
 31. Lonetto, M. A., K. L. Brown, K. E. Rudd, and M. J. Buttner. 1994. Analysis of the *Streptomyces coelicolor sigE* gene reveals the existence of a subfamily of eubacterial RNA polymerase σ factors involved in the regulation of extracytoplasmic functions. *Proc. Natl. Acad. Sci. USA* **91**:7573–7577.
 32. Majdalani, N., C. Cunniff, D. Sledjeski, T. Elliott, and S. Gottesman. 1998. DsrA RNA regulates translation of RpoS message by an anti-antisense mechanism, independent of its action as an antisilencer of transcription. *Proc. Natl. Acad. Sci. USA* **95**:12462–12467.
 33. Massé, E., F. E. Escorcia, and S. Gottesman. 2003. Coupled degradation of a small regulatory RNA and its mRNA targets in *Escherichia coli*. *Genes Dev.* **17**:2374–2383.
 34. Massé, E., and S. Gottesman. 2002. A small RNA regulates the expression of genes involved in iron metabolism in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **99**:4620–4625.
 35. McDowell, J. C., J. W. Roberts, D. J. Jin, and C. Gross. 1994. Determination of intrinsic transcription initiation by RNA polymerase elongation rate. *Science* **266**:822–825.
 36. Mecsas, J., P. E. Rouviere, J. W. Erickson, T. J. Donohue, and C. A. Gross. 1993. The activity of σ^E , an *Escherichia coli* heat-inducible σ -factor, is modulated by expression of outer membrane proteins. *Genes Dev.* **7**:2618–2628.
 37. Mecsas, J., R. Welch, J. W. Erickson, and C. A. Gross. 1995. Identification and characterization of an outer membrane protein, OmpX, in *Escherichia coli* that is homologous to a family of outer membrane proteins including Ail of *Yersinia enterocolitica*. *J. Bacteriol.* **177**:799–804.
 38. Miller, J. H. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 39. Molloy, M. P., B. R. Herbert, M. B. Slade, T. Rabilloud, A. S. Nouwens, and K. L. Williams. 2000. Proteomic analysis of the *Escherichia coli* outer membrane. *Eur. J. Biochem.* **267**:2871–2881.
 40. Nitta, T., H. Nagamitsu, M. Murata, H. Izu, and M. Yamada. 2000. Function of the σ^E regulon in dead-cell lysis in stationary phase *Escherichia coli*. *J. Bacteriol.* **182**:5231–5237.
 41. Norioka, S., K. Ramakrishnan, K. Ikenaka, and M. Inouye. 1986. Interaction of a transcriptional activator, OmpR, with reciprocally osmoregulated genes, *ompF* and *ompC*, of *Escherichia coli*. *J. Biol. Chem.* **261**:17113–17119.
 42. Papenfort, K., V. Pfeiffer, F. Mika, S. Lucchini, J. C. D. Hinton, and J. Vogel. 2006. σ^E -dependent small RNAs of *Salmonella* respond to membrane stress by accelerating global *omp* mRNA decay. *Mol. Microbiol.* **62**:1674–1688.
 43. Peters, J. E., and N. L. Craig. 2000. Tn7 transposes proximal to DNA double-strand breaks and into regions where chromosomal DNA replication terminates. *Mol. Cell* **6**:573–582.
 44. Raivio, T. L., and T. J. Silhavy. 1997. Transduction of envelope stress in *Escherichia coli* by the Cpx two-component system. *J. Bacteriol.* **179**:7724–7733.
 45. Rhodius, V., N. Savary, A. Kolb, and S. Busby. 2001. Assays for transcription factor activity. *Methods Mol. Biol.* **148**:451–464.
 46. Rhodius, V. A., W. C. Suh, G. Nonaka, J. West, and C. A. Gross. 2006. Conserved and variable functions of the σ^E stress response in related genomes. *PLoS Biol.* **4**:0043–0059.
 47. Rivas, E., R. J. Klein, T. A. Jones, and S. R. Eddy. 2001. Computational identification of noncoding RNAs in *E. coli* by comparative genomics. *Curr. Biol.* **11**:1369–1373.
 48. Rouvière, P., A. De Las Penas, J. Mecsas, C. Zen Lu, K. E. Rudd, and C. A. Gross. 1995. *rpoE*, the gene encoding the second heat-shock sigma factor, σ^E , in *Escherichia coli*. *EMBO J.* **14**:1032–1042.
 49. Silhavy, T. J., M. L. Berman, and L. W. Enquist. 1984. Experiments with gene fusions. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
 50. Simons, R. W., F. Houtman, and N. Kleckner. 1987. Improved single and multicopy *lac*-based cloning vectors for protein and operon fusions. *Gene* **53**:85–96.
 51. Sledjeski, D. D., and S. Gottesman. 1996. The small RNA, DsrA, is essential for the low temperature expression of RpoS during exponential growth in *Escherichia coli*. *EMBO J.* **15**:3993–4000.
 52. Straus, D. B., W. A. Walter, and C. A. Gross. 1989. The activity of σ^{32} is reduced under conditions of excess heat shock protein production in *Escherichia coli*. *Genes Dev.* **3**:2003–2010.
 53. Udekwi, K. I., and E. G. Wagner. 2007. Sigma E controls biogenesis of the antisense RNA MicA. *Nucleic Acids Res.* **35**:1279–1288.
 54. Ulbrandt, N. D., J. A. Newitt, and H. D. Bernstein. 1997. The *E. coli* signal recognition particle is required for the insertion of a subset of inner membrane proteins. *Cell* **88**:187–196.
 55. Valentin-Hansen, P., M. Eriksen, and C. Udesen. 2004. The bacterial Sm-like protein Hfq: a key player in RNA transactions. *Mol. Microbiol.* **51**:1525–1533.
 56. Vanderpool, C. K., and S. Gottesman. 2004. Involvement of a novel transcriptional activator and small RNA in post-transcriptional regulation of the glucose phosphoenolpyruvate phosphotransferase system. *Mol. Microbiol.* **54**:1076–1089.
 57. Vogel, J., V. Bartels, H. H. Tang, G. Churakov, J. G. Slagter-Jager, A. Huttenhofer, and E. G. H. Wagner. 2003. RNomics in *Escherichia coli* detects new sRNA species and indicates parallel transcriptional output in bacteria. *Nucleic Acids Res.* **31**:6435–6443.
 58. Vogel, J., and K. Papenfort. 2006. Small non-coding RNAs and the bacterial outer membrane. *Curr. Opin. Microbiol.* **9**:605–611.
 59. Vollmer, W., M. von Rechenberg, and J.-V. Holtje. 1999. Demonstration of molecular interaction between the murein polymerase PBPI, the lytic transglycosylase MltA, and the scaffolding protein MipA of *Escherichia coli*. *J. Biol. Chem.* **274**:6726–6734.

60. **Wagner, E. G. H., and R. W. Simons.** 1994. Antisense RNA control in bacteria, phages, and plasmids. *Annu. Rev. Microbiol.* **48**:713–742.
61. **Walsh, N. P., B. M. Alba, B. Bose, C. A. Gross, and R. T. Sauer.** 2003. OMP peptide signals initiate the envelope-stress response by activating DegS protease via relief of inhibition mediated by its PDZ domain. *Cell* **113**:61–71.
62. **Wassarman, K. M., F. Repoila, C. Rosenow, G. Storz, and S. Gottesman.** 2001. Identification of novel small RNAs using comparative genomics and microarrays. *Genes Dev.* **15**:1637–1651.
63. **Xiong, A., A. Gottman, C. Park, M. Baetens, S. Pandza, and A. Martin.** 2000. The EmrR protein represses the *Escherichia coli emrRAB* multidrug resistance operon by directly binding to its promoter region. *Antimicrob. Agents Chemother.* **44**:2905–2907.
64. **Yoshida, M., L. Qin, L. A. Egger, and M. Inouye.** 2006. Transcription regulation of *ompF* and *ompC* by a single transcription factor, OmpR. *J. Biol. Chem.* **281**:17114–17123.
65. **Young, B. A., L. C. Anthony, T. M. Gruber, T. M. Arthur, E. Heyduk, C. Z. Lu, M. M. Sharp, A. Heydorn, R. R. Burgess, and C. A. Gross.** 2001. A coiled-coil from the RNA polymerase β' subunit allosterically induces selective nontemplate strand binding by $\sigma 70$. *Cell* **105**:935–944.
66. **Zhou, Y.-N., and S. Gottesman.** 1998. Regulation of proteolysis of the stationary-phase sigma factor RpoS. *J. Bacteriol.* **180**:1154–1158.