

# Functional Interactions between Coexisting Toxin-Antitoxin Systems of the *ccd* Family in *Escherichia coli* O157:H7<sup>∇</sup>

Myriam Wilbaux,<sup>1</sup> Natacha Mine,<sup>1</sup> Anne-Marie Guérout,<sup>2</sup>  
 Didier Mazel,<sup>2</sup> and Laurence Van Melderen<sup>1\*</sup>

Laboratoire de Génétique des Procaryotes, Institut de Biologie et Médecine Moléculaires, Université Libre de Bruxelles, 12 Rue des Professeurs Jeener et Brachet, 6041 Gosselies, Belgium,<sup>1</sup> and Unité Plasticité du Génome Bactérien, Département Génomes et Génétique, Institut Pasteur, 25 Rue du Dr. Roux, 75724 Paris, France<sup>2</sup>

Received 30 October 2006/Accepted 16 January 2007

**Toxin-antitoxin (TA) systems are widely represented on mobile genetic elements as well as in bacterial chromosomes. TA systems encode a toxin and an antitoxin neutralizing it. We have characterized a homolog of the *ccd* TA system of the F plasmid (*ccd<sub>F</sub>*) located in the chromosomal backbone of the pathogenic O157:H7 *Escherichia coli* strain (*ccd<sub>O157</sub>*). The *ccd<sub>F</sub>* and the *ccd<sub>O157</sub>* systems coexist in O157:H7 isolates, as these pathogenic strains contain an F-related virulence plasmid carrying the *ccd<sub>F</sub>* system. We have shown that the chromosomal *ccd<sub>O157</sub>* system encodes functional toxin and antitoxin proteins that share properties with their plasmidic homologs: the CcdB<sub>O157</sub> toxin targets the DNA gyrase, and the CcdA<sub>O157</sub> antitoxin is degraded by the Lon protease. The *ccd<sub>O157</sub>* chromosomal system is expressed in its natural context, although promoter activity analyses revealed that its expression is weaker than that of *ccd<sub>F</sub>*. *ccd<sub>O157</sub>* is unable to mediate postsegregational killing when cloned in an unstable plasmid, supporting the idea that chromosomal TA systems play a role(s) other than stabilization in bacterial physiology. Our cross-interaction experiments revealed that the chromosomal toxin is neutralized by the plasmidic antitoxin while the plasmidic toxin is not neutralized by the chromosomal antitoxin, whether expressed ectopically or from its natural context. Moreover, the *ccd<sub>F</sub>* system is able to mediate postsegregational killing in an *E. coli* strain harboring the *ccd<sub>O157</sub>* system in its chromosome. This shows that the plasmidic *ccd<sub>F</sub>* system is functional in the presence of its chromosomal counterpart.**

Toxin-antitoxin (TA) proteic systems were originally discovered on low-copy-number plasmids (for reviews on TA systems, see references 11, 22, 24, and 29). They are composed of two genes organized in an operon encoding a toxin and an antitoxin that antagonizes it. The expression of the TA genes is autoregulated at the transcriptional level; the antitoxin acts as a repressor and the toxin often as a corepressor. The antitoxin is an unstable protein degraded by an ATP-dependent protease, while the toxin is a stable protein that inhibits an essential cellular process (e.g., replication and translation). TA systems contribute to plasmid stability by a mechanism called postsegregational killing (PSK). PSK relies on the differential stabilities of the antitoxin and toxin proteins and leads to the killing of daughter bacteria that did not receive a plasmid copy at cell division (31, 50, 53).

Recent computational analyses have shown that TA systems are widely represented in eubacterial and archaeobacterial chromosomes, suggesting a role for horizontal gene transfer in the spread of these genes (5, 6, 38). The localization of chromosomal TA systems is quite varied. Some are localized within exogenous DNA islands like phages (*relBE*<sub>K-12</sub> in the cryptic lambdoid Qin prophage of *Escherichia coli* MG1655) (40),

transposons (*relBE* homolog in Tn5401 of *Bacillus thuringiensis* (23), and superintegrons (*relBE*, *parDE*, *phd-doc*, and *higAB* homologs in the superintegron of *Vibrio cholerae* [17, 38, 41]). Others, such as *mazEF* (*chpA*) and *chpB* of *E. coli* K-12, are apparently settled in the chromosome, flanked by metabolic or regulatory genes (35, 36). The chromosomal TA systems were shown to be activated under stress conditions, although the outcomes of activation appear to be different for the various systems. On the one hand, the *mazEF* system was shown to be a suicide module leading to cell death under various stressful conditions (e.g., overproduction of ppGpp, DNA damage, high temperatures, and oxidative stress) (2, 4, 30, 46, 47). On the other hand, the *mazEF* and *relBE*<sub>K-12</sub> systems have been described as growth modulators under conditions of amino acid starvation (15, 16, 39). The function of TA systems localized in transposons or superintegrons could be similar to that of plasmidic TA systems, i.e., they could stabilize these exogenous DNA islands within the bacterial genome (17, 41).

We have performed a functional analysis of the chromosomal *ccd<sub>O157</sub>* system of *E. coli* O157:H7, which is homologous to the *ccd* TA system of the F plasmid (*ccd<sub>F</sub>*) (for reviews on *ccd*, see references 19 and 52). The *ccd<sub>F</sub>* system is composed of the unstable CcdA<sub>F</sub> antitoxin, which is degraded by the Lon ATP-dependent protease (53, 54), and of the stable CcdB<sub>F</sub> toxin. Both CcdA<sub>F</sub> and CcdB<sub>F</sub> are required for autoregulation (1, 42, 49). CcdB<sub>F</sub> targets the DNA gyrase and leads to replication and transcription inhibition, SOS induction, and ultimately, to cell death (8, 9, 20, 31, 34). The *ccd<sub>F</sub>* and the *ccd<sub>O157</sub>* systems coexist in *E. coli* O157:H7 isolates; the *ccd<sub>F</sub>* system is

\* Corresponding author. Mailing address: Laboratoire de Génétique des Procaryotes, Institut de Biologie et Médecine Moléculaires, Université Libre de Bruxelles, 12 Rue des Professeurs Jeener et Brachet, 6041 Gosselies, Belgium. Phone: 32 2 650 97 78. Fax: 32 2 650 97 70. E-mail: lvmelder@ulb.ac.be.

<sup>∇</sup> Published ahead of print on 26 January 2007.

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Genotype or plasmid properties <sup>a</sup>	Reference or source
<b>Strains</b>		
B462	DH2 <i>lacI</i> <sup>q</sup> <i>gyrA462</i> <i>zei::Tn10</i>	Laboratory collection
CSH50	<i>ara</i> $\Delta$ ( <i>lac-pro</i> ) <i>strA thi</i>	37
SG22622	MC4100 <i>cpsB::lacZ</i> $\Delta$ <i>ara malP::lacI</i> <sup>q</sup>	S. Gottesman
SG22622 <i>gyrA462</i> <i>zei::Tn10</i>	SG22622 containing the <i>gyrA462</i> <i>CcdB</i> <sub>F</sub> resistance mutation	3
SG22623	SG22622 $\Delta$ <i>lon-510</i>	S. Gottesman
O55:H7 TB182A	Enterohemorrhagic <i>E. coli</i>	STEC center
O55:H7 $\Delta$ <i>ccd</i> <sub>O157</sub>	O55:H7 TB182A $\Delta$ <i>ccd</i> <sub>O157</sub>	This work
<b>Plasmids</b>		
pBAD33	p15A, Cm <sup>r</sup> , pBAD promoter	28
pBAD33- <i>ccdB</i> <sub>F</sub>	pBAD33 derivative containing the <i>ccdB</i> <sub>F</sub> gene under the control of the pBAD promoter	This work
pBAD33- <i>ccdB</i> <sub>O157</sub>	pBAD33 derivative containing the <i>ccdB</i> <sub>O157</sub> gene under the control of the pBAD promoter	This work
pCP20	pSC101 ts derivative containing the FLP gene under the control of the $\lambda$ cI857 repressor	13
pJL207	p15A, Cm <sup>r</sup> , <i>lacZ</i>	33
pJL- <i>ccd</i> <sub>F</sub> (pULB2600)	pJL207 containing the OP of the <i>ccd</i> <sub>F</sub> operon	42
pJL- <i>ccd</i> <sub>O157</sub>	pJL207 containing the OP of the <i>ccd</i> <sub>O157</sub> operon	This work
pKD4	pANTS $\gamma$ derivative containing an FRT-flanked kanamycin resistance gene, Amp <sup>r</sup>	21
pKK223-3	ColE1, Amp <sup>r</sup> , pTac promoter	10
pKK- <i>ccdA</i> <sub>F</sub> (pULB2709)	pKK223-3 derivative containing the <i>ccdA</i> <sub>F</sub> gene under the control of the Tac promoter	53
pKK- <i>ccdA</i> <sub>O157</sub>	pKK223-3 derivative containing the <i>ccdA</i> <sub>O157</sub> gene under the control of the Tac promoter	This work
pKOBEG	pSC101 ts derivative containing an arabinose-inducible $\lambda$ red $\gamma$ $\beta$ $\alpha$ operon, Cm <sup>r</sup>	12
pKT279	pBR322 derivative, Tet <sup>r</sup>	48
pKT- <i>ccd</i> <sub>F</sub> (pULB2707)	pKT279 containing the <i>ccd</i> <sub>F</sub> operon	42
pKT- <i>ccd</i> <sub>O157</sub>	pKT279 containing the <i>ccd</i> <sub>O157</sub> operon	This work
pMLO59	pGB2 ts derivative, Spec <sup>r</sup>	M. Labocka
pMLO- <i>ccd</i> <sub>F</sub> (pULB2710)	pMLO59 containing the <i>ccd</i> <sub>F</sub> operon	53
pMLO- <i>ccd</i> <sub>O157</sub>	pMLO59 containing the <i>ccd</i> <sub>O157</sub> operon	This work

<sup>a</sup> FRT, FLP recombination target; ts, temperature sensitive.

present on an F-related virulence plasmid (pO157), while the *ccd*<sub>O157</sub> system is present on the chromosome. Therefore, we have tested whether the components of these systems interact and showed that the *ccd*<sub>F</sub> system is functional in the presence of its chromosomal homolog.

## MATERIALS AND METHODS

**Bacterial strains, plasmids, and media.** Strains and plasmids used in this work are listed in Table 1.

**Construction of the O55:H7 $\Delta$ *ccd*<sub>O157</sub> strain.** The pKOBEG plasmid (12) was used as described at the website <http://www.pasteur.fr/recherche/unites/Ggb/3SPCRprotocol.html>. The kanamycin resistance cassette of pKD4 was amplified by PCR with the following primers: P1 (5' ATA CTA GAC GTA TAA ATT GTA CAG GAG CAC GAT ATC GTG TAG GCT GGA GCT GCT TC) and P2 (5' AAG GAT TTG GGT GAG GGA GAG GCG GTC GCG TCT TAA CAT ATG AAT ATC CTC CTT AG). Deletion of the *ccd*<sub>O157</sub> system was constructed by following the method described in reference 21. The deletion and the flanking regions were checked by DNA sequencing.

**Construction of plasmids. (i) Expression plasmids.** The expression plasmids are isogenic, i.e., all the open reading frames (ORFs) were cloned using the same restriction sites in the expression vectors, and all the regulatory sequences were added by PCR (the Shine-Dalgarno box and the sequence between the Shine-Dalgarno box and the ATG of the ORFs were identical).

To construct the pBAD-*ccdB*<sub>F</sub> plasmid, the *ccdB* gene from the F plasmid was amplified by PCR using pULB2250 as a template (8) and the following primers: 5' CcdB-XbaI (5'TCT AGA AGG AGG GTG AAA TGC AGT TTA AGG) and 3' CcdB-PstI (5'AGT CTC TGC AGT TAT ATT CCC CAG AAC). The PCR product was cloned into the TOPO-XL vector (Invitrogen). The resulting plasmid was then digested by XbaI and PstI. The fragment containing *ccdB*<sub>F</sub> was inserted into pBAD33 that was opened by the same enzymes. The ligation mixture was transformed in strain B462.

To construct the pBAD-*ccdB*<sub>O157</sub> plasmid, the *ccdB* gene from *E. coli* O157:H7 was amplified by PCR using *E. coli* O157:H7 chromosomal DNA

(ATCC 700927) as a template and the following primers: 5' CcdB<sub>O157</sub>-XbaI (5'TCT AGA AGG AGG TAG CGA TGC AAT TTA CGG) and 3' CcdB<sub>O157</sub>-PstI (5'AGT CTC TGC AGT TAA ATC CCG TCG AGC). The PCR product was digested by XbaI and PstI and inserted into pBAD33 that was digested with the same enzymes. The ligation mixture was transformed in strain B462.

To construct the pKK-*ccdA*<sub>O157</sub> plasmid, the *ccdA* gene from *E. coli* O157:H7 was amplified by PCR using *E. coli* O157:H7 chromosomal DNA (ATCC 700927) as a template and the following primers: 5' CcdA<sub>O157</sub>-EcoRI (5'TTG TGA ATT CTA TGA CTG CAA AAC GTA CCA) and 3' CcdA<sub>O157</sub>-PstI (5'AGT CTC TGC AGC TAG AAG CTC CGG TAC TC). The PCR product was cloned into the TOPO-XL vector (Invitrogen). The resulting plasmid was then digested by EcoRI and PstI. The fragment containing *ccdA*<sub>O157</sub> was inserted into pKK223-3 that was opened by the same enzymes.

**(ii) Promoter activity plasmids.** To construct the pJL-OP*ccd*<sub>O157</sub> plasmid, the operator/promoter region of the *ccd*<sub>O157</sub> operon was amplified by PCR using the Topo-*folAccd*<sub>O157</sub> plasmid (see below) as a template and the following primers: 5'O/P2-*ccd*<sub>O157</sub> (5'GGT ATT CAG CGA ATT CCA CGA CGC TG) and 3'O/P2-*ccd*<sub>O157</sub> (5'TCA GCA TTG AGC GCA ACC GTA AGG G). The PCR product was cloned into the TOPO-XL vector (Invitrogen). The resulting plasmid was then digested by HindIII and PstI (sites from the TOPO-XL vector). The fragment containing the operator/promoter region of the *ccd*<sub>O157</sub> operon was inserted into pJL207 that was opened by the same enzymes.

The Topo-*folAccd*<sub>O157</sub> plasmid was constructed by amplifying the chromosomal region comprising the *folA* gene and the *ccd*<sub>O157</sub> operon by PCR using *E. coli* O157:H7 chromosomal DNA (ATCC 700927) as a template and the following primers: 5'*folA* (5'CCC TCA TCC TAA TAA AGA GTG ACG) and 3' operon-*ccd*<sub>O157</sub> (5'CGA ACC GGC ATA AGG ATT TGG GTG AGG G). The PCR product was cloned into the TOPO-XL vector (Invitrogen).

To construct the pKT-*ccd*<sub>O157</sub> plasmid, the *ccd*<sub>O157</sub> operon was amplified by PCR using *E. coli* O157:H7 chromosomal DNA (ATCC 700927) as a template and the following primers: 5' operon-*ccd*<sub>O157</sub> (5'GAG ATT CTG GAG CGG CGG TAA TTT TG) and 3' operon-*ccd*<sub>O157</sub> (5'CGA ACC GGC ATA AGG ATT TGG GTG AGG G). The PCR product was cloned into the TOPO-XL vector (Invitrogen). The resulting plasmid was digested by EcoRI and then inserted into the pKT279 plasmid digested by EcoRI.






FIG. 2. The *ccd*<sub>O157</sub> system carries a toxin-antitoxin gene pair. SG22622/pKK223-3/pBAD-*ccd*<sub>B<sub>F</sub></sub> (filled squares), SG22622/pKK223-3/pBAD-*ccd*<sub>B<sub>O157</sub></sub> (filled triangles), SG22622/pKK-*ccd*<sub>A<sub>F</sub></sub>/pBAD-*ccd*<sub>B<sub>F</sub></sub> (open squares), and SG22622/pKK-*ccd*<sub>A<sub>O157</sub></sub>/pBAD-*ccd*<sub>B<sub>O157</sub></sub> (open triangles) were grown as described in Materials and Methods. After the addition of 1% arabinose, serial dilutions of the cultures were plated without arabinose and incubated overnight at 37°C. Values correspond to the means of results of three independent experiments.

ing the various constructs were grown exponentially, and *ccd*<sub>B<sub>F</sub></sub> and *ccd*<sub>B<sub>O157</sub></sub> transcription was induced at time zero by the addition of arabinose (Fig. 2). As observed for CcdB<sub>F</sub>, the production of CcdB<sub>O157</sub> results in a dramatic decrease in viable counts (by about 2 logs) after 10 min of induction. Viability is not affected in the presence of the cognate antitoxins (CcdA<sub>F</sub> and CcdA<sub>O157</sub>). This shows that the chromosomal *ccd*<sub>O157</sub> system encodes functional toxin and antitoxin proteins.

CcdB<sub>F</sub> kills plasmid-free segregant bacteria by poisoning the DNA gyrase, an essential topoisomerase II (8, 9). To determine whether the CcdB<sub>O157</sub> toxin also targets the DNA gyrase, the pBAD-*ccd*<sub>B<sub>O157</sub></sub> plasmid was transformed in a wild-type strain and in the isogenic strain carrying the CcdB<sub>F</sub>-resistant mutation *gyrA462*. Table 2 shows that the transformation efficiency for the pBAD-*ccd*<sub>B<sub>F</sub></sub> and pBAD-*ccd*<sub>B<sub>O157</sub></sub> plasmids in the wild-type strain is very low in the presence of 1% arabinose (<10<sup>-4</sup>), while it is comparable to that of the vector in the CcdB<sub>F</sub>-resistant strain. This shows that the DNA gyrase is the cellular target of CcdB<sub>O157</sub>. The key amino acids of the toxic active site of CcdB<sub>F</sub> have been previously identified as glycine100 and isoleucine101 (7). These two carboxy-terminal amino acids are conserved in CcdB<sub>O157</sub> (Fig. 1). We introduced the G100R or I101K mutations in CcdB<sub>O157</sub> and found that these mutations completely abolished the toxic activity of CcdB<sub>O157</sub> (data not shown), indicating that the active site of CcdB<sub>F</sub> is conserved in CcdB<sub>O157</sub>.

CcdA<sub>F</sub> is unstable and degraded by the Lon ATP-dependent protease. Figure 3 shows that the half-life of CcdA<sub>O157</sub> in a wild-type strain is comparable to that of CcdA<sub>F</sub> (~30 min) and

TABLE 2. Like CcdB<sub>F</sub>, the CcdB<sub>O157</sub> toxin targets the DNA gyrase

Strain	Efficiency of transformation <sup>a</sup>		
	pBAD33	pBAD- <i>ccd</i> <sub>B<sub>F</sub></sub>	pBAD- <i>ccd</i> <sub>B<sub>O157</sub></sub>
SG22622	1.1	6.7 × 10 <sup>-5</sup>	7.1 × 10 <sup>-5</sup>
SG22622 <i>gyrA462</i>	1.1	0.9	1

<sup>a</sup> Efficiency of transformation was calculated as the ratio of the number of transformants obtained on 1% arabinose plates to the number of transformants obtained on plates without arabinose. This experiment was performed at least in triplicate.




FIG. 3. Turnover of the CcdA<sub>F</sub> and CcdA<sub>O157</sub> antitoxins. SG22622 (wild type [wt]) and SG22623 (*Δlon* mutant) carrying either the pKK-*ccd*<sub>A<sub>F</sub></sub> or the pKK-*ccd*<sub>A<sub>O157</sub></sub> plasmid were grown to early log phase in LB at 37°C. Spectinomycin was added (100 μg/ml) to block protein synthesis. Culture samples were removed at the times indicated in the figure. Protein extraction and Western blot analysis were performed as described in Materials and Methods.

that both antitoxins are stabilized in a *Δlon* strain. Thus, CcdA<sub>O157</sub> is a Lon substrate.

The putative promoter region of the *ccd*<sub>O157</sub> system was cloned in a plasmid carrying a promoter-free *lacZ* gene (pJL-OP*ccd*<sub>O157</sub>). Table 3 shows that the promoter activity of the *ccd*<sub>O157</sub> system is fivefold weaker than that of the *ccd*<sub>F</sub> system. Regions of various lengths upstream and encompassing the 5' end of the *ccd*<sub>O157</sub> gene were tested, and their transcriptional activities are comparable to that of pJL-OP*ccd*<sub>O157</sub> (data not shown). We analyzed the autoregulation property of the *ccd*<sub>O157</sub> system and found that its expression is autoregulated by CcdA<sub>O157</sub> and CcdB<sub>O157</sub> proteins expressed in *trans* (Table 3). However, the autoregulation appears to be less efficient than that of the *ccd*<sub>F</sub> system (60% versus 90% of repression).

**Interactions between the *ccd*<sub>F</sub> and the *ccd*<sub>O157</sub> systems.** As the *ccd*<sub>O157</sub> and the *ccd*<sub>F</sub> systems coexist in *E. coli* O157:H7 isolates, we tested the cross-interactions between the components of the two systems. The ability of each antitoxin to counteract the toxic activity of its noncognate toxin was assayed in *E. coli* K-12. The expression of CcdB<sub>F</sub> and CcdB<sub>O157</sub> from the pBAD promoter results in a dramatic loss of viable counts. Basal expression of CcdA<sub>F</sub> from the pTac promoter in *trans* restores the viability of bacteria producing either CcdB<sub>O157</sub> or CcdB<sub>F</sub> (Fig. 4A). CcdA<sub>F</sub> is thus able to efficiently counteract the toxic activity of CcdB<sub>O157</sub> as its cognate toxin. However, expression of CcdA<sub>O157</sub> from the pTac promoter is unable to counteract CcdB<sub>F</sub> toxicity (Fig. 4B), even under overproduction conditions (in the presence of 1 mM IPTG) (data not shown).

TABLE 3. Promoter activity and autoregulation of the *ccd*<sub>F</sub> and *ccd*<sub>O157</sub> systems

OP:: <i>lacZ</i> fusion and corresponding operon	β-Galactosidase sp act (Miller units) (mean ± SD) <sup>a</sup>
pJL-OP <i>ccd</i> <sub>F</sub> pKT279	233 ± 11.83 (1)
pJL-OP <i>ccd</i> <sub>F</sub> pKT <i>ccd</i> <sub>F</sub>	22 ± 0.76 (0.09)
pJL-OP <i>ccd</i> <sub>O157</sub> pKT279	46 ± 0.97 (1)
pJL-OP <i>ccd</i> <sub>O157</sub> pKT <i>ccd</i> <sub>O157</sub>	17 ± 1.96 (0.37)

<sup>a</sup> β-galactosidase-specific activity of the OP::*lacZ* fusions in the presence of the corresponding operon in *trans* was normalized to that obtained with the control vector (shown in parentheses). Values correspond to the means of results for three experiments ± standard deviations.




FIG. 4. Interactions between the *ccd<sub>F</sub>* and the *ccd<sub>O157</sub>* systems. All strains were grown as described in Materials and Methods and treated as described in the legend for Fig. 2. Values correspond to the means of results of three independent experiments. Data shown were obtained from the same experiments and are represented in two panels for clarity. (A) Ability of CcdA<sub>F</sub> to counteract the toxic activity of CcdB<sub>O157</sub> SG22622/pKK223-3/pBAD-*ccdB<sub>F</sub>* (filled squares), SG22622/pKK223-3/pBAD-*ccdB<sub>O157</sub>* (filled triangles), SG22622/pKK-*ccdA<sub>F</sub>*/pBAD-*ccdB<sub>F</sub>* (open squares), and SG22622/pKK-*ccdA<sub>F</sub>*/pBAD-*ccdB<sub>O157</sub>* (open triangles). (B) Ability of CcdA<sub>O157</sub> to counteract the toxic activity of CcdB<sub>F</sub> SG22622/pKK223-3/pBAD-*ccdB<sub>F</sub>* (filled squares), SG22622/pKK223-3/pBAD-*ccdB<sub>O157</sub>* (filled triangles), SG22622/pKK-*ccdA<sub>O157</sub>*/pBAD-*ccdB<sub>F</sub>* (open squares), and SG22622/pKK-*ccdA<sub>O157</sub>*/pBAD-*ccdB<sub>O157</sub>* (open triangles).

We also tested the cross-interactions between the *ccd<sub>F</sub>* and the *ccd<sub>O157</sub>* systems in an *E. coli* strain carrying the *ccd<sub>O157</sub>* system in its chromosome. For that purpose, we could not use the O157:H7 strain, since it contains the pO157 plasmid carrying the *ccd<sub>F</sub>* system.

Evolutionary analyses have shown that O55:H7 strains are genetically closely related to O157:H7 strains (55, 56), although they are devoid of the pO157 plasmid. We screened O55:H7 isolates by PCR using two sets of primers, one specific to *ccd<sub>O157</sub>* and the other to *ccd<sub>F</sub>*, and confirmed that these strains carry *ccd<sub>O157</sub>* but lack *ccd<sub>F</sub>*. DNA sequencing revealed that the *ccd<sub>O157</sub>* system and its flanking regions are identical to those of O157:H7 (data not shown). A deletion of the *ccd<sub>O157</sub>* system was constructed in O55:H7 (O55:H7Δ*ccd<sub>O157</sub>*; see Materials and Methods).

Strains O55:H7 and O55:H7Δ*ccd<sub>O157</sub>* were transformed with the pBAD-*ccdB<sub>F</sub>* and the pBAD-*ccdB<sub>O157</sub>* plasmids. The viabilities of both strains were measured upon expression of the CcdB<sub>F</sub> and CcdB<sub>O157</sub> toxins. Figure 5 shows that the viability of O55:H7 is only slightly affected after 30 min of CcdB<sub>O157</sub> induction, while it is largely reduced by that of CcdB<sub>F</sub> (by about 2 logs). This shows that CcdA<sub>O157</sub> is produced from its natural location at a basal level that is sufficient to at least partially counteract the toxic activity of CcdB<sub>O157</sub> produced in *trans*. These results also confirm that CcdA<sub>O157</sub> is unable to counteract CcdB<sub>F</sub> produced in *trans*. On the other hand, the viability of O55:H7Δ*ccd<sub>O157</sub>* is drastically affected by the expression of both toxins (Fig. 5).




FIG. 5. Expression of the chromosomal *ccd<sub>O157</sub>* system in the O55:H7 *E. coli* strain. O55:H7 (squares) and O55:H7Δ*ccd<sub>O157</sub>* (triangles) containing pBAD-*ccdB<sub>F</sub>* (filled symbols) or pBAD-*ccdB<sub>O157</sub>* (open symbols) plasmids were grown as described in Materials and Methods. After addition of 0.25% arabinose, serial dilutions of the cultures were plated without arabinose and incubated ON at 37°C. Values correspond to the means of results for three independent experiments.

#### The *ccd<sub>F</sub>* system is able to mediate postsegregational killing in the presence of its chromosomal homolog *ccd<sub>O157</sub>*.

The chromosomal CcdA<sub>O157</sub> antitoxin, whether expressed ectopically or in its natural context, is not able to counteract the plasmidic CcdB<sub>F</sub> toxin. Therefore, we tested whether the *ccd<sub>F</sub>* system is able to mediate PSK in the presence of *ccd<sub>O157</sub>*. The O55:H7 strain that carries the *ccd<sub>O157</sub>* system in its chromosome was transformed with a conditionally replicating (thermosensitive) plasmid carrying the *ccd<sub>F</sub>* system (pMLO59-*ccd<sub>F</sub>*). Figure 6 shows that, after 180 min of culture at 42°C, the ability of O55:H7/pMLO59-*ccd<sub>F</sub>* to form colonies decreases in comparison with that of the control strain O55:H7/pMLO59. It only slightly increases during the next 135 min of the experiment, showing that the loss of pMLO59-*ccd<sub>F</sub>* mediates PSK. Thus, the *ccd<sub>F</sub>* system is functional for plasmid stabilization in an *E. coli* strain carrying the *ccd<sub>O157</sub>* system in its chromosome.

The *ccd<sub>O157</sub>* system is unable to mediate postsegregational killing. TA systems located in mobile genetic elements such as plasmids (e.g., *ccd<sub>F</sub>* and *relBE<sub>p307</sub>*), prophages (*relBE<sub>K-12</sub>*), and superintegrons (*higBA*) have been shown to be able to mediate PSK (17, 26, 27, 31). We tested the capacity of the chromosomal *ccd<sub>O157</sub>* system to mediate PSK using the same system




FIG. 6. The *ccd<sub>F</sub>* system mediates postsegregational killing in the O55:H7 *E. coli* strain. O55:H7 containing the pMLO59 vector (diamond) and its derivative carrying the *ccd<sub>F</sub>* system (squares) were grown as described in Materials and Methods. Exponential cultures in LB at 42°C were sampled at the times indicated in the figure. Surviving bacteria were scored as the number of CFU on LB plates at 30°C. Values correspond to the means of results for three independent experiments.




FIG. 7. The *ccd*<sub>O157</sub> system is unable to mediate postsegregational killing. O55:H7Δ*ccd*<sub>O157</sub> containing the pMLO59 vector (squares) and its derivatives carrying either the *ccd*<sub>F</sub> (triangles) or the *ccd*<sub>O157</sub> (diamonds) system were grown as described in Materials and Methods. Exponential cultures in LB at 42°C were sampled at the times indicated in the figure. Surviving bacteria were scored as the number of CFU on LB plates at 30°C. The values correspond to the means of results for three independent experiments.

described above. The O55:H7Δ*ccd*<sub>O157</sub> strain was transformed with the pMLO59-*ccd*<sub>F</sub> or the pMLO59-*ccd*<sub>O157</sub> plasmid. As expected, the loss of pMLO59-*ccd*<sub>F</sub> decreases the ability of the O55:H7Δ*ccd*<sub>O157</sub> strain to form colonies after 180 min of culture at 42°C. On the contrary, the loss of pMLO59-*ccd*<sub>O157</sub> does not affect the ability of the O55:H7Δ*ccd*<sub>O157</sub> strain to form colonies, even after 315 min of culture at 42°C (Fig. 7). As a control, we confirmed that the loss of both plasmids is comparable to that of the pMLO59 vector control (data not shown).

## DISCUSSION

TA systems are abundant in bacterial chromosomes and on mobile genetic elements such as plasmids, phages, and transposons (24). Computational analyses have grouped TA systems in families based on proteic sequence similarities and sequence profile analysis (5, 38). Members of a given family often coexist in the same bacteria. For instance, the *relBE* and the *vapBC* families are represented at 9 and 13 copies, respectively, in the chromosome of *Nitrosomonas europaea* (38). Situations in which plasmidic and chromosomal systems belonging to the same family are present in the same bacteria also exist. We have identified such a situation in the pathogenic *E. coli* O157:H7 strains, since they contain the plasmidic *ccd*<sub>F</sub> system located on an F-derivative virulence plasmid (pO157) and a chromosomal homolog of this plasmidic system, the *ccd*<sub>O157</sub> system.

The *ccd*<sub>O157</sub> chromosomal system is located between the *folA* and *apaH* genes in the chromosomes of 5 of the 17 sequenced *E. coli* strains. This heterogeneous distribution among the *E. coli* species suggests that the *ccd*<sub>O157</sub> system has been acquired by horizontal gene transfer. In the nine *E. coli* strains lacking it, a palindromic unit sequence is found in the *folA-apaH* intergenic region. Since palindromic units are specific targets for IS1397 insertion (18), an attractive hypothesis is that a composite transposon carrying an ancestral *ccd* system and other genes may have hopped in this specific site. DNA rearrangements as well as coevolution with the host genome may have occurred (e.g., GC content of *ccd*<sub>O157</sub> [49%] is close

to that of the chromosome of O157:H7 [50%]), leading to the actual *ccd*<sub>O157</sub> system. The *folA-apaH* intergenic region seems to be subject to insertion events, as a DNA segment containing an ORF coding for a hypothetical protein of unknown function and a sequence homologous to the C-terminal part of CcdA<sub>O157</sub> has been detected in 2 of the 17 *E. coli* strains (data not shown).

The chromosomal *relBE*<sub>K-12</sub> and *mazEF* systems of *E. coli* K-12 have been described as having a function in bacterial physiology (2, 14, 15, 16, 30, 47). However, each system seems to have a different function and/or a different function under different physiological conditions. *mazEF* is activated by numerous stresses and leads to cell death (22, 30), while activation of *relBE*<sub>K-12</sub> appears to be more specific to amino acid starvation and results in growth inhibition (15, 39). Although we do not yet have data involving the *ccd*<sub>O157</sub> system in the *E. coli* physiology, we have shown that it is expressed in its natural context, i.e., in O55:H7 *E. coli* isolates. We have shown that, unlike the *ccd*<sub>F</sub> system, the *ccd*<sub>O157</sub> system is unable to mediate postsegregational killing. The weak promoter activity of *ccd*<sub>O157</sub> in comparison with that of *ccd*<sub>F</sub>, rather than the toxicity of CcdB<sub>O157</sub>, is likely to be responsible for its inability to mediate PSK, since CcdB<sub>O157</sub>, when expressed ectopically, causes a reduction in viability comparable to that of CcdB<sub>F</sub>. A low transcriptional activity might reflect the adaptation of the *ccd*<sub>O157</sub> system to its chromosomal location. It is likely that chromosomal TA systems might have to adapt to their hosts to be maintained throughout evolution. Therefore, their regulation might be more complex and integrated than that of plasmidic TA systems. For instance, the alarmone ppGpp regulates negatively the transcription of the *mazEF* system in *E. coli* K-12 (2).

The *ccd*<sub>F</sub> and *ccd*<sub>O157</sub> systems naturally coexist in O157:H7 isolates, as these strains contain an F-derivative virulence plasmid (pO157), which is *ccd*<sub>F</sub><sup>+</sup>. Our cross-talk experiments have shown that the chromosomal CcdA<sub>O157</sub> antitoxin is not able to counteract the plasmidic CcdB<sub>F</sub> toxin, even under overproduction conditions, while the plasmidic CcdA<sub>F</sub> antitoxin is able to counteract the chromosomal CcdB<sub>O157</sub> toxin. Moreover, we have shown that the *ccd*<sub>F</sub> system is able to mediate PSK in an *E. coli* O55:H7 strain harboring the *ccd*<sub>O157</sub> system in its chromosome, showing that the *ccd*<sub>F</sub> system is functional in the presence of its chromosomal counterpart. A similar observation was made for the *pem* (*parD*) TA system located on the R100 (R1) plasmid of *E. coli*. The two antitoxins of its chromosomal homologs (*mazEF* and *chpB*) are unable to counteract the toxicity of the plasmid-encoded toxin (PemK) (35), explaining how the *pem* system is able to stabilize plasmids in *E. coli* (51). However, mutants of the antitoxins of the *mazEF* and *chpB* systems are able to counteract PemK toxicity (44, 45), most likely due to an overproduction of the chromosomal antitoxins. In our experiments, overproduction of the chromosomal CcdA<sub>O157</sub> antitoxin does not counteract the plasmidic CcdB<sub>F</sub> toxin (data not shown). Thus, only plasmidic TA systems able to evade cross-interaction seem to coexist with their chromosomal homologs. This raises the hypothesis that chromosomal TA systems might serve as “exclusion” systems to protect bacteria from being loaded with an excess of exogenous DNA carrying identical TA systems (plasmids, transposons, phages) (18a, 41), a function somewhat reminiscent of that of

restriction-modification (RM) systems (for a review on RM, see reference 32). RM and TA systems share several properties. They are located in plasmids as well as in chromosomes, they seem to move from one location to another through horizontal gene transfer, and they are able to mediate PSK when cloned on an unstable plasmid. Further experiments are needed to test whether certain TA systems might promote the "exclusion" of incoming homologous TA systems.

#### ACKNOWLEDGMENTS

We are grateful to Susan Gottesman for helpful suggestions and critical reading of the manuscript. We thank Christophe Beloin and Jean-Marc Ghigo for providing us with the pKOBEG plasmid. We also thank Beth Whittam from the STEC Center (Michigan State University) for kindly providing us with the O55:H7 strain.

This work was supported by the FNRS (FRSM 3.4510.02), the Institut Pasteur, the CNRS (URA 2171), MENESR, and the European Union (STREP LSH-2004-2.1.2-2). L.V.M. is Chercheur Qualifié at the FNRS. M.W. is supported by the FRIA.

#### REFERENCES

- Afif, H., N. Allali, M. Couturier, and L. Van Melderen. 2001. The ratio between CcdA and CcdB modulates the transcriptional repression of the ccd poison-antidote system. *Mol. Microbiol.* **41**:73–82.
- Aizenman, E., H. Engelberg-Kulka, and G. Glaser. 1996. An *Escherichia coli* chromosomal "addiction module" regulated by guanosine [corrected] 3',5'-bispyrophosphate: a model for programmed bacterial cell death. *Proc. Natl. Acad. Sci. USA* **93**:6059–6063.
- Allali, N., H. Afif, M. Couturier, and L. Van Melderen. 2002. The highly conserved TldD and TldE proteins of *Escherichia coli* are involved in microcin B17 processing and in CcdA degradation. *J. Bacteriol.* **184**:3224–3231.
- Amitai, S., Y. Yassin, and H. Engelberg-Kulka. 2004. MazF-mediated cell death in *Escherichia coli*: a point of no return. *J. Bacteriol.* **186**:8295–8300.
- Anantharaman, V., and L. Aravind. 2003. New connections in the prokaryotic toxin-antitoxin network: relationship with the eukaryotic nonsense-mediated RNA decay system. *Genome Biol.* **4**:R81.
- Arcus, V. L., P. B. Rainey, and S. J. Turner. 2005. The PIN-domain toxin-antitoxin array in mycobacteria. *Trends Microbiol.* **13**:360–365.
- Bahassi, E. M., M. A. Salmon, L. Van Melderen, P. Bernard, and M. Couturier. 1995. F plasmid CcdB killer protein: ccdB gene mutants coding for non-cytotoxic proteins which retain their regulatory functions. *Mol. Microbiol.* **15**:1031–1037.
- Bernard, P., and M. Couturier. 1992. Cell killing by the F plasmid CcdB protein involves poisoning of DNA-topoisomerase II complexes. *J. Mol. Biol.* **226**:735–745.
- Bernard, P., K. E. Kezdy, L. Van Melderen, J. Steyaert, L. Wyns, M. L. Pato, P. N. Higgins, and M. Couturier. 1993. The F plasmid CcdB protein induces efficient ATP-dependent DNA cleavage by gyrase. *J. Mol. Biol.* **234**:534–541.
- Brosius, J., and A. Holy. 1984. Regulation of ribosomal RNA promoters with a synthetic lac operator. *Proc. Natl. Acad. Sci. USA* **81**:6929–6933.
- Buts, L., J. Lah, M. H. Dao-Thi, L. Wyns, and R. Loris. 2005. Toxin-antitoxin modules as bacterial metabolic stress managers. *Trends Biochem. Sci.* **30**:672–679.
- Chaveroche, M. K., J. M. Ghigo, and C. d'Enfert. 2000. A rapid method for efficient gene replacement in the filamentous fungus *Aspergillus nidulans*. *Nucleic Acids Res.* **28**:E97.
- Cherepanov, P. P., and W. Wackernagel. 1995. Gene disruption in *Escherichia coli*: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. *Gene* **158**:9–14.
- Christensen, S. K., and K. Gerdes. 2004. Delayed-relaxed response explained by hyperactivation of RelE. *Mol. Microbiol.* **53**:587–597.
- Christensen, S. K., M. Mikkelsen, K. Pedersen, and K. Gerdes. 2001. RelE, a global inhibitor of translation, is activated during nutritional stress. *Proc. Natl. Acad. Sci. USA* **98**:14328–14333.
- Christensen, S. K., K. Pedersen, F. G. Hansen, and K. Gerdes. 2003. Toxin-antitoxin loci as stress-response-elements: ChpAK/MazF and ChpBK cleave translated RNAs and are counteracted by tmRNA. *J. Mol. Biol.* **332**:809–819.
- Christensen-Dalsgaard, M., and K. Gerdes. 2006. Two higBA loci in the *Vibrio cholerae* superintegron encode mRNA cleaving enzymes and can stabilize plasmids. *Mol. Microbiol.* **62**:397–411.
- Clément, J.-M., C. Wilde, S. Bachelier, P. Lambert, and M. Hofnung. 1999. IS1397 is active for transposition into the chromosome of *Escherichia coli* K-12 and inserts specifically into palindromic units of bacterial interspersed mosaic elements. *J. Bacteriol.* **181**:6929–6936.
- Cooper, T. F., and J. A. Heinemann. 2000. Postsegregational killing does not increase plasmid stability but acts to mediate the exclusion of competing plasmids. *Proc. Natl. Acad. Sci. USA* **97**:12643–12648.
- Couturier, M., E. M. Bahassi, and L. Van Melderen. 1998. Bacterial death by DNA gyrase poisoning. *Trends Microbiol.* **6**:269–275.
- Critchlow, S. E., M. H. O'Dea, A. J. Howells, M. Couturier, M. Gellert, and A. Maxwell. 1997. The interaction of the F plasmid killer protein, CcdB, with DNA gyrase: induction of DNA cleavage and blocking of transcription. *J. Mol. Biol.* **273**:826–839.
- Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**:6640–6645.
- Engelberg-Kulka, H., R. Hazan, and S. Amitai. 2005. mazEF: a chromosomal toxin-antitoxin module that triggers programmed cell death in bacteria. *J. Cell Sci.* **118**:4327–4332.
- Gerdes, K. 2000. Toxin-antitoxin modules may regulate synthesis of macromolecules during nutritional stress. *J. Bacteriol.* **182**:561–572.
- Gerdes, K., S. K. Christensen, and A. Lobner-Olesen. 2005. Prokaryotic toxin-antitoxin stress response loci. *Nat. Rev. Microbiol.* **3**:371–382.
- Glandsdorff, N. 1965. Topography of cotransducible arginine mutations in *Escherichia coli* K-12. *Genetics* **51**:167–179.
- Goffredsen, M., and K. Gerdes. 1998. The *Escherichia coli* relBE genes belong to a new toxin-antitoxin gene family. *Mol. Microbiol.* **29**:1065–1076.
- Gronlund, H., and K. Gerdes. 1999. Toxin-antitoxin systems homologous with relBE of *Escherichia coli* plasmid P307 are ubiquitous in prokaryotes. *J. Mol. Biol.* **285**:1401–1415.
- Guzman, L.-M., D. Belin, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose P<sub>BAD</sub> promoter. *J. Bacteriol.* **177**:4121–4130.
- Hayes, F. 2003. Toxins-antitoxins: plasmid maintenance, programmed cell death, and cell cycle arrest. *Science* **301**:1496–1499.
- Hazan, R., B. Sat, and H. Engelberg-Kulka. 2004. *Escherichia coli* mazEF-mediated cell death is triggered by various stressful conditions. *J. Bacteriol.* **186**:3663–3669.
- Jaffé, A., T. Ogura, and S. Hiraga. 1985. Effects of the ccd function of the F plasmid on bacterial growth. *J. Bacteriol.* **163**:841–849.
- Kobayashi, I. 2001. Behavior of restriction-modification systems as selfish mobile elements and their impact on genome evolution. *Nucleic Acids Res.* **29**:3742–3756.
- Light, J., and S. Molin. 1982. The sites of action of the two copy number control functions of plasmid R1. *Mol. Gen. Genet.* **187**:486–493.
- Maki, S., S. Takiguchi, T. Miki, and T. Horiuchi. 1992. Modulation of DNA supercoiling activity of *Escherichia coli* DNA gyrase by F plasmid proteins. Antagonistic actions of LetA (CcdA) and LetD (CcdB) proteins. *J. Biol. Chem.* **267**:12244–12251.
- Masuda, Y., K. Miyakawa, Y. Nishimura, and E. Ohtsubo. 1993. *chpA* and *chpB*, *Escherichia coli* chromosomal homologs of the *pem* locus responsible for stable maintenance of plasmid R100. *J. Bacteriol.* **175**:6850–6856.
- Metzger, S., I. B. Dror, E. Aizenman, G. Schreiber, M. Toone, J. D. Friesen, M. Cashel, and G. Glaser. 1988. The nucleotide sequence and characterization of the relA gene of *Escherichia coli*. *J. Biol. Chem.* **263**:15699–15704.
- Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Pandey, D. P., and K. Gerdes. 2005. Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. *Nucleic Acids Res.* **33**:966–976.
- Pedersen, K., S. K. Christensen, and K. Gerdes. 2002. Rapid induction and reversal of a bacteriostatic condition by controlled expression of toxins and antitoxins. *Mol. Microbiol.* **45**:501–510.
- Perna, N. T., J. D. Glasner, V. Burland, and G. Plunkett III. 2002. The genomes of *Escherichia coli* K-12 and pathogenic *E. coli*, p. 3–53. In M. S. Donnenberg, (ed.), *Escherichia coli*: virulence mechanisms of a versatile pathogen. Academic Press, San Diego, CA.
- Rowe-Magnus, D. A., A. M. Guerout, L. Biskri, P. Bouige, and D. Mazel. 2003. Comparative analysis of superintegrons: engineering extensive genetic diversity in the Vibrionaceae. *Genome Res.* **13**:428–442.
- Salmon, M. A., L. Van Melderen, P. Bernard, and M. Couturier. 1994. The antidote and autoregulatory functions of the F plasmid CcdA protein: a genetic and biochemical survey. *Mol. Gen. Genet.* **244**:530–538.
- Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Santos-Sierra, S., R. Giraldo, and R. Diaz-Orejas. 1997. Functional interactions between homologous conditional killer systems of plasmid and chromosomal origin. *FEMS Microbiol. Lett.* **152**:51–56.
- Santos Sierra, S., R. Giraldo, and R. Diaz Orejas. 1998. Functional interactions between *chpB* and *parD*, two homologous conditional killer systems found in the *Escherichia coli* chromosome and in plasmid R1. *FEMS Microbiol. Lett.* **168**:51–58.
- Sat, B., R. Hazan, T. Fisher, H. Khaner, G. Glaser, and H. Engelberg-Kulka. 2001. Programmed cell death in *Escherichia coli*: some antibiotics can trigger mazEF lethality. *J. Bacteriol.* **183**:2041–2045.
- Sat, B., M. Reches, and H. Engelberg-Kulka. 2003. The *Escherichia coli* mazEF suicide module mediates thymineless death. *J. Bacteriol.* **185**:1803–1807.

48. **Talmadge, K., and W. Gilberg.** 1980. Construction of plasmid vectors with unique PstI cloning sites in a signal sequence coding region. *Gene* **12**:235–241.
49. **Tam, J. E., and B. C. Kline.** 1989. The F plasmid *ccd* autorepressor is a complex of CcdA and CcdB proteins. *Mol. Gen. Genet.* **219**:26–32.
50. **Tsuchimoto, S., Y. Nishimura, and E. Ohtsubo.** 1992. The stable maintenance system *pem* of plasmid R100: degradation of PemI protein may allow PemK protein to inhibit cell growth. *J. Bacteriol.* **174**:4205–4211.
51. **Tsuchimoto, S., H. Ohtsubo, and E. Ohtsubo.** 1988. Two genes, *pemK* and *pemI*, responsible for stable maintenance of resistance plasmid R100. *J. Bacteriol.* **170**:1461–1466.
52. **Van Melderren, L.** 2002. Molecular interactions of the CcdB poison with its bacterial target, the DNA gyrase. *Int. J. Med. Microbiol.* **291**:537–544.
53. **Van Melderren, L., P. Bernard, and M. Couturier.** 1994. Lon-dependent proteolysis of CcdA is the key control for activation of CcdB in plasmid-free segregant bacteria. *Mol. Microbiol.* **11**:1151–1157.
54. **Van Melderren, L., M. H. Thi, P. Lecchi, S. Gottesman, M. Couturier, and M. R. Maurizi.** 1996. ATP-dependent degradation of CcdA by Lon protease. Effects of secondary structure and heterologous subunit interactions. *J. Biol. Chem.* **271**:27730–27738.
55. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Ørskov, I. Ørskov, and R. A. Wilson.** 1993. Clonal relationships among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**:1619–1629.
56. **Wick, L. M., W. Qi, D. W. Lacher, and T. S. Whittam.** 2005. Evolution of genomic content in the stepwise emergence of *Escherichia coli* O157:H7. *J. Bacteriol.* **187**:1783–1791.
57. **Zhou, Y., and S. Gottesman.** 1998. Regulation of proteolysis of the stationary-phase sigma factor RpoS. *J. Bacteriol.* **180**:1154–1158.