

Genome Sequence of *Staphylococcus aureus* Strain Newman and Comparative Analysis of Staphylococcal Genomes: Polymorphism and Evolution of Two Major Pathogenicity Islands[∇]

Tadashi Baba,^{1*} Taeok Bae,² Olaf Schneewind,³ Fumihiko Takeuchi,⁴ and Keiichi Hiramatsu¹

Department of Microbiology and Infection Control Science, Juntendo University, Tokyo 113-8421, Japan¹; Microbiology and Immunology, Indiana University School of Medicine, Northwest 3400 Broadway, Med. Ed. 3056, Gary, Indiana 46408²; Department of Microbiology, The University of Chicago, 920 E. 58th Street, Chicago, Illinois 60637³; and Wellcome Trust Sanger Institute, Hinxton, Cambridge CB10 1SA, United Kingdom⁴

Received 22 June 2007/Accepted 9 October 2007

Strains of *Staphylococcus aureus*, an important human pathogen, display up to 20% variability in their genome sequence, and most sequence information is available for human clinical isolates that have not been subjected to genetic analysis of virulence attributes. *S. aureus* strain Newman, which was also isolated from a human infection, displays robust virulence properties in animal models of disease and has already been extensively analyzed for its molecular traits of staphylococcal pathogenesis. We report here the complete genome sequence of *S. aureus* Newman, which carries four integrated prophages, as well as two large pathogenicity islands. In agreement with the view that *S. aureus* Newman prophages contribute important properties to pathogenesis, fewer virulence factors are found outside of the prophages than for the highly virulent strain MW2. The absence of drug resistance genes reflects the general antibiotic-susceptible phenotype of *S. aureus* Newman. Phylogenetic analyses reveal clonal relationships between the staphylococcal strains Newman, COL, NCTC8325, and USA300 and a greater evolutionary distance to strains MRSA252, MW2, MSSA476, N315, Mu50, JH1, JH9, and RF122. However, polymorphism analysis of two large pathogenicity islands distributed among these strains shows that the two islands were acquired independently from the evolutionary pathway of the chromosomal backbones of staphylococcal genomes. Prophages and pathogenicity islands play central roles in *S. aureus* virulence and evolution.

Staphylococcus aureus is a human pathogen that causes both nosocomial and community-acquired infections. The emergence of strains resistant to many antibiotics (methicillin-resistant *S. aureus* [MRSA]) and of highly virulent community-acquired MRSA that can cause fatal infections such as necrotizing pneumonia is of considerable concern even in countries with well-developed health surveillance systems (24, 30). In order to study mechanisms of staphylococcal antibiotic resistance and virulence, whole genome sequences of several different *S. aureus* strains have been determined. MRSA strains N315 and Mu50 were the first staphylococcal genomes to be sequenced (18), which were followed by nine additional strains (1, 5, 9, 10, 13, 14). All staphylococcal genomes are approximately 2.8 Mbp in size with a relatively low G+C content. Comparative analysis revealed that most regions of the staphylococcal genome are well conserved, whereas several large sequence blocks display high variability. *S. aureus* strains likely acquired these genomic islands horizontally and, at least initially, their integration into the genome must have required dedicated DNA recombination (integrase) genes. Furthermore, variable blocks of genome sequence frequently carry virulence and antibiotic resistance determinants that aid in the development of staphylococcal diseases. Variable regions can

be classified as prophages, pathogenicity islands, or staphylococcal cassette chromosomes. The overall combination of variable sequence elements and the encoded spectrum of virulence properties varies from strain to strain and appears to be reflective of the overall large spectrum of clinical disease manifestations in humans (1, 2).

S. aureus strain Newman was isolated in 1952 from a human infection (6) and has been used extensively in animal models of staphylococcal disease due to its robust virulence phenotypes. Thirty genes that are required for staphylococcal pathogenesis were identified in *S. aureus* Newman after a screen of 1,736 *bursa aurealis* mutants with transposon insertions in different genes. Both well-characterized virulence genes and genes with unknown function were shown to be involved in the pathogenesis of staphylococcal infections (4). Additional benefits of systematic insertional mutagenesis are the identification of genes that are dispensable for staphylococcal growth under laboratory conditions. Subsequent work identified four prophages, ϕ NM1 to ϕ NM4, in the genome of strain Newman genome. Indeed, six paralogous groups of virulence determinants that were identified via *bursa aurealis* mutagenesis are encoded by these prophages (3). *S. aureus* Newman variants that lacked either ϕ NM3 or ϕ NM1, ϕ NM2, and ϕ NM4, or all four prophages (ϕ NM1 to ϕ NM4) displayed dramatic reductions in their ability to form organ specific abscesses after intravenous infection of mice, suggesting that the prophages ϕ NM1 to ϕ NM4 play important roles during the pathogenesis of staphylococcal infections.

To further unravel molecular mechanisms of the physiology

* Corresponding author. Mailing address: Department of Microbiology and Infection Control Science, Juntendo University, Tokyo 113-8421, Japan. Phone: (81) 3-5802-1041. Fax: (81) 3-5684-7830. E-mail: tbaba@med.juntendo.ac.jp.

[∇] Published ahead of print on 19 October 2007.

and pathogenesis of disease caused by *S. aureus* Newman, all of its genes must be known. This experimental goal was achieved, as we report here the complete genome sequence of *S. aureus* Newman. In contrast to other staphylococcal strains, which carry some virulence genes in mobile pathogenicity islands or genomic islets, virulence determinants of *S. aureus* Newman strain are conspicuous in prophages (2). Strain Newman carries a similar combination of major pathogenicity islands, ν Sa α and ν Sa β , as *S. aureus* strains COL, NCTC8325, and USA300. In contrast to hospital-acquired MRSA, *S. aureus* Newman harbors only a small number of insertion sequences (IS) and lacks known antibiotic resistance determinants.

MATERIALS AND METHODS

Shotgun sequencing and contig assembly. The whole-genome sequence was determined as described previously (1, 18). Shotgun sequencing was carried out by Hitachi High-Tech Fielding Co. (Tokyo, Japan) and Takara Bio, Inc. (Otsu, Japan). Sequences were also assembled as described previously. We have entered the whole-genome sequence of Newman in the DNA Database of Japan, with accession number AP009351.

Determination of open reading frames and structural RNA. Determination of open reading frames, structural RNAs and annotations were performed as described previously (1, 18). Briefly, open reading frames were initially extracted with Genome GAMBLER program (Xanagen, Kawasaki, Japan) based on GLIMMER and rbsfinder software. The predicted open reading frames were then individually reviewed with GAMBLER. We searched a nonredundant protein database with the determined open reading frames using BLAST software for annotation. tRNA and tmRNA genes were identified by tRNAscan-SE (22) and with web-based software (<http://www.indiana.edu/~tmrna/>), respectively. Illustration of G+C contents on a genome map was drawn by using Insilico molecular cloning software (Insilico Biology, Yokohama, Japan). This software was also used for comparative genome analysis among strains.

***S. aureus* genomes used for comparative analysis.** Sequences of *S. aureus* strains MW2 and N315 (accession numbers are BA000033 and BA000018, respectively) were used for whole-genome comparative analysis with strain Newman. The genome sequences of strains Mu50 (BA000017) (18), NCTC8325 (CP000253) (10), COL (CP000046) (9), MRSA252 (BX571856) (14), MSSA476 (BX571857) (14), RF122 (AJ938182) (13), USA300 (CP000255) (5), JH1 (CP000736) (A. Copeland et al., unpublished data), and JH9 (CP000703) (Copeland et al., unpublished) were also used for comparison among pathogenicity islands ν Sa α and ν Sa β .

Calculation of phylogenetic relationship among strains. From genomes sequences of 10 different *S. aureus* strains nucleotide sequences of carbamate kinase, shikimate dehydrogenase, glycerol kinase, guanylate kinase, phosphate acetyltransferase, triosephosphate isomerase and acetyl coenzyme A acetyltransferase were used for multilocus sequence typing (MLST) analysis (7). Nucleotide sequences were combined and then aligned with CLUSTAL X (26). For phylogenetic tree display, results from the CLUSTAL X calculation were visualized with TreeView (25).

RESULTS

Overview and genomic islands of the *S. aureus* Newman genome. The whole-genome sequence of *S. aureus* Newman was determined as described previously (1, 17). Briefly, shotgun cloning of *S. aureus* strain Newman genomic DNA allowed for DNA sequencing of random fragments and data assembly into contiguous genome segments (contigs). Sequence gaps between assembled contigs were read by series of PCRs, using primers based on predetermined sequences. We encountered assembly difficulties for genome segments surrounding prophages due to the high degree of sequence homology between prophages, in particular for ϕ NM1 and ϕ NM2 (a nearly 40-kb identical sequence). This obstacle was overcome by deriving DNA sequences from isolated ϕ NM1 or ϕ NM2 phage particles that had been isolated distinctively by mitomycin treat-

ment of strain Newman (3). Indeed, sequences of ϕ NM1 and ϕ NM2 showed clear differences only in their attachment core sequences and integrases that recognize the attachment sequence and had little uniqueness in other domains. The clear differences in the attachment sites and the integrase sequences strongly supported that ϕ NM1 and ϕ NM2 were distinct prophages from each other, and this was confirmed by identifying unique chromosomal locus where insertion of each phage occurred in contiguous sequences upon shotgun assembly along with the different integrase sequences of ϕ NM1 and ϕ NM2. In conjunction with sequences for phage integration sites, this eventually permitted assembly of a circular chromosomal DNA sequence. The length of the *S. aureus* Newman chromosome is 2,878,897 bp, and it encodes 2,614 open reading frames (Table 1). Plasmids sequences were not identified in our experiments involving *S. aureus* Newman.

We compared the Newman genome with other *S. aureus* chromosomes thus far sequenced (Table 1). Although the strains are categorized as a single species, *S. aureus*, the chromosomes from different strains had unique features. The G+C contents did not vary drastically; however, the lengths of chromosomes differed by more than 5% when the longest chromosome from strain JH9 was compared to the shortest RF122, with lengths ranging 2.74 to 2.91 Mbp. Chromosomes were classified into two groups according to the number of rRNA genes. Importantly, the ribosomal gene numbers did not correlate with genomic island subtypes, as seen in SCCmec and ν Sa islands (see below). A number of genomic islands also had large varieties among different strains. At least one prophage was found in each genome, and Newman has as many as four, which is maximum number thus far identified in a genome. There was also wide variability among strains possessing other classes of islands and IS, indicating that these genetic elements play key roles in conferring chromosomal diversity to *S. aureus* strains.

Figure 1 displays a circular map of *S. aureus* Newman chromosomal DNA. Genomic islands including prophages and pathogenicity islands are shown as green lines. The integration of four different prophages is unique to strain Newman; for example, *S. aureus* MW2 and N315 harbor either two or only one prophage, respectively (Fig. 2). Two IS1181 insertion sequences and ten remnants of IS were found in strain Newman. One of the IS1181 was inserted into the corresponding site of IS1181-6 in strain N315 (17). Major transposons such as Tn554 were not found in the *S. aureus* Newman chromosome, in contrast to strain N315 with many IS, as well as five copies of Tn554 (Table 1). SCCmec (16) was not found in the chromosome of strain Newman, a finding in agreement with the observation that this strain is susceptible to methicillin and other β -lactam antibiotics (data not shown).

Open reading frames are indicated in second (found in the forward strand) and third (found in the reverse strand) circles as either red (virulence determinants) or blue (others) bars. The open reading frame orientation showed clear contrast according to movement of replication fork, which is in agreement with a tendency seen in other strains previously sequenced (1, 18). The G+C skew value distribution was also asymmetrical across the axle of replication origin termination sites. G+C contents tended to be relatively high not only in the loci where structural RNA genes were concentrated but also

TABLE 1. Overview of *S. aureus* strain Newman genome in comparison with other strains

Parameter	Strain											
	Newman	MW2	N315	Mu50	NCTC8325	MRSA252	MSSA476	COL	RF122	USA300	JH1	JH9
Length of sequence (bp)	2,878,897	2,826,402	2,814,816	2,878,529	2,821,361	2,902,619	2,799,802	2,809,422	2,742,531	2,872,769	2,906,507	2,906,700
G+C content (%)	32.9	32.8	32.8	32.9	32.9	32.8	32.9	32.8	32.8	32.8	33.0	33.0
Open reading frames ^a	2,614	2,632	2,593	2,714	2,892	2,744	2,619	2,673	2,589	2,560	2,747	2,697
No. of protein coding regions	83.4	83.5	83.4	83.8	85.1	90.0	85.6	82.9	83.9	82.1	83.7	83.4
% Coding												
rRNAs												
16S	5	6	5	5	5	5	6	6	5	5	6	6
23S	5	6	5	5	5	5	6	6	5	5	6	6
5S	6	7	6	6	6	6	7	7	6	6	7	7
tRNAs	56	61	62	60	61	60	60	53	60	53	60	59
tmRNA	1	1	1	1	1	1	1	1	1	1	1	1
No. of insertion sequences ^b												
IS1/81	2	0	8	10	2	0	0	3	0	2	8	8
IS431	0	1	2	2	0	2	1	1	2	2	1	1
IS1272	0	1	0	0	1	9	0	0	0	1	1	1
Others (including remnant)	10	4	10	11	5	19	4	6	9	7	7	7
Transposon Tn554	0	0	5	2	0	3	0	0	0	0	2	2
Genomic islands												
Prophages	4	2	1	2	3	2	2	1	1	2	4	4
SCC _{mec} type	None	IV	II	II	None	II	None ^c	I	None	IV	II	II
vSaα type ^d	I	II	I	I	I	III	II	I	IV	I	I	I
vSaβ type ^d	II	II	I	I	II	III	II	II	II	II	I	I
Other islands ^e	2	3	2	3	2	2	2	3	3	4	1	1

^a Based on published records and therefore the criteria for open reading frame adoption differs from one genome to another.

^b Based on published records or BLAST searches using IS Finder (<http://www-is.biotoul.fr/is.html>), where genetic elements that e values were 0.0 to known IS were adopted.

^c SCC_{Isd} is present instead of SCC_{mec}.

^d See Fig. 3 and 4.

^e That is, islands without pathogenic features are also included.




FIG. 1. Circular display of *S. aureus* strain Newman chromosome. Green bars inside the first (outer) scale circle indicate the positions of genomic islands. The second circle shows open reading frames oriented in the forward direction, whereas the third circle indicates those oriented in the reverse direction. The fourth and fifth circles show genes for rRNAs and transfer RNAs, respectively. The sixth circle represents G+C content values. Purple indicates domains with G+C contents higher than 50%. The seventh circle shows G+C skew, in which purple indicates positive values.

where genomic islands were located, a finding in agreement with the general hypothesis that horizontal gene transfer causes acquisition of the genomic islands.

Genomic island *vSa* and the nomenclature. The term *vSa* refers to nonphage and non-SCC genomic islands that are exclusively present in *S. aureus*, often (but not always) encode for virulence determinants, are inserted at a specific locus in chromosome, and are associated with either intact or remnant DNA recombinase (1). The feature of a *vSa* genomic island possessing DNA recombinase also supports the hypothesis that staphylococcal pathogenicity islands are acquired by horizontal gene transfer. Due to the allelicity of the islands among dif-

ferent strains, the designation of the islands based on their structures or genetic content may create multiple island names that differ from strain to strain, regardless of the fact that they are inserted in identical loci of *S. aureus* chromosomes. We therefore propose that the term *vSa* does not designate an island with specific structure or a particular genetic content in a strain, but rather a locus where the island is inserted in the *S. aureus* chromosome. Hence, a term indicating a specific pathogenicity island such as “SaPI” should be used as well as “*vSa*.”

Among all of the *S. aureus* strains sequenced thus far, two major pathogenicity islands *vSa* α and *vSa* β are present in strain Newman, and these islands appear to be allelic among




FIG. 2. Homology alignment of whole genomes between strains Newman and MW2 (A) and strains Newman and N315 (B). Gaps are highlighted with circles in green (unique elements to Newman), purple (unique to MW2 or N315), or yellow (the same elements but with differences). The $\nu\text{Sa}\beta$ in panel A and $\nu\text{Sa}\alpha$ in panel B are homologous each other, but only the pathogenicity island names are indicated in parentheses.

different strains (2) (see also Fig. 3 and 4). The previously identified $\nu\text{Sa}\gamma$, which has been shown to be present in all sequenced *S. aureus* strains, encoding exfoliative toxin and exotoxins (9), was also found in strain Newman. A fourth class

of νSa island, $\nu\text{Sa}4$, was located downstream of $\phi\text{NM}3$. This island was inserted at the corresponding site into the chromosome of strain N315, where a $\nu\text{Sa}4$ island coding for 20 open reading frames, including three superantigen genes, is present in N315. Unlike the $\nu\text{Sa}4$ in N315, the one found in strain Newman lacks known virulence determinants and encodes for only integrase and three functionally unknown proteins. Therefore, the $\nu\text{Sa}4$ island in strain Newman probably lacks the features of a pathogenicity island and is structurally similar to the $\nu\text{Sa}4$ island in strain MW2. The difference in $\nu\text{Sa}4$ in strain Newman or strain MW2 compared to that in N315 also indicates that $\nu\text{Sa}4$ shows polymorphism among strains as $\nu\text{Sa}\alpha$ and $\nu\text{Sa}\beta$.

Comparative analysis of staphylococcal genomes. Figure 2 shows a comparison of the *S. aureus* Newman genome with those of strains MW2 and N315. Depiction of homologous regions as dots or lines revealed that the overall size of the chromosome and the order of its genes are conserved among all three strains. Major homology gaps are caused by the insertion of four prophages into the genome of strain Newman. Similar to the insertion of $\phi\text{NM}3$ in strain Newman, *S. aureus* MW2 and N315 also carry prophages in the *hlyB* gene (encoding beta-hemolysin); however, sequence homology between these phages is low, and differences are readily detectable in the plots in Fig. 2. *SCCmec* elements are found in strains N315 and MW2; however, this element is absent in *S. aureus* Newman. The observed gap size is larger in *S. aureus* N315 than in strain MW2, since *SCCmec* in strain N315 carries not only β -lactam resistance but also determinants for resistance to other antibiotics, whereas MW2 possesses only the β -lactam resistance gene *mecA* (23). Additional sequence gaps between staphylococcal strains are mainly due to differences in pathogenicity islands. The gap at 0.45 Mbp (Fig. 2A) is due to differences in pathogenicity island $\nu\text{Sa}\alpha$ between Newman and MW2, whereas the gap at 1.9 Mbp in Fig. 2B reflects the differences in the $\nu\text{Sa}\beta$ pathogenicity island between strains Newman and N315. These data suggest that *S. aureus* Newman $\nu\text{Sa}\alpha$ is similar to $\nu\text{Sa}\alpha$ in N315 but not to $\nu\text{Sa}\alpha$ in MW2. Further, *S. aureus* Newman $\nu\text{Sa}\beta$ is similar to $\nu\text{Sa}\beta$ in MW2 but not to $\nu\text{Sa}\beta$ in N315.

In *S. aureus* Newman, prophage $\phi\text{NM}3$ is inserted into *hlyB*, and similar insertions have been observed for *hlyB*-converting phages of *S. aureus* strains N315, MW2, Mu50, NCTC8325, MSSA476, MRSA252, USA300, JH1, and JH9. The genetic content of *hlyB*-converting phages is, however, variable, especially with regard to virulence determinants. For example, *hlyB*-converting phages of *S. aureus* N315 and Newman carry genes for staphylococcal complement inhibitor and chemotaxis inhibitory protein (31); the latter is absent in the *S. aureus* MW2 prophage. In contrast, the MW2 *hlyB*-converting phage carries genes for enterotoxins K2 and Q that are not found in phages of strains N315 and Newman. Despite these differences, the integrase gene of $\phi\text{NM}3$ and those of related phages are virtually identical, suggesting that all *hlyB*-converting phages evolved from a common ancestor. Other prophages of strain Newman, $\phi\text{NM}1$, $\phi\text{NM}2$, and $\phi\text{NM}4$, are absent in *S. aureus* MW2 and N315 (Fig. 2). However, $\phi\text{NM}1$ is inserted at the same integration site as $\phi 11$ in *S. aureus* NCTC8325 (15), and $\phi 11$ and $\phi\text{NM}1$ harbor the same integrase gene. The integrase of $\phi\text{NM}4$ is identical to that of $\phi\text{L}54a$ in *S. aureus* COL and,




FIG. 3. Comparison of major pathogenicity islands vSaα (A) and vSaβ (B) in Newman with other *S. aureus* strains. Arrows represent open reading frames and their orientations. The two islands for strain Newman are shown on the top of each panel, whereas each type of vSaα and vSaβ is shown with representative strains below that for Newman: vSaα of N315, MW2, MRSA252, and RF122 are shown as types I through IV (A), whereas vSaβ of N315, MW2, and MRSA252 are shown as types I through III (B), respectively. Refer to Fig. 4 for the classes of the islands carried by other strains. Note that HsdS sequences for strains are polymorphic and that a common type of the pathogenicity island always harbors identical HsdS sequences regardless of the strains. Another pathogenicity island, SaPIbov (8), is inserted downstream of *guaA* gene in type IV vSaα of strain RF122, and only the insertion position is indicated.

similarly, φL54a and φNM4 insert into the same locus (*geh*). Thus, it seems highly likely that site-specific integration of phages into the staphylococcal genome is associated with different classes of integrase genes.

Virulence determinants. Table 2 summarizes major virulence-related genes found in strain Newman compared to strains MW2 and N315. Twelve additional virulence-related genes that belong to six paralogous groups identified in previous studies (3, 4) were encoded by the four Newman prophages, in addition to genes in other loci of the chromosome.

The functionally unknown phage genes were conspicuously present in Newman, whereas most of them were absent in prophages of strains MW2 and N315, suggesting that the phage genes play an important role in the virulence of strain Newman.

Most of the exoenzymes and adhesins were commonly present in the three strains shown in Table 2, although a lipase encoded in *geh* gene was truncated due to the insertion of φNM4, and extra genes were present in the *spl* serine protease cluster in vSaβ in strain Newman. It has also been noted that




FIG. 4. Tree view showing the phylogenetic relationships among 12 *S. aureus* strains calculated based on the sequence diversity of seven housekeeping genes and the classification of the genomic islands (Fig. 3) $vSa\alpha$ (A) and $vSa\beta$ (B) using ellipses. The genomic island classification is based on its structural differences and HsdS sequences (1): when the HsdS sequences in either $vSa\alpha$ or $vSa\beta$ island from different strains are identical, the islands are categorized into a common class, whose genetic content is strongly associated with the HsdS allotype in the island, as shown in Fig. 3. If strains harbor a common class of $vSa\alpha$ (A) or $vSa\beta$ (B), their names are covered by an ellipse with a unique color. Both HsdS in $vSa\alpha$ and $vSa\beta$ of RF122 are truncated, although $vSa\beta$ of the strain is classified into type II. The scale indicates the relative distance on the phylogenetic tree view.

fibronectin-binding proteins encoded by *fnbA* and *fnbB* genes were present in strain Newman, but they lack C-terminal cell wall sorting signals (27), as reported previously (11). The three strains in Table 2 share the same sets of hemolysin components and LukDE leukocidin genes. However, unlike *S. aureus* N315 and MW2, genes for staphylococcal superantigens were not found in the genome in strain Newman with the single exception of enterotoxin A (*sea*), which is encoded by ϕ NM3. Since other sequenced *S. aureus* strains normally harbor at least two enterotoxin genes in their chromosomes (data not shown), strain Newman is characteristic in its possession of a smaller number of enterotoxin genes. In *S. aureus* MW2, genes for enterotoxin H (*seh*) and collagen-binding protein (*cna*) are located within genome islets (1) and Panton-Valentine leukocidin component genes, whose product is responsible for fatal outcomes in humans such as necrotizing pneumonia (20) in a prophage; however, these virulence genes are not present in *S. aureus* Newman.

Major pathogenicity islands of *S. aureus* Newman. Prophages, staphylococcal cassette chromosome, and pathogenicity islands (8, 21) are categorized as genomic islands and carry characteristic integrase genes (DNA recombinases) (1). Unlike most other genomic islands, the pathogenicity islands $vSa\alpha$ and $vSa\beta$ are present in all *S. aureus* genomes sequenced thus far; however, $vSa\alpha$ and $vSa\beta$ harbor only remnants of their integrase genes. It therefore seems reasonable to assume that $vSa\alpha$ and $vSa\beta$ are no longer mobile and that these pathogenicity islands must have played a major role in the evolution of this pathogen (2). For example, even though the genome of *Staphylococcus epidermidis* is highly homologous to that of *S. aureus*,

the genomic islands $vSa\alpha$ and $vSa\beta$ are absent from the genome of this or any other coagulase-negative staphylococci (9, 19, 29, 33). These findings are in agreement with a general hypothesis that acquisition of $vSa\alpha$ and $vSa\beta$ into a primordial staphylococcal genome may have been associated with subsequent evolution of *S. aureus* as a major human pathogen.

$vSa\alpha$ and $vSa\beta$ display polymorphisms among strains and can be classified into three to four groups based on their structural differences and HsdS subtypes generated in each of the two islands. Tandem arrays of exotoxin and lipoprotein genes are characteristic features of $vSa\alpha$ in Newman (Fig. 3). $vSa\beta$ carries genes responsible for lantibiotic biosynthesis in addition to genes for leukocidin components (*lukD* and *lukE*) and a serine protease gene (*spl*) cluster. Compared to other strains, $vSa\alpha$ for Newman belongs to the same type as $vSa\alpha$ in strains N315, Mu50, NCTC8325, COL, JH1, JH9, and USA300 (type I in Fig. 3A), whereas $vSa\beta$ belongs to the same type as $vSa\beta$ in strains NCTC8325, COL, MW2, MSSA476, and RF122 (type II in Fig. 3B). Therefore, the combination of $vSa\alpha$ and $vSa\beta$ in *S. aureus* Newman (blue and brown ellipses in Fig. 4A and B, respectively) is the same as in strains NCTC8325, COL, and USA300 (see also Table 1). It is noteworthy that different types of $vSa\alpha$ and $vSa\beta$ correlate with sequence variation in *hsdS*, whose product determines the sequence specificity of DNA methylation and restriction via staphylococcal restriction-modification systems (17). Figure 4, however, also shows that the type distribution of the two major pathogenicity islands in the strains does not correlate with phylogenetic relationships among strains when a phylogenetic tree image is drawn based on allelic distribution of seven essential housekeeping

TABLE 2. Major virulence-related genes in strains Newman, MW2, and N315 and their locations^a

Type	Strain								
	Newman			MW2			N315		
	Locus	Gene	Island	Locus	Gene	Island	Locus	Gene	Island
Toxins									
Enterotoxin	NWMN1883	<i>sea</i>	ϕNM3	MW1889	<i>sea</i>	ϕSa3mw	SA1761	<i>sep</i>	ϕSa3n
				MW1938	<i>sek2</i>	ϕSa3mw	SA1817	<i>sec3</i>	νSa4
				MW1937	<i>seq</i>	ϕSa3mw	SA1816	<i>sel</i>	νSa4
				MW0759	<i>sec4</i>	νSa3	SA1642	<i>seg</i>	νSaβ
				MW0760	<i>sel2</i>	νSa3	SA1643	<i>sen</i>	νSaβ
				MW0051	<i>seh</i>		SA1646	<i>sei</i>	νSaβ
							SA1647	<i>sem</i>	νSaβ
							SA1648	<i>seo</i>	νSaβ
							SA0382	<i>set6</i>	νSaα
							SA0383	<i>set7</i>	νSaα
Exotoxin	NWMN0388	<i>set1nm</i>	νSaα	MW0382	<i>set16</i>	νSaα	SA0382	<i>set6</i>	νSaα
	NWMN0389	<i>set2nm</i>	νSaα	MW0383	<i>set17</i>	νSaα	SA0383	<i>set7</i>	νSaα
	NWMN0390	<i>set3nm</i>	νSaα	MW0384	<i>set18</i>	νSaα	SA0384	<i>set8</i>	νSaα
	NWMN0391	<i>set4nm</i>	νSaα	MW0385	<i>set19</i>	νSaα	SA0385	<i>set9</i>	νSaα
	NWMN0392	<i>set5nm</i>	νSaα	MW0386	<i>set20</i>	νSaα	SA0386	<i>set10</i>	νSaα
	NWMN0393	<i>set6nm</i>	νSaα	MW0387	<i>set21</i>	νSaα	SA0387	<i>set11</i>	νSaα
	NWMN0394	<i>set7nm</i>	νSaα	MW0388	<i>set22</i>	νSaα	SA0388	<i>set12</i>	νSaα
	NWMN0395	<i>set8nm</i>	νSaα	MW0389	<i>set23</i>	νSaα	SA0389	<i>set13</i>	νSaα
	NWMN0396	<i>set9nm</i>	νSaα	MW0390	<i>set24</i>	νSaα	SA0390	<i>set14</i>	νSaα
	NWMN0397	<i>set10nm</i>	νSaα	MW0391	<i>set25</i>	νSaα	SA0393	<i>set15</i>	νSaα
	NWMN0400	<i>set11nm</i>	νSaα	MW0394	<i>set26</i>	νSaα			
Toxic shock syndrome toxin						SA1819	<i>tst</i>	νSa4	
Exfoliative toxin	NWMN1082	<i>eta</i>	νSaγ	MW1054	<i>eta</i>	νSaγ	SA1016	<i>eta</i>	νSaγ
Alpha-hemolysin	NWMN1073	<i>hly</i>	νSaγ	MW0955	<i>hly</i>	νSaγ	SA1007	<i>hly</i>	νSaγ
Beta-hemolysin	Truncated	<i>hlyB</i>	ϕNM3	Truncated	<i>hlyB</i>	ϕSa3mw	Truncated	<i>hlyB</i>	ϕSa3n
Delta-hemolysin	NWMN1942	<i>hld</i>		MW1959	<i>hld</i>		SAS065	<i>hld</i>	
Gamma-hemolysin component	NWMN2318	<i>hlgA</i>		MW2342	<i>hlgA</i>		SA2207	<i>hlgA</i>	
	NWMN2319	<i>hlgC</i>		MW2343	<i>hlgC</i>		SA2208	<i>hlgC</i>	
	NWMN2320	<i>hlgB</i>		MW2344	<i>hlgB</i>		SA2209	<i>hlgB</i>	
LukDE leukocidin component	NWMN1717	<i>lukD</i>	νSaβ	MW1767	<i>lukD</i>	νSaβ	SA1637	<i>lukD</i>	νSaβ
Panton-Valentine leukicidin component	NWMN1718	<i>lukE</i>	νSaβ	MW1768	<i>lukE</i>	νSaβ	SA1638	<i>lukE</i>	νSaβ
				MW1379	<i>lukS-PV</i>	ϕSa2mw			
				MW1378	<i>lukF-PV</i>	ϕSa2mw			
Exoenzymes									
Serine protease	NWMN1706	<i>splA</i>	νSaβ	MW1755	<i>splA</i>	νSaβ	SA1631	<i>splA</i>	νSaβ
	NWMN1705	<i>splB</i>	νSaβ	MW1754	<i>splB</i>	νSaβ	SA1630	<i>splB</i>	νSaβ
	NWMN1704	<i>splC</i>	νSaβ	MW1753	<i>splC</i>	νSaβ	SA1629	<i>splC</i>	νSaβ
	NWMN1703	<i>splD</i>	νSaβ	MW1752	<i>splF</i>	νSaβ	SA1628	<i>splD</i>	νSaβ
	NWMN1702	<i>splE</i>	νSaβ				SA1627	<i>splF</i>	νSaβ
	NWMN1701	<i>splF</i>	νSaβ						
	NWMN0892	<i>htrA</i>		MW0903	<i>htrA</i>		SA0879	<i>htrA</i>	
	NWMN0918	<i>sspA</i>		MW0932	<i>sspA</i>		SA0901	<i>sspA</i>	
Serine V8 protease	NWMN0917	<i>sspB</i>		MW0931	<i>sspB</i>		SA0900	<i>sspB</i>	
	NWMN0916	<i>sspC</i>		MW0930	<i>sspC</i>		SA0899	<i>sspC</i>	
Cysteine protease	Truncated ^e	<i>geh</i>	ϕNM4	MW0297	<i>geh</i>		SA0309	<i>geh</i>	
	NWMN2569	<i>lip</i>		MW2590	<i>lip</i>		SA2463	<i>lip</i>	
Lipase	NWMN0624	<i>lipA</i>		MW0617	<i>lipA</i>		SA0610	<i>lipA</i>	
Lipase/esterase	NWMN2106	<i>hysA</i>		MW2129	<i>hysA</i>		SA2003	<i>hysA</i>	
Hyaluronate lyase	NWMN1236	<i>nuc</i>		MW1211	<i>nuc</i>		SA1160	<i>nuc</i>	
Thermonuclease									
Immunomodulators									
Staphylokinase	NWMN1880	<i>sak</i>	ϕNM3	MW1885	<i>sak</i>	ϕSa3mw	SA1758	<i>sak</i>	ϕSa3n
Chemotaxis inhibiting protein	NWMN1877	<i>chp</i>	ϕNM3				SA1755	<i>chp</i>	ϕSa3n
	NWMN1876	<i>scn</i>	ϕNM3	MW1884	<i>scn</i>	ϕSa3mw	SA1754	<i>scn</i>	ϕSa3n
Complement inhibitor	NWMN0055	<i>spa</i>		MW0084	<i>spa</i>		SA0107	<i>spa</i>	
Immunoglobulin G binding protein A	NWMN2317	<i>sbi</i>		MW2341	<i>sbi</i>		SA2206	<i>sbi</i>	
Immunoglobulin G binding protein sbi									
Adhesins									
Collagen-binding protein				MW2612	<i>cna</i>				
Fibronectin-binding protein	NWMN2399	<i>fnbA^b</i>		MW2421	<i>fnbA</i>		SA2291	<i>fnbA</i>	
	NWMN2397	<i>fnbB^b</i>		MW2420	<i>fnbB</i>		SA2290	<i>fnbB</i>	

Continued on following page

Downloaded from <http://jpb.asm.org/> on September 15, 2019 by guest

TABLE 2—Continued

Type	Strain								
	Newman			MW2			N315		
	Locus	Gene	Island	Locus	Gene	Island	Locus	Gene	Island
SD-rich fibrinogen-binding protein	NWMN0756	<i>clfA</i>		MW0764	<i>clfA</i>		SA0742	<i>clfA</i>	
	NWMN2529	<i>clfB</i>		MW2551	<i>clfB</i>		SA2423	<i>clfB</i>	
	NWMN0523	<i>sdrC</i>		MW0516	<i>sdrC</i>		SA0519	<i>sdrC</i>	
	NWMN0524	<i>sdrD</i>		MW0517	<i>sdrD</i>		SA0520	<i>sdrD</i>	
	NWMN0525	<i>sdrE</i>		MW0518	<i>sdrE</i>		SA0521	<i>sdrE</i>	
	NWMN1940	<i>sdrH</i>		MW1956	<i>sdrH</i>		SA1839	<i>sdrH</i>	
Elastin-binding protein	NWMN1389	<i>ebpS</i>		MW1369	<i>ebpS</i>		SA1312	<i>ebpS</i>	
Other virulence-related phage proteins ^c									
Functionally unknown protein ^A	NWMN0270 ^d		ϕNM4						
Functionally unknown protein ^B	NWMN0273		ϕNM4						
Functionally unknown protein ^C	NWMN0280		ϕNM4	MW1420		ϕSa2mw	SA1786		ϕSa3n
Functionally unknown protein	NWMN0284		ϕNM4						
Functionally unknown protein	NWMN0995		ϕNM2						
Functionally unknown protein ^B	NWMN1001		ϕNM2						
Functionally unknown protein ^C	NWMN1008		ϕNM2						
Functionally unknown protein ^C	NWMN1800		ϕNM1						
Functionally unknown protein ^B	NWMN1807		ϕNM1						
Functionally unknown protein ^A	NWMN1809 ^e		ϕNM1						
Functionally unknown protein ^C	NWMN1905 ^f		ϕNM3						
Functionally unknown protein	NWMN1912		ϕNM3	MW1924		ϕSa3mw	SA1795		ϕSa3n

^a Among virulence-related genes shown as red bars in Fig. 1, those involved in capsular polysaccharide biosynthesis and iron uptake are not listed.

^b The C terminus, including cell wall anchoring signal, is truncated, and therefore the product is nonfunctional (11).

^c Identified by *bursa aurealis* mutagenesis of strain Newman followed by nematode killing assay employing the mutants (4). Unknown proteins with the same superscript capital letter are paralogs.

^d Located between NWMN0270 and NWMN0271.

^e Located between NWMN1809 and NWMN1810.

^f Located between NWMN1905 and NWMN1906.

^g Truncated due to insertion of the indicated prophage.

genes used for MLST analysis (7). In addition, the distribution of ν Sa α types among strains differs from that of ν Sa β , suggesting that the two pathogenicity islands have been acquired by the strains independently of each other and differently from housekeeping genes that have presumably evolved in a vertical fashion.

Due to their general distribution in *S. aureus* strains and absence in other staphylococci, the two major pathogenicity islands are considered to play important roles in virulence for their human hosts. The molecular mechanisms that implement such putative strategies are, however, still unknown, and future work will need to unravel how pathogenicity islands are involved in staphylococcal virulence during host infection.

DISCUSSION

Following the first sequencing of *S. aureus* N315 (18), 11 additional *S. aureus* genomes have been determined and de-

posited into the databases. Here we add the whole genome sequence of *S. aureus* Newman to this rapidly growing list. Genome sequencing projects for multiple isolates of a bacterial pathogen are of considerable scientific value because the generated data reveal not only gene content but also conservation and variability between different strains and their associated human or animal diseases. Staphylococcal diversity is mainly due to polymorphisms that occur in genomic islands, which also carry many virulence and antibiotic resistance determinants. Nevertheless, some genes, such as the staphylocoagulase gene, are located outside of genomic islands and are known to be polymorphic (32). One can add to this list certain combinations of virulence genes, for example, *seh* (enterotoxin H) and *cna* (collagen-binding protein), which are present only in certain types of *S. aureus* strains. A hallmark of the *S. aureus* classification is the ability of these microbes to ferment mannitol and to produce characteristic proteins such as DNase, coagulase, and protein A. *S. aureus* strains differ from one

another in virulence and drug resistance features that are carried in or outside of genomic islands.

Previous works (3, 4) revealed virulence genes or candidate virulence genes within four prophages that have integrated into the genome of *S. aureus* Newman. Our determination of the whole genome sequence for strain Newman showed that many virulence-related genes are encoded by prophages. One superantigen, staphylococcal enterotoxin A (*sea*), is located in ϕ NM3; however, unlike other staphylococcal strains, additional superantigen genes were not found. Furthermore, *S. aureus* Newman carries a small pathogenicity island but lacks known virulence genes. We also failed to identify the collagen adhesin gene that is present in strains MW2, MRSA252, and MSSA476. Therefore, it is likely that virulence caused by strain Newman largely relies on prophages, in addition to the contribution by other virulence determinants present in all *S. aureus* strains, and the nonprophage regions of strain Newman genome seem to form the basic backbone of pathogenic *S. aureus*. While en bloc transfer of virulence genes via prophages and pathogenicity islands appears to be important for *S. aureus* acquisition of virulence properties, stepwise incorporation of additional genes and/or mutations may play an additional role in the evolution of clones with similar, yet discretely different strategies for the pathogenesis of human disease.

As shown in Fig. 4, analysis of two major pathogenicity islands in 12 different *S. aureus* genome sequences revealed that these strains do not always share the same combinations of ν Sa α and ν Sa β classes. Moreover, the classes do not correlate with phylogenetic relationship based on the allelic distribution of seven housekeeping genes upon MLST analysis (7). This clearly shows that these two pathogenicity islands were horizontally acquired and must have evolved independently of *S. aureus* genomes, whereas housekeeping genes are considered to evolve in a vertical fashion. Interestingly, sequences of *hsdS* gene products that determine the site specificity of methylation and restriction in restriction-modification systems vary depending on the type of pathogenicity islands that encodes them. The reasons why modification subunits of the R-M system are present in ν Sa α and ν Sa β and have sequence variations remain unknown. One possible explanation is that sequence diversity in pathogenicity islands requires its distinct restriction modification site determined by HsdS: since self-DNA protection by modification system is promoted by sequence-specific methylation on DNA, sequence diversity in genomic islands should coincide with the methylation site determined by HsdS. DNA methylation of pathogenicity islands may further influence expression of the virulence gene and thereby affect the pathogenesis of infectious diseases caused by this organism. Recent studies have revealed that type I RM system activity and modification site specificity are related to changes in the surface antigenic protein in *Mycoplasma pulmonis*, depending on the organism's infection sites (12, 28). This suggests that the RM system in *S. aureus* also plays a direct role in virulence.

Some of the genes located within the major pathogenicity islands, ν Sa α and ν Sa β , are presumed to be involved in virulence. However, their molecular contributions to pathogenicity are still unclear. It should also be noted that the presence of any one gene does not result in its expression. In order to

reveal the mechanisms of virulence further, microarray experiments could be used to reveal their expression.

The overall spectrum and individual combinations of virulence genes, as they are diversely encoded by different genomic islands, appears to be the major factor in determining clinical symptoms after *S. aureus* infection and may even dictate the severity of diseases caused by this pathogen. Together with an analysis of transposon insertion mutants (4), our work here may provide experimental strategies for better understanding the pathogenicity and physiology of *S. aureus*.

ACKNOWLEDGMENTS

This study was supported by a Grant-in-Aid for 21st Century COE, a Grant-in-Aid for Scientific Research on Priority Areas (no. 13226114), and a Grant-in-Aid for Scientific Research B (no. 14370097) from the Ministry of Education, Science, Sports, Culture, and Technology of Japan.

REFERENCES

- Baba, T., F. Takeuchi, M. Kuroda, H. Yuzawa, K. Aoki, A. Oguchi, Y. Nagai, N. Iwama, K. Asano, T. Naimi, H. Kuroda, L. Cui, K. Yamamoto, and K. Hiramatsu. 2002. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* **359**:1819–1827.
- Baba, T., F. Takeuchi, M. Kuroda, T. Ito, H. Yuzawa, and K. Hiramatsu. 2003. The genome of *Staphylococcus aureus*, p. 66–153. In D. Al'Aladeen and K. Hiramatsu (ed.), *The Staphylococcus aureus: molecular and clinical aspects*. Ellis Harwood, London, United Kingdom.
- Bae, T., T. Baba, K. Hiramatsu, and O. Schneewind. 2006. Prophages of *Staphylococcus aureus* Newman and their contribution to virulence. *Mol. Microbiol.* **62**:1035–1047.
- Bae, T., A. K. Banger, A. Wallace, E. M. Glass, F. Aslund, O. Schneewind, and D. M. Missiakas. 2004. *Staphylococcus aureus* virulence genes identified by *bursa aurealis* mutagenesis and nematode killing. *Proc. Natl. Acad. Sci. USA* **101**:12312–12317.
- Diep, B. A., S. R. Gill, R. F. Chang, T. H. Phan, J. H. Chen, M. G. Davidson, F. Lin, J. Lin, H. A. Carleton, E. F. Mongodin, G. F. Sensabaugh, and F. Perdreau-Remington. 2006. Complete genome sequence of USA300, an epidemic clone of acquired methicillin-resistant *Staphylococcus aureus*. *Lancet* **367**:731–739.
- Duthie, E. S., and L. L. Lorenz. 1952. Staphylococcal coagulase: mode of action and antigenicity. *J. Gen. Microbiol.* **6**:95–107.
- Enright, M. C., N. P. J. Day, C. E. Davies, S. J. Peacock, and B. G. Spratt. 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *Clin. Microbiol.* **38**:1008–1015.
- Fitzgerald, J. R., S. R. Monday, T. J. Foster, G. A. Bohach, P. J. Hartigan, W. J. Meaney, and C. J. Smyth. 2001. Characterization of a putative pathogenicity island from bovine *Staphylococcus aureus* encoding multiple superantigens. *J. Bacteriol.* **183**:63–70.
- Gill, S. R., D. E. Fouts, G. L. Archer, E. F. Mongodin, R. T. Deboy, J. Ravel, I. T. Paulsen, J. F. Kolonay, L. Brinkac, M. R. J. Beanan, S. C. Daugherty, R. Madupu, S. V. Angiuoli, A. S. Durkin, D. H. Haft, J. Vamathevan, H. Khouri, T. Utterback, C. Lee, G. Dimitrov, L. Jiang, H. Qin, J. Weidman, K. Tran, K. Kang, I. R. Hance, K. E. Nelson, and C. M. Fraser. 2005. Insights on evolution of virulence and resistance from the complete genome analysis of an early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing methicillin-resistant *Staphylococcus epidermidis* strain. *J. Bacteriol.* **187**:2426–2438.
- Gillaspay, A. F., V. Worrell, J. Orvis, B. A. Roe, D. W. Dyer, and J. J. Iandolo. 2006. *Staphylococcus aureus* NCTC8325 genome, p. 381–412. In V. Fischetti, R. Novick, J. Ferretti, D. Portnoy, and J. Rood (ed.), *Gram-positive pathogens*. ASM Press, Washington, DC.
- Grundmeier, M., M. Hussain, P. Becker, C. Heilmann, G. Peters, and B. Sinha. 2004. Truncation of fibronectin-binding proteins in *Staphylococcus aureus* strain Newman leads to deficient adherence and host cell invasion due to loss of the cell wall anchor function. *Infect. Immun.* **72**:7155–7163.
- Gumulak-Smith, J., A. Teachman, A. H. Tu, J. W. Simecka, J. R. Lindsey, and K. Dybvig. 2001. Variations in the surface proteins and restriction enzyme systems of *Mycoplasma pulmonis* in the respiratory tract of infected rats. *Mol. Microbiol.* **40**:1037–1044.
- Herron, L. L., R. Chakravarty, C. Dwan, J. R. Fitzgerald, J. M. Musser, E. Retzel, and V. Kapur. 2002. Genome sequence survey identifies unique sequences and key virulence genes with unusual rates of amino acid substitution in bovine *Staphylococcus aureus*. *Infect. Immun.* **70**:3978–3981.
- Holden, M. T., E. J. Feil, J. A. Lindsay, S. J. Peacock, N. P. Day, M. C. Enright, T. J. Foster, C. E. Moore, L. Hurst, R. Atkin, A. Barron, N. Bason,

- S. D. Bentley, C. Chillingworth, T. Chillingworth, C. Churcher, L. Clark, C. Corton, A. Cronin, J. Doggett, L. Dowd, T. Feltwell, Z. Hance, B. Harris, H. Hauser, S. Holroyd, K. Jagels, K. D. James, N. Lennard, A. Line, R. Mayes, S. Moule, K. Mungall, D. Ormond, M. A. Quail, E. Rabinowitch, K. Rutherford, M. Sanders, S. Sharp, M. Simmonds, K. Stevens, S. Whitehead, B. G. Barrell, B. G. Spratt, and J. Parkhill. 2004. Complete genomes of two clinical *Staphylococcus aureus* strains: evidence for the rapid evolution of virulence and drug resistance. *Proc. Natl. Acad. Sci. USA* **101**:9786–9791.
15. Iandolo, J. J., V. Worrell, K. H. Groicher, Y. Qian, R. Tian, S. Kenton, A. Dorman, H. Ji, S. Lin, P. Loh, S. Qi, H. Zhu, and B. A. Roe. 2002. Comparative analysis of the genomes of the temperate bacteriophages phi 11, phi 12, and phi 13 of *Staphylococcus aureus* 8325. *Gene* **289**:109–118.
 16. Katayama, Y., T. Ito, and K. Hiramatsu. 2001. Genetic organization of the chromosome region surrounding *mecA* in clinical staphylococcal strains: role of IS*A31*-mediated *mecI* deletion in expression of resistance in *mecA*-carrying, low-level methicillin-resistant *Staphylococcus haemolyticus*. *Antimicrob. Agents Chemother.* **45**:1955–1963.
 17. Kobayashi, I. 2001. Behavior of restriction-modification systems as selfish mobile elements and their impact on genome evolution. *Nucleic Acids Res.* **29**:3742–3756.
 18. Kuroda, M., T. Ohta, I. Uchiyama, T. Baba, H. Yuzawa, I. Kobayashi, L. Cui, A. Oguchi, K. Aoki, Y. Nagai, J.-Q. Lian, T. Ito, M. Kanamori, H. Matsumaru, A. Maruyama, H. Murakami, A. Hosoyama, Y. Mizutani-Ui, K. Kobayashi, T. Tanaka, T. Sawano, R. Inoue, C. Kaito, K. Sekimizu, K. Hirakawa, S. Kuhara, S. Goto, J. Yabuzaki, M. Kanehisa, A. Yamashita, K. Oshima, K. Furuya, C. Yoshino, T. Shiba, M. Hattori, N. Ogasawara, H. Hayashi, and K. Hiramatsu. 2001. Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* **357**:1225–1240.
 19. Kuroda, M., A. Yamashita, H. Hirakawa, M. Kumano, K. Morikawa, M. Higashide, A. Maruyama, Y. Inose, K. Matoba, H. Toh, S. Kuhar, M. Hattori, and T. Ohta. 2005. Whole genome sequence of *Staphylococcus saprophyticus* reveals the pathogenesis of uncomplicated urinary tract infection. *Proc. Natl. Acad. Sci. USA* **102**:13272–13277.
 20. Labandeira-Rey, M., F. Couzon, S. Boisset, E. L. Brown, M. Bes, Y. Benito, E. M. Barbu, V. Vazquez, M. Hook, J. Etienne, F. Vandenesch, and M. G. Bowden. 2007. *Staphylococcus aureus* Panton-Valentine leukocidin causes necrotizing pneumonia. *Science* **315**:1130–1133.
 21. Lindsay, J. A., A. Ruzin, H. F. Ross, N. Kurepina, and R. P. Novick. 1998. The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Mol. Microbiol.* **29**:527–543.
 22. Lowe, T. M., and S. R. Eddy. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**:955–964.
 23. Ma, X. X., T. Ito, C. Tiensasitorn, M. Jamklang, P. Chongtrakool, S. Boyle-Vavra, R. S. Daum, and K. Hiramatsu. 2002. Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob. Agents Chemother.* **46**:1147–1152.
 24. Naimi, T. S., K. H. LeDell, D. J. Boxrud, A. V. Groom, C. D. Steward, S. K. Johnson, J. M. Besser, C. O'Boyle, R. N. Danila, J. E. Cheek, M. T. Osterholm, K. A. Moore, and K. E. Smith. 2001. Epidemiology and clonality of community-acquired methicillin-resistant *Staphylococcus aureus* in Minnesota, 1996–1998. *Clin. Infect. Dis.* **33**:990–996.
 25. Page, R. D. M. 1996. TREEVIEW: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**:357–358.
 26. Ramu, C., H. Sugawara, T. Koike, R. Lopez, T. J. Gibson, D. G. Higgins, and J. D. Thompson. 2003. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* **31**:3497–3500.
 27. Schneewind, O., P. Model, and V. A. Fischetti. 1992. Sorting of protein A to the staphylococcal cell wall. *Cell* **70**:267–281.
 28. Sitaraman, R., A. M. Deniso, and K. Dybvig. 2002. A unique, bifunctional site-specific DNA recombinase from *Mycoplasma pulmonis*. *Mol. Microbiol.* **46**:1033–1040.
 29. Takeuchi, F., S. Watanab, T. Baba, H. Yuzawa, T. Ito, Y. Morimoto, M. Kuroda, L. Cui, M. Takahashi, A. Ankai, S. Baba, S. Fukui, J. C. Lee, and K. Hiramatsu. 2005. Whole-genome sequencing of *Staphylococcus haemolyticus* uncovers the extreme plasticity of its genome and the evolution of human-colonizing staphylococcal species. *J. Bacteriol.* **187**:7292–7308.
 30. Torell, E., D. Molin, E. Tano, C. Ehrenborg, and C. Ryden. 2005. Community-acquired pneumonia and bacteraemia in a healthy young woman caused by methicillin-resistant *Staphylococcus aureus* (MRSA) carrying the genes encoding Panton-Valentine leukocidin (PVL). *Scand. J. Infect. Dis.* **37**:902–904.
 31. van Wamel, W. J., S. H. Rooijackers, M. Ruyken, K. P. van Kessel, and J. A. van Strijp. 2006. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of *Staphylococcus aureus* are located on β -hemolysin-converting bacteriophages. *J. Bacteriol.* **188**:1310–1315.
 32. Watanabe, S., T. Ito, F. Takeuchi, M. Endo, E. Okuno, and K. Hiramatsu. 2005. Structural comparison of ten serotypes of staphylocoagulases in *Staphylococcus aureus*. *J. Bacteriol.* **187**:3698–3707.
 33. Zhang, Y. Q., S. X. Ren, H. L. Li, Y. X. Wang, G. Fu, J. Yang, Z. Q. Qin, Y. G. Miao, W. Y. Wang, R. S. Chen, Y., Shen, Z. Chen, Z. H. Yuan, G. P. Zhao, D. Qu, A. Danchin, and Y. M. Wen. 2003. Genome-based analysis of virulence genes in a non-biofilm-forming *Staphylococcus epidermidis* strain (ATCC 12228). *Mol. Microbiol.* **49**:1577–1593.