

## CDP-Alcohol Hydrolase, a Very Efficient Activity of the 5'-Nucleotidase/UDP-Sugar Hydrolase Encoded by the *ushA* Gene of *Yersinia intermedia* and *Escherichia coli*<sup>∇†</sup>

Isabel Alves-Pereira,<sup>1,2</sup> José Canales,<sup>1</sup> Alicia Cabezas,<sup>1</sup> Paloma Martín Cordero,<sup>3</sup> María Jesús Costas,<sup>1</sup> and José Carlos Cameselle<sup>1\*</sup>

*Grupo de Enzimología, Departamento de Bioquímica y Biología Molecular y Genética, Facultad de Medicina, Universidad de Extremadura, Badajoz, Spain*<sup>1</sup>; *Departamento de Química, Universidade de Évora, Évora, Portugal*<sup>2</sup>; and *Servicio de Microbiología, Complejo Hospitalario Universitario de Badajoz, Servicio Extremeño de Salud, Badajoz, Spain*<sup>3</sup>

Received 11 May 2008/Accepted 6 July 2008

Nucleoside 5'-diphosphate-X hydrolases are interesting enzymes to study due to their varied activities and structure-function relationships and the roles they play in the disposal, assimilation, and modulation of the effects of their substrates. Few of these enzymes with a preference for CDP-alcohols are known. In *Yersinia intermedia* suspensions prepared from cultures on Columbia agar with 5% sheep blood, we found a CDP-alcohol hydrolase liberated to Triton X-100-containing medium. Growth at 25°C was deemed optimum in terms of the enzyme-activity yield. The purified enzyme also displayed 5'-nucleotidase, UDP-sugar hydrolase, and dinucleoside-polyphosphate hydrolase activities. It was identified as the protein product (UshA<sub>Yi</sub>) of the *Y. intermedia ushA* gene (*ushA<sub>Yi</sub>*) by its peptide mass fingerprint and by PCR cloning and expression to yield active enzyme. All those activities, except CDP-alcohol hydrolase, have been shown to be the properties of UshA of *Escherichia coli* (UshA<sub>Ec</sub>). Therefore, UshA<sub>Ec</sub> was expressed from an appropriate plasmid and tested for CDP-alcohol hydrolase activity. UshA<sub>Ec</sub> and UshA<sub>Yi</sub> behaved similarly. Besides being the first study of a UshA enzyme in the genus *Yersinia*, this work adds CDP-alcohol hydrolase to the spectrum of UshA activities and offers a novel perspective on these proteins, which are viewed here for the first time as highly efficient enzymes with  $k_{cat}/K_m$  ratios near the theoretical maximum level of catalytic activities. The results are discussed in the light of the known structures of UshA<sub>Ec</sub> conformers and the respective homology models constructed for UshA<sub>Yi</sub>, and also in relation to possible biological functions. Interestingly, every *Yersinia* species with a sequenced genome contains an intact *ushA* gene, except *Y. pestis*, which in all its sequenced biovars contains a *ushA* gene inactivated by frameshift mutations.

Nucleoside 5'-diphosphate-X (NDP-X) hydrolases with different specificities for the nucleoside (N) and X moieties of the NDP-X substrates are widespread among all the kingdoms of life. They possess interesting properties by virtue of their varied activities and relationships between structure and function and the roles they play in the disposal, assimilation, or modulation of the effects of their substrates. Many NDP-X hydrolases belong to the Nudix protein superfamily, characterized by a conserved amino acid motif and by their reputation for being housecleaning enzymes (5, 28). However, there are also non-Nudix NDP-X hydrolases, including mammalian enzymes, like nucleotide pyrophosphatases/phosphodiesterases (6), dinucleoside triphosphatase (42), and ADP-ribose/CDP-alcohol pyrophosphatase (ADPRibase-Mn) (10, 11), or, relevant to this work, the bacterial 5'-nucleotidase/UDP-sugar hydrolase (UshA) (see below).

The *Escherichia coli* 5'-nucleotidase/UDP-sugar hydrolase (UshA<sub>Ec</sub>) is a periplasmic metallophosphoesterase expressed from the *ushA* gene as an immature precursor which, upon

export to the periplasm, undergoes maturation by proteolytic cleavage of a signal sequence (7, 8, 13). The substrate specificity of UshA<sub>Ec</sub> is complex, because it hydrolyzes many nucleotides and NDP-X compounds, but not without discrimination (Fig. 1A to C), so that different types of activities have been defined: 5'-nucleotidase (including apyrase-like activity), UDP-sugar hydrolase, and dinucleoside polyphosphate hydrolase activities (18, 20, 32, 38). Extensive structural studies of mature UshA<sub>Ec</sub> have been performed, including solving the structures of several enzyme conformers in complex with substrate (ATP), substrate-analog inhibitor ( $\alpha,\beta$ -methylene-ADP), or products (adenosine and phosphate) (24–26, 39, 40). However, the structural basis of its substrate specificity is still unsolved.

Few NDP-X hydrolases with a preference for CDP-X substrates are known. Besides the above-mentioned ADPRibase-Mn, there are two bacterial Nudix enzymes: the bifunctional YZGD from *Paenibacillus thioaminolyticus*, which is both a pyridoxal phosphatase and an NDP-X hydrolase with some resemblance to the specificity of ADPRibase-Mn (46), and a CDP-choline hydrolase identified in a survey of the Nudix hydrolases present in the *Bacillus subtilis* genome (48).

Here, starting from a CDP-alcohol hydrolase activity recovered from bacterial suspensions of *Yersinia intermedia*, we found that the UshA ortholog in this species (UshA<sub>Yi</sub>), and UshA<sub>Ec</sub> itself, hydrolyzes CDP-alcohols (Fig. 1D) with very high  $k_{cat}/K_m$  ratios. Besides being the first study of a UshA

\* Corresponding author. Mailing address: Departamento de Bioquímica y Biología Molecular y Genética, Facultad de Medicina, Apartado 108, 06080 Badajoz, Spain. Phone: 34-924289470. Fax: 34-924289468. E-mail: cameselle@unex.es.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

∇ Published ahead of print on 18 July 2008.




FIG. 1. Major groups of substrates for the activities of UshA<sub>Ec</sub> and UshA<sub>Yi</sub> enzymes. The structures shown are typical substrates hydrolyzed by the known 5'-nucleotidase and apyrase-like (AMP, ADP, and ATP are shown) (A), UDP-sugar hydrolase (UDP-glucose is shown) (B), or dinucleoside polyphosphate hydrolase (Ap<sub>2</sub>A, Ap<sub>3</sub>A, and Ap<sub>4</sub>A are shown) (C) activities and by the novel CDP-alcohol hydrolase activity (CDP-ethanolamine, CDP-glycerol, and CDP-choline are shown) (D) reported in this work.

enzyme in the genus *Yersinia*, this work makes the unexpected addition of CDP-alcohol hydrolase to the spectrum of UshA activities and offers a novel perspective on these proteins, which appear to be highly efficient enzymes acting near catalytic perfection.

## MATERIALS AND METHODS

**Bacteria, media, and other products for bacterial cultures.** The isolation of the *Y. intermedia* strain BA-1 was accidental, since it was obtained as a contaminant of chromatographic fractions of rat lung supernatant. *Y. intermedia* colonies were originally isolated from cultures in Columbia agar with 5% sheep blood (COS). They were positively identified with MicroScan Combo Gram Negative 1S panels from Siemens (Madrid, Spain) and ID 32 GN galleries from bioMérieux (Madrid, Spain). The strain is deposited in the Colección Española de Cultivos Tipo (Burjassot, Valencia, Spain) with accession no. CECT 7230. For this work, we used glycerol cultures kept frozen at  $-80^{\circ}\text{C}$  in our laboratory. *E. coli* BL21 and BL21(DE3) were purchased as transformation-competent, Gold-type preparations from Stratagene (Cultek, Madrid, Spain).

Precast COS plates were from bioMérieux. Liquid Luria-Bertani (LB) medium contained 10 g/liter Bacto tryptone, 5 g/liter yeast extract (both from Difco, Francisco Soria Melguizo, Madrid, Spain), and 5 g/liter NaCl. For solid LB medium, the liquid medium was supplemented with 15 g/liter Bacto agar (from Difco). For culture of plasmid-transformed *E. coli* cells, 0.1 g/liter ampicillin (from Sigma-Aldrich Química, Madrid, Spain) was added to LB media. For induction experiments, isopropylthiogalactoside (IPTG) (from Roche, Barcelona, Spain) was used at the concentrations indicated.

**Chemicals and biochemicals.** All the inorganic salts and acids, EDTA, and sodium dodecyl sulfate (SDS) were from Merck. Triton X-100, all the nucleotide compounds, and other enzyme substrates tested were from Sigma-Aldrich. Tris was from Roche. Auxiliary enzymes for assays of the activities or the reaction products were from Sigma-Aldrich (glycerol kinase and lactate dehydrogenase) or from Roche (alkaline phosphatase, glucose-6-phosphate dehydrogenase, glycerol-3-phosphate dehydrogenase, hexokinase, phosphoglucomutase, pyrophosphatase, and pyruvate kinase).

**Products for molecular biology.** Enzymes used for cloning included NotI and SalI (Roche), T4 DNA ligase (New England Biolabs, Izasa, Barcelona, Spain), and *Pfu* ultra-high-fidelity DNA polymerase (Stratagene). Plasmid pGEX-6P-3 was purchased from GE Healthcare (Barcelona, Spain), and pLM-2, containing the *E. coli ushA* gene (*ushA<sub>Ec</sub>*), was prepared as described by McMillen et al. (29).

**Purification of endogenous CDP-alcohol hydrolase of *Y. intermedia*.** Bacterial cells from glycerol cultures kept at  $-80^{\circ}\text{C}$  were seeded on 10-cm COS plates and grown for 36 h at  $25^{\circ}\text{C}$ . The cells were scraped from the surfaces of four plates with a stainless steel spatula, resuspended in 24 ml of buffer tMg (20 mM Tris-HCl, pH 7.5, 0.5 mM EDTA, 5 mM MgCl<sub>2</sub>), sedimented by centrifugation at  $10,000 \times g$  ( $25^{\circ}\text{C}$ ), and resuspended in buffer tMg supplemented with 5 mg/ml Triton X-100. After standing for 30 min at  $25^{\circ}\text{C}$ , the cells were sedimented by centrifugation for 10 min at  $10,000 \times g$  at  $4^{\circ}\text{C}$ . The supernatant of this centrifugation constituted step 1 of CDP-alcohol hydrolase purification, and it was chromatographed in a Sephacryl S-200 column (83 cm by 2.4 cm; GE Healthcare) equilibrated and eluted with buffer tMg supplemented with 0.1 M KCl at a flow rate of 17 ml/h (purification step 2). Fractions of 4.5 ml were collected and assayed for CDP-ethanolamine hydrolase activity. Those corresponding to the single activity peak recovered at an elution volume of  $\sim 240$  ml were pooled; dialyzed for 24 h against 1 liter of 20 mM Tris-HCl, pH 7.5, 5 mM MgCl<sub>2</sub> (with one buffer change); and applied to a Q-Sepharose column (25 cm by 1.6 cm; GE Healthcare) equilibrated in dialysis buffer. After application of the sample, the column was washed at 25 ml/h with 150 ml of the same buffer, followed by a linear 200-ml gradient of 0 to 400 mM KCl in the same buffer. The enzyme was recovered as a single peak of CDP-alcohol hydrolase activity centered around the gradient fraction with  $\sim 160$  mM KCl. This was the preparation used for enzyme characterization (step 3) (see Table S1 in the supplemental material). However, for peptide mass fingerprinting (PMF) of protein bands potentially associated with CDP-alcohol hydrolase activity, an additional purification step 4 was performed. The pooled fractions of the step 3 peak were diluted 1:1 in 5 mM MgCl<sub>2</sub> and applied to a Bio-Gel HTP column (25 cm by 1.6 cm; Bio-Rad) equilibrated in 5 mM MgCl<sub>2</sub>. The unbound protein was washed with the same salt solution, and the enzyme was eluted with a 100-ml linear gradient of 0 to 400 mM sodium phosphate, pH 7. Due to the high phosphate concentration, the chromatographic profile of CDP-ethanolamine hydrolase could not be determined using the P<sub>i</sub> liberation assay (see below). Instead, the enzyme was assayed by measuring the formation of the yellow product 4-nitrophenol using bis-*p*-nitrophenylphosphate as the substrate. The active fractions were pooled, dialyzed twice against 350 volumes of buffer tMg, assayed for CDP-ethanolamine hydrolase activity, and immediately submitted to SDS-polyacrylamide gel electrophoresis (PAGE) and then processed for PMF analysis (Servicio de Proteómica, UAM, Madrid, Spain).

**PCR cloning of the *Y. intermedia* ortholog of the *ushA* gene.** Genomic DNA was isolated from *Y. intermedia* CECT 7230 using the DNA Isolation Kit for Cells and Tissues according to the manufacturer's instructions (Roche). The *ushA* gene ortholog open reading frame (ORF) was amplified with primers accorded to the sequence that in the genome of *Y. intermedia* ATCC 29909 codes for the hypothetical protein ZP\_00834762 (the candidate pinpointed by the PMF of the endogenous CDP-alcohol hydrolase of *Y. intermedia*). The forward primer, GG ATAGTCGACATGCGTTTTTCATTACCGACC, contained a SalI site (underlined) followed by the first seven ORF codons; the reverse primer, GTCAGGC GGCCGCTTACTTATAAACAATCTC, contained a NotI site (underlined) followed by the reverse complement of the last six codons, including the stop codon. A product of the expected size (1,653 bp) was obtained, cut with SalI and NotI, and inserted into the corresponding sites of the pGEX-6P-3 plasmid to obtain plasmid pGEX-6P-3-Yi-*ushA*. Both strands of the insert were sequenced in the Servicio de Secuenciación Automática (IIB, CSIC-UAM, Madrid, Spain).

**Expression and purification of recombinant UshA proteins from *Y. intermedia* and *E. coli*.** To express the product of the *Y. intermedia ushA* gene, liquid LB cultures of BL21 cells transformed with pGEX-6P-3-Yi-*ushA* were grown overnight at  $37^{\circ}\text{C}$ . Five milliliters of the overnight culture was inoculated in 100 ml of fresh medium and cultured at  $28^{\circ}\text{C}$  until the  $A_{600}$  reached 0.8; then, IPTG (0.5 mM) was added and incubation continued for 2 h. The postsonication supernatant obtained in 10 ml of 20 mM Tris-HCl, pH 7.5, 0.5 mM EDTA, 50 mM KCl containing one dissolved tablet of protease inhibitor cocktail (Complete Mini; Roche) was concentrated to 2.7 ml by ultrafiltration. For purification, this preparation was fractionated by gel filtration in a Sephadex G-75 column (135 cm by 0.9 cm; GE Healthcare) equilibrated and eluted with buffer tMg with 50 mM KCl at a flow rate of 2 ml/h. Fractions (1 ml) were collected and assayed for CDP-ethanolamine hydrolase and 5'-nucleotidase activities. A single peak with both activities was collected at a  $V_e$  of  $\sim 58$  ml. The active fractions were pooled, dialyzed for 24 h against 100 volumes of 20 mM Tris-HCl, pH 7.5, 5 mM MgCl<sub>2</sub>, and chromatographed in a Q-Sepharose column as described for the endogenous CDP-alcohol hydrolase of *Y. intermedia* (see above).

To overexpress the product of the *E. coli ushA* gene, liquid LB cultures of BL21(DE3) cells transformed with pLM-2 were processed as described above, except that 1 mM IPTG was used for induction and, rather than lysing the bacteria by sonication, the periplasm was obtained (29) and used for enzyme




FIG. 2. Recovery of CDP-ethanolamine hydrolase and 5'-nucleotidase activities from *Y. intermedia* suspensions. CECT 7230 cells were collected from the surfaces of four COS plates and processed as described in Materials and Methods for the purification of endogenous CDP-alcohol hydrolase, except that (i) COS cultures were grown at different temperatures and (ii) after centrifugation to obtain the step 1 supernatant, the cell precipitates were resuspended in 10 ml of buffer tMg with Triton X-100 and were lysed by sonication. Enzyme activities were assayed in the supernatants (white portions of the bars) and in the lysed precipitates (black portions of the bars). The activities are expressed relative to the numbers of cells in the suspensions as estimated by  $A_{600}$  measurements.

purification as described for the endogenous CDP-alcohol hydrolase of *Y. intermedia* up to step 3 (see above).

**Enzyme activity assays.** Unless otherwise indicated, phosphohydrolytic activities were determined at 37°C by discontinuous assay of  $P_i$  liberation, either directly from substrates with a terminal phosphate (chain) or coupled to alkaline phosphatase for substrates without terminal phosphates. The standard reaction mixtures (50 to 400  $\mu$ l) contained 50 mM Tris-HCl, pH 7.5, 5 mM  $MgCl_2$ , 1 mg/ml bovine serum albumin, an amount of enzyme sample within the linearity range, 0.5 mM substrate, and, when needed, 9 units/ml alkaline phosphatase. The incubations were stopped by the addition of one of the  $P_i$  reagents indicated below.

**Assays of enzyme reaction products.**  $P_i$  was assayed colorimetrically with a dodecyl sulfate-ascorbate-molybdate reagent prepared in two different versions depending on the sensitivity required (35): either 0.7 ml of the standard reagent was added to 50- to 100- $\mu$ l samples or 0.2 ml of a concentrated reagent was added to 0.4-ml samples.

The amount of glycerol was assayed spectrophotometrically with glycerol kinase coupled to phosphoenolpyruvate kinase and lactate dehydrogenase (14), L-glycerol-3-phosphate with glycerol-3-phosphate dehydrogenase (30), glucose with hexokinase coupled to glucose-6-phosphate dehydrogenase (3), and glucose-1-phosphate with phosphoglucosyltransferase coupled to glucose-6-phosphate dehydrogenase (4).

Nucleosides and nucleotides were assayed by ion pair reverse-phase high-performance liquid chromatography in a Hypersil-ODS column (15 cm by 0.4 cm) protected by a guard column of the same material (1 cm by 0.4 cm) and connected to an Agilent HP 1100 chromatograph. Samples (20  $\mu$ l) were manually injected, and the elutions were performed at a flow rate of 0.5 ml/min: a 5-min isocratic elution with 1 mM sodium phosphate, pH 7, 20 mM tetrabutylammonium bromide, 10% (vol/vol) methanol, followed by a 20-min gradient elution with 1 to 100 mM sodium phosphate at constant tetrabutylammonium bromide and methanol levels. Chromatograms were recorded at 250 nm. Product peaks were identified by their retention times, compared with authentic standards, and integrated. Since in every reaction studied, the substrate and nucleotide product had equivalent molar extinction coefficients, the product peak areas were converted to molar amounts using the area of the substrate peak as an internal standard.

**Nucleotide sequence accession number.** The sequence of the 1,653-nucleotide insert ORF of plasmid pGEX-6P-3-Yi-*ushA* was deposited in GenBank (accession no. EF472003).




FIG. 3. Endogenous CDP-alcohol hydrolase from *Y. intermedia*. (A) SDS-PAGE analysis of partially purified preparations. The lanes show analyses of purification steps 3 (III) and 4 (IV) shown in Table S1 in the supplemental material. The gels were stained with silver. The arrows on the right mark the protein bands analyzed by PMF. The band identified as belonging to UshA<sub>Yi</sub> is the ~63-kDa band. (B) Activation by divalent cations. For this experiment, the enzyme (purification step 3 [see Table S1 in the supplemental material]) was dialyzed against 250 volumes of 20 mM Tris-HCl, pH 7.5, for 20 h, with one buffer change. The initial rates of CDP-ethanolamine hydrolysis were measured in 0.1-ml reaction mixtures using the standard assay, except that different divalent chloride salts were used at the indicated concentrations and bovine serum albumin was omitted. Equal amounts of dialyzed enzyme were used for the assay with each divalent cation.

## RESULTS

**Purification and characterization of an endogenous CDP-alcohol hydrolase of *Y. intermedia* and identification of the ortholog of *ushA* as a candidate for cloning.** Strain CECT 7230 of *Y. intermedia* was isolated in the laboratory as a contaminant of a partially purified rat lung fraction in which studies of CDP-alcohol hydrolases were in progress. From those experiments, it became clear that *Y. intermedia* produced a CDP-ethanolamine hydrolase. This activity appeared in the extracellular medium of pure *Y. intermedia* suspensions prepared with bacteria collected from the surfaces of COS plates and suspended in a buffer containing Triton X-100. The amount of enzyme in that medium was strongly dependent on the temperature at which the COS cultures were grown. Maximum activity was obtained from cultures grown at 25°C, while the activity declined significantly when the bacteria were grown at 28°C or higher (Fig. 2). The enzyme was purified 160-fold from *Y. intermedia* suspensions prepared from bacteria grown at 25°C (see Table S1 in the supplemental material). The last purification step contained two protein bands of ~63 kDa and ~23 kDa (Fig. 3A).

TABLE 1. Substrate specificities of the endogenous CDP-alcohol hydrolase of *Y. intermedia* (UshA<sub>Yi</sub>) and overexpressed UshA<sub>Ec</sub>

Substrate	Activity <sup>a</sup>	
	UshA <sub>Yi</sub> <sup>b</sup>	UshA <sub>Ec</sub> <sup>c</sup>
AMP	302	169
ADP	146–291 <sup>d</sup>	153–306 <sup>d</sup>
ATP	118–353 <sup>d</sup>	101–303 <sup>d</sup>
CMP	213	139
CDP	105–210 <sup>d</sup>	139–278 <sup>d</sup>
GMP	388	
UMP	147	
L-Glycerol-3-phosphate	<1	<1
D-Glucose-1-phosphate	<1	<1
Phosphoethanolamine	<1	<1
Phosphocholine	<1	<1
2'-AMP	<1	<1
CDP-glycerol	237	137
CDP-ethanolamine	100	100
CDP-choline	28	64
CDP-glucose	4	1
UDP-galactose	55	24
UDP-N-acetylglucosamine	40	48
UDP-glucose	38	20
UDP-glucuronate	36	16
UDP-N-acetylgalactosamine	36	18
UDP-galacturonate	10	3
ADP-glucose	<1	<1
dTDP-glucose	3	1
GDP-glucose	<1	<1
GDP-mannose	<1	<1
ADP-ribose	33	5
Ap <sub>2</sub> A	3	2
Ap <sub>3</sub> A	12	2
Ap <sub>4</sub> A	<1	<1
Ap <sub>5</sub> A	<1	<1
NAD <sup>+</sup>	<1	2
NADH	<1	1
FAD	<1	2
bis-p-Nitrophenylphosphate	4	1
Glycerophosphocholine	2	<1
Glycerophosphoethanolamine	<1	<1
4-Nitrophenylphosphocholine	<1	<1
4-Nitrophenyl-dTMP	<1	<1
2',3'-cyclic AMP	<1	1
3',5'-cyclic AMP	<1	<1

<sup>a</sup> Percentage of that measured with CDP-ethanolamine.

<sup>b</sup> Assayed with enzyme from purification step 3 (see Table S1 in the supplemental material).

<sup>c</sup> Overexpressed activity in lysate supernatants of *E. coli* BL21 cells transformed with the cloned *ushA* gene of *E. coli* (plasmid pLM-2) (29).

<sup>d</sup> Assuming that the number of P<sub>i</sub> moles formed per mole of substrate split was 1 to 3 (ATP) or 1 to 2 (ADP and CDP).

A study of enzyme specificity showed, besides CDP-alcohol hydrolase activity on CDP-glycerol, CDP-ethanolamine, and CDP-choline (but not CDP-glucose), a set of activities typical of UshA<sub>Ec</sub>: 5'-nucleotidase, UDP-sugar hydrolase, and dinucleoside polyphosphate hydrolase (Table 1). All were measured with Mg<sup>2+</sup> as the activating cation, but the enzyme was also activated by Co<sup>2+</sup> or Mn<sup>2+</sup>, but not by Ca<sup>2+</sup> (Fig. 3B). UshA<sub>Ec</sub> is known to be activated by all these cations, including Ca<sup>2+</sup> (20, 29, 32, 38).

The study of the reactions with CDP-alcohols and UDP-glucose revealed in both cases the same product pattern (see Table S2 in the supplemental material). The nucleotide CMP or UMP was not detected either as a final or an intermediate product. Instead, the nucleoside cytidine or uridine was appar-

ently formed directly from the substrate, together with an equimolar amount of P<sub>i</sub>. In addition, the hydrolysis of UDP-glucose gave an equimolar amount of D-glucose-1-phosphate, and the hydrolysis of CDP-glycerol yielded 0.5 mol of L-glycerol-3-phosphate per mol of cytidine and P<sub>i</sub> (since CDP-glycerol is a racemic mixture, the formation of another 0.5 mol of D-glycerol-3-phosphate, which does not react in the enzymatic assay used, can be assumed). Glycerol and D-glucose were not direct reaction products, but when the reaction was coupled to alkaline phosphatase, they were detected at the rate of 1 mol per mol of nucleoside, together with 2 mol of P<sub>i</sub>. These results correspond to the following general reaction: NDP-X + 2H<sub>2</sub>O → N + P<sub>i</sub> + P-X, where N is the nucleoside cytidine or uridine and P-X is the corresponding phosphoalcohol or phosphosugar (phosphoethanolamine, phosphocholine, DL-glycerol-3-phosphate, or D-glucose-1-phosphate). This reaction is identical to the hydrolysis of UDP-glucose by UshA<sub>Ec</sub>: the phosphoanhydride and the nucleoside 5'-ester linkages are hydrolyzed without detectable signs of nucleoside monophosphate (NMP) as an intermediate, possibly because it is hydrolyzed as soon as it is formed without leaving the active center (20).

The native molecular weight of the CDP-alcohol hydrolase, estimated by gel filtration chromatography, was about 52,000 (not shown). This left unclear whether the enzyme is a monomer of the ~63-kDa band or an oligomer of the ~23-kDa band seen by SDS-PAGE (Fig. 3A). In this respect, it is relevant that the molecular weight of mature UshA<sub>Ec</sub>, calculated from its amino acid sequence, is 58,100 (8), but during gel filtration chromatography, it behaves as a 52-kDa protein (38).

To investigate the molecular identity of the ~63-kDa and ~23-kDa protein bands, their tryptic PMFs were obtained and used to run Mascot searches (<http://www.matrixscience.com>) against the NCBI nr database. The best hits corresponded to hypothetical proteins conceptually translated from *Yersinia* genomic DNA sequences. The ~23-kDa protein was related with low significance to *Yersinia enterocolitica* and *Y. intermedia* superoxide dismutases. The ~63-kDa protein pinpointed, with a strong score, a *Y. intermedia* (strain ATCC 29909) hypothetical protein (GenBank accession no. ZP\_00834762) that showed 75% amino acid identity with UshA<sub>Ec</sub>.

**PCR cloning and expression of the *Y. intermedia* CECT 7230 *ushA* gene (*ushA<sub>Yi</sub>*) and biochemical characterization of its recombinant product.** The evidence supported the notion that CDP-alcohol hydrolase was likely an activity of UshA<sub>Yi</sub>. Therefore, we cloned *ushA<sub>Yi</sub>* by PCR, constructed plasmid pGEX-6P-3-Yi-*ushA*, and sequenced its 1,653-nucleotide insert ORF. With respect to the genome of *Y. intermedia* ATCC 29909, it showed four variations: C412T, T1056C, G1257A, and G1490A, with only the last changing the encoded amino acid (Ser497Asn).

The expression of plasmid pGEX-6P-3-Yi-*ushA* was expected to yield an 87.8-kDa GST-UshA<sub>Yi</sub> fusion protein with affinity for glutathione-Sepharose. SDS-PAGE analysis of the lysate supernatants of transformed *E. coli* cells showed three bands of about 80 kDa, 60 kDa, and 30 kDa responsive to induction with IPTG (Fig. 4A). Of these, only the ~30-kDa band was also induced in control cells transformed with vector pGEX-6P-3, where the expression of the GST protein alone was expected. The same supernatants showed the IPTG-induced overexpression of a set of Mg<sup>2+</sup>-dependent enzyme activities: CDP-alcohol hydrolase, 5'-nucleotidase, UDP-sugar




FIG. 4. Expression and purification of the recombinant UshA<sub>Yi</sub> protein. (A) Proteins induced by IPTG in *E. coli* cells transformed with plasmid pGEX-6P-3-Yi-ushA. The cells transformed with the empty vector pGEX-6P-3 (lanes 1 and 2) or with pGEX-6P-3-Yi-ushA (lanes 3 and 4), either with (+) or without (-) IPTG induction, were lysed and centrifuged. Shown is the SDS-PAGE analysis (Coomassie blue staining) of 12- $\mu$ l (lanes 1 and 2) or 8- $\mu$ l (lanes 3 and 4) samples of lysate supernatants. The arrows point to the three proteins induced by IPTG, two of which are specific to cells transformed with the *ushA* gene of *Y. intermedia*. The ~80-kDa band may correspond to the 87.8-kDa GST-UshA<sub>Yi</sub> fusion protein. The ~60-kDa band may correspond to the UshA<sub>Yi</sub> protein, either including (60.3 kDa) or not including (58.0 kDa) the predicted signal sequence (see Fig. S1 in the supplemental material). (B) Copurification of the activities overexpressed from pGEX-6P-3-Yi-ushA with an ~60-kDa protein band but not with the GST-UshA<sub>Yi</sub> fusion protein. Shown are (G-75) the chromatographic profiles of purification step 2, including enzyme activities (■, 5'-nucleotidase on AMP; ●, CDP-ethanolamine hydrolase), A<sub>280</sub> (▲), and SDS-PAGE analyses of selected fractions as indicated and (QS) the analysis of the enzymatically active fractions after purification step 3 (see Table S3 in the supplemental material). The arrows mark the gel positions of the ~80-kDa band (putative GST-UshA<sub>Yi</sub> fusion protein) and the ~60-kDa band. The PMF of the latter demonstrated unambiguously the presence of the UshA<sub>Yi</sub> protein (not shown).

TABLE 2. Hydrolytic activities overexpressed in *E. coli* cells transformed with *ushA*<sub>Yi</sub>

Substrate	Activity (U/mg)		
	pGEX-6P-3	pGEX-6P-3-Yi-ushA	Overexpressed <sup>a</sup>
AMP	0.08	18.81	18.73 (427)
UDP-glucose	0.08	2.15	2.07 (47)
CDP-glycerol	0.09	7.04	6.95 (158)
CDP-ethanolamine	0.06	4.45	4.39 (100)
CDP-choline	0.09	2.1	2.01 (46)
CDP-glucose	0.11	0.26	0.15 (3.4)
ADP-ribose	0.02	1.0	0.98 (22.3)
Ap <sub>3</sub> A	0.01	0.56	0.55 (12.5)
2',3'-cyclic AMP	0.07	0.09	0.02 (0.5)
3',5'-cyclic AMP	0.15	0.13	-0.02 (-0.5)
3',5'-cyclic AMP	0.09	0.09	0.00 (0.0)

<sup>a</sup> Percentages are in parentheses.

hydrolase, and dinucleoside polyphosphate hydrolase (Table 2), in agreement with the specificity of the endogenous enzyme of *Y. intermedia* (Table 1). Despite several attempts, neither the ~80-kDa nor the ~60-kDa protein band nor the IPTG-induced enzyme activities were adsorbed onto glutathione-Sepharose. In addition, when the lysate supernatant of *E. coli* cells transformed with *ushA*<sub>Yi</sub> and induced with IPTG was submitted to gel filtration chromatography, very little 5'-nucleotidase or CDP-alcohol hydrolase activity coeluted with the ~80-kDa band. Instead, they coeluted with the ~60-kDa protein, which by its size could correspond to the UshA<sub>Yi</sub> moiety separated from GST (Fig. 4B). To test this point, and to eliminate significant interference by the UshA<sub>Ec</sub> protein of the expression host, the PMF of the 60-kDa protein band was obtained (not shown). It confirmed unambiguously the presence of UshA<sub>Yi</sub> without signs of UshA<sub>Ec</sub>-specific peptides (for a possible explanation for the recovery of UshA<sub>Yi</sub> separated from GST upon expression of plasmid pGEX-6P-3-Yi-ushA, see below). Recombinant UshA<sub>Yi</sub> was then purified to near homogeneity (Fig. 4B, lane QS) by an additional ion-exchange chromatography step (the purification results are summarized in Table S3 in the supplemental material). This purified preparation displayed the same substrate specificity (not shown) displayed by the IPTG-induced set of activities in *E. coli* cells transformed with pGEX-6P-3-Yi-ushA (Table 2).

**Presence of a signal sequence in UshA<sub>Yi</sub>.** To ascertain whether UshA<sub>Yi</sub> contains a periplasmic signal sequence like that of UshA<sub>Ec</sub> (7, 8, 13), both proteins were submitted to informatics analysis. First, four different programs available online (19, 27, 31, 50) predicted a periplasmic location both for UshA<sub>Yi</sub> and UshA<sub>Ec</sub>, in agreement with experimental data for UshA<sub>Ec</sub>. Second, UshA<sub>Yi</sub> and UshA<sub>Ec</sub> were subjected to predictive analysis for the presence of a signal sequence with the SignalP program (15). For UshA<sub>Ec</sub>, the predictions indicated a signal sequence with a peptidase site between amino acids 25 and 26, in agreement with experimental evidence (7, 8, 13). For UshA<sub>Yi</sub>, SignalP also pointed to a strong signal sequence with a peptidase site between either amino acids 23 and 24 or 25 and 26. The signal sequences at the N ends of UshA<sub>Ec</sub> and UshA<sub>Yi</sub> are indicated in Fig. S1 in the supplemental material. Tryptic fingerprinting can provide evidence of the presence or absence of signal peptides. In the case of UshA<sub>Yi</sub>, the

TABLE 3. Kinetic parameters and catalytic efficiencies of UshA<sub>Yi</sub> and UshA<sub>Ec</sub>

Substrate	From this work <sup>a</sup>						From others
	$k_{\text{cat}}$ (s <sup>-1</sup> )		$K_m$ (μM)		$k_{\text{cat}}/K_m$ (M <sup>-1</sup> s <sup>-1</sup> )		$k_{\text{cat}}/K_m$ (M <sup>-1</sup> s <sup>-1</sup> )
	UshA <sub>Yi</sub>	UshA <sub>Ec</sub>	UshA <sub>Yi</sub>	UshA <sub>Ec</sub>	UshA <sub>Yi</sub>	UshA <sub>Ec</sub>	UshA <sub>Ec</sub>
AMP	735	372	9.6	1.8	$7.7 \times 10^7$	$2.1 \times 10^8$	$1.0 \times 10^{7b}$ $5.4 \times 10^{7c}$
CDP-glycerol	399	324	11.6	2.9	$3.4 \times 10^7$	$1.1 \times 10^8$	
CDP-ethanolamine	330	261	8.1	1.7	$4.1 \times 10^7$	$1.5 \times 10^8$	
CDP-choline	172	231	10.3	2.4	$1.7 \times 10^7$	$9.6 \times 10^7$	
UDP-glucose	97	71	9.6	10.4	$1.0 \times 10^7$	$6.8 \times 10^6$	$1.0 \times 10^{7d}$
ADP-ribose	72		5.8		$1.2 \times 10^7$		
Ap <sub>3</sub> A							$1.0 \times 10^{7e}$

<sup>a</sup> UshA<sub>Yi</sub> was the recombinant enzyme of *Y. intermedia* expressed in *E. coli* and purified (see Table S3 in the supplemental material). UshA<sub>Ec</sub> was the overexpressed enzyme of *E. coli* BL-21 cells transformed with plasmid pLM-2; the enzyme was purified by the same procedure as for UshA<sub>Yi</sub>, and a nearly homogeneous preparation was obtained (not shown). The standard enzyme assay was used, with a concentrated P<sub>i</sub> reagent to measure initial rates at different substrate concentrations. The data were adjusted to the Michaelis-Menten equation (12).

<sup>b</sup> Calculated as follows, with Co<sup>2+</sup> as the activating cation:  $k_{\text{cat}} = 750 \text{ s}^{-1}$  and  $K_m = 74 \text{ μM}$  (39).

<sup>c</sup> Calculated as follows, with Co<sup>2+</sup>-Ca<sup>2+</sup> as the activating cation:  $k_{\text{cat}} = 2,440 \text{ s}^{-1}$  and  $K_m = 45 \text{ μM}$  (38).

<sup>d</sup> Calculated as follows, with Mn<sup>2+</sup> as the activating cation:  $k_{\text{cat}} = 504 \text{ s}^{-1}$  and  $K_m = 45 \text{ μM}$  (38).

<sup>e</sup> Calculated as follows, with Mn<sup>2+</sup> as the activating cation:  $k_{\text{cat}} = 360 \text{ s}^{-1}$  and  $K_m = 35 \text{ μM}$  (38).

expected tryptic peptide formed by amino acids 3 to 28, with a calculated mass of 2,679.237 Da, is diagnostic of the presence of the signal sequence. Interestingly, experimental data for the endogenous and the recombinant proteins differed in this respect: the PMF of endogenous UshA<sub>Yi</sub> included the diagnostic peptide, while recombinant UshA<sub>Yi</sub> did not (not shown). It was concluded that the endogenous protein corresponded, at least in part, to a UshA<sub>Yi</sub> precursor. This can be explained by the liberation of immature UshA<sub>Yi</sub> from *Y. intermedia* in the cell suspensions prepared in the presence of Triton X-100. On the other hand, the recombinant protein appeared as the mature form of UshA<sub>Yi</sub>, indicating that the signal peptidase of *E. coli* acted on recombinant UshA<sub>Yi</sub>. Two pathways could account for the unexpected expression of mature UshA<sub>Yi</sub> from a plasmid encoding a fusion of GST with immature UshA<sub>Yi</sub> (pGEX-6P-3-Yi-*ushA*): either the translated fusion protein was processed directly by the *E. coli* signal peptidase, or part of the mRNA transcribed from plasmid pGEX-6P-3-Yi-*ushA* was translated from the immature UshA<sub>Yi</sub> initiation codon and the immature protein was processed by the signal peptidase.

**CDP-alcohol hydrolase activity of UshA<sub>Ec</sub>.** CDP-alcohol hydrolase activity has never been reported for a UshA enzyme. To elucidate whether it is a peculiarity of UshA<sub>Yi</sub> or a more general feature, *ushA<sub>Ec</sub>* was expressed in *E. coli* from plasmid pLM-2 (29). In addition to the known 5'-nucleotidase, UDP-sugar hydrolase, and dinucleoside polyphosphate hydrolase activities, CDP-alcohol hydrolase was strongly overexpressed (Table 1).

**The catalytic efficiencies of UshA<sub>Ec</sub> and UshA<sub>Yi</sub> activities, including CDP-alcohol hydrolase, are near the limits of enzymatic perfection.** Since the catalytic efficiency ratio,  $k_{\text{cat}}/K_m$ , cannot be greater than the diffusion-controlled enzyme and substrate association rate constant, the upper limit for this ratio is  $10^8$  to  $10^9 \text{ M}^{-1} \text{ s}^{-1}$ . Values of  $\geq 10^7 \text{ M}^{-1} \text{ s}^{-1}$  are typical of so-called "perfect" enzymes, which are limited by the diffusion rather than the chemical steps of their mechanisms (17). To our knowledge, there are two studies reporting  $k_{\text{cat}}$  and  $K_m$  values for some UshA<sub>Ec</sub> substrates, but they have not been

explicitly used to derive  $k_{\text{cat}}/K_m$  ratios (38, 39). In the present work, the saturation kinetics of UshA<sub>Ec</sub> and UshA<sub>Yi</sub> were studied for a selection of substrates with Mg<sup>2+</sup> as the activating cation (Table 3). Considering the  $k_{\text{cat}}/K_m$  ratios, the 5'-nucleotidase activity on AMP was the most efficient of those studied, followed very closely by CDP-alcohol hydrolase, and then by UDP-glucose and ADP-ribose hydrolase activities. The 5'-nucleotidase and CDP-alcohol hydrolase activities of UshA<sub>Ec</sub> displayed  $k_{\text{cat}}/K_m$  ratios of  $\geq 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , while UDP-glucose hydrolase displayed a 10-fold-lower value, similar to those calculated for the Mn<sup>2+</sup>-dependent UDP-glucose and diadenosine triphosphate (Ap<sub>3</sub>A) hydrolase activities from published saturation parameters (38, 39). A similar picture was found for UshA<sub>Yi</sub>, except that the  $k_{\text{cat}}/K_m$  values for 5'-nucleotidase and CDP-alcohol hydrolase activities were somewhat lower than those for UshA<sub>Ec</sub>, but still higher than  $10^7 \text{ M}^{-1} \text{ s}^{-1}$ .

**Theoretical analysis of UshA<sub>Yi</sub> structure in comparison to that of UshA<sub>Ec</sub>.** The structures of several UshA<sub>Ec</sub> conformers have been solved by X-ray diffraction and revealed a monomeric protein with two domains linked by an  $\alpha$ -helix. The N-terminal domain displayed the structural features typical of the metallo-dependent phosphatase superfamily, and it also contained the catalytic site. However, the substrate first binds far away from this site, in the C-terminal domain of the so-called "open" conformer. This inactive conformer is converted to the "closed" active conformer by a hinge-bending C-domain rotation of 96° that brings the bound substrate close to the catalytic center (24–26). To evaluate whether UshA<sub>Yi</sub> could adopt structures similar to those of UshA<sub>Ec</sub> and work by similar mechanisms, we adopted a theoretical approach. The UshA<sub>Yi</sub> protein (accession no. ABO69612) has the same length as the immature precursor of UshA<sub>Ec</sub> (accession no. NP\_415013). They align with 75% sequence identity and 86% similarity, including identity (see Fig. S1 in the supplemental material). Therefore, theoretical structures for UshA<sub>Yi</sub> were obtained by automatic homology modeling on the SWISS-MODEL server (41), using as models the X-ray structures of the two UshA<sub>Ec</sub> conformers that represent the extreme stages

of the 96° rotation. The homology models of UshA<sub>Yi</sub> showed the two-domain organization, the structural features of metallophosphoesterases in the N-terminal domain, and the possibility of adopting open and closed conformations, similarly to UshA<sub>Ec</sub> (see Fig. S2 in the supplemental material). In fact, comparing UshA<sub>Yi</sub> with UshA<sub>Ec</sub> structures, a full sequence and spatial conservation of the amino acids of the metallophosphoesterase signatures, the substrate-binding sites, and the catalytic centers was observed (see Fig. S1 and S3 in the supplemental material).

## DISCUSSION

**Major advances in regard to UshA enzymes.** In this study, a UshA protein was cloned, expressed, and characterized for the first time from bacteria of the genus *Yersinia*. This led to the finding of two properties shared by UshA<sub>Yi</sub> and UshA<sub>Ec</sub> that had remained unnoticed in this type of enzyme. One is the possession of CDP-alcohol hydrolase activity, not to a minor degree, but to the extent of being the second highest activity of the enzyme, not much lower than the 5'-nucleotidase and higher than the UDP-sugar hydrolase activity that lends its name to the *ushA* gene and its encoded protein. In retrospect, the reason why the high CDP-alcohol hydrolase activity of UshA<sub>Ec</sub> remained hidden may have been the low activity toward CDP-glucose compared to UDP-glucose, a feature that, after its early description (20), may have discouraged testing CDP-alcohols as substrates. The other novel feature found in this work for UshA enzymes is a catalytic efficiency ( $k_{cat}/K_m$ ) so high that it allows their inclusion in the category of "perfect" enzymes. This is particularly noteworthy considering that the catalytic cycle of the enzyme involves a 96° rotation of its C domain forward and back (see Fig. S2 in the supplemental material). It seems a considerable feat for this large motion, together with the chemical steps, to occur at a rate near that of the diffusion-controlled encounter of the enzyme and substrate in solution.

**The specificity patterns of UshA activities may depend on an interplay of substrate interactions with different protein domains.** The phosphohydrolytic behavior of UshA enzymes is rather complex, because although they hydrolyze a broad spectrum of phosphorylated compounds, they can be sorted into groups that depict an apparently multifunctional enzyme: 5'-nucleotidase (including apyrase-like activity), CDP-alcohol hydrolase, UDP-sugar hydrolase, and dinucleoside polyphosphate hydrolase. The apparent multifunctionality arises from the fact that, considering NDP-X as the prototypical substrate of the enzyme, those activities differ in their abilities to discriminate either among N groups or among X groups. The picture is not complete, because only a limited set of N/X combinations is available to the experimenter in the form of NDP-X compounds. However, the rather extensive list tested (Table 1) (18, 20, 32, 38) allows some conclusions to be drawn. The UDP-sugar hydrolase activity discriminates strongly for U against the other nitrogen bases (A, G, C, and T) so that UDP-glucose, UDP-galactose, UDP-N-acetylglucosamine, and UDP-N-acetylgalactosamine are hydrolyzed at similar rates, while ADP-glucose, GDP-glucose, CDP-glucose, and dTDP-glucose are not or are only marginally hydrolyzed. In contrast, the 5'-nucleotidase/apyrase activity or activities show little dis-

criminatory ability among nitrogen bases: all the NMP, NDP, and nucleoside triphosphate substrates are hydrolyzed at relatively similar rates. The CDP-alcohol hydrolase activity reported here adds further complexity to the UshA specificity pattern. Namely, the strong discrimination for uracil versus cytosine in the N moiety of the substrate, observed when glucose occupies the X position, is apparently overridden when glycerol or ethanolamine substitutes for glucose, as CDP-glycerol and CDP-ethanolamine are very good substrates, hydrolyzed with higher efficiencies than UDP-glucose.

The influence that the X group of the substrate exerts on the enzyme's ability to discriminate among N groups (or vice versa) can be accounted for by interactions of the NMP and P-X moieties of the substrate with different parts of the active site that are located in different protein domains. The structures of UshA complexes with the substrate ATP (see Fig. S2 in the supplemental material) (inactive conformer), the substrate-analog inhibitor  $\alpha,\beta$ -methylene-ADP (see Fig. S2 in the supplemental material) (active conformer), or the reaction products adenosine and P<sub>i</sub> (not shown) (inactive conformer) show that the NMP moiety binds to a C-domain pocket fitting AMP through several contacts (see Fig. S3A in the supplemental material). On the other hand, the  $\beta$  and  $\gamma$  phosphate groups of ATP (i.e., its P-X moiety) do not make any contact with the open UshA conformer and protrude freely toward the solvent (see Fig. S2 in the supplemental material), while the  $\beta$  phosphonate of  $\alpha,\beta$ -methylene-ADP, the nearest equivalent to a substrate studied in complex with UshA, makes contacts with the N domain of the closed conformer (see Fig. S2 and S3A in the supplemental material). Assuming that the NMP and P-X moieties of UDP-sugars and CDP-alcohols interact with the C and N domains of the closed conformer, we suggest that the specificity pattern of UshA enzymes results from a subtle interplay between NMP and P-X interactions, which might have consequences for the positioning of the scissile bonds of different compounds with respect to catalytic groups. The details of such interplay remain to be studied.

**The biological roles of UshA enzymes.** Because of its periplasmic location, UshA<sub>Ec</sub> has a role in the utilization of extracellular nucleotides. *E. coli* mutants isolated by a deficiency in 5'-nucleotidase activity are unable to take up and use AMP as a carbon source, which requires periplasmic conversion of AMP to adenosine (1, 49). Very recently, *ushA* knockouts of *E. coli* have been shown to be unable to grow on AMP as the only carbon source and are significantly delayed or impaired in their growth on other purine nucleotides. Growth on nonadenine purine nucleotides is made possible by an acid phosphatase encoded by the *aphA* gene, which at neutral pH acts as an efficient 5' nucleotidase of purine nucleotides, except AMP. Double knockouts in *ushA* and *aphA* do not grow on nucleotides (22).

In the gram-positive *Corynebacterium glutamicum*, but not in *E. coli*, the *ushA* gene is part of the phosphate starvation regulon, increasing its expression under conditions of phosphate deprivation (21), and its encoded UshA protein, with 5'-nucleotidase and UDP-sugar hydrolase activities, is required for growth on AMP or UDP-glucose as sole sources of phosphorus (36).

The resemblance to UshA<sub>Ec</sub>, both structurally and enzymatically, indicates that UshA<sub>Yi</sub> may play a role similar to that of

UshA<sub>Ec</sub> in regard to extracellular nucleotides. Besides *Y. intermedia*, with the exception of *Yersinia pestis*, all the *Yersinia* species for which sequenced genomes are available (*Y. enterocolitica*, *Yersinia pseudotuberculosis*, *Yersinia mollaretii*, *Yersinia berkoverii*, and *Yersinia frederiksenii*) contain genes coding for closely related UshA proteins (85% identity and 92% similarity, including identity) with conserved amino acids at all the positions of the substrate binding site and the dimetallic center (BlastP and alignment results not shown). The exceptionality of *Y. pestis* comes from two frameshift mutations: a single-nucleotide insertion between 252 and 253 and the deletion of nucleotide 967 in the *ushA* genes of the seven sequenced biovar genomes of the species (reference 34; nucleotide Blast and alignment results not shown). Most likely, the frameshifts prevent the expression of a functional UshA protein in *Y. pestis* (33). In addition, BlastN and TblastN searches of the genomes of *Yersinia* species indicated they lack genes orthologous to *aphA*. Therefore, it is possible that wild-type *Y. pestis* is unable to use nucleotides as a carbon source, similar to the *ushA* and *aphA* double knockouts of *E. coli* (see above).

The following factors seem to correlate the *ushA* genes of the genus *Yersinia* with the polysaccharidic O-antigen gene cluster: (i) the cluster contains genes related to the metabolism of NDP sugars and the biosynthesis of O antigen; (ii) *ushA* genes are located downstream and near the O-antigen clusters of *Y. pestis*, *Y. pseudotuberculosis*, and *Y. enterocolitica* serovar O:8 (16, 34, 43, 45); (iii) the lack of expression of O antigen in *Y. pestis* has been attributed to frameshift mutations in a few genes of the O-antigen cluster, and as noted above, this includes the nearby *ushA* (34, 44); (iv) the expression of the O-antigen cluster of *Y. enterocolitica* O:8 is markedly higher at near-ambient temperature than at 37°C (2), a behavior very similar to UshA<sub>Yi</sub> enzyme activities in our experiments (Fig. 2). These correlations make us wonder about a hypothetical role of *Yersinia* UshA proteins in the modulation of UDP sugars as glycosidic precursors.

Concerning the novel CDP-alcohol hydrolase activities of UshA<sub>Ec</sub> and UshA<sub>Yi</sub> on CDP-glycerol, CDP-ethanolamine, and CDP-choline, one would consider whether their very high catalytic efficiencies (Table 3) could be related to some of the known physiological roles of these compounds. (i) CDP-glycerol is the precursor from which gram-positive bacteria synthesize polyglycerolphosphate, one of the major forms of cell wall teichoic acid (47). (ii) CDP-ethanolamine and CDP-choline are well known as key intermediates of mammalian pathways for the synthesis of phosphatidylethanolamine and phosphatidylcholine; although its operation in bacteria is far from general, there is at least one well-documented case of use of the CDP-choline pathway by *Treponema denticola* (23). (iii) CDP-choline is the putative precursor for the “decoration” of cell surface lipopolysaccharides or (lipo)teichoic acids with phosphocholine (9, 37). Any of these CDP-alcohol-dependent processes could be affected by UshA enzymes with CDP-alcohol hydrolase activity, like UshA<sub>Ec</sub> and UshA<sub>Yi</sub>. To our knowledge, there is no evidence for CDP-alcohol use by *Yersinia* or *E. coli*; therefore, we ran TblastN searches, using the amino acid sequence of UshA<sub>Ec</sub> as the probe, against the sequenced genomes of *T. denticola* and other bacterial species selected for their putative abilities to use either CDP-glycerol to synthesize teichoic acid or CDP-choline to decorate their surfaces with

phosphocholine. In this way, we found nearly 30 bacterial species potentially using a CDP-alcohol for one of the above-mentioned purposes and containing a possible UshA ortholog ( $10^{-20} > E \text{ value} \geq 10^{-256}$ ) in their genomes (not shown).

Finally, the CDP-alcohol hydrolase activities of UshA proteins in the periplasmic space could act on CDP-alcohols from the infected host. In cases of intracellular infection, the CDP-alcohol hydrolase could disrupt the normal functioning of phospholipid biosynthetic routes by the hydrolysis of CDP-ethanolamine and CDP-choline.

#### ACKNOWLEDGMENTS

We are grateful to Ifor Beacham and Dennis Burns, Griffith University, Australia, for their kind gift of plasmid pLM-2.

This research was supported by grant BFU2006-00510 from the Ministerio de Educación y Ciencia, Spain, cofinanced by FEDER; grants GRU06031 and GRU07066 from the Consejería de Infraestructuras y Desarrollo Tecnológico, Junta de Extremadura Spain, cofinanced by FSE and FEDER; and grant A7-09 from the Vicerrectorado de Investigación, Desarrollo e Innovación, UEX.

#### REFERENCES

1. Beacham, I. R., R. Kahana, L. Levy, and E. Yagil. 1973. Mutants of *Escherichia coli* K-12 “cryptic,” or deficient in 5'-nucleotidase (uridine diphosphate-sugar hydrolase) and 3'-nucleotidase (cyclic phosphodiesterase) activity. *J. Bacteriol.* **116**:957–964.
2. Bengoechea, J. A., L. Zhang, P. Toivanen, and M. Skurnik. 2002. Regulatory network of lipopolysaccharide O-antigen biosynthesis in *Yersinia enterocolitica* includes cell envelope-dependent signals. *Mol. Microbiol.* **44**:1045–1062.
3. Bergmeyer, H. U., E. Bernt, F. Schmidt, and H. Stork. 1974. D-Glucose. Determination with hexokinase and glucose 6-phosphate dehydrogenase, p. 1196–1201. In H. U. Bergmeyer (ed.), *Methods of enzymatic analysis*, 2nd ed., vol. 3. Academic Press, New York, NY.
4. Bergmeyer, H. U., and G. Michal. 1974. D-Glucose-1-phosphate, p. 1233–1237. In H. U. Bergmeyer (ed.), *Methods of enzymatic analysis*, 2nd ed., vol. 3. Academic Press, New York, NY.
5. Bessman, M. J., D. N. Frick, and S. F. O'Handley. 1996. The MutT proteins or “Nudix” hydrolases, a family of versatile, widely distributed, “housecleaning” enzymes. *J. Biol. Chem.* **271**:25059–25062.
6. Bollen, M., R. Gijssbers, H. Ceulemans, W. Stalmans, and C. Stefan. 2000. Nucleotide pyrophosphatases/phosphodiesterases on the move. *Crit. Rev. Biochem. Mol. Biol.* **35**:393–432.
7. Burns, D. M., L. J. Abraham, and I. R. Beacham. 1983. Characterization of the *ush* gene of *Escherichia coli* and its protein products. *Gene* **25**:343–353.
8. Burns, D. M., and I. R. Beacham. 1986. Nucleotide sequence and transcriptional analysis of the *E. coli ushA* gene, encoding periplasmic UDP-sugar hydrolase (5'-nucleotidase): regulation of the *ushA* gene, and the signal sequence of its encoded protein product. *Nucleic Acids Res.* **14**:4325–4342.
9. Campbell, H. A., and C. Kent. 2001. The CTP:phosphocholine cytidylyltransferase encoded by the *licC* gene of *Streptococcus pneumoniae*: cloning, expression, purification, and characterization. *Biochim. Biophys. Acta* **1534**:85–95.
10. Canales, J., A. Fernández, J. M. Ribeiro, A. Cabezas, J. R. Rodrigues, J. C. Cameselle, and M. J. Costas. 2008. Mn<sup>2+</sup>-dependent ADP-ribose/CDP-alcohol pyrophosphatase: a novel metallophosphoesterase family preferentially expressed in rodent immune cells. *Biochem. J.* **413**:103–113.
11. Canales, J., R. M. Pinto, M. J. Costas, M. T. Hernández, A. Miró, D. Bernet, A. Fernández, and J. C. Cameselle. 1995. Rat liver nucleoside diphosphate or diphosphoalcohol pyrophosphatases different from nucleotide pyrophosphatase or phosphodiesterase I: substrate specificities of Mg<sup>2+</sup>- and/or Mn<sup>2+</sup>-dependent hydrolases acting on ADP-ribose. *Biochim. Biophys. Acta* **1246**:167–177.
12. Cleland, W. W. 1967. The statistical analysis of enzyme kinetic data. *Adv. Enzymol. Relat. Areas Mol. Biol.* **29**:1–32.
13. Cowman, A., and I. R. Beacham. 1980. Molecular cloning of the gene (*ush*) from *Escherichia coli* specifying periplasmic UDP-sugar hydrolase (5'-nucleotidase). *Gene* **12**:281–286.
14. Eggstein, M., and E. Kulhlmann. 1974. Triglycerides and glycerol. Determination after alkaline hydrolysis, p. 1825–1831. In H. U. Bergmeyer (ed.), *Methods of enzymatic analysis*, 2nd ed., vol. 4. Academic Press, New York, NY.
15. Emanuelsson, O., S. Brunak, G. von Heijne, and H. Nielsen. 2007. Locating proteins in the cell using TargetP, SignalP and related tools. *Nat. Protoc.* **2**:953–971.
16. Eppinger, M., M. J. Rosovitz, W. F. Fricke, D. A. Rasko, G. Kokorina, C. Fayolle, L. E. Lindler, E. Carniel, and J. Ravel. 2007. The complete genome



- sequence of *Yersinia pseudotuberculosis* IP31758, the causative agent of Far East scarlet-like fever. *PLoS Genet.* **3**:e142.
17. Fehrst, A. 1998. Structure and mechanism in protein science. A guide to enzyme catalysis and protein folding. W. Freeman & Co., New York, NY.
  18. Garcia, L., L. Chayet, A. M. Kettlun, L. Collados, M. Chiong, A. Traverso-Cori, M. Mancilla, and M. A. Valenzuela. 1997. Kinetic characteristics of nucleoside mono-, di- and triphosphatase activities of the periplasmic 5'-nucleotidase of *Escherichia coli*. *Comp. Biochem. Physiol. B* **117**:135–142.
  19. Gardy, J. L., M. R. Laird, F. Chen, S. Rey, C. J. Walsh, M. Ester, and F. S. Brinkman. 2005. PSORTb v. 2.0: expanded prediction of bacterial protein subcellular localization and insights gained from comparative proteome analysis. *Bioinformatics* **21**:617–623.
  20. Glaser, L., A. Melo, and R. Paul. 1967. Uridine diphosphate sugar hydrolase. Purification of enzyme and protein inhibitor. *J. Biol. Chem.* **242**:1944–1954.
  21. Ishige, T., M. Krause, M. Bott, V. F. Wendisch, and H. Sahn. 2003. The phosphate starvation stimulon of *Corynebacterium glutamicum* determined by DNA microarray analyses. *J. Bacteriol.* **185**:4519–4529.
  22. Kakehi, M., Y. Usuda, Y. Tabira, and S. Sugimoto. 2007. Complete deficiency of 5'-nucleotidase activity in *Escherichia coli* leads to loss of growth on purine nucleotides but not of their excretion. *J. Mol. Microbiol. Biotechnol.* **13**:96–104.
  23. Kent, C., P. Gee, S. Y. Lee, X. Bian, and J. C. Fenno. 2004. A CDP-choline pathway for phosphatidylcholine biosynthesis in *Treponema denticola*. *Mol. Microbiol.* **51**:471–481.
  24. Knöfel, T., and N. Sträter. 2001. *E. coli* 5'-nucleotidase undergoes a hinge-bending domain rotation resembling a ball-and-socket motion. *J. Mol. Biol.* **309**:255–266.
  25. Knöfel, T., and N. Sträter. 2001. Mechanism of hydrolysis of phosphate esters by the dimetal center of 5'-nucleotidase based on crystal structures. *J. Mol. Biol.* **309**:239–254.
  26. Knöfel, T., and N. Sträter. 1999. X-ray structure of the *Escherichia coli* periplasmic 5'-nucleotidase containing a dimetal catalytic site. *Nat. Struct. Biol.* **6**:448–453.
  27. Lu, Z., D. Szafron, R. Greiner, P. Lu, D. S. Wishart, B. Poulin, J. Anvik, C. Macdonell, and R. Eisner. 2004. Predicting subcellular localization of proteins using machine-learned classifiers. *Bioinformatics* **20**:547–556.
  28. McLennan, A. G. 2006. The Nudix hydrolase superfamily. *Cell Mol. Life Sci.* **63**:123–143.
  29. McMillen, L., I. R. Beacham, and D. M. Burns. 2003. Cobalt activation of *Escherichia coli* 5'-nucleotidase is due to zinc ion displacement at only one of two metal-ion-binding sites. *Biochem. J.* **372**:625–630.
  30. Michal, G., and G. Lang. 1974. 1-(-)-Glycerol-3-phosphate, p. 1415–1418. In H. U. Bergmeyer (ed.), *Methods of enzymatic analysis*, 2nd ed., vol. 3. Academic Press, New York, NY.
  31. Nair, R., and B. Rost. 2005. Mimicking cellular sorting improves prediction of subcellular localization. *J. Mol. Biol.* **348**:85–100.
  32. Neu, H. C. 1967. The 5'-nucleotidase of *Escherichia coli*. I. Purification and properties. *J. Biol. Chem.* **242**:3896–3904.
  33. Pieper, R., S. T. Huang, D. J. Clark, J. M. Robinson, P. P. Parmar, H. Alami, C. L. Bunai, R. D. Perry, R. D. Fleischmann, and S. N. Peterson. 2008. Characterizing the dynamic nature of the *Yersinia pestis* periplasmic proteome in response to nutrient exhaustion and temperature change. *Proteomics* **8**:1442–1458.
  34. Prior, J. L., J. Parkhill, P. G. Hitchen, K. L. Mungall, K. Stevens, H. R. Morris, A. J. Reason, P. C. Oyston, A. Dell, B. W. Wren, and R. W. Titball. 2001. The failure of different strains of *Yersinia pestis* to produce lipopolysaccharide O-antigen under different growth conditions is due to mutations in the O-antigen gene cluster. *FEMS Microbiol. Lett.* **197**:229–233.
  35. Ribeiro, J. M., A. Carloto, M. J. Costas, and J. C. Cameselle. 2001. Human placenta hydrolases active on free ADP-ribose: an ADP-sugar pyrophosphatase and a specific ADP-ribose pyrophosphatase. *Biochim. Biophys. Acta* **1526**:86–94.
  36. Rittmann, D., U. Sorger-Herrmann, and V. F. Wendisch. 2005. Phosphate starvation-inducible gene *ushA* encodes a 5' nucleotidase required for growth of *Corynebacterium glutamicum* on media with nucleotides as the phosphorus source. *Appl. Environ. Microbiol.* **71**:4339–4344.
  37. Rock, C. O., R. J. Heath, H. W. Park, and S. Jackowski. 2001. The *licC* gene of *Streptococcus pneumoniae* encodes a CTP:phosphocholine cytidyltransferase. *J. Bacteriol.* **183**:4927–4931.
  38. Ruiz, A., C. Hurtado, J. Meireles Ribeiro, A. Sillero, and M. A. Günther Sillero. 1989. Hydrolysis of bis(5'-nucleosidyl) polyphosphates by *Escherichia coli* 5'-nucleotidase. *J. Bacteriol.* **171**:6703–6709.
  39. Schultz-Heienbrok, R., T. Maier, and N. Sträter. 2005. A large hinge bending domain rotation is necessary for the catalytic function of *Escherichia coli* 5'-nucleotidase. *Biochemistry* **44**:2244–2252.
  40. Schultz-Heienbrok, R., T. Maier, and N. Sträter. 2004. Trapping a 96 degrees domain rotation in two distinct conformations by engineered disulfide bridges. *Protein Sci.* **13**:1811–1822.
  41. Schwede, T., J. Kopp, N. Guex, and M. C. Peitsch. 2003. SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res.* **31**:3381–3385.
  42. Sillero, M. A., R. Villalba, A. Moreno, M. Quintanilla, C. D. Lobatón, and A. Sillero. 1977. Dinucleosidetriphosphatase from rat liver. Purification and properties. *Eur. J. Biochem.* **76**:331–337.
  43. Skurnik, M., and J. A. Bengoechea. 2003. The biosynthesis and biological role of lipopolysaccharide O-antigens of pathogenic *Yersiniae*. *Carbohydr. Res.* **338**:2521–2529.
  44. Skurnik, M., A. Peippo, and E. Ervela. 2000. Characterization of the O-antigen gene clusters of *Yersinia pseudotuberculosis* and the cryptic O-antigen gene cluster of *Yersinia pestis* shows that the plague bacillus is most closely related to and has evolved from *Y. pseudotuberculosis* serotype O:1b. *Mol. Microbiol.* **37**:316–330.
  45. Thomson, N. R., S. Howard, B. W. Wren, M. T. Holden, L. Crossman, G. L. Challis, C. Churcher, K. Mungall, K. Brooks, T. Chillingworth, T. Feltwell, Z. Abdellah, H. Hauser, K. Jagels, M. Maddison, S. Moule, M. Sanders, S. Whitehead, M. A. Quail, G. Dougan, J. Parkhill, and M. B. Prentice. 2006. The complete genome sequence and comparative genome analysis of the high pathogenicity *Yersinia enterocolitica* strain 8081. *PLoS Genet.* **2**:e206.
  46. Tirrell, I. M., J. L. Wall, C. J. Daley, S. J. Denial, F. G. Tennis, K. G. Galens, and S. F. O'Handley. 2006. YZGD from *Paenibacillus thiaminolyticus*, a pyridoxal phosphatase of the HAD (haloacid dehalogenase) superfamily and a versatile member of the Nudix (nucleoside diphosphate x) hydrolase superfamily. *Biochem. J.* **394**:665–674.
  47. Ward, J. B. 1981. Teichoic and teichuronic acids: biosynthesis, assembly, and location. *Microbiol. Rev.* **45**:211–243.
  48. Xu, W., C. A. Dunn, C. R. Jones, G. D'Souza, and M. J. Bessman. 2004. The 26 Nudix hydrolases of *Bacillus cereus*, a close relative of *Bacillus anthracis*. *J. Biol. Chem.* **279**:24861–24865.
  49. Yagil, E., and I. R. Beacham. 1975. Uptake of adenosine 5'-monophosphate by *Escherichia coli*. *J. Bacteriol.* **121**:401–405.
  50. Yu, C. S., C. J. Lin, and J. K. Hwang. 2004. Predicting subcellular localization of proteins for Gram-negative bacteria by support vector machines based on *n*-peptide compositions. *Protein Sci.* **13**:1402–1406.