

The BlcC (AttM) Lactonase of *Agrobacterium tumefaciens* Does Not Quench the Quorum-Sensing System That Regulates Ti Plasmid Conjugative Transfer^{∇†}

Sharik R. Khan and Stephen K. Farrand*

Department of Microbiology, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801

Received 16 September 2008/Accepted 31 October 2008

The conjugative transfer of *Agrobacterium* plasmids is controlled by a quorum-sensing system consisting of TraR and its acyl-homoserine lactone (HSL) ligand. The acyl-HSL is essential for the TraR-mediated activation of the Ti plasmid *Tra* genes. Strains A6 and C58 of *Agrobacterium tumefaciens* produce a lactonase, BlcC (AttM), that can degrade the quorumone, leading some to conclude that the enzyme quenches the quorum-sensing system. We tested this hypothesis by examining the effects of the mutation, induction, or mutational derepression of *blcC* on the accumulation of acyl-HSL and on the conjugative competence of strain C58. The induction of *blc* resulted in an 8- to 10-fold decrease in levels of extracellular acyl-HSL but in only a twofold decrease in intracellular quorumone levels, a measure of the amount of active intracellular TraR. The induction or mutational derepression of *blc* as well as a null mutation in *blcC* had no significant effect on the induction of or continued transfer of pTiC58 from donors in any stage of growth, including stationary phase. In matings performed in developing tumors, wild-type C58 transferred the Ti plasmid to recipients, yielding transconjugants by 14 to 21 days following infection. *blcC*-null donors yielded transconjugants 1 week earlier, but by the following week, transconjugants were recovered at numbers indistinguishable from those of the wild type. Donors mutationally derepressed for *blcC* yielded transconjugants in planta at numbers 10-fold lower than those for the wild type at weeks 2 and 3, but by week 4, the two donors showed no difference in recoverable transconjugants. We conclude that BlcC has no biologically significant effect on Ti plasmid transfer or its regulatory system.

Two isolates of *Agrobacterium tumefaciens*, strains C58 and A6, produce a lactonase, BlcC (AttM), that degrades acyl-homoserine lactone (HSL) quorum-sensing signals (3, 57). *blcC*, which encodes the lactonase, is part of the three-gene *blcABC* operon (Fig. 1) that confers on the bacteria the ability to utilize γ -butyrolactone (GBL) as the sole source of carbon and energy (3, 6). The operon is located on the 543-kbp plasmid pAtC58 in strain C58 (3) and on the linear chromosome of strain A6 (57). The expression of the operon is controlled by BlcR, a repressor coded for by a gene located just upstream of and oriented in reverse to *blcABC* (Fig. 1) (3, 6, 57). The operon is strongly induced by two intermediates of the catabolic pathway, γ -hydroxybutyrate (GHB) and succinyl semialdehyde (SSA), very weakly by GBL, and not at all by acyl-HSLs (3, 6, 57). In strain A6, the operon also has been reported to be under the control of the stringent response system and induced during entry into stationary phase (52, 57, 58).

The conjugative transfer of pTiC58 and pTiA6, the Ti plasmids found in strains C58 and A6, is controlled by the quorum-sensing transcription factor TraR, which itself is encoded by the Ti plasmids (18, 38; reviewed in reference 54). The activator induces the expression of three Ti plasmid transfer operons, *traAFB*, *traCDG*, and *traI-trb* (19, 27). Functional TraR

requires the acyl-HSL quorum-sensing signal *N*-(3-oxo-octanoyl)-L-homoserine lactone (60), synthesized by TraI, which is also encoded by the Ti plasmid (21), as a ligand. Signal production is essential; *traI* mutants do not transfer the Ti plasmid unless the cultures are supplemented with exogenous quorumone (45). The quorum-dependent activation of Ti plasmid conjugative transfer systems is itself controlled in a hierarchical manner by a second signaling system. The Ti plasmids confer on their bacterial hosts the ability to induce plant cancers called crown galls. The galls arise following the transfer and integration of a specific small segment of the Ti plasmid, called the T region, from the bacterium into nuclear DNA of target plant cells (42). The resulting tumors produce metabolites, called opines, that the bacteria utilize as sole sources of carbon and energy (9, 15). The opines, which are synthesized by the transformed plant cells using enzymes coded for by genes on the integrated T-DNA, also induce conjugative transfer functions in the bacteria by directly controlling the expression of *traR* (15). In strain C58, agrocinopines, produced by tumors induced by this strain, control the expression of *traR* through the opine-modulated repressor AccR on pTiC58 (2, 39). In octopine-type Ti plasmids such as pTiR10 and pTiA6, octopine regulates expression of *traR* through OccR, an opine-modulated activator encoded on the plasmid (18, 19, 53). The quorum-dependent expression of the Ti plasmid *tra* regulon thus requires the exposure of the bacteria to the conjugative opine followed by an accumulation of the acyl-HSL signal necessary to generate active TraR. Quite clearly, the development of conjugative competence in *A. tumefaciens* has evolved to be intimately linked to the niche provided by the crown gall tumor (24, 25, 34, 35, 46).

* Corresponding author. Mailing address: Department of Microbiology, University of Illinois at Urbana-Champaign, B103 Chemical and Life Sciences Laboratory, 601 South Goodwin Avenue, Urbana, IL 61801. Phone: (217) 333-1524. Fax: (217) 244-6697. E-mail: stephenf@uiuc.edu.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

[∇] Published ahead of print on 14 November 2008.




FIG. 1. Genetic organization of the *blc* operon of pAtC58 and the pathway for degradation of GBL conferred by this operon. Gene organization is from the genome sequence of pAtC58 (GenBank accession number NC_003064). The main pathway and gene-enzyme relationships were described previously by Chai et al. (6). The shunt by which GABA enters the pathway is speculative.

The expression of BlcC results in greatly diminished levels of accumulation of the acyl-HSL signal in culture supernatants (3, 7, 57), and the lactonase has been reported to interfere with the acyl-HSL-dependent quorum-sensing system that controls Ti plasmid conjugative transfer (57). Such inhibition constitutes one example of a phenomenon called quorum quenching (reviewed in references 1, 10, and 11), and BlcC, based on its ability to degrade acyl-HSLs synthesized by the donor bacteria, has been called a quorum-quenching lactonase (3, 7, 57, 58).

While it is clear that the expression of the lactonase can affect levels of extracellular quorumone in culture, the role of this enzyme in quenching the quorum-dependent induction of the Ti plasmid conjugative transfer systems has not been rigorously assessed. Little is known concerning the regulation of the *blc* operon in planta, and no studies have reported the effect of this enzyme on the development or maintenance of conjugative transfer either in culture or, more importantly, in situ. Moreover, given the dependence of the regulatory circuitry controlling the induction of the conjugative transfer system on the development of the crown gall tumors and their subsequent production of opines, it is not at all clear that the bacteria will express the *blc* operon and the *tra* regulon at the same time in the infected plant.

In this study, we tested the hypothesis that BlcC is a quorum-quenching lactonase by assessing the influence of the enzyme on both extracellular and intracellular quorumone accumulation and on the development and maintenance of Ti plasmid conjugative transfer both in culture and in situ on developing crown gall tumors.

MATERIALS AND METHODS

Bacterial strains and plasmids. *Agrobacterium tumefaciens* strain C58 is a classical nopaline/agrocinopine, biovar 1 pathogen. Strain NTL4 (30) is a derivative of strain C58 that lacks pTiC58 but still harbors pAtC58, which codes for the *blcABC* operon. Strain C58C1RS (37) lacks pTiC58, is resistant to rifampin

and streptomycin, and contains a derivative of pAtC58 with an uncharacterized deletion that has removed the *blc* operon. This strain fails to utilize GBL (data not shown). Strain C58C1, the parent of C58C1RS, harbors an apparently full-sized pAtC58, contains a copy of *blc*, and grows with GBL as the sole carbon source (data not shown). We constructed a derivative of this strain, C58NTRS, that is resistant to rifampin and streptomycin by selecting sequentially for mutants that are resistant to the two antibiotics. The new strain is otherwise indistinguishable from C58C1 (data not shown). Strain AB153, a reconstructed derivative of C58 that harbors pTiC58 and pAtC58 (33), was used in several experiments. pTiC58 Δ *accR* contains a deletion in *accR* that removes opine control and renders the plasmid transfer constitutive (*tra*⁺) (2). pKPC12 is a derivative of pTiC58 that contains an inactive mutant allele of *traR* (39). This Ti plasmid fails to transfer and produces only small amounts of acyl-HSL. Ti plasmids were marked with a kanamycin resistance cassette as previously described (45). The nonpolar *tral* mutant pTiC58 Δ *traIKm* was constructed as previously described (45). *Agrobacterium* and *Rhizobium* spp. listed in Table 1 are from our collection.

Media, growth conditions, and reagents. *Agrobacterium* and *Rhizobium* spp. were grown at 28°C. LB or MG/L medium (28) was used as the rich liquid medium, while nutrient agar (Difco), yeast extract mannitol agar, or TY (55) agar was used as the rich solid medium. AB medium (8) supplemented with mannitol (0.2%) (ABM) or with GBL (26 mM; Sigma Chemical Co.) was used as the minimal medium for culturing *Agrobacterium* spp. *Rhizobium* spp. were grown on the minimal medium described previously by Ramírez-Romano et al. (41). Antibiotics were included in medium at the following concentrations: rifampin (Rf) at 50 μ g per ml, streptomycin (Sm) at 200 μ g per ml, kanamycin (Km) at 50 μ g per ml, and gentamicin at 25 μ g per ml. Growth in liquid cultures was monitored turbidometrically using a Klett colorimeter (red filter) or a Perkin-Elmer spectrophotometer at 600 nm and quantified by viable-colony counts (45). Succinic semialdehyde was obtained from Sigma Chemical Co. (St. Louis, MO).

Mutant and reporter constructions. A 2,098-bp PCR-generated fragment containing the *blcC* gene from pAtC58 was cloned into the BamHI site of pSF208, a derivative of pBluescript SK(+) in which the EcoRV and SmaI sites had been removed. A 251-bp internal fragment containing the first 224 bp of *blcC* was removed from this construct by digestion with EcoRV and ScaI and replaced with a gentamicin resistance cassette from pMGM (32). This allele of *blcC* was marker exchanged into pAtC58 by homologous recombination to produce pAtC58 Δ *blcC*. To construct a *blcR* mutant, a PCR fragment containing 699 bp of sequence downstream of *blcR* was fused with a 976-bp PCR fragment containing the first 204 bp of *blcR* of the *blcR*-*blcA* intergenic region and a 5' 718-bp segment of *blcA*, and this fragment was cloned into pBluescript II SK(+). The resulting construct is missing the 3' half of *blcR* but contains almost 700 bp of

pAtC58 DNA directly downstream of the lactonase gene. The gentamicin resistance gene cassette from pMGM was inserted at the BamHI fusion site, and the construct was marker exchanged into pAtC58 to produce pAtC58 Δ blcR.

A *blcC::lacZ* reporter fusion in pAtC58 was constructed by cloning a 272-bp internal fragment of the lactonase gene as an EcoRI-XbaI fragment into pVIK111 (22), generating a translational fusion between the fragment of *blcC* and *lacZYA*. The introduction of this nonreplicating plasmid into *Agrobacterium* strains yielded genomic reporters by Campbell insertions (22).

Expression constructs. The construction and properties of pSRKGm::*traR*, in which the expression of *traR*_{C58} is tightly regulated and inducible with IPTG (isopropyl- β -D-thiogalactopyranoside), were previously described (26). For the construction of pSRKTc::*traR*, The entire *traR* expression cassette from pSRKGm::*traR* was excised as a 2.5-kb BstBI fragment and cloned into BstBI-digested pSRKTc (26). In the absence of IPTG, this construct does not induce the transfer of wild-type pTiC58 ($<10^{-8}$ transconjugants per input donor). A PCR-generated fragment containing the entire *blcC* gene from pAtC58 was cloned into pRK415 (23) to yield pRK415::*blcC*. In this construct, *blcC* is constitutively expressed from the *lac* promoter of the vector.

Cloning and purification of BlcC. The *blcC* gene was amplified from genomic DNA of C58 by PCR, and the amplicon was cloned into pQE30 (Qiagen), generating a version of the gene with an in-frame His₆ sequence at its 5' end. In this construct, the expression of *blcC* is under the control of the T5 promoter. The His-tagged BlcC protein was overexpressed in *Escherichia coli* M15 cells (Qiagen) grown in LB induced with IPTG at 20°C, and the soluble product was purified by Ni-nitrilotriacetic acid affinity chromatography and dialyzed into 50 mM phosphate buffer (pH 7.5) containing 10% glycerol.

Purification of TraR. The active dimer TraR expressed from pETR in *E. coli* BL21(DE3) cells grown with 25 nM 3-oxo-C₈-HSL was purified as previously described (40). The purified protein is stable in solution and tightly retains the acyl-HSL ligand (40).

Quantitation of acyl-HSLs. Acyl-HSLs in culture supernatants, in cell pellets, or from enzyme assay buffer were extracted with ethyl acetate and quantified on thin-layer chromatography plates as previously described by using strain NTL4(pZLR4) as the bioreporter (17, 31, 45).

Matings. All matings were conducted by the spot plate method using C58CIRS or C58NTRS as a recipient as described previously (37, 45). In this method, volumes of dilutions of donors are spotted onto selective medium onto which a culture of the recipient has been spread as a lawn. Each colony appearing within a spot represents a transconjugant derived from mating with a single donor cell. Since the donors cannot transcribe genes or translate message on the mating plates, this method assesses the conjugative competence of the donor population at the instant that the cells are spotted onto the lawn of recipients. Frequencies are expressed as transconjugant colonies arising per number of donor cells spotted.

In planta matings. Donors and recipients were grown in MG/L medium to late exponential phase, and population densities were normalized to ca. 10^8 CFU per ml. Wounds measuring 2.5 cm were produced between the first and second nodes on 6- to 10-week-old tomato plants (Sunny Hybrid; Asgrow) grown in standard potting mix in the greenhouse. The wounds were inoculated with 10- μ l suspensions of donors, of recipients, or of a mixture of the two. Inoculated plants were maintained in the greenhouse without pesticides. At 2 weeks after infection and at weekly intervals thereafter, three plants from each treatment were chosen at random, the wounded segment was dissected, and each segment was macerated in 10 ml of sterile 0.9% saline. Volumes of 0.1 and 1.0 ml of the macerates were spread onto selection plates: ABM with Km for donors, ABM with Rf and Sm for recipients, and ABM with Km, Rf, and Sm for transconjugants. Tumors were first visible between weeks 2 and 3 following inoculation.

β -Galactosidase assays. β -Galactosidase activity, expressed as units per 10^9 cells, was quantified as described previously (28).

Southern analysis. Genomic DNA was extracted from bacterial cultures (29) and digested with EcoRV, HindIII, or BglII, and fragments were separated by electrophoresis and transferred onto nylon membranes (29). The membranes were incubated with the *blcC* gene labeled with digoxigenin under conditions of moderate stringency, and hybridizing fragments were detected by chemiluminescence as described in the digoxigenin application manual (Roche Diagnostics).

RESULTS

BlcC is inducible by pathway intermediates but not by GBL or entry into stationary phase. There are differences in the literature with respect to conditions under which the *blc* operon expresses (3, 6, 7, 52, 57, 58). For our studies, it was

important to verify under what conditions this gene set is induced. We monitored the expression of the operon in strain C58 grown in minimal mannitol medium under several conditions by monitoring levels of β -galactosidase from a *blcC::lacZ* fusion recombined into pAtC58. The reporter was induced about 20-fold by SSA but only twofold in cultures supplemented with GBL (see Fig. S1A in the supplemental material). Moreover, no change in the basal level of expression was detectable in unsupplemented cultures at any phase of growth, including late exponential and early stationary, mid-stationary, and late stationary phases (see Fig. S1B and S1C in the supplemental material). Induction by SSA was rapid, peaking after about 5 h of incubation. As previously reported (7), growth with γ -aminobutyric acid (GABA) also induced the operon (see Fig. S1D in the supplemental material). SSA yielded the highest levels of induction at concentrations as low as 500 μ M, while growth with GABA even at 10 mM yielded induction levels about twofold lower (see Fig. S1D in the supplemental material). Consistent with previous studies (3, 6, 57), a strain harboring pAtC58 Δ blcR *blcC::lacZ* in which the repressor gene *blcR* is deleted constitutively expressed the operon at high levels and in all growth phases (see Fig. S1C in the supplemental material).

The finding that GBL does not significantly induce the *blcC* reporter suggests that the lactone is not a true inducer of the operon. However, *blcC* is mutated in our reporter, and GBL has been reported to induce the wild-type operon (3). These observations suggest that the lactonase is required to convert GBL to an inducing intermediate, either GHB or SSA. We tested this hypothesis by examining the effect of GBL on the induction of the *blcC::lacZ* reporter in a strain that also expresses a cloned copy of *blcC* in *trans*. When grown with lactone, reporter strains also expressing *blcC* expressed the reporter fusion at levels 10-fold higher than did strains grown without supplement (see Fig. S2 in the supplemental material). We conclude that in strain C58, the *blc* operon is controlled strictly by BlcR in response to inducing substrates including SSA and GABA and that the operon does not autoinduce to any significant level as the cells enter stationary phase.

Induction of BlcC strongly affects extracellular but not intracellular levels of acyl-HSL. We assessed the influence of the lactonase on the levels of accumulation of the acyl-HSL in cultures of strain NTL4 harboring pTiC58 Δ accR, a Ti plasmid that is constitutive for transfer and for the synthesis of the quorumone. When cultured in the absence of an inducer, extracellular acyl-HSL accumulated to a maximum level of about 0.6 μ M at about 20 h of growth and plateaued as the cells entered stationary phase (Fig. 2A). In cultures in which the expression of *blcC* was induced with SSA, the signal was detectable but accumulated to levels four- to eightfold lower than that in the uninduced culture (Fig. 2A). A strain deleted for *blcC* accumulated acyl-HSL to levels about twice that of the *blcC*⁺ parent (Fig. 2A).

We also determined the amount of the acyl-HSL retained by the cells as a measure of levels of active intracellular TraR (5, 31). In contrast to the greatly decreased levels of extracellular signal, bacteria grown with SSA retained significant amounts of the acyl-HSL, about half the level detected in cells grown without inducer (Fig. 2B). The Δ blcC mutant accumulated intracellular quorumone at levels two- to fourfold higher than




FIG. 2. Expression of *blcC* strongly affects extracellular but not intracellular levels of acyl-HSL. Strain NTL4(pTiC58 Δ accRKm) harboring pAtC58 (\square and \diamond) or pAtC58 Δ blcC (\circ) was grown in ABM alone (\square and \circ) or ABM with SSA at 500 μ M (\diamond). Samples were removed at the indicated times and assayed for levels of extracellular (A) and intracellular (B) quorumone. A growth curve representative of the three cultures is shown (\blacksquare). The experiment was repeated once, with qualitatively similar results.

that of the *blcC*⁺ parent during exponential phase (Fig. 2B). However, the intracellular signal in the *blcC* mutant dropped to levels approximating those of the *blcC*⁺ parent as the cells transitioned into stationary phase.

BlcC does not significantly affect conjugative transfer of pTiC58 Δ accR. If the lactonase quenches quorum sensing, the induction of the *blc* operon should affect conjugative transfer frequencies. To test this hypothesis, we examined the influence of BlcC on the transfer properties of our *tra*^c Ti plasmid. Donors harboring pTiC58 Δ accRKm were cultured in minimal mannitol medium with or without SSA, samples were taken during growth, and cells were tested for conjugative transfer. Donors incubated with SSA transferred their plasmids at frequencies indistinguishable from those in which the *blc* operon was not induced at all time points tested (Fig. 3A). We also compared transfer frequencies of donors in which *blcC* or *blcR* had been deleted. Donors harboring pAtC58 Δ blcC or those harboring pAtC58 Δ blcR that constitutively expresses the *blc* operon transferred the Ti plasmid at frequencies similar to that of donors harboring wild-type pAtC58 (Fig. 3A and B). We conclude from these experiments that either the induced or constitutive expression of the *blc* operon has no significant effect on conjugative transfer from donors in which the *tra* system is already induced.

BlcC does not degrade the acyl-HSL bound by TraR. The induction of *blcC* had only a modest effect on quorumone retention by the cells (Fig. 2B) and failed to significantly affect transfer in transfer-constitutive donors (Fig. 3), suggesting that the acyl-HSL bound by TraR is not degraded by the lactonase. We tested this hypothesis by incubating the ligand-bound dimer form of TraR with a purified preparation of BlcC. As a control, we incubated the lactonase with authentic 3-oxo-C₈-




FIG. 3. BlcC lactonase does not affect conjugative transfer of a transfer-constitutive mutant of pTiC58. (A) Wild type compared with a *blcC* mutant. (B) Wild type compared with a *blcR* mutant. Cultures of strain NTL4(pTiC58 Δ accRKm) harboring pAtC58 (\square and \diamond), pAtC58 Δ blcC (\blacktriangle and \triangle) or pAtC58 Δ blcR (\bullet) were grown in ABM alone (\diamond , \blacktriangle , and \bullet) or in ABM supplemented with SSA at 500 μ M (\square and \triangle). Samples were removed at the indicated times, and frequencies of transfer to C58C1RS, expressed as transconjugants per donor, were measured by drop-plate mating. A growth curve (\circ) representative of the tested donor cultures is shown. Measurements were performed in triplicate at each time point, and the experiment was repeated once.

HSL in solution. While the enzyme rapidly degraded the free signal, the amount of acyl-HSL bound by TraR was not detectably decreased following the addition of lactonase (see Fig. S3 in the supplemental material).

Induction of BlcC has no significant influence on development of conjugative competence. Although BlcC had no effect on conjugation from donors already committed for transfer, the lactonase might interfere with the induction of the transfer process by preventing the accumulation of the quorumone to levels required to activate TraR. The transfer of pTiC58 is induced by the agropinopine opines, which control the transcription of *traR* (39). Since this opine is not available, we constructed a donor in which *traR* is under the control of a very tightly regulated *lac* promoter (26). In the absence of the *lac* inducer IPTG, the Ti plasmid does not transfer from such donors, while the addition of IPTG results in transfer at frequencies similar to that of pTiC58 Δ accR (26). In this system, IPTG functions as the surrogate conjugative opine.

To test the influence of the lactonase on the activation of the transfer system, we designed a system in which we could independently induce the expression of *blcABC* and *traR*. Donors harboring wild-type pAtC58; Ti plasmid pKPC12, in which *traR* is mutant; and pSRKGm::*traR* were cultured in ABM supplemented with IPTG to induce *traR* and also SSA to induce the *blc* operon of pAtC58. Samples were removed at intervals, and the donors were tested for conjugative competence by mating with strain C58C1RS. Donors in which the *blc*




FIG. 4. Preinduction but not simultaneous induction of *BlcC* lactonase transiently and modestly affects induction of transfer of pTiC58. (A) Simultaneous induction. A culture of NTL4(pKPC12, pSRKGm::*traR*) in ABM was split into two subcultures. IPTG was added to both cultures (■ and △), SSA (500 μ M) was added to one culture (△), and the two cultures were reincubated. (B) Preinduction. Two cultures of NTL4(pKPC12, pSRKGm::*traR*), one in ABM and the other in ABM with SSA (500 μ M), were grown for 4 h. IPTG was added to both (time zero), and the subculture containing IPTG only was further divided into two subcultures, one left untreated (□) and the other supplemented with SSA (●). The culture with SSA was similarly divided into two subcultures. The four cultures were incubated for 4 h, at which time SSA was added to one of the two subcultures that had been pretreated with SSA (▲) (arrow), and growth was continued. In both sets of experiments, donors sampled at the indicated times were tested for conjugative competence in drop-plate matings with C58C1RS. A growth curve representative of the tested donor cultures is shown (○). Measurements were performed in triplicate at each time point, and the experiment was repeated once.

operon was induced simultaneously with the *tra* regulon developed conjugative competence and transferred their Ti plasmids at rates and frequencies indistinguishable from those of donors in which only the transfer system was induced (Fig. 4A).

We considered the possibility that the preinduction of the *blc* operon is required to produce enough lactonase to significantly affect the production of the acyl-HSL and the consequent development of the conjugative transfer system. To test this hypothesis, we precultured the donor in medium containing SSA for 4 h to induce the *blc* operon and then added IPTG to induce *traR*. Four hours after the induction of *traR*, we split the culture and added a second dose of SSA to one subculture to ensure the continued expression of the lactonase. In parallel, we cultured the donor under conditions in which only *traR* was induced. As described above, samples were removed at intervals after the addition of IPTG, and donors were tested for conjugative competence. The preinduction of the *blc*




FIG. 5. Effects of *blcC* on induction of transfer depend on levels of expression of *traR*. Cultures of NTL4(pKPC12, pSRKTc::*traR*), one with pAtC58 (closed symbols) and the other with pAtC58 Δ *blcR* (open symbols), were grown to early exponential phase in ABM. At time zero, IPTG was added to induce *traR* at concentrations of 50 μ M (◆ and ◇), 100 μ M (■ and □), and 1 mM (● and ○). Growth was continued, samples were removed at the indicated times, and donors were tested for conjugative competence. Measurements were performed in triplicate at each time point, and the experiment was repeated once.

operon resulted in an initial frequency of transfer about three-fold lower than that in donors in which the operon was not induced (Fig. 4B). This difference was reduced to about 1.5-fold by 5 h, and by 10 h, both populations of donors transferred the Ti plasmid with the same efficiencies. The addition of a second dose of SSA to the preinduced donor culture did not further affect transfer frequencies (Fig. 4B). In all three cultures, transfer frequencies remained high even after the donors had transitioned into stationary phase.

As a most stringent test of the hypothesis, we assessed the effect of the constitutive expression of the lactonase on the induction of conjugative competence. Donors harboring our inducible *traR* system and either wild-type pAtC58 or pAtC58 Δ *blcR*, in which the *blc* operon is constitutively expressed at high levels (see Fig. S1C in the supplemental material), were cultured in fresh ABM, IPTG was added at several concentrations to induce expression of *traR* to different levels (26), and incubation of the culture was continued. Samples were removed at intervals, and the donors were tested for conjugative transfer by matings with C58C1RS. Growth with IPTG at 1 mM resulted in a rapid induction of conjugative competence, with transconjugants first appearing in both donors at 2 h (Fig. 5). Donors harboring wild-type pAtC58 transferred the Ti plasmid at frequencies almost 2 orders of magnitude higher than those containing the *blcR* mutation. Donors harboring pAtC58 Δ *blcR* incubated with lower, suboptimal concentrations of IPTG showed delays in the development of conjugative transfer (Fig. 5). However, in all cases, conjugative transfer from donors constitutively expressing the *blc* operon rose to levels of the corresponding wild-type donors by the second or third sampling time (Fig. 5).

BlcC does not significantly affect transfer of the Ti plasmid in planta. The regulation of Ti plasmid transfer is tuned via opines to the habitat of the crown gall tumors (9, 15). We assessed the influence of the lactonase on the quorum-sensing system in the natural habitat by measuring Ti plasmid transfer occurring on developing tumors induced by the donors. Tomato plants were inoculated with suspensions of donor cells,




FIG. 6. *blcC* does not significantly affect in planta transfer of pTiC58. Tomato plants were wounded and inoculated with mixtures of NTL4(pTiC58Km,pAtC58) (◆) or NTL4(pTiC58Km, pAtC58Δ*blcC*) (○) as tumorigenic donors and C58C1RS as the recipient. Beginning at week 2, wound sections from three plants were removed at weekly intervals for each mating set and macerated individually, and macerates were plated onto medium selective for donors, recipients, and transconjugants as described in Materials and Methods. (A) Numbers of transconjugants obtained per plant. Each symbol, which represents the average of data for three determinations, represents the datum from one plant at each sampling time point. (B) Frequency of transfer expressed as number of transconjugants recovered per donor.

either the wild type or mutant for *blcC*; with recipient cells; and with mixtures of one or the other donor and recipients. At intervals, segments of stems at the inoculation sites were removed, and macerates were plated onto medium selective for donors, for recipients, and for transconjugants. Plants infected with the pathogenic donors developed visible tumors between 2 and 3 weeks after inoculation.

In plants infected with wild-type donors, total numbers of recoverable donors and recipients increased by about 20-fold over the first week, plateauing at levels of about 10⁷ and 10⁶ CFU, respectively (see Fig. S4A and S4B in the supplemental material). Transconjugants were detectable 2 weeks postinoculation in plants infected with the *blcC* donor and after 3 weeks in plants infected with the wild-type donor (Fig. 6A). At 3 weeks postinfection, two plants infected with the wild-type donor yielded no detectable transconjugants (<50 transconjugants per tumor), while the third plant yielded the same number of transconjugants as the plants infected with the *blcC* donor. From week 4 on, the numbers of transconjugants that were isolatable from plants infected with either donor were not significantly different (Fig. 6A). Expressing the data as number of transconjugants per donor yielded the same pattern; al-




FIG. 7. Constitutive expression of the *blc* operon exerts a modest early and transient effect on in planta transfer of pTiC58. Tomato plants were infected with mixtures of NTL4(pTiC58Km, pAtC58) (◆) or NTL4(pTiC58Km, pAtC58Δ*blcR*) (○) as tumorigenic donors and C58C1RS as the recipient, and the wound sites were sampled and analyzed as described in the legend of Fig. 6. (A) Numbers of transconjugants obtained per plant. Each symbol, which represents the average of data from three determinations, represents the datum from one plant at each sampling time point. (B) Frequency of transfer expressed as number of transconjugants recovered per donor.

though initially delayed by 1 week, from week 4 on, the efficiency of transfer from *blcC*⁺ donors in planta was indistinguishable from that seen with donors that do not produce the lactonase (Fig. 6B).

It is conceivable that tomato plants do not produce inducers of the *blc* operon. To address this possibility, we conducted an in planta mating as described above using the Δ*blcR* mutant that constitutively expresses the operon as one of the donors. While the number of recoverable wild-type donors remained relatively constant over the course of the experiment, the numbers of Δ*blcR* donors decreased more than 10-fold at week 2 postinfection (see Fig. S5A in the supplemental material). However, by week 4, Δ*blcR* donors were recoverable at numbers equal to those of the wild type. The number of recoverable recipients remained relatively constant over the course of the experiment (see Fig. S5B in the supplemental material). Transconjugants were first detectable 2 weeks after infection in crosses with both wild-type and Δ*blcR* donors, and their numbers increased throughout the 6 weeks of the experiment (Fig. 7A). Compared to the Δ*blcR* donors, wild-type donors yielded up to 10 times as many transconjugants at weeks 2 and 3 postinfection. However, by weeks 4 and 5, there was little difference in the numbers of transconjugants recovered from




FIG. 8. Conjugative transfer in planta requires the acyl-HSL signal. Tomato plants were infected with mixtures of NTL4(pTiC58Km) (■) or NTL4(pTiC58ΔtraIKm) (□) as donors and C58C1RS as the recipient. The wound sites were sampled and analyzed for the conjugative competence of the *Agrobacterium* donors as described in the legend of Fig. 6. The data represent the average of the total numbers of transconjugants recovered from three plants of each mating at each time point.

tumors representing either mating (Fig. 7A). Transfer efficiencies, expressed as the number of transconjugants per donor, showed similar patterns (Fig. 7B).

Our standard recipient strain, C58C1RS, harbors a copy of pAtC58 that contains an uncharacterized deletion that has removed the *blc* operon. The use of this strain allowed us to focus on the role of the lactonase produced by the donors. However, in these matings, we could not address the possibility that the BlcC lactonase produced by a *blc*⁺ recipient could influence the development of conjugative competence. To test this hypothesis, we conducted a set of in planta matings between wild-type C58 and, as recipients, strain C58C1RS (*ΔblcRABC*) and the *blc*⁺ strain C58NTRS. In the two sets of matings, transconjugants were recoverable at similar numbers at 2 weeks postinfection (see Fig. S6 in the supplemental material). From week 4 on, matings with the *blc* wild-type recipient yielded six- to eightfold more transconjugants than did matings with the *Δblc* mutant (see Fig. S6 in the supplemental material).

The quorumone is essential for conjugative transfer in planta. It is possible that tomato plants produce a lactonase-resistant agonistic mimic of acyl-HSL (47). To test this hypothesis, we compared plasmid transfer from wild-type donors with that from donors harboring a Ti plasmid with a nonpolar deletion of *traI*, which encodes the acyl-HSL synthase (45). In tumors induced by the wild-type donor, transconjugants were detectable at week 3, with their numbers increasing almost 100-fold over the next 2 weeks (Fig. 8). However, no transconjugants were detected in tumors induced by the *ΔtraI* donor at any time point (Fig. 8).

The *blc* operon is not widely distributed in the genus *Agrobacterium*. If it is a dedicated component of the TraR-type quorum-sensing system, the *blc* operon should be widely distributed among bacteria that harbor plasmids in which transfer is controlled by this regulatory system. We tested this hypothesis by examining a representative group of field isolates of

Agrobacterium spp. as well as two members of the genus *Rhizobium* that harbor plasmids in which conjugation is known to be controlled by TraR-mediated quorum sensing (48, 55). As assessed by Southern analysis, 8 of the 20 agrobacterial isolates tested, with representatives from all three biovars of the genus, contain a detectable homolog of *blcC* (Table 1). Growth studies showed a strong correlation between the presence of this homolog and utilization of GBL as a sole carbon source. Among the two *Rhizobium* isolates tested, *R. leguminosarum* contains the entire *blc* operon and can utilize GBL, while *R. etli* lacks this gene system and fails to grow with this substrate (Table 1).

DISCUSSION

Quorum quenching as a natural phenomenon should result in quantitatively and temporally significant changes in quorum-regulated phenotypes as a result of conditions that induce the inhibitory mechanism. In the case of Ti plasmid transfer, quorum quenching by BlcC should cause demonstrably significant and lasting changes in transfer frequencies of the Ti plasmid under conditions in which the lactonase is produced by the donor cells. There is little from our results to support the hypothesis that this enzyme exerts such effects on the regulation of Ti plasmid transfer. While the expression of lactonase significantly decreased levels of extracellular quorumone (Fig. 2), there was little effect of this enzyme on the endpoint of the process, the conjugative proficiency of Ti plasmid donors. Inducing the expression of *blcC* did not significantly affect the induction of transfer in donors harboring wild-type Ti plasmids (Fig. 4) or steady-state rates of transfer from donors harboring *tra*^c plasmids in culture (Fig. 3). Where normal levels of induced expression of *blcC* did influence transfer, the effect was modest (≤ 10 -fold), early, and transient (Fig. 4 and 5).

More to the point, the role of its conjugative transfer system is to disseminate the Ti plasmid to appropriate recipients in the environs of the crown gall tumor (24, 25; reviewed in references 9 and 15). *blcC* either regulated or constitutively expressed in the donor significantly influenced neither the numbers of Ti plasmid-containing transconjugants nor the rate at which they appeared in planta (Fig. 6 and 7). Similarly, the expression of the lactonase by recipients had no inhibitory effect on Ti plasmid transfer from donors coresident on the tumors (see Fig. S6 in the supplemental material). The greatest effect, again transient, was observed in matings on tumors induced by donors in which *blcR* had been deleted (Fig. 7). However, under these conditions, *blcC* is constitutively expressed at levels significantly higher than those observed in wild-type cells grown with optimum amounts of inducers such as SSA and GABA (see Fig. S1D in the supplemental material). It is unlikely that the wild-type operon can be induced to such high levels, either in culture or in planta, casting doubt on the biological relevance of even this relatively low level of inhibition. Moreover, while such *blcR* donors transferred their Ti plasmids at frequencies up to 10-fold lower than those of *blcR*⁺ donors early in the infection, by 5 weeks after inoculation, numbers of transconjugants had reached wild-type levels (Fig. 7). In its wild-type state, BlcC had at best only a modest early effect on the development of conjugative competence in the tumorigenic donors and no demonstrable effect on the

TABLE 1. Distribution of *blcC* among *Agrobacterium* and *Rhizobium* isolates

Strain ^a	Biovar ^f	TraR-QS (reference) ^b	C58 <i>blcC</i> homolog(s) ^c	Growth on ^d :	
				Glucose	GBL
<i>Agrobacterium</i> spp.					
C58*	1	E, S (39)	SA ⁺ , G ⁺	+++++	+++++
A6*	1	E, S (59)	SA ⁺	+++++	+++++
15955*	1	E, S (16)	SA ⁺	+++++	+++
R10	1	E, S (18)	SA ⁺	+++++	+++++
Ach5	1	E, S (59)	SA ⁺	+++++	+++++
IIBv7	1	ND	SA ⁺	+++++	++++
Chry5*	1	E, S (34)	SA ⁺	+++++	++
Ant4*	1	E (51)	SA ⁺	+++++	+++
Bo542*	1	E, S (34)	SA ⁻	+++++	-
T37	1	E (14)	SA ⁻	+++++	+
K599*	1	S (4)	SA ⁻	+++++	+
B6S3 ^e	1	E, S (46)	SA ⁻	+++++	-
B6-Schil ^c	1	E, S (46)	SA ⁻	+++++	-
K84*	2	E, S (35)	SA ⁻ , G ⁻	+++++	-
J73	2	ND	SA ⁻	+++++	+
AB2-73*	2	E (50)	SA ⁻	+++++	-
A4*	2	E, S (36)	SA ⁻	+++++	-
K299*	2	E (35)	SA ⁻	+++++	+
8196	2	E (36)	SA ⁻	+++++	-
S4	3	S	SA ⁻	+++++	-
TM4	3	ND	SA ⁻	+++++	-
<i>R. leguminosarum</i> 3841	NA	E, S (55)	G ⁺	+++++	++++
<i>R. etli</i> CFN42	NA	E, S (48)	G ⁻	+++++	+

^a All listed strains are held in our collection. Strains marked with an asterisk have been reported to produce *N*-(3-oxooctanoyl)-L-homoserine lactone (4).

^b TraR-QS, presence of a plasmid coding for a TraR-type quorum-sensing system; E, experimentally determined; S, determined by sequence analysis; ND, no data available.

^c Determined by moderate-stringency Southern analysis using a *blcC* probe (SA⁺ and SA⁻) and by Blastn, Blastp, and Tblastn analyses of genome sequences available in public databases (G⁺ and G⁻).

^d Growth on solid minimal medium with the indicated substrate as the sole carbon source was assessed visually and graded from +++++ (luxuriant growth) to - (no detectable growth).

^e Two isolates of strain B6 with different plasmid complements (20).

^f NA, not applicable.

long-term emergence of transconjugants in the habitat of the tumor. Finally, the finding that donors that are unable to synthesize an acyl-HSL do not transfer their Ti plasmids at detectable frequencies in planta (Fig. 8) shows that the quorumone is essential and that tomato plants do not produce acyl-HSL mimics that might compensate for signal degradation by the lactonase.

Although greatly reducing the amount of extracellular quorumone, the expression of the lactonase only modestly affected the levels of intracellular signal (Fig. 2), taken to be a gauge of the amount of active TraR in the cell (5, 31). In culture, 3-oxo-C₈-HSL fully activates Ti plasmid transfer at an extracellular concentration as low as 1 nM (45). Assuming free diffusion of the quorumone and an average cell volume of 1.5 μm³, this concentration represents between 10 and 20 molecules of signal per cell, corresponding to between 5 and 10 copies of TraR dimer. In culture, signal can accumulate to μM levels (Fig. 2) (57), an amount about 1,000-fold higher than that needed to fully induce the quorum-sensing system. This requirement for only a few molecules of signal per cell may explain why BlcC, even when induced to maximum levels, does not significantly affect the quorum-sensing system.

Zhang et al. (57) previously reported that the expression of the *blc* operon of strains A6 and C58 is induced during entry into stationary phase. Since they observed a concurrent de-

crease in the amount of extracellular quorumone, they concluded that such induction could serve to shut down conjugation during entry into stationary phase. However, consistent with a previous report by Carlier et al. (3), we did not observe an increase in the expression of *blc* in stationary-phase cells of C58 in the absence of an inducer (see Fig. S1 in the supplemental material). In addition, quorumone levels stayed high in stationary-phase cultures in which the *blc* operon was not induced with exogenous substrate. Ligand-bound TraR is stable (40, 63), and the acyl-HSL bound by the dimer is not accessible to degradation by BlcC (see Fig. S2 in the supplemental material). These observations make it unlikely that the lactonase plays a role in turning over active TraR in any stage of growth. Moreover, if the role of the lactonase is to degrade the signal, the hypothesis predicts that conjugative proficiency would rapidly decay since the signal would become limiting. The transfer of the Ti plasmid from *A. tumefaciens* B6 apparently decreases substantially in stationary phase (46), an observation consistent with the hypothesis. However, the BlcC lactonase cannot be responsible for this decay; strain B6 lacks the *blcC* gene and fails to grow with GBL (Table 1). Moreover, a null mutation in or the overexpression of *blcC* had no significant effect on the conjugative competence of strain C58 in any stage of growth including stationary phase (Fig. 3 and 4). This observation is consistent with recent studies showing that C58 retains conju-

gative competence well into stationary phase even when extracellular quorumone is removed (45). These results strongly suggest that BlcC does not play a role in returning the quorum-sensing system to its off state at any stage of the culture cycle.

More likely, *blc* has evolved for the catabolism of butyryl compounds. The operon responds to and confers the utilization of GBL as well as its catabolic intermediates GHB and SSA but not acyl-HSLs (3). Interestingly, GBL is itself at best a very poor inducer of the operon (see Fig. S1 in the supplemental material). Induction following growth with this substrate apparently requires its conversion to SSA, which is a strong inducer (6). Although GBL has not been reported to be present in plants, GABA, which does accumulate in plants, induces the *blc* operon in C58 (see Fig. S1 in the supplemental material) (7), presumably following its conversion to SSA by a GABA-T-like transaminase (Fig. 1). While such an activity has not been reported for *Agrobacterium*, the genome of strain C58 encodes a protein, Atu3300, that is 66% similar to GABA-T from tomato at the amino acid sequence level. To date, gene sets orthologous in sequence and organization to the *blcABC* operon and its *blcR* regulator have been identified in the genomes of only three other bacteria, *Yersinia intermedia* ATCC 29909, *Azorhizobium caulinodans* ORS571, and *Rhizobium leguminosarum* 3841. The operon is not present in another *Rhizobium* sp., *R. etli* CNF42, or in other members of the family *Rhizobiaceae* for which a complete or draft genome sequence is available. Within the agrobacteria, sequences homologous to *blcC* were detected by Southern analysis in 8 of the 20 isolates that we examined (Table 1). However, of the 12 isolates that lack these sequences, 10 harbor plasmids for which there is either direct or circumstantial evidence for transfer systems controlled by a TraR-dependent quorum-sensing system. While there is a strong association between the presence of the operon and the utilization of GBL, there is no such correlation with a functional TraR-type regulatory system (Table 1). Interestingly, all of the *blc*⁺ agrobacteria are members of the biovar 1 subgroup, although not all biovar 1 strains contain the gene. Similarly, while the two *Rhizobium* isolates examined harbor plasmids with TraR-dependent transfer systems (48, 55), only *R. leguminosarum* contains the *blc* operon (Table 1). Moreover, only this isolate catabolizes GBL. Based upon these observations, we propose that the biologically relevant function of the *blc* operon concerns the catabolism of its butyryl substrates.

Quite clearly, engineered quorum-quenching systems can impact the quorum-sensing systems of targeted bacteria, including pathogens of plants and animals (12, 13, 43, 44, 49, 62). Directed quorum quenching could well be used as an intervention strategy for controlling disease. Moreover, most likely, niches will be found in which quorum-quenching factors produced by one member of a natural consortium influence the signaling systems of other members of the microbial flora (1, 56, 61). It also is conceivable that bacteria that use a quorum-quenching mechanism to modulate their own quorum-sensing system will be found. However, our studies clearly show that the BlcC lactonase produced by *Agrobacterium* isolates does not quench Ti plasmid quorum sensing to a biologically significant level and is not a dedicated component of the Ti plasmid quorum-sensing regulatory system. Any influence of BlcC on conjugative transfer and its regulation by TraR requires the

high-level, continuous expression of the *blc* operon and is transient, of marginal magnitude, and apparently inadvertent. Given that the *blcC* system is the only example described to date, the existence of such lactonases or other acyl-HSL-degrading enzymes that have evolved to moderate quorum-sensing systems is an interesting hypothesis still in need of a supporting biological system.

ACKNOWLEDGMENTS

This work was supported by grant no. R01 GM52465 from the NIH to S.K.F.

We appreciate valuable comments and suggestions from members of the laboratory and from Allen Kerr.

ADDENDUM IN PROOF

In a recent paper, Shephard and Lindow (Appl. Environ. Microbiol. 75:45–53, 2009) reported that *Pseudomonas syringae* B728a produces two potent acylases, HacA and HacB, that degrade acyl-HSL quorum-sensing signals. However, they showed that expression of these two enzymes does not affect phenotypes known to be controlled by the acyl-CHSL-dependent quorum-sensing system of this bacterium.

REFERENCES

- Bauer, W. D., and J. B. Robinson. 2002. Disruption of bacterial quorum sensing by other organisms. *Curr. Opin. Biotechnol.* **13**:234–237.
- Beck von Bodman, S., G. T. Hayman, and S. K. Farrand. 1992. Opine catabolism and conjugal transfer of the nopaline Ti plasmid pTiC58 are coordinately regulated by a single repressor. *Proc. Natl. Acad. Sci. USA* **89**:643–647.
- Carlier, A., R. Chevrot, Y. Dessaux, and D. Faure. 2004. The assimilation of γ -butyrolactone in *Agrobacterium tumefaciens* C58 interferes with the accumulation of the *N*-acyl-homoserine lactone signal. *Mol. Plant-Microbe Interact.* **17**:951–957.
- Cha, C., P. Gao, Y.-C. Chen, P. D. Shaw, and S. K. Farrand. 1998. Production of acyl-homoserine lactone quorum-sensing signals by gram-negative, plant-associated bacteria. *Mol. Plant-Microbe Interact.* **11**:1119–1129.
- Chai, Y., and S. C. Winans. 2004. Site-directed mutagenesis of a LuxR-type quorum-sensing transcription factor: alteration of autoinducer specificity. *Mol. Microbiol.* **51**:765–776.
- Chai, Y., C. S. Tsai, H. Cho, and S. C. Winans. 2007. Reconstitution of the biochemical activities of the AttJ repressor and the AttK, AttL, and AttM catabolic enzymes of *Agrobacterium tumefaciens*. *J. Bacteriol.* **189**:3674–3679.
- Chevrot, R., R. Rosen, E. Haudecoeur, A. Cirou, B. J. Shelp, E. Ron, and D. Faure. 2006. GABA controls the level of quorum-sensing signal in *Agrobacterium tumefaciens*. *Proc. Natl. Acad. Sci. USA* **103**:7460–7464.
- Chilton, M. D., T. C. Currier, S. K. Farrand, A. J. Bendich, M. P. Gordon, and E. W. Nester. 1974. *Agrobacterium tumefaciens* and bacteriophage PS8 DNA not detected in crown gall tumors. *Proc. Natl. Acad. Sci. USA* **71**:3672–3676.
- Dessaux, Y., A. Petit, S. K. Farrand, and P. J. Murphy. 1998. Opines and opine-like molecules involved in plant-*Rhizobiaceae* interactions, p. 173–197. In H. P. Spaink, A. Kondorosi, and P. J. J. Hooykaas (ed.), *The Rhizobiaceae*, molecular biology of model plant-associated bacteria. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Dong, Y.-H., and L.-H. Zhang. 2005. Quorum sensing and quorum-quenching enzymes. *J. Microbiol.* **43**:101–109.
- Dong, Y.-H., L. Y. Wang, and L.-H. Zhang. 2007. Quorum-quenching microbial infections: mechanisms and implications. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **362**:1201–1211.
- Dong, Y.-H., S.-F. Zhang, J.-L. Xu, and L.-H. Zhang. 2004. Insecticidal *Bacillus thuringiensis* silences *Erwinia carotovora* virulence by a new form of microbial antagonism, signal interference. *Appl. Environ. Microbiol.* **70**:954–960.
- Dong, Y.-H., L. H. Wang, J.-L. Xu, H.-D. Zhang, X.-F. Zhang, and L.-H. Zhang. 2001. Quenching quorum-sensing dependent bacterial infection by an *N*-acyl homoserine lactonase. *Nature* **411**:813–817.
- Ellis, J. G., A. Kerr, A. Petit, and J. Tempé. 1982. Conjugal transfer of nopaline and agropine Ti-plasmids—the role of agrocinopines. *Mol. Gen. Genet.* **186**:269–274.
- Farrand, S. K. 1998. Conjugal plasmids and their transfer, p. 199–233. In H. P. Spaink, A. Kondorosi, and P. J. J. Hooykaas (ed.), *The Rhizobiaceae*, molecular biology of model plant-associated bacteria. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Farrand, S. K., C. I. Kado, and C. R. Ireland. 1981. Suppression of the tumorigenicity by the IncW R plasmid pSa in *Agrobacterium tumefaciens*. *Mol. Gen. Genet.* **181**:44–51.

17. Farrand, S. K., Y. Qin, and P. Oger. 2002. Quorum-sensing system of *Agrobacterium* plasmids: analysis and utility. *Methods Enzymol.* **358**:452–484.
18. Fuqua, W. C., and S. C. Winans. 1994. A LuxR-LuxI-type regulatory system activates *Agrobacterium* Ti plasmid conjugal transfer in the presence of a plant tumor metabolite. *J. Bacteriol.* **176**:2796–2806.
19. Fuqua, W. C., and S. C. Winans. 1996. Localization of OccR-activated and TraR-activate promoters that express two ABC-type permeases and the *traR* gene of Ti plasmid pTiR10. *Mol. Microbiol.* **20**:1199–1210.
20. Hamada, S. E., and S. K. Farrand. 1980. Diversity among B6 strains of *Agrobacterium tumefaciens*. *J. Bacteriol.* **141**:1127–1133.
21. Hwang, I., P.-L. Li, L. Zhang, K. R. Piper, D. M. Cook, M. E. Tate, and S. K. Farrand. 1994. TraI, a LuxI homologue, is responsible for production of conjugation factor, the Ti plasmid *N*-acylhomoserine lactone autoinducer. *Proc. Natl. Acad. Sci. USA* **91**:4639–4643.
22. Kalogeraki, V., and S. C. Winans. 1997. Suicide plasmids containing promoterless reporter genes can simultaneously disrupt and create fusions to target genes of diverse bacteria. *Gene* **188**:69–75.
23. Keen, N. T., S. Tamaki, D. Kobayashi, and D. Trollinger. 1988. Improved broad-host-range plasmids for DNA cloning in gram-negative bacteria. *Gene* **70**:191–197.
24. Kerr, A. 1969. Transfer of virulence between isolates of *Agrobacterium*. *Nature* **223**:1175–1176.
25. Kerr, A., P. Manigault, and J. Tempé. 1977. Transfer of virulence in vivo and in vitro by *Agrobacterium*. *Nature* **265**:560–561.
26. Khan, S. R., J. Gaines, R. M. Roop II, and S. K. Farrand. 2008. Broad host range expression vectors with tightly regulated promoters and their use for examining the influence of TraR and TraM expression on Ti plasmid quorum sensing. *Appl. Environ. Microbiol.* **74**:5053–5062.
27. Li, P.-L., and S. K. Farrand. 2000. The replicator of the nopaline-type Ti plasmid pTiC58 is a member of the *repABC* family and is influenced by the TraR-dependent quorum-sensing regulatory system. *J. Bacteriol.* **182**:179–188.
28. Luo, Z.-Q., and S. K. Farrand. 1999. Signal-dependent DNA binding and functional domains of the quorum-sensing activator TraR as identified by repressor activity. *Proc. Natl. Acad. Sci. USA* **96**:9009–9014.
29. Luo, Z.-Q., and S. K. Farrand. 2001. The *Agrobacterium tumefaciens rml* homolog is required for TraR-mediated quorum-dependent activation of Ti plasmid *tra* gene expression. *J. Bacteriol.* **183**:3919–3930.
30. Luo, Z.-Q., T. E. Clemente, and S. K. Farrand. 2001. Construction of a derivative of *Agrobacterium tumefaciens* C58 that does not mutate to tetracycline resistance. *Mol. Plant-Microbe Interact.* **14**:98–103.
31. Luo, Z.-Q., S. Su, and S. K. Farrand. 2003. In situ activation of the quorum-sensing transcription factor TraR by cognate and noncognate acyl-homoserine lactone ligands: kinetics and consequences. *J. Bacteriol.* **185**:5665–5672.
32. Murillo, J., H. Shen, D. Gerhold, A. Sharma, D. A. Cooksey, and N. T. Keen. 1994. Characterization of pPT23B, the plasmid involved in syringolide production by *Pseudomonas syringae* pv. *Tomato* PT23. *Plasmid* **31**:275–287.
33. Nair, G. R., Z. Liu, and A. N. Binns. 2003. Reexamining the role of the accessory plasmid pAtC58 in the virulence of *Agrobacterium tumefaciens* C58. *Plant Physiol.* **133**:989–999.
34. Oger, P., and S. K. Farrand. 2001. Co-evolution of the agrocinopine opines and the agrocinopine-mediated control of TraR, the quorum-sensing activator of the Ti plasmid conjugation system. *Mol. Microbiol.* **41**:1173–1185.
35. Oger, P., and S. K. Farrand. 2002. Two opines control conjugative transfer of an *Agrobacterium* plasmid by regulating expression of separate copies of the quorum-sensing activator gene *traR*. *J. Bacteriol.* **184**:1121–1131.
36. Petit, A., C. David, G. Dahl, J. G. Ellis, P. Guyon, F. Casse-Delbart, and J. Tempé. 1983. Further extension of the opine concept: plasmids in *Agrobacterium rhizogenes* cooperate for opine degradation. *Mol. Gen. Genet.* **190**: 204–214.
37. Piper, K. R., and S. K. Farrand. 2000. Quorum sensing but not autoinduction of Ti plasmid conjugal transfer requires control by the opine regulon and the antiactivator TraM. *J. Bacteriol.* **182**:1080–1088.
38. Piper, K. R., S. Beck von Bodman, and S. K. Farrand. 1993. Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction. *Nature* **362**:448–450.
39. Piper, K. R., S. Beck von Bodman, I. Hwang, and S. K. Farrand. 1999. Hierarchical gene regulatory systems arising from fortuitous gene associations: controlling quorum sensing by the opine regulon in *Agrobacterium*. *Mol. Microbiol.* **32**:1077–1089.
40. Qin, Y., S. Su, and S. K. Farrand. 2007. Molecular basis of transcriptional antiactivation. TraM disrupts the TraR-DNA complex through stepwise interactions. *J. Biol. Chem.* **282**:19979–19991.
41. Ramírez-Romano, M. A., I. Masulis, M. A. Cevallos, V. González, and G. Dávila. 2006. The *Rhizobium etli* sigma70 (SigA) factor recognizes a lax consensus promoter. *Nucleic Acid Res.* **34**:1470–1480.
42. Ream, W. 1989. *Agrobacterium tumefaciens* and interkingdom genetic exchange. *Annu. Rev. Phytopathol.* **27**:583–618.
43. Romero, M., S. P. Diggle, S. Heeb, M. Cámara, and A. Otero. 2008. Quorum quenching activity in *Anabaena* sp. PCC 7120: identification of AiiC, a novel AHL-acylase. *FEMS Microbiol. Lett.* **280**:73–80.
44. Sio, C. F., L. G. Otten, R. H. Cool, S. P. Diggle, P. G. Braun, R. Bos, M. Daykin, M. Cámara, P. Williams, and W. J. Quax. 2006. Quorum quenching by an *N*-acyl-homoserine lactone acylase from *Pseudomonas aeruginosa* PAO1. *Infect. Immun.* **74**:1673–1682.
45. Su, S., S. R. Khan, and S. K. Farrand. 2008. Induction and loss of Ti plasmid conjugative competence in response to the acyl-homoserine lactone quorum-sensing signal. *J. Bacteriol.* **190**:4398–4407.
46. Tempé, J., C. Estrade, and A. Petit. 1978. The biological significance of opines. II. The conjugative activities of the Ti plasmid of *Agrobacterium tumefaciens*, p. 153–160. In M. Ridé (ed.), *Proceedings of the 4th International Conference on Plant Pathogenic Bacteria*. INRA Press, Angers, France.
47. Teplitski, M., J. B. Robinson, and W. D. Bauer. 2000. Plants secrete substances that mimic bacterial *N*-acyl homoserine lactone signal activities and affect population density-dependent behaviors in associated bacteria. *Mol. Plant-Microbe Interact.* **13**:637–648.
48. Tun-Garrido, C., P. Bustos, V. González, and S. Brom. 2003. The conjugative transfer of p42a from *Rhizobium etli* CFN42, which is required for mobilization of the symbiotic plasmid, is regulated by quorum sensing. *J. Bacteriol.* **185**:1681–1692.
49. Ulrich, R. L. 2004. Quorum quenching: enzymatic disruption of *N*-acylhomoserine lactone-mediated bacterial communication in *Burkholderia thailandensis*. *Appl. Environ. Microbiol.* **70**:6173–6180.
50. Unger, L., S. F. Ziegler, G. A. Huffman, V. C. Knauf, R. Peet, L. W. Moore, M. P. Gordon, and E. W. Nester. New class of limited-host-range *Agrobacterium* mega-tumor-inducing plasmids lacking homology to the transferred DNA of a wide-host-range, tumor-inducing plasmid. *J. Bacteriol.* **164**:723–730.
51. Vaudequin-Dransart, V., A. Petit, W. S. Chilton, and Y. Dessaux. The cryptic plasmid of *Agrobacterium tumefaciens* cointegrates with the Ti plasmid and cooperates for opine degradation. *Mol. Plant-Microbe Interact.* **7**:583–591.
52. Wang, C., H.-B. Zhang, L.-H. Wang, and L.-H. Zhang. 2006. Succinic semi-aldehyde couples stress response to quorum-sensing signal decay in *Agrobacterium tumefaciens*. *Mol. Microbiol.* **62**:45–56.
53. Wang, C., H.-B. Zhang, G. Chen, L. Chen, and L.-H. Zhang. 2006. Dual control of quorum sensing by two TraM-type antiactivators in *Agrobacterium tumefaciens* octopine strain A6. *J. Bacteriol.* **188**:2435–2445.
54. White, C. E., and S. C. Winans. 2007. Cell-cell communication in the plant pathogen *Agrobacterium tumefaciens*. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **362**:1135–1148.
55. Wilkinson, A., V. Danino, F. Wisniewski-Dyé, J. K. Lithgow, and J. A. Downie. 2002. *N*-Acyl-homoserine lactone inhibition of rhizobial growth is mediated by two quorum-sensing genes that regulate plasmid transfer. *J. Bacteriol.* **184**:4510–4519.
56. Wopperer, J., S. T. Cardona, B. Huber, C. A. Jacobi, M. A. Valvano, and L. Eberl. 2006. A quorum-quenching approach to investigate the conservation of quorum-sensing-regulated functions within the *Burkholderia cepacia* complex. *Appl. Environ. Microbiol.* **72**:1579–1587.
57. Zhang, H.-B., K.-H. Wang, and L.-H. Zhang. 2002. Genetic control of quorum-sensing signal turnover in *Agrobacterium tumefaciens*. *Proc. Natl. Acad. Sci. USA* **99**:4638–4643.
58. Zhang, H.-B., C. Wang, and L.-H. Zhang. 2004. The quorumone degradation system of *Agrobacterium tumefaciens* is regulated by starvation signal and stress alarmone(p) ppGpp. *Mol. Microbiol.* **52**:1389–1401.
59. Zhang, L., and A. Kerr. 1991. A diffusible compound can enhance transfer of the Ti plasmid in *Agrobacterium tumefaciens*. *J. Bacteriol.* **173**:1867–1872.
60. Zhang, L., P. J. Murphy, A. Kerr, and M. E. Tate. 1993. *Agrobacterium* conjugation and gene regulation by *N*-acyl-homoserine lactones. *Nature* **362**:446–448.
61. Zhang, L.-H. 2003. Quorum-quenching and proactive host defense. *Trends Plant Sci.* **8**:238–244.
62. Zhu, H., Y. L. Shen, D. Z. Wei, and J. W. Zhu. 2008. Inhibition of quorum sensing in *Serratia marcescens* H30 by molecular regulation. *Curr. Microbiol.* **56**:645–650.
63. Zhu, J., and S. C. Winans. 1999. Autoinducer binding by the quorum-sensing regulator TraR increases affinity for target promoters in vitro and decreases TraR turnover rates in whole cells. *Proc. Natl. Acad. Sci. USA* **98**:1507–1512.