

# Genome Sequence of *Fulvimarina pelagi* HTCC2506<sup>T</sup>, a Mn(II)-Oxidizing Alphaproteobacterium Possessing an Aerobic Anoxygenic Photosynthetic Gene Cluster and Xanthorhodopsin<sup>∇</sup>

Ilnam Kang,<sup>1</sup> Hyun-Myung Oh,<sup>1</sup> Seung-Il Lim,<sup>1</sup> Steve Ferreira,<sup>2</sup>  
 Stephen J. Giovannoni,<sup>3</sup> and Jang-Cheon Cho<sup>1\*</sup>

Division of Biology and Ocean Sciences, Inha University, Incheon 402-751, Republic of Korea<sup>1</sup>; J. Craig Venter Institute, 9704 Medical Center Drive, Rockville, Maryland 20850<sup>2</sup>; and Department of Microbiology, Oregon State University, Corvallis, Oregon 97331<sup>3</sup>

Received 30 June 2010/Accepted 7 July 2010

***Fulvimarina pelagi* is a Mn(II)-oxidizing marine heterotrophic bacterium in the order Rhizobiales. Here we announce the draft genome sequence of *F. pelagi* HTCC2506<sup>T</sup>, which was isolated from the Sargasso Sea by using dilution-to-extinction culturing. The genome sequence contained a xanthorhodopsin gene as well as a photosynthetic gene cluster, which suggests the coexistence of two different phototrophic mechanisms in a single microorganism.**

Besides being mediated by the well-known process of aerobic oxygenic photosynthesis, utilization of light energy in the marine carbon and nutrient cycling processes is usually mediated by aerobic anoxygenic phototrophic bacteria (AAPB) or prokaryotes containing microbial rhodopsin family proteins (18). AAPB comprise about 10% of total microbial cells in the euphotic zone of diverse marine regimes, with alpha-, beta-, and gammaproteobacteria as major constituents (8, 10). Rhodopsin family proteins, including bacteriorhodopsin, proteorhodopsin, and xanthorhodopsin (XR), have been shown to exist in 7 to 70% of marine prokaryotes and contribute phototrophy in diverse phylogenetic groups such as *Flavobacteria*, *Proteobacteria*, and *Archaea* (15, 19). However, no single microorganism has been reported to contain the genes for both aerobic anoxygenic phototrophy (AAnP) and rhodopsin family proteins.

*F. pelagi* HTCC2506<sup>T</sup> was cultivated through a dilution-to-extinction approach (3) from the western Sargasso Sea and identified as a novel genus and species in the order *Rhizobiales* of the *Alphaproteobacteria* by polyphasic taxonomy (2). Later, three *F. pelagi* strains, including HTCC2506<sup>T</sup>, were shown to have Mn(II)-oxidizing activity (1). The draft genome sequence of HTCC2506<sup>T</sup> was determined by shotgun sequencing at the J. Craig Venter Institute as part of the Moore Foundation Microbial Genome Sequencing Project and analyzed by the GenDB annotation program (17) at the Center for Genome Research and Biocomputing at Oregon State University and the Joint Genome Institute IMG system (<http://img.jgi.doe.gov>) (14).

The draft genome was 3,802,689 bp in length, distributed in 20 contigs with 61.2% G+C content, and contained 3,754 pro-

tein-coding genes, three copies of 16S-23S-5S rRNA genes, and 54 tRNA genes. Remarkably, HTCC2506<sup>T</sup> possessed XR and a complete gene set for AAnP together. A gene cluster encoding the AAnP apparatus was composed of *bchIDO-crtCDF-bchCXYZ-pufBALMC-puhE-acsF-puhCBA-lhaA-bchMLHBNF-aerR-ppsR-ubiA-pucC-bchP-hemT*. Mu-like prophage sequences were located closely adjacent to the AAnP gene cluster, which might imply the possibility of the lateral gene transfer of the AAnP gene cluster. The XR gene was followed by *blh*, a gene involved in retinal biosynthesis (16). To our knowledge, this is the first report of a microbe possessing both an AAnP apparatus and a rhodopsin family protein, although the two gene sets in HTCC2506<sup>T</sup> had been separately reported to be present (5, 13). The HTCC2506<sup>T</sup> genome also encoded a bacteriophytochrome (6, 7), a light-regulated signal transduction histidine kinase. Overall, the existence of a diverse repertoire of genes for sensing or harvesting of light energy implicates the importance of phototrophic metabolism for HTCC2506<sup>T</sup>.

In addition to genes for phototrophy, the genome contained several genes with diverse metabolic potential. A lithotrophic mode of energy acquisition was predicted from the presence of the form II *coxSLM* genes, which encode aerobic-type carbon monoxide dehydrogenase (9). As expected from the Mn(II)-oxidizing activity of HTCC2506<sup>T</sup>, the genome also encoded a multicopper oxidase (MCO) enzyme, suggesting a potential lithotrophy. In terms of carbon assimilation, genes encoding RuBisCO and phosphoribulokinase of the Calvin-Benson-Bassham cycle (11) were predicted, suggesting the possibility of autotrophic CO<sub>2</sub> fixation in HTCC2506<sup>T</sup>. Serine transhydroxymethylase for formaldehyde assimilation (12) was predicted. A gene encoding dimethylsulfoniopropionate (DMSP) lyase, which conveys the production of dimethylsulfide from DMSP (4), was also found in the genome.

Although HTCC2506<sup>T</sup> was originally isolated as a chemoheterotroph (2), the genome sequence clearly shows the phototrophic potential of this bacterium. This finding, combined

\* Corresponding author. Mailing address: Division of Biology and Ocean Sciences, Inha University, Incheon 402-751, Republic of Korea. Phone: 82-32-860-7711. Fax: 82-32-232-0541. E-mail: chojc@inha.ac.kr.

<sup>∇</sup> Published ahead of print on 16 July 2010.

with the prediction of many genes related to lithotrophy, carbon fixation,  $C_1$  compound assimilation, and DMSP lysis, suggests that *F. pelagi* HTCC2506<sup>T</sup> may use a wide range of potential metabolic functions to survive in the marine euphotic environment.

**Nucleotide sequence accession number.** The draft genome sequence of HTCC2506<sup>T</sup> was deposited in GenBank under the accession number AATP00000000. The GenDB-generated data can be also accessed at Marine Microbial Genomics at Oregon State University (<http://bioinfo.cgrb.oregonstate.edu/microbes/>).

The initial phase of sequencing, assembly, and annotation efforts were carried out at the OSU High Throughput Culturing Laboratory with support from a Gordon and Betty Moore Foundation investigator award and the Gordon and Betty Moore Foundation Marine Microbial Sequencing Project. Subsequent data analysis was conducted at Inha University and also supported by the 21C Frontier Program of Microbial Genomics and Applications.

We thank Justin Johnson, Jason Miller, Granger Sutton, Saul A. Kravitz, Robert Friedman, and Yu-Hui Rogers for their technical and scientific support of the Gordon and Betty Moore Foundation-funded project.

#### REFERENCES

- Anderson, C. R., G. J. Dick, M.-L. Chu, J.-C. Cho, R. E. Davis, S. L. Bräuer, and B. M. Tebo. 2009. *Aurantimonas manganoxydans*, sp. nov. and *Aurantimonas litoralis*, sp. nov.: Mn(II) oxidizing representatives of a globally distributed clade of alpha-Proteobacteria from the order Rhizobiales. *Geomicrobiol. J.* **26**:189–198.
- Cho, J.-C., and S. J. Giovannoni. 2003. *Fulvimarina pelagi* gen. nov., sp. nov., a marine bacterium that forms a deep evolutionary lineage of descent in the order 'Rhizobiales.' *Int. J. Syst. Evol. Microbiol.* **53**:1853–1859.
- Connon, S. A., and S. J. Giovannoni. 2002. High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Appl. Environ. Microbiol.* **68**:3878–3885.
- Curson, A. R. J., R. Rogers, J. D. Todd, C. A. Brearley, and A. W. B. Johnston. 2008. Molecular genetic analysis of a dimethylsulfoniopropionate lyase that liberates the climate-changing gas dimethylsulfide in several marine  $\alpha$ -proteobacteria and *Rhodobacter sphaeroides*. *Environ. Microbiol.* **10**:757–767.
- Giovannoni, S. J., D. H. Hayakawa, H. J. Tripp, U. Stingl, S. A. Givan, J.-C. Cho, H.-M. Oh, J. B. Kitner, K. L. Vergin, and M. S. Rappé. 2008. The small genome of an abundant coastal ocean methylophile. *Environ. Microbiol.* **10**:1771–1782.
- Giraud, E., and A. Verméglio. 2008. Bacteriophytochromes in anoxygenic photosynthetic bacteria. *Photosynth. Res.* **97**:141–153.
- Jaubert, M., L. Vuillet, L. Hannibal, J.-M. Adriano, J. Fardoux, P. Bouyer, K. Bonaldi, D. Fleischman, E. Giraud, and A. Verméglio. 2008. Control of peripheral light-harvesting complex synthesis by a bacteriophytochrome in the aerobic photosynthetic bacterium *Bradyrhizobium* strain BTai1. *J. Bacteriol.* **190**:5824–5831.
- Jiao, N., Y. Zhang, Y. Zeng, N. Hong, R. Liu, F. Chen, and P. Wang. 2007. Distinct distribution pattern of abundance and diversity of aerobic anoxygenic phototrophic bacteria in the global ocean. *Environ. Microbiol.* **9**:3091–3099.
- King, G. M., and C. F. Weber. 2007. Distribution, diversity and ecology of aerobic CO-oxidizing bacteria. *Nat. Rev. Microbiol.* **5**:107–118.
- Kolber, Z. S., F. G. Plumley, A. S. Lang, J. T. Beatty, R. E. Blankenship, C. L. VanDover, C. Vetriani, M. Koblizek, C. Rathgeber, and P. G. Falkowski. 2001. Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* **292**:2492–2495.
- Kusian, B., and B. Bowien. 1997. Organization and regulation of *ccb* CO<sub>2</sub> assimilation genes in autotrophic bacteria. *FEMS Microbiol. Rev.* **21**:135–155.
- Lidstrom, M. 2006. Aerobic methylotrophic prokaryotes, p. 618–634. *In* M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer, and E. Stackebrandt (ed.), *The prokaryotes*, vol. 2. Springer-Verlag, New York, NY.
- Liotenberg, S., A.-S. Steunou, M. Picaud, F. Reiss-Husson, C. Astier, and S. Ouchane. 2008. Organization and expression of photosynthesis genes and operons in anoxygenic photosynthetic proteobacteria. *Environ. Microbiol.* **10**:2267–2276.
- Markowitz, V. M., I.-M. A. Chen, K. Palaniappan, K. Chu, E. Szeto, Y. Grechkin, A. Ratner, I. Anderson, A. Lykidis, K. Mavromatis, N. N. Ivanova, and N. C. Kyrpides. 2010. The integrated microbial genomes system: an expanding comparative analysis resource. *Nucleic Acids Res.* **38**:D382–D390.
- Martinez, A., A. S. Bradley, J. R. Waldbauer, R. E. Summons, and E. F. DeLong. 2007. Proteorhodopsin photosystem gene expression enables photophosphorylation in a heterologous host. *Proc. Nat. Acad. Sci. U. S. A.* **104**:5590–5595.
- McCarren, J., and E. F. DeLong. 2007. Proteorhodopsin photosystem gene clusters exhibit co-evolutionary trends and shared ancestry among diverse marine microbial phyla. *Environ. Microbiol.* **9**:846–858.
- Meyer, F., A. Goesmann, A. McHardy, D. Bartels, T. Bekel, J. Clausen, J. Kalinowski, B. Linke, O. Rupp, R. Giegerich, and A. PuEhler. 2003. GenDB—an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res.* **31**:2187–2195.
- Moran, M. A., and W. L. Miller. 2007. Resourceful heterotrophs make the most of light in the coastal ocean. *Nat. Rev. Microbiol.* **5**:792–800.
- Riedel, T., J. Tomasch, I. Buchholz, J. Jacobs, M. Kollenberg, G. Gerdt, A. Wichels, T. Brinkhoff, H. Cypionka, and I. Wagner-Dobler. 2010. Constitutive expression of the proteorhodopsin gene by a *Flavobacterium* strain representative of the proteorhodopsin-producing microbial community in the North Sea. *Appl. Environ. Microbiol.* **76**:3187–3197.