

DosS Responds to a Reduced Electron Transport System To Induce the *Mycobacterium tuberculosis* DosR Regulon[∇]

Ryan W. Honaker,¹ Rakesh K. Dhiman,² Prabakaran Narayanasamy,²
 Dean C. Crick,² and Martin I. Voskuil^{1*}

University of Colorado Denver, School of Medicine, Department of Microbiology, P18-9115, 12800 East 19th Avenue, P.O. Box 6511, Aurora, Colorado 80045,¹ and Colorado State University, Mycobacteria Research Laboratories, Department of Microbiology, Immunology and Pathology, 1682 Campus Delivery, Fort Collins, Colorado 80523²

Received 18 August 2010/Accepted 1 October 2010

The DosR regulon in *Mycobacterium tuberculosis* is involved in respiration-limiting conditions, its induction is controlled by two histidine kinases, DosS and DosT, and recent experimental evidence indicates DosS senses either molecular oxygen or a redox change. Under aerobic conditions, induction of the DosR regulon by DosS, but not DosT, was observed after the addition of ascorbate, a powerful cytochrome *c* reductant, demonstrating that DosS responds to a redox signal even in the presence of high oxygen tension. During hypoxic conditions, regulon induction was attenuated by treatment with compounds that occluded electron flow into the menaquinone pool or decreased the size of the menaquinone pool itself. Increased regulon expression during hypoxia was observed when exogenous menaquinone was added, demonstrating that the menaquinone pool is a limiting factor in regulon induction. Taken together, these data demonstrate that a reduced menaquinone pool directly or indirectly triggers induction of the DosR regulon via DosS. Biochemical analysis of menaquinones upon entry into hypoxic/anaerobic conditions demonstrated the disappearance of the unsaturated species and low-level maintenance of the mono-saturated menaquinone. Relative to the unsaturated form, an analog of the saturated form is better able to induce signaling via DosS and rescue inhibition of menaquinone synthesis and is less toxic. The menaquinone pool is central to the electron transport system (ETS) and therefore provides a mechanistic link between the respiratory state of the bacilli and DosS signaling. Although this report demonstrates that DosS responds to a reduced ETS, it does not rule out a role for oxygen in silencing signaling.

Mycobacteria are strict aerobes, but *Mycobacterium tuberculosis* encounters microaerobic to anaerobic environments during the course of infection. Oxygen-limited microenvironments occur in mature granulomas, which are known to be avascular, inflammatory, and necrotic (1, 5, 20, 40, 56). Recent reports have detected mycobacterial DNA in visibly normal lung tissue (17), as well as adipose tissue (32). Interestingly, adipose tissue has been associated with hypoxia as well as the presence of nitric oxide (NO), both of which inhibit respiration (38, 51, 57, 60). Further, *M. tuberculosis* is able to survive for long periods of time in a nonreplicating and nonrespiring state (15, 55).

The DosR regulon is expressed in response to hypoxia, NO, and carbon monoxide (CO) and is thought to be important for early adaptation to these stimuli as well as long-term survival in the host (2, 25, 26, 46, 47, 53, 58). Despite induction by other gases, the presence of oxygen itself inhibits induction of the regulon (26, 41, 43, 48). The DosR regulon is regulated by the response regulator DosR (DevR; Rv3133c) and was recently shown to be positively regulated by PhoP (Rv0757) to a basal level during aerobic growth (16). DosR is activated by two sensor histidine kinases, DosT (Rv2027c) and DosS (DevS; Rv3132c). The activation of DosR occurs through autophos-

phorylation of either sensor followed by transfer of the phosphate to DosR (39).

DosS and DosT are modular proteins that contain C-terminal histidine kinase domains, an ATP binding domain, and two tandem GAF (cGMP, adenylyl cyclase, and FhlA) domains (GAF A and GAF B). GAF domains are small molecule-binding regulatory domains found throughout prokaryotes and eukaryotes. Within these domains, both DosT and DosS bind heme, and in DosS the location of heme binding within the GAF A domain is known. The heme moiety itself binds the divalent gases (NO, CO, and O₂) responsible for modulating activity of the sensor (6, 22, 26, 41, 43, 48), and specifics of the hydrogen bond network required for ligand discrimination have been described recently (59).

The current biochemical literature on DosS is in disagreement regarding its signal for activation. One body of work indicates that DosS directly binds oxygen, suppressing the sensor's activity. In this way DosS functions as an oxygen sensor: when oxygen becomes limited and is no longer bound, the sensor autophosphorylates. Several hydrogen bond specifics of the oxygen interaction have been described, and the different hydrogen bond networks involved in discrimination between NO/CO and O₂ have been delineated, as have aspects of the kinetics of oxygen binding as well as bond affinity (21, 22, 43, 48, 59). Two recent reports proposed a second model for DosT/S signaling, wherein DosS is a redox sensor (6, 26). This conclusion was based in part on *in vitro* oxidation rates and the stability of the oxy complex reported by these groups. Further evidence was derived from the interaction of DosS with cy-

* Corresponding author. Mailing address: University of Colorado Denver, School of Medicine, Department of Microbiology, P18-9115, 12800 East 19th Avenue, P.O. Box 6511, Aurora, CO 80045. Phone: (303) 724-4219. Fax: (303) 724-4226. E-mail: martin.voskuil@ucdenver.edu.

[∇] Published ahead of print on 15 October 2010.

nide as well as anaerobic autokinase activity assays (26). However, similar experiments resulting in contrary data have been published in two other reports by proponents of oxygen as the direct signal (21, 48). Support for the redox sensor hypothesis comes from data that demonstrate that under aerobic conditions DosS exists in an inactive Fe³⁺ form as a result of its rapid oxidation (but not direct binding) by O₂ (6, 26); however, when DosS is reduced to Fe²⁺, its autokinase activity is greatly increased; thus, it uses its oxidation state as a signal for activation (6, 26). Lastly, it was demonstrated that flavin nucleotides are capable of directly reducing the heme iron of DosS (6).

Despite these various biochemical reports, little work has been conducted *in vivo*. In this report we investigate the stimulus involved in DosS signaling within bacterial cells.

MATERIALS AND METHODS

Growth conditions and strains. All cultures were grown in Dubos Tween albumin medium (Becton Dickinson). For anaerobic container experiments, BD BBL GasPak Plus envelopes were used to achieve anaerobic conditions, and anaerobic indicators were used to ensure anaerobiosis. Vented culture flasks were placed in sealed GasPak containers (BD, Franklin Lakes, NJ) and stirred. All experiments had a starting culture optical density at 600 nm (OD₆₀₀) of 0.2. The anaerobic dormancy model was performed as previously described (18). CFU were calculated as previously described (19). Additives were given at day 12 of the anaerobic dormancy model, 6 days after anaerobiosis (29). Tubes were opened for the addition of treatments in an anaerobic chamber to maintain the anaerobic status of the media and then further incubated for 8 days, after which they were plated. Additive concentrations were the same as those used for real-time analysis. Statistical significance was determined by a paired two-tailed Student *t* test. For direct oxygen concentration measurements, a 2-mm O₂ sensor and TBR4100 free radical analyzer (World Precision Instruments) were used. Culture conditions were identical to those used for the experimental data, with the exception of the strain, which was an unmarked derivative of 6030 (42). Data represent the percent oxygen of untreated media and are the averages of four experiments. Oxyrase was used according to the manufacturer's guidelines.

Culture chemical treatments. For aerobic experiments, treatments were added for 1 h. For hypoxic conditions, cultures were placed in anaerobic GasPak containers for 4 h. Treatments were then added to the media anaerobically (in an anaerobic chamber) and incubated for two additional hours for all treatments, with the exception of chlorpromazine and CSU-20. For these experiments, cultures were aerobically pretreated for 4 h and then placed in GasPak containers for four additional hours. Four hours in an anaerobic container was previously demonstrated to create a hypoxic medium environment (18). Treatment concentrations were as follows, with their respective vehicle in parentheses: 0.1 mM NO consisting of NO donor diethylenetriamine/NO adduct (water), 20 mM ascorbate (no vehicle; powder alone added), 400 μM vitamin K1 (ethanol), 400 μM vitamin K2 (DMSO), 50 μg/ml chlorpromazine (ethanol), 40 μM CSU-20 (ethanol), 1 mM 2,4-dinitrophenol (DMSO), and 10 mM nitrate (water). Vehicles for each treatment were added separately as controls. All chemicals were from Sigma, with the exception of CSU-20, which was synthesized as previously described (27).

Real-time PCR. RNA was extracted, and cDNA prepared and labeled as previously described (53). To help stabilize RNA and avoid harvesting-induced artifacts, a solution of 5 M guanidine thiocyanate, 0.5% sodium *N*-lauryl sarcosine, 50 mM tri-sodium citrate, and 0.1 M 2-mercaptoethanol was used to harvest cultures at a 1:1 ratio with bacterial culture (44). Reverse transcription and quantitative real-time PCR were performed with gene-specific primers. Fluorescent probes were designed with the program FastPCR (23) and synthesized by Integrated DNA technologies. Primer sequences were as follows: Rv0239 forward (TGATTCGAACGCAAGTCCAGCTC), reverse (AGATCCG CACCATGTGCTCCA), and probe (TCGCGCACGAGCACGAAATG); Rv2626c forward (CCGCGACATTGTGATCAAAG), reverse (GCTCTGAGA TGACCGGAACAC), and probe (CGAACGCAAGCATCCAGGAGATGC); and Rv1738 forward (CACTGGACCGTCGACATATCG), reverse (CGGTCCG GCCGGATTG), and probe (CCAACGCAGCCGTGCCTTCG). Real-time PCR was performed on the Roche LightCycler 480. A reverse transcriptase negative reaction was used to account for residual DNA, and copy numbers were

normalized to the level of Rv0239 transcript (18). These copy numbers were then used to determine the copy number per bacterium.

Microarrays. Oligonucleotide-based microarrays were printed, microarrays were processed, RNA was extracted, and cDNA was prepared and labeled as previously reported (18, 53). Microarrays were scanned with a GenePix 4000b scanner (Molecular Devices), and spot intensities were obtained by using GenePix Pro 6.0 (Molecular Devices). Aerobic cultures for microarray analysis were treated for 1 h with 20 mM ascorbate; the reference was untreated aerobic culture. Experiments were repeated in triplicate for the Δ *dosT* mutant and in duplicate for the Δ *dosS* mutant. Data analysis was performed as previously described (53). Significance Analysis of Microarrays (SAM) (<http://www-stat.stanford.edu/~tibs/SAM/>) was used to determine statistically significant regulated genes in the Δ *dosT* mutant experiments (52). Genes in the Δ *dosT* strain included in Table 1 conformed to highly stringent selection criteria, exhibiting at least a 4-fold induction ratio and a corresponding SAM false discovery *q* value of zero. Microarray data are available at the NCBI Gene Expression Omnibus website (<http://www.ncbi.nlm.nih.gov/geo/>).

Menquinone analysis. Bacterial cultures were concentrated by centrifugation, and 6 ml chloroform-methanol (2:1) containing 5 μg of vitamin K1 (Sigma) were added; the sample was stored in the dark at -20°C until processed. Samples were then placed on a rotating mixer (Glas-Col) for 3 h, and 1 ml of water was added to generate a phase separation. The lower, organic phase was washed once with water-methanol-chloroform (48:47:3 [vol/vol/vol]) and transferred to new tubes. The solvent was evaporated under a stream of N₂, and the sample was dissolved in chloroform, loaded on a silicic acid column, and eluted with chloroform to remove the polar lipids. The elution solvent was removed from the resulting material under a N₂ stream, and the sample was dissolved in ethanol in a volume normalized to the original culture OD₆₀₀ and subjected to high-pressure liquid chromatography (HPLC) mass spectrometry (MS) on an HP1100 series HPLC connected to a 2000 Finnigan LCQ-DUO ion trap mass spectrometer with an APPI interface. HPLC separation was achieved using an XBridge column (Waters) and a gradient running from 100% methanol, for 2 min, to methanol-isopropanol (50:50 [vol/vol]) over 50 min at 0.4 ml min⁻¹ with a column temperature of 40°C. Eluted molecules were subjected to positive ion MS using APPI as the ionization interface. Capillary temperature was 150°C, and APPI vaporizer temperature was 350°C.

RESULTS

DosR regulon genes respond to a reduced electron transport system. To examine DosR regulon induction, two representative regulon genes, Rv2626c and Rv1738, were selected for real-time PCR analysis. A *dosT* mutant was used for all experiments, except when noted, to ensure all regulation observed was solely from DosS. Ascorbate, a reductant of cytochrome *c*, was added to elucidate the role in DosR signaling via a reduced electron transport system (ETS). Robust induction of the two DosR regulon genes was observed when ascorbate was added to the Δ *dosT* strain maintained under aerobic conditions; however, no induction of the genes was observed with a Δ *dosS* strain (Fig. 1A). This indicated that a reductant of the ETS is capable of inducing the DosR regulon even in the presence of oxygen via DosS, but not DosT. When ascorbate was added to hypoxic cultures, increased induction of the regulon genes was also observed (Fig. 1C). However, the induction ratio was attenuated compared to that observed during aerobic conditions due to normal hypoxic induction of the regulon in the absence of ascorbate. The ETS is relatively more reduced during hypoxic conditions than during aerobic conditions due to the lack of oxygen and resulting diminution of aerobic respiration. To ensure observed signaling was not the result of hypoxia caused by the addition of ascorbate, oxygen levels were directly measured with an oxygen probe; ascorbate did not alter medium oxygen levels (Fig. 2). Oxyrase was used as a control for decreased oxygen levels, and a reduced amount of oxygen was observed. Furthermore, the lack of induction via

TABLE 1. Induction ratios for $\Delta dosT$ and $\Delta dosS$

Gene no. ^a	Gene	Induction ratio		Gene product
		$\Delta dosT$ strain	$\Delta dosS$ strain	
DosR				
Rv0079		27.6	0.4	Hypothetical protein
Rv0080		12.5	1.2	Conserved hypothetical protein
Rv0081		3.1	1.0	Probable transcriptional regulatory protein
Rv0082		7.8	1.4	Probable oxidoreductase
Rv0569		14.9	0.8	Conserved hypothetical protein
Rv0570	<i>nrdZ</i>	5.9	0.5	Probable ribonucleoside-diphosphate reductase
Rv0572c		6.7	0.7	Hypothetical protein
Rv1733c		5.8	0.7	Probable conserved transmembrane protein
Rv1736c	<i>narX</i>	11.8	0.9	Probable nitrate reductase
Rv1737c	<i>narK2</i>	18.6	0.5	Possible nitrate/nitrite transporter
Rv1738		32.4	0.7	Conserved hypothetical protein
Rv1813c		7.7	4.3	Conserved hypothetical protein
Rv1996		6.4	0.7	Conserved hypothetical protein
Rv2003c		6.6	0.7	Conserved hypothetical protein
Rv2004c		7.1	0.9	Conserved hypothetical protein
Rv2005c		12.1	0.5	Conserved hypothetical protein
Rv2007c	<i>fdxA</i>	82.0	0.9	Probable ferredoxin
Rv2028c		13.2	0.7	Conserved hypothetical protein
Rv2029c	<i>pfkB</i>	20.8	0.9	Probable phosphofructokinase
Rv2030c		67.1	0.8	Conserved hypothetical protein
Rv2031c	<i>hspX</i>	76.2	0.5	14-kD antigen, heat shock protein Hsp20 family
Rv2032	<i>acg</i>	30.8	0.5	Conserved hypothetical protein
Rv2623		1.7	0.7	Conserved hypothetical protein
Rv2624c		6.2	1.2	Conserved hypothetical protein
Rv2626c		36.5	0.8	Conserved hypothetical protein
Rv2627c		16.3	0.3	Conserved hypothetical protein
Rv2628		8.9	0.8	Hypothetical protein
Rv2629		8.1	0.9	Conserved hypothetical protein
Rv2630		4.2	1.6	Hypothetical protein
Rv3127		44.3	1.2	Conserved hypothetical protein
Rv3130c	<i>tgsl</i>	45.7	1.7	Triacylglycerol synthase
Rv3131		41.0	0.9	Conserved hypothetical protein
Rv3132c	<i>dosS</i>	16.7	0.4	Sensor histidine kinase
Rv3133c	<i>dosR</i>	11.1	0.6	Two-component response regulator
Rv3134c		26.0	0.7	Conserved hypothetical protein
Non-DosR				
Rv0467	<i>icl</i>	7.6	4.5	Isocitrate lyase
Rv0791c		5.0	3.5	Conserved hypothetical protein
Rv0792c		4.4	3.2	Probable GNTR family regulatory protein
Rv0793		4.9	2.6	Conserved hypothetical protein
Rv2025c		7.4	2.7	Possible conserved membrane protein
Rv2108	<i>PPE36</i>	4.1	2.4	PPE family protein
Rv2172c		4.4	2.2	Conserved hypothetical protein
Rv3083		27.7	20.4	Probable monooxygenase (hydroxylase)
Rv3084	<i>lipR</i>	10.3	9.9	Probable acetyl-hydrolase/esterase
Rv3086	<i>adhD</i>	7.3	8.8	Probable zinc-type alcohol dehydrogenase
Rv3087		6.7	4.0	Possible triacylglycerol synthase
Rv3088		6.9	13.4	Possible triacylglycerol synthase
Rv3089	<i>fadD13</i>	7.1	4.3	Probable fatty-acid coenzyme A (CoA) ligase
Rv3841	<i>bfrB</i>	17.7	15.3	Possible bacterioferritin

^a Genes in the $\Delta dosT$ strain that were included in the table had an induction ratio of at least 4-fold and a corresponding SAM *q* value of 0.

DosT, a known oxygen sensor, indicates that oxygen levels were not responsible for induction via DosS by ascorbate.

Additional experiments were conducted to further delineate the evident role of the ETS in DosS signaling. Chlorpromazine (CPZ), a type II oxidoreductase inhibitor, and CSU-20, a menaquinone synthesis inhibitor (27), were added to hypoxic cultures (Fig. 1C). Inhibition of oxidoreductases attenuates electron flow through the ETS, as does decreasing the size of the menaquinone pool by inhibiting menaquinone synthesis. The addition of both CPZ and CSU-20 to hypoxic cultures

resulted in a marked reduction in signaling, and no effect on aerobic cultures was observed (Fig. 1B and C). When nitrate was added to hypoxic cultures, a corresponding decrease in signaling occurred (Fig. 1C). This was expected, as nitrate addition should result in oxidation of the menaquinone pool because of the ability of nitrate to function as an alternative electron acceptor. These findings corroborate the ascorbate data by correlating a decrease in electron flow into and through the menaquinone pool of the ETS (CPZ and CSU-20) or a decrease in reduction of the menaquinone pool (nitrate)




FIG. 1. Induction of DosR regulon genes during aerobic cultures (A and B) and hypoxic conditions (C). The data shown represent fold induction of untreated or vehicle-treated cultures of the transcript copy numbers of two DosR regulon genes, Rv2626c and Rv1738, normalized to levels of Rv0239. All experiments were performed with a $\Delta dosT$ strain except when noted.

with a decline in signaling. They also imply that either the menaquinone pool itself or a reduced component of the ETS is the signal monitored by DosS.

To test this further, vitamin K2, a functional unsaturated menaquinone analog, was added to cultures. Aerobically no effect was observed, but hypoxically an increased induction of the DosR regulon was observed, indicating that menaquinone levels were limiting for regulon induction during hypoxic conditions (Fig. 1B and C). When CSU-20 was added simultaneously with ascorbate to again correlate menaquinone levels and signaling, a decrease in signaling was observed compared to ascorbate-treated cultures, as expected (Fig. 1A).

Hypoxia could diminish overall proton motive force (PMF). Decreased PMF may also occur as a result of excess electrons




FIG. 2. Oxygen levels of ascorbate-treated media. To ensure that observed transcriptional findings were not the result of diminished oxygen levels from the addition of ascorbate, an oxygen sensor was used for direct measurement of cultures. Results represent percent oxygen detected of untreated media. Bacterial culture is white, bacterial culture plus ascorbate is black, and media treated with Oxyrase is striped.

in the ETS from the addition of ascorbate. To ensure that a decreased PMF is not the signal for induction of the DosR regulon, a protonophore, 2,4-dinitrophenol (DNP), was added to aerobic cultures. Addition of DNP resulted in no induction of the DosR regulon genes (shown in Fig. 1B and described previously in reference 3). Finally, nitric oxide was added to both aerobic and hypoxic conditions as a positive control for DosR regulon induction (53), and a 10-fold induction was observed in both conditions.

Microarray analysis of ascorbate-treated aerobic cultures. To confirm the real-time PCR data as well as determine the overall effect of ascorbate on aerobic cultures, microarray analysis was performed on aerobic $\Delta dosT$ and $\Delta dosS$ mutant cultures treated with ascorbate. In the $\Delta dosT$ strain, 49 genes were significantly induced and had induction ratios greater than 4-fold (Table 1). Of these 49 genes, 35 were part of the DosR regulon and had an average induction ratio of 21-fold. Contrarily, the $\Delta dosS$ strain had an average induction ratio of the DosR genes of 0.9-fold. Fourteen additional genes were induced in both mutants.

Change in menaquinone pool during entry into anaerobiosis. To elucidate the kinetics of the various species of menaquinone during this model as well as to show a correlation between menaquinone levels and signaling, the status of the menaquinone pool at various points during adaptation to anaerobic dormancy was next examined. The presence of oxygen in this system was as follows: time zero represents aerobic growth, day 4 represents hypoxic growth, and by day 6 the cultures were anaerobic (29). A rapid decrease in both species, MK-9(II-H₂) as well as MK-9 (see Fig. 5 for structure references), was observed by the time the cultures became hypoxic (Fig. 3 and 4). However, at that point, MK-9(II-H₂) ceased to decrease as rapidly and reached a fairly stable concentration, while MK-9 declined to below the limit of detection.

To examine the possible differential effect of menaquinone saturation status on DosS signaling, vitamin K1 and vitamin K2 were added to cultures (Fig. 1B). These compounds differ in




FIG. 3. Representative mass spectral analysis of neutral lipids extracted from *M. tuberculosis*. (A) Total ion chromatogram. (B) Extracted ion chromatogram for 787.6315 Da, which corresponds to the molecular ion of MK-9(II-H₂). (C) Extracted ion chromatogram for 785.6158 Da, which corresponds to the molecular ion of MK-9. Arrows indicate the peaks assigned to MK-9 (B) and MK-9(II-H₂) (C).

the saturation status from the second isoprenoid group (from the head group) to the end (Fig. 5); vitamin K1 is saturated, and vitamin K2 is unsaturated. Therefore, they were added as surrogates for MK-9 and MK-9(II-H₂), respectively, because the saturation status at the second isoprenoid group (from the head group) is equivalent (Fig. 5). Increased transcript levels for Rv2626 and Rv1738 were detected after the addition of vitamin K1 and were 2-fold higher relative to the addition of vitamin K2 (induction levels of the two genes versus those of the uninduced were 6.7/4.7 for K1 and 4.7/2.7 for K2) (Fig. 1B).

Differential survival effects of supplemental menaquinone analogs. Under aerobic conditions it was previously demonstrated that vitamin K2 is better able than K1 (Fig. 5, structural information) to prevent killing caused by the menaquinone synthesis inhibitor CSU-20 (8). Changes in the saturation state of the menaquinone pools during anaerobic dormancy led us




FIG. 4. Menaquinone content of *M. tuberculosis* H37Rv cells in the RAD dormancy model. Cultures were terminated at the indicated time point (days). Day 0 corresponds to logarithmic aerobic growth. Harvested cells were extracted with chloroform-methanol (2:1), and the nonpolar lipids were isolated using silicic acid column chromatography eluted with chloroform. Samples were normalized to the culture OD, subjected to LC-MS, and quantitated using vitamin K1 as an internal recovery standard to derive the relative detector response. MK-9(II-H₂) is in black, and MK-9 is diagonally striped. By day 4, the levels of MK-9 had fallen to near the limit of detection.

to determine the effects of these two menaquinones in rescuing killing by CSU-20 during anaerobic conditions. All treatments were added at day 12 of the anaerobic dormancy model, after the cultures were anaerobic for approximately 1 week, and were plated 8 days later. When treated with CSU-20, a decrease in survival of three orders of magnitude was observed (Fig. 6). Vitamin K1 had a statistically significant increased ability to rescue killing by CSU-20 compared to vitamin K2 ($P < 0.0001$), converse to what was reported under aerobic conditions (8), further indicating a preferred role for MK-9(II-H₂) during dormancy conditions. In addition, supplementation with either menaquinone alone resulted in decreased survival of the anaerobic cultures. Vitamin K2 addition resulted in a statistically significant increase in killing compared to vitamin K1 ($P < 0.05$). As shown in Fig. 1, the addition of either vitamin K1 or K2 also resulted in increased DosR regulon expression. Therefore, ascorbate was tested as a control to




FIG. 5. Structures and calculated monoisotopic masses of menaquinone analogs.




FIG. 6. Vitamin K rescue of CSU-20-treated cultures. Cultures were treated on day 12 of the anaerobic dormancy model and incubated for 8 days. Values shown are CFU (A and B) and percent survival (C and D) derived from CFU of treated cultures divided by untreated or vehicle-treated cultures. (A and C) Ascorbate is black, vitamin K1 is vertically striped, vitamin K2 is horizontally striped, vehicles are gridded, and untreated is white. (B and D) CSU-20 is thinly diagonally striped, CSU-20 plus vitamin K1 is double diagonally striped, and CSU-20 plus vitamin K2 is thickly diagonally striped.

determine if overexpression of the DosR regulon resulted in killing, which it did not (Fig. 6).

DISCUSSION

The findings in this report demonstrate a role for the ETS in DosS control of the DosR regulon. Data herein indicate that DosS, but not DosT, initiates signaling when treated with ascorbate, a cytochrome *c* reductant, even in the presence of oxygen. Interestingly, the effect of ascorbate is quite specific to the DosR regulon, although a small group of other genes is moderately affected by ascorbate in a DosS-independent manner (Table 1). This provides a link between electron flow and DosS signaling, a link further supported by the observation that inhibition of electron flow decreased signaling, as did the inhibition of menaquinone synthesis.

Regardless of the numerous routes of electron flow through the ETS, the menaquinone pool is always utilized, and thus, monitoring its redox status provides a mechanism for the bacterium to accurately sense the state of respiration. The concentration of menaquinones decreased during hypoxic entry into anaerobiosis. The unsaturated form disappeared, and the concentration of the saturated form initially declined but was maintained at a lower level. Experiments with saturated and unsaturated analogs of these species demonstrated differences in their role in signaling and survival and indicate that specific concentrations are maintained for optimal performance.

This work demonstrates that DosS responds to the redox status of the ETS; however, the direct interaction between the sensor and the ETS is as yet unknown. Recent biochemical work provides some intriguing possibilities. *In vitro* a ferredoxin/ferredoxin reductase pair was demonstrated to activate DosS (21). Ferredoxins are often reduced by electrons from the quinone pool; thus, there is a possible link to the ETS. The genome of *M. tuberculosis* contains several annotated ferredoxins, including *fdxA*, which is part of the DosR regulon and is thus upregulated during conditions in which DosS activates the DosR regulon (7, 53). A second report demonstrated that flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD), cofactors used by many respiratory and ETS enzymes, were able to reduce DosS *in vitro* (6). The *in vivo* relevance of these interactions is as yet unknown, but their involvement with the ETS makes them interesting candidates for interaction with DosS (6). Interestingly, this same report did not observe a direct interaction between reduced menaquinone and DosS, implying that an intermediate is required. It is also worth mentioning that it was previously demonstrated with *Mycobacterium smegmatis* that DevS, which is a homolog to DosT (not to DosS), was insensitive to changes in redox and the menaquinone pool (28).

Several bacterial systems are known to sense components of the ETS in order to respond to a changing respiratory environment. In *Escherichia coli*, Aer responds to redox changes in the ETS to facilitate aerotaxis, although the mechanism of interaction is still unknown (11, 50). Also in *E. coli*, ArcB senses the redox state of the quinone pool (14, 30). In *Helicobacter pylori*, TlpD senses a decrease in electron flow and responds by altering its taxis away from electron-flow limiting conditions (45). *Rhodobacter sphaeroides* uses PrrBA to sense and respond to electron flow through the *cbb*₃ oxidase (33, 34). PhoR in *Bacillus subtilis* responds to the reduced quinones, and several other bacterial sensors containing redox-active cysteine residues have been reported (12). Thus, it is apparent that monitoring the state of the ETS is an important function in many bacteria.

In *M. tuberculosis*, the DosR regulon is important for conditions that inhibit respiration, and several of these conditions result in its induction. Aside from reduction of the ETS, the slowing of aerobic respiration affects other systems which could function as signals for DosR regulon induction. Proton motive force may decrease under conditions that inhibit aerobic respiration; however, the addition of a protonophore does not induce the DosR regulon (shown in Fig. 1B and described previously in reference 3). A decreased level of ATP is likely not the signal, because while hypoxia does lower the overall amount of ATP, other additives that decrease ATP do not induce the regulon (3). Changes in respiration may have an effect on the NADH/NAD ratio; however, no correlation was shown between this ratio and DosR regulon induction via the addition of a wide variety of different additives (3). Cyanide blocks regulon induction by hypoxia and NO (53), likely by directly binding the heme moieties in DosT and DosS, an interaction shown for DosS (22). However, cyanide, a cytochrome *c* oxidase inhibitor, does not induce the regulon itself (3, 53). Azide, another cytochrome oxidase inhibitor, also does not affect regulon induction (22); these results imply that reduced cytochrome oxidase is not directly sensed by DosS.

Interestingly, W/Beijing strains of *M. tuberculosis* constitutively express the DosR regulon, and some lineages have numerous duplications and deletions, including a duplication of *dosR* as well as a portion of the DosR regulon (9, 13, 37). These strains also have an inactivating frameshift mutation in *DosT* (13, 37). It was observed in these reports that neither the *DosT* mutation nor the duplication of the regulon was responsible for the constitutive expression levels. In light of the findings in the present work, it seems feasible that these strains may have a metabolic abnormality which results in a constitutively reduced ETS, which could result in overexpression of the regulon via *DosS*. The plausibility of this is corroborated by the finding of additional gene duplications in these strains, specifically of the *nuo* gene cluster, responsible for encoding the type I oxidoreductase, as well as duplication of the succinate dehydrogenase gene (9). Both of the enzyme clusters encoded by these genes are involved in the ETS. We hypothesize that the mutation in *DosT* may be an evolutionary attempt to attenuate *DosR* regulon expression in certain conditions in order to counteract its overexpression.

Oxygen-limited, nonreplicating bacilli require electron flow via menaquinones, an intact proton motive force, and ATP production for survival (8, 24, 36). The major menaquinone species in mycobacteria is MK-9(II-H₂), a 1,4-naphthoquinone derivative which has nine isoprene units, the second of which is saturated (Fig. 5) (31). Mono-saturated menaquinones of this type have been described for various organisms (35). Several other species of menaquinone exist in mycobacteria, although they are less prevalent, including a fully unsaturated form of the major species, MK-9 (8, 10). Functional differences between these two types of quinone species are presently unclear in the literature. Treatment with CSU-20 was shown to decrease the amount of both species of menaquinone [MK-9(II-H₂) and MK-9] via *menA* inhibition; it was also shown to inhibit oxygen consumption (8). In this report we observed that the level of menaquinone correlated with the amount of *DosR* regulon expression. When oxygen became limited, the levels of menaquinone decreased, while restoring menaquinone to the system resulted in increased signaling. The decrease in menaquinone levels during anaerobic dormancy appears to result in the gradual decrease in *DosR* regulon expression observed in long-term models (54).

Aerobically, it was previously reported that the menaquinone analogs vitamin K1 and vitamin K2 [analogous to MK-9(II-H₂) and MK-9, respectively, because of the saturation status of the second isoprenoid group; Fig. 5] were both able to rescue killing and decreased oxygen consumption caused by CSU-20; further, K2 rescued more effectively than did K1 (8). Interestingly, results reported herein demonstrate that the opposite rescuing effect occurs during anaerobic conditions: K1 rescued more effectively than K2 (Fig. 6). Differential effects of vitamin K1 and K2 as well as other related naphthoquinones have previously been reported for bacterial extracts from *Mycobacterium phlei*, further corroborating biological significance of the saturation status (4).

Overall, the survival data indicate that the bacilli maintain an optimal level of menaquinones during anaerobiosis, given that inhibition of synthesis results in death, but supplementation does as well. Also, it appears that not only is the amount important, but the species is as well, since addition of vitamin

K2 resulted in more death than vitamin K1. The role of MK-9(II-H₂) appears to be specifically important during anaerobiosis. This is based on the observed decrease of MK-9 to the level of detection and the maintenance of MK-9(II-H₂) during the hypoxic adaptation period (Fig. 4). The increased anaerobic rescuing effect by K1 could be partially due to the fact that the *DosR* regulon is more induced by K1 than K2 (Fig. 1), and previously we reported a correlation between regulon induction and anaerobic survival (18). Thus, to summarize the menaquinone results, MK-9(II-H₂) is maintained at a low level. Its analog, vitamin K1, induces the *DosR* regulon more effectively, rescues killing by CSU-20 more effectively, and is less toxic relative to vitamin K2, the analog of MK-9.

A recent report examined the effects of ascorbate-induced hypoxia on the *DosR* regulon (49). As previously reported (18), differential effects and response to oxygen levels were observed for single-sensor mutants of *DosS* and *DosT*. However, the results in our current work clearly separate the effects of oxygen and redox sensing by the two sensors. Ascorbate experiments in this work were performed at a time point when oxygen was not sufficiently limited to induce either sensor (Fig. 1 and 2 and Table 1), unlike the previously mentioned report. This enabled the differences between *DosS* and *DosT* sensing to be observed, differences which were confirmed by a variety of other additives which modulate the ETS.

This report demonstrates that induction of the *DosR* regulon via *DosS* is in response to a reduced ETS. Despite this role of *DosS* in redox monitoring, under aerobic conditions oxygen maintains *DosS* in an inactive, oxidized form. Thus, it appears that both bodies of research on *DosS* signaling are correct: *DosS* functions as both an oxygen sensor (off switch) and redox sensor (on switch). The menaquinone pool is explicitly implicated in this sensing. Menaquinone levels in general decrease, and the saturated form is preferentially maintained at low but functional levels under anaerobic conditions. Relative to the unsaturated form, this saturated form is less toxic, better able to induce signaling via *DosS* during hypoxia, and better able to rescue a drug-mediated decrease in the menaquinone pool. These data indicate that the saturated form of menaquinone is best utilized during oxygen-limited conditions. The menaquinone pool is dynamic, its composition changes with the respiratory environment, and its centrality in the ETS makes it a viable target for monitoring the redox status of the ETS.

ACKNOWLEDGMENTS

This research project was funded by NIH grants RO1 AI061505, entitled the *Mycobacterium tuberculosis* Dormancy Program and awarded to M.I.V., AI049151 awarded to D.C.C., and T32 AI052066-06 awarded to R.W.H.

REFERENCES

1. Anonymous. 1953. Report of panel discussion on survival and revival of tubercle bacilli in healed tuberculous lesions. *Am. Rev. Tuberc.* **68**:477–495.
2. Balási, G., A. P. Heath, L. Shi, and M. L. Gennaro. 2008. The temporal response of the *Mycobacterium tuberculosis* gene regulatory network during growth arrest. *Mol. Syst. Biol.* **4**:225.
3. Boshoff, H. I., T. G. Myers, B. R. Copp, M. R. McNeil, M. A. Wilson, and C. E. Barry III. 2004. The transcriptional responses of *Mycobacterium tuberculosis* to inhibitors of metabolism: novel insights into drug mechanisms of action. *J. Biol. Chem.* **279**:40174–40184.
4. Brodie, A. F., and J. Ballantine. 1960. Oxidative phosphorylation in fractionated bacterial systems. III. Specificity of vitamin K reactivation. *J. Biol. Chem.* **235**:232–237.
5. Canetti, G. 1955. The tubercle bacillus in the pulmonary lesion of man. Springer, New York, NY.

6. Cho, H. Y., H. J. Cho, Y. M. Kim, J. I. Oh, and B. S. Kang. 2009. Structural insight into the heme-based redox sensing by DosS from *Mycobacterium tuberculosis*. *J. Biol. Chem.* **284**:13057–13067.
7. Cole, S. T., R. Brosch, J. Parkhill, T. Garnier, C. Churcher, D. Harris, S. V. Gordon, K. Eiglmeier, S. Gas, C. E. Barry III, F. Tekoa, K. Badcock, D. Basham, D. Brown, T. Chillingworth, R. Connor, R. Davies, K. Devlin, T. Feltwell, S. Gentles, N. Hamlin, S. Holroyd, T. Hornsby, K. Jagels, A. Krogh, J. McLean, S. Moule, L. Murphy, K. Oliver, J. Osborne, M. A. Quail, M. A. Rajandream, J. Rogers, S. Rutter, K. Seeger, J. Skelton, R. Squares, S. Squares, J. E. Sulston, K. Taylor, S. Whitehead, and B. G. Barrell. 1998. TubercuList. *Nature* **393**:537–544.
8. Dhiman, R. K., S. Mahapatra, R. A. Slayden, M. E. Boyne, A. Lenaerts, J. C. Hinshaw, S. K. Angala, D. Chatterjee, K. Biswas, P. Narayanasamy, M. Kurosu, and D. C. Crick. 2009. Menaquinone synthesis is critical for maintaining mycobacterial viability during exponential growth and recovery from non-replicating persistence. *Mol. Microbiol.* **72**:85–97.
9. Domenech, P., G. S. Kolly, L. Leon-Solis, A. Fallow, and M. B. Reed. 2010. Massive gene duplication event among clinical isolates of the *Mycobacterium tuberculosis* W/Beijing family. *J. Bacteriol.* **192**:4562–4570.
10. Dunphy, P. J., P. G. Phillips, and A. F. Brodie. 1971. Separation and identification of menaquinones from microorganisms. *J. Lipid Res.* **12**:442–449.
11. Edwards, J. C., M. S. Johnson, and B. L. Taylor. 2006. Differentiation between electron transport sensing and proton motive force sensing by the Aer and Tsr receptors for aerotaxis. *Mol. Microbiol.* **62**:823–837.
12. Eldakak, A., and F. M. Hulett. 2007. Cys303 in the histidine kinase PhoR is crucial for the phosphotransfer reaction in the PhoPR two-component system in *Bacillus subtilis*. *J. Bacteriol.* **189**:410–421.
13. Fallow, A., P. Domenech, and M. B. Reed. 2010. Strains of the East Asian (W/Beijing) lineage of *Mycobacterium tuberculosis* are DosS/DosT-DosR two-component regulatory system natural mutants. *J. Bacteriol.* **192**:2228–2238.
14. Georgellis, D., O. Kwon, and E. C. Lin. 2001. Quinones as the redox signal for the arc two-component system of bacteria. *Science* **292**:2314–2316.
15. Gomez, J. E., and J. D. McKinney. 2004. *M. tuberculosis* persistence, latency, and drug tolerance. *Tuberculosis (Edinb.)* **84**:29–44.
16. Gonzalo-Asensio, J., S. Mostowy, J. Harders-Westerveen, K. Huygen, R. Hernandez-Pando, J. Thole, M. Behr, B. Gicquel, and C. Martin. 2008. PhoP: a missing piece in the intricate puzzle of *Mycobacterium tuberculosis* virulence. *PLoS One* **3**:e3496.
17. Hernández-Pando, R., M. Jeyanathan, G. Mengistu, D. Aguilar, H. Orozco, M. Harboe, G. A. Rook, and G. Bjune. 2000. Persistence of DNA from *Mycobacterium tuberculosis* in superficially normal lung tissue during latent infection. *Lancet* **356**:2133–2138.
18. Honaker, R. W., R. L. Leistikow, I. L. Bartek, and M. I. Voskuil. 2009. Unique roles of DosT and DosS in DosR regulon induction and *Mycobacterium tuberculosis* dormancy. *Infect. Immun.* **77**:3258–3263.
19. Honaker, R. W., A. Stewart, S. Schittone, A. Izzo, M. R. Klein, and M. I. Voskuil. 2008. *Mycobacterium bovis* BCG vaccine strains lack *narK2* and *narX* induction and exhibit altered phenotypes during dormancy. *Infect. Immun.* **76**:2587–2593.
20. Imboden, P., and G. K. Schoolnik. 1998. Construction and characterization of a partial *Mycobacterium tuberculosis* cDNA library of genes expressed at reduced oxygen tension. *Gene* **213**:107–117.
21. Ioanoviciu, A., Y. T. Meharena, T. L. Poulos, and P. R. Ortiz de Montellano. 2009. DevS oxy complex stability identifies this heme protein as a gas sensor in *Mycobacterium tuberculosis* dormancy. *Biochemistry* **48**:5839–5848.
22. Ioanoviciu, A., E. T. Yukl, P. Moenne-Loccoz, and P. R. de Montellano. 2007. DevS, a heme-containing two-component oxygen sensor of *Mycobacterium tuberculosis*. *Biochemistry* **46**:4250–4260.
23. Kalendar, R. 2006. FastPCR, PCR primer design, DNA and protein tools, repeats and own database searches program. Helsinki, Finland. <http://www.biocenter.helsinki.fi/bi/programs/fastpcr.htm>.
24. Koul, A., L. Vranckx, N. Dendouga, W. Balemans, I. V. Den Wyngaert, K. Vergaunen, H. W. Goehlmann, R. Willebrords, A. Poncellet, J. Guillemont, D. Bald, and K. Andries. 2008. Diarylquinolines are bactericidal for dormant mycobacteria as a result of disturbed ATP homeostasis. *J. Biol. Chem.* **283**:25273–25280.
25. Kumar, A., J. S. Deshane, D. K. Crossman, S. Bolisetty, B. S. Yan, I. Kramnik, A. Agarwal, and A. J. Steyn. 2008. Heme oxygenase-1 derived carbon monoxide induces the *Mycobacterium tuberculosis* dormancy regulon. *J. Biol. Chem.* **283**:18032–18039.
26. Kumar, A., J. C. Toledo, R. P. Patel, J. R. Lancaster, Jr., and A. J. Steyn. 2007. *Mycobacterium tuberculosis* DosS is a redox sensor and DosT is a hypoxia sensor. *Proc. Natl. Acad. Sci. U. S. A.* **104**:11568–11573.
27. Kurosu, M., P. Narayanasamy, K. Biswas, R. Dhiman, and D. C. Crick. 2007. Discovery of 1,4-dihydroxy-2-naphthoate [corrected] prenyltransferase inhibitors: new drug leads for multidrug-resistant gram-positive pathogens. *J. Med. Chem.* **50**:3973–3975.
28. Lee, J. M., H. Y. Cho, H. J. Cho, I. J. Ko, S. W. Park, H. S. Baik, J. H. Oh, C. Y. Eom, Y. M. Kim, B. S. Kang, and J. I. Oh. 2008. O₂ and NO sensing mechanism through the DevSR two-component system in *Mycobacterium smegmatis*. *J. Bacteriol.* **190**:6795–6804.
29. Leistikow, R. L., R. A. Morton, I. L. Bartek, I. Frimpong, K. Wagner, and M. I. Voskuil. 2010. The *Mycobacterium tuberculosis* DosR regulon assists in metabolic homeostasis and enables rapid recovery from non-respiring dormancy. *J. Bacteriol.* **192**:1662–1670.
30. Malpica, R., B. Franco, C. Rodriguez, O. Kwon, and D. Georgellis. 2004. Identification of a quinone-sensitive redox switch in the ArcB sensor kinase. *Proc. Natl. Acad. Sci. U. S. A.* **101**:13318–13323.
31. Minnikin, D. E., S. M. Minnikin, M. Goodfellow, and J. L. Stanford. 1982. The mycolic acids of *Mycobacterium chelonae*. *J. Gen. Microbiol.* **128**:817–822.
32. Neyrolles, O., R. Hernandez-Pando, F. Pietri-Rouxel, P. Fornes, L. Tailleux, J. A. Pagan, E. Pivert, Y. Bordat, D. Aguilar, M. C. Prevost, C. Petit, and B. Gicquel. 2006. Is adipose tissue a place for *Mycobacterium tuberculosis* persistence? *PLoS One* **1**:e43.
33. Oh, J. I., and S. Kaplan. 2000. Redox signaling: globalization of gene expression. *EMBO J.* **19**:4237–4247.
34. Oh, J. I., I. J. Ko, and S. Kaplan. 2004. Reconstitution of the *Rhodospirillum rubrum* cbb3-PrrBA signal transduction pathway in vitro. *Biochemistry* **43**:7915–7923.
35. Phillips, P. G., P. J. Dunphy, K. L. Servis, and A. F. Brodie. 1969. A new menaquinone series differing in the degree of unsaturation of the side chain. *Biochemistry* **8**:2856–2861.
36. Rao, S. P., S. Alonso, L. Rand, T. Dick, and K. Pethe. 2008. The protonmotive force is required for maintaining ATP homeostasis and viability of hypoxic, nonreplicating *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. U. S. A.* **105**:11945–11950.
37. Reed, M. B., S. Gagneux, K. Deriemer, P. M. Small, and C. E. Barry III. 2007. The W-Beijing lineage of *Mycobacterium tuberculosis* overproduces triglycerides and has the DosR dormancy regulon constitutively upregulated. *J. Bacteriol.* **189**:2583–2589.
38. Ribiere, C., A. M. Jaubert, N. Gaudiot, D. Sabourault, M. L. Marcus, J. L. Boucher, D. Denis-Henriot, and Y. Giudicelli. 1996. White adipose tissue nitric oxide synthase: a potential source for NO production. *Biochem. Biophys. Res. Commun.* **222**:706–712.
39. Roberts, D. M., R. P. Liao, G. Wisedchaisri, W. G. Hol, and D. R. Sherman. 2004. Two sensor kinases contribute to the hypoxic response of *Mycobacterium tuberculosis*. *J. Biol. Chem.* **279**:23082–23087.
40. Russel, W. F., S. H. Dressler, G. Middlebrook, and J. Denst. 1955. Implications of the phenomenon of open cavity healing for the chemotherapy of pulmonary tuberculosis. *Am. Rev. Tuberc.* **71**:441–446.
41. Saini, D. K., V. Malhotra, and J. S. Tyagi. 2004. Cross talk between DevS sensor kinase homologue, Rv2027c, and DevR response regulator of *Mycobacterium tuberculosis*. *FEBS Lett.* **565**:75–80.
42. Sambandamurthy, V. K., S. C. Derrick, T. Hsu, B. Chen, M. H. Larsen, K. V. Jalapathy, M. Chen, J. Kim, S. A. Porcelli, J. Chan, S. L. Morris, and W. R. Jacobs, Jr. 2006. *Mycobacterium tuberculosis* DeltaRD1 DeltapanCD: a safe and limited replicating mutant strain that protects immunocompetent and immunocompromised mice against experimental tuberculosis. *Vaccine* **24**:6309–6320.
43. Sardiwal, S., S. L. Kendall, F. Movahedzadeh, S. C. Rison, N. G. Stoker, and S. Djordjevic. 2005. A GAF domain in the hypoxia/NO-inducible *Mycobacterium tuberculosis* DosS protein binds haem. *J. Mol. Biol.* **353**:929–936.
44. Schnappinger, D., S. Ehrh, M. I. Voskuil, Y. Liu, J. A. Mangan, I. M. Monahan, G. Dolganov, B. Efron, P. D. Butcher, C. Nathan, and G. K. Schoolnik. 2003. Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages: insights into the phagosomal environment. *J. Exp. Med.* **198**:693–704.
45. Schweinitzer, T., T. Mizote, N. Ishikawa, A. Dudnik, S. Inatsu, S. Schreiber, S. Suerbaum, S. Aizawa, and C. Josenhans. 2008. Functional characterization and mutagenesis of the proposed behavioral sensor TlpD of *Helicobacter pylori*. *J. Bacteriol.* **190**:3244–3255.
46. Sherman, D. R., M. Voskuil, D. Schnappinger, R. Liao, M. I. Harrell, and G. K. Schoolnik. 2001. Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha-crystallin. *Proc. Natl. Acad. Sci. U. S. A.* **98**:7534–7539.
47. Shiloh, M. U., P. Manzanillo, and J. S. Cox. 2008. *Mycobacterium tuberculosis* senses host-derived carbon monoxide during macrophage infection. *Cell Host Microbe* **3**:323–330.
48. Sousa, E. H., J. R. Tuckerman, G. Gonzalez, and M. A. Gilles-Gonzalez. 2007. DosT and DevS are oxygen-switched kinases in *Mycobacterium tuberculosis*. *Protein Sci.* **16**:1708–1719.
49. Taneja, N. K., S. Dhingra, A. Mittal, M. Naresh, and J. S. Tyagi. 2010. *Mycobacterium tuberculosis* transcriptional adaptation, growth arrest and dormancy phenotype development is triggered by vitamin C. *PLoS One* **5**:e10860.
50. Taylor, B. L. 2007. Aer on the inside looking out: paradigm for a PAS-HAMP role in sensing oxygen, redox and energy. *Mol. Microbiol.* **65**:1415–1424.
51. Trayhurn, P., and I. S. Wood. 2004. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Br. J. Nutr.* **92**:347–355.

52. Tusher, V. G., R. Tibshirani, and G. Chu. 2001. Significance analysis of microarrays applied to the ionizing radiation response. *Proc. Natl. Acad. Sci. U. S. A.* **98**:5116–5121.
53. Voskuil, M. L., D. Schnappinger, K. C. Visconti, M. I. Harrell, G. M. Dolganov, D. R. Sherman, and G. K. Schoolnik. 2003. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *J. Exp. Med.* **198**:705–713.
54. Voskuil, M. L., K. C. Visconti, and G. K. Schoolnik. 2004. *Mycobacterium tuberculosis* gene expression during adaptation to stationary phase and low-oxygen dormancy. *Tuberculosis (Edinb.)* **84**:218–227.
55. Wayne, L. G., and C. D. Sohaskey. 2001. Nonreplicating persistence of *Mycobacterium tuberculosis*. *Annu. Rev. Microbiol.* **55**:139–163.
56. Wolstenholme, G. E. W., M. P. Cameron, and C. M. O'Connor (ed.). 1955. Ciba Foundation symposium on experimental tuberculosis, bacillus and host. Little, Brown, and Company, Boston, MA.
57. Yan, H., E. Aziz, G. Shillabeer, A. Wong, D. Shanghavi, A. Kermouni, M. Abdel-Hafez, and D. C. Lau. 2002. Nitric oxide promotes differentiation of rat white preadipocytes in culture. *J. Lipid Res.* **43**:2123–2129.
58. Yuan, Y., D. D. Crane, R. M. Simpson, Y. Q. Zhu, M. J. Hickey, D. R. Sherman, and C. E. Barry III. 1998. The 16-kDa alpha-crystallin (Acr) protein of *Mycobacterium tuberculosis* is required for growth in macrophages. *Proc. Natl. Acad. Sci. U. S. A.* **95**:9578–9583.
59. Yuki, E. T., A. Ioanoviciu, M. M. Nakano, P. R. Ortiz de Montellano, and P. Moenne-Loccoz. 2008. A distal tyrosine residue is required for ligand discrimination in DevS from *Mycobacterium tuberculosis*. *Biochemistry* **47**:12532–12539.
60. Yun, Z., H. L. Maecker, R. S. Johnson, and A. J. Giaccia. 2002. Inhibition of PPAR gamma 2 gene expression by the HIF-1-regulated gene DEC1/Stra13: a mechanism for regulation of adipogenesis by hypoxia. *Dev. Cell* **2**:331–341.