

Genome Sequence of the Marine Photoheterotrophic Bacterium *Erythrobacter* sp. Strain NAP1

Michal Koblížek,^{1*} Jan Janouškovec,² Miroslav Oborník,^{1,2,3} Justin H. Johnson,⁴
Steven Ferriera,⁴ and Paul G. Falkowski⁵

Institute of Microbiology CAS, Opatovický mlýn, 379 81 Třeboň, Czech Republic¹; University of South Bohemia, Faculty of Science, Branišovská 31, 37005 České Budějovice, Czech Republic²; Institute of Parasitology CAS, Branišovská 31, 37005 České Budějovice, Czech Republic³; J. Craig Venter Institute, 9704 Medical Center Drive, Rockville, Maryland 20850⁴; and Institute of Marine and Coastal Sciences, Rutgers University, 71 Dudley Road, New Brunswick, New Jersey 08901⁵

Received 25 July 2011/Accepted 4 August 2011

Here we report the full genome sequence of marine phototrophic bacterium *Erythrobacter* sp. strain NAP1. The 3.3-Mb genome contains a full set of photosynthetic genes organized in one 38.9-kb cluster; however, it does not contain genes for CO₂ or N₂ fixation, thereby confirming that the organism is a photoheterotroph.

Aerobic anoxygenic phototrophic (AAP) bacteria were discovered in the late 1970s in Tokyo Bay, Japan (2, 9). The AAP bacteria contain bacteriochlorophyll *a* as the main light-harvesting pigment and photochemically functional reaction centers; however, in contrast to purple nonsulfur photosynthetic bacteria, they are obligate aerobes. They perform photoheterotrophic metabolism based on an obligatory supply of organic substrates for growth but derive a significant portion of their energy requirements from light (4, 5, 11). One of the first isolates (OCh 101) was later classified as *Erythrobacter longus* (10), establishing the first genus of marine AAP bacteria. However, the importance of AAP bacteria in the marine environment was fully recognized only later, after Kolber et al. reported their widespread occurrence in the euphotic zone of the oceans (6, 7).

Erythrobacter sp. strain NAP1 (order *Sphingomonadales*, family *Alphaproteobacteria*) was isolated by plating on minimal medium from a water sample collected in the Northwest Atlantic on 16 April 2000 (3, 7). The genome was sequenced using the whole-genome random shotgun method (1) by the J. Craig Venter Institute. The automatic annotation was done with NCBI Prokaryotic Genomes Automatic Annotation Pipeline, which combines hidden Markov model (HMM)-based gene prediction methods with a sequence similarity-based approach.

The complete genome of *Erythrobacter* sp. NAP1 consists of a single circular chromosome of 3,264,238 bp with an average G+C content of 61%. The predicted 3,177 putative genes cover 92% of the genome. The average gene size is 974.67 bp. There is one copy for each of the rRNA genes and 45 genes for tRNAs. The genome contains one continuous 38.9-kb-long photosynthetic gene cluster. The organization of the genes in the cluster is *bchIDO-crtCDF-bchCXYZ-pufBALM-tspO-bchP-ORF-bchG-ppsR-ppaA-bchFNBHLM-lhaA-puhABC-acsF-puhE-hemA-cycA*. While the photosynthetic gene cluster contains a

complete set of genes for bacteriochlorophyll biosynthesis and reaction center proteins, most of the carotenoid biosynthesis genes are located outside the cluster. The gene cluster also contains homologues of regulatory genes *ppaA* (cobalamin binding) and *ppsR* (*crtJ* homologue).

Erythrobacter sp. NAP1 lacks the genes of any autotrophic CO₂ fixation pathway. The catabolic pathway contains genes encoding enzymes for glycolysis and the tricarboxylic acid cycle. The genome does not contain nitrogenase or nitrate reductase, which is consistent with its inability to grow on nitrate. There is a full set of genes for heme biosynthesis; δ -aminolevulinic acid is synthesized through the Shemin (C₄) pathway. There is a complete set of genes for the siroheme biosynthesis pathway, but the final enzymes of cobalamin (vitamin B₁₂) biosynthesis are missing.

The presence of photosynthetic genes but the absence of autotrophic fixation pathways in *Erythrobacter* sp. NAP1 is consistent with its photoheterotrophic metabolism in culture. The provided genome offers a valuable source of information for further studies focused on AAP metabolism and physiological nature. This valuable insight could also provide a comparison with the published genome of the closely related but heterotrophic strain *Erythrobacter litoralis* HTCC2594 (8).

Nucleotide sequence accession number. The genome is available in NCBI GenBank under accession number NZ_AAMW000000000.

The genome sequencing was supported by the Gordon and Betty Moore Foundation as a part of its Marine Microbial Sequencing Project (www.moore.org/marinemicro.aspx). The manual annotation and publication were supported by Czech project GAČR P501/10/0221, Algatech (CZ.1.05/2.1.00/03.0110), and the institutional research concepts MSM6007665808 and AV0Z50200510.

REFERENCES

1. Goldberg, S. M., et al. 2006. A Sanger/pyrosequencing hybrid approach for the generation of high-quality draft assemblies of marine microbial genomes. *Proc. Natl. Acad. Sci. U. S. A.* **103**:11240–11245.
2. Harashima, K., T. Shiba, T. Totsuka, U. Simidu, and N. Taga. 1978. Occurrence of bacteriochlorophyll *a* in a strain of an aerobic heterotrophic bacterium. *Agric. Biol. Chem.* **42**:1627–1628.
3. Koblížek, M., et al. 2003. Isolation and characterization of *Erythrobacter* sp. strains from the upper ocean. *Arch. Microbiol.* **180**:327–338.

* Corresponding author. Mailing address: Institute of Microbiology CAS, Opatovický mlýn, 379 81 Třeboň, Czech Republic. Phone: 420-384340432. Fax: 420-384340415. E-mail: koblizek@alga.cz.

4. **Koblížek, M., J. Mlčoušková, Z. Kolber, and J. Kopecký.** 2010. On the photosynthetic properties of marine bacterium COL2P belonging to Roseobacter clade. *Arch. Microbiol.* **192**:41–49.
5. **Koblížek, M.** 2011. Role of photoheterotrophic bacteria in the marine carbon cycle, p. 49–51. *In* N. Jiao, F. Azam, and S. Sanders (ed.), *Microbial carbon pump in the ocean*. Science/AAAS, Washington, DC.
6. **Kolber, Z. S., C. L. Van Dover, R. A. Niederman, and P. G. Falkowski.** 2000. Bacterial photosynthesis in surface waters of the open ocean. *Nature* **407**: 177–179.
7. **Kolber, Z. S., et al.** 2001. Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* **292**:2492–2495.
8. **Oh, H.-M., S. J. Giovannoni, S. Ferreira, J. Johnson, and J.-C. Cho.** 2009. Complete genome sequence of *Erythrobacter litoralis* HTCC2594. *J. Bacteriol.* **191**:2419–2420.
9. **Shiba, T., U. Simidu, and N. Taga.** 1979. Distribution of aerobic bacteria which contain bacteriochlorophyll a. *Appl. Environ. Microbiol.* **38**:43–45.
10. **Shiba, T., and U. Simidu.** 1982. *Erythrobacter longus* gen. nov., sp. nov., an aerobic bacterium which contains bacteriochlorophyll a. *Int. J. Syst. Bacteriol.* **32**:211–217.
11. **Yurkov, V. V., and J. T. Csotonyi.** 2009. New light on aerobic anoxygenic phototrophs, p. 31–55. *In* C. N. Hunter, F. Daldal, M. C. Thurnauer, and J. T. Beatty (ed.), *The purple phototrophic bacteria*. Springer Verlag, Dordrecht, The Netherlands.