

Complete Genome Sequence of *Helicobacter cinaedi* Strain PAGU611, Isolated in a Case of Human Bacteremia

Takatsugu Goto,^a Yoshitoshi Ogura,^{b,c} Hideki Hirakawa,^d Junko Tomida,^e Yuji Morita,^e Takaaki Akaike,^f Tetsuya Hayashi,^{b,c} and Yoshiaki Kawamura^e

Division of Anaerobe Research, Life Science Research Center, Gifu University, Gifu, Japan^a; Division of Bioenvironmental Science, Frontier Science Research Center, University of Miyazaki, Miyazaki, Japan^b; Division of Microbiology, Department of Infectious Diseases, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan^c; Department of Plant Genome Research, Kazusa DNA Research Institute, Chiba, Japan^d; Department of Microbiology, School of Pharmacy, Aichi Gakuin University, Aichi, Japan^e; and Department of Microbiology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan^f

We report the complete genome sequence of *Helicobacter cinaedi* strain PAGU611, isolated in a case of human bacteremia. The PAGU611 genome comprises a 2,078,348-bp chromosome and a 23,054-bp plasmid. The chromosome contains a unique genomic island, encoding a type VI secretion system and clustered regularly interspaced short palindromic repeat (CRISPR) loci.

Helicobacter cinaedi causes cellulitis, diarrhea, and bacteremia frequently in immunocompromised patients and occasionally in immunocompetent individuals (7, 10, 11, 20). Cases of nosocomial *H. cinaedi* infection (10, 12) and results of epidemiological studies (9) have recently been reported. Despite the clinical importance of *H. cinaedi*, its genome sequence and virulence mechanism have not been completely defined. We report the first complete genome sequence of *H. cinaedi* species.

The genome of PAGU611, an *H. cinaedi* strain isolated in a case of human bacteremia, was sequenced using the Roche 454 GS FLX Titanium system with an 8-kb paired-end library (341,681 reads, 30.8-fold genome coverage). Sequence reads were assembled using Celera Assembler 5.3, and gaps were closed by sequencing PCR products spanning the gaps. Sequence errors were corrected by mapping with Illumina Miseq 150-bp paired-end reads (727,442 reads, 52-fold coverage). The genome sequence was automatically annotated using the Microbial Genome Annotation Pipeline (18) and manually corrected using *in silico* Molecular Cloning Genomics Edition software (In Silico Biology, Inc., Yokohama, Japan) (13).

The PAGU611 genome comprises a 2,078,348-bp chromosome and a 23,054-bp plasmid, pHci1, with average G+C contents of 38.6% and 31.6%, respectively. The chromosome contains 2,096 predicted protein-coding sequences (CDSs), 39 tRNA genes, and 2 rRNA operons. The plasmid encodes 29 predicted CDSs, of which 27 are hypothetical proteins. The genome has no prophage or insertion sequence element.

We performed reciprocal best hit analysis of PAGU611 chromosomal CDSs against 4 sequenced *Helicobacter* species using BLASTP (1) (E value cutoff, 10^{-10}), and significant homology was observed: homologies of 66.1% to *Helicobacter hepaticus* (17), 49.6% to *Helicobacter mustelae* (14), 47.4% to *Helicobacter felis* (2), and 47.3% to *Helicobacter pylori* 26695 (19). Synteny plots of all orthologs indicated that *H. cinaedi* is most similar to *H. hepaticus*.

Synteny plots also identified a unique *H. cinaedi* genomic island—HciGI1. The HciGI1 island (approximately 132 kb, 32.3% G+C content, spanning from HCN_1310 to HCN_1482) contains 173 CDSs, including 147 hypothetical protein genes and 12 genes to assemble a type VI secretion system (T6SS). These T6SS genes are homologous to those of *H. hepaticus* (33.7 to 83.4% amino acid sequence identity), which limit the colonization of *H.*

hepaticus to a germfree murine intestine and prevent intestinal inflammation during colonization (6). The *H. cinaedi* chromosome encodes 2 clustered regularly interspaced short palindromic repeats (CRISPR) and CRISPR-associated (Cas) systems (3, 8), which confer immunity to host cells against foreign DNA. Of these, one is encoded on HciGI1. The *H. mustelae* chromosome encodes a CRISPR/Cas system similar to that in HciGI1. However, the 3 other sequenced *Helicobacter* chromosomes have no CRISPR/Cas system.

Additionally, the PAGU611 chromosome encodes 2 known virulence factors, cytolethal distending toxin (16) and alkyl hydroperoxide reductase (4), and several potential virulence-related proteins, such as neutrophil activation protein (5) and a homolog of *Campylobacter jejuni* invasion antigen B (15).

Nucleotide sequence accession numbers. The complete chromosome and plasmid sequences of *H. cinaedi* PAGU611 have been deposited in DDBJ/EMBL/GenBank under accession no. AP012344 and AP012345, respectively.

ACKNOWLEDGMENTS

This work was supported in part by a Grant-in-Aid for Scientific Research (Challenging Exploratory Research) from the Ministry of Education, Culture, Science, Sports, and Technology (MEXT), Japan, and in part by funds from the Association for Research on Lactic Acid Bacteria in 2012.

REFERENCES

1. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.
2. Arnold IC, et al. 2011. Comparative whole genome sequence analysis of the carcinogenic bacterial model pathogen *Helicobacter felis*. *Genome Biol. Evol.* 3:302–308.
3. Bhaya D, Davison M, Barrangou R. 2011. CRISPR-Cas systems in bacteria and archaea: versatile small RNAs for adaptive defense and regulation. *Annu. Rev. Genet.* 45:273–297.
4. Charoenlap N, et al. 2012. Alkyl hydroperoxide reductase is required for *Helicobacter cinaedi* intestinal colonization and survival under oxidative

Received 17 April 2012 Accepted 24 April 2012

Address correspondence to Yoshiaki Kawamura, kawamura@dpc.agu.ac.jp.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00645-12

- stress in BALB/c and BALB/c interleukin-10^{-/-} mice. *Infect. Immun.* 80:921–928.
5. Choli-Papadopoulou T, Kottakis F, Papadopoulos G, Pendas S. 2011. *Helicobacter pylori* neutrophil activating protein as target for new drugs against *H. pylori* inflammation. *World J. Gastroenterol.* 17:2585–2591.
 6. Chow J, Mazmanian SK. 2010. A pathobiont of the microbiota balances host colonization and intestinal inflammation. *Cell Host Microbe* 7:265–276.
 7. Fox JG. 1997. The expanding genus of *Helicobacter*: pathogenic and zoonotic potential. *Semin. Gastrointest. Dis.* 8:124–141.
 8. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res.* 35:W52–W57. doi:10.1093/nar/gkm360.
 9. Iwashita H, et al. 2008. Identification of the major antigenic protein of *Helicobacter cinaedi* and its immunogenicity in humans with *H. cinaedi* infections. *Clin. Vaccine Immunol.* 15:513–521.
 10. Kitamura T, et al. 2007. *Helicobacter cinaedi* cellulitis and bacteremia in immunocompetent hosts after orthopedic surgery. *J. Clin. Microbiol.* 45:31–38.
 11. Matsumoto T, et al. 2007. Multicenter study to evaluate bloodstream infection by *Helicobacter cinaedi* in Japan. *J. Clin. Microbiol.* 45:2853–2857.
 12. Minauchi K, et al. 2010. The nosocomial transmission of *Helicobacter cinaedi* infections in immunocompromised patients. *Intern. Med.* 49:1733–1739.
 13. Ohyama A, et al. 2006. Bioinformatics tool for genomic era; a step towards the *in silico* experiments—focused on molecular cloning. *J. Comp. Aid. Chem.* 7:102–115.
 14. O'Toole PW, et al. 2010. Comparative genomics and proteomics of *Helicobacter mustelae*, an ulcerogenic and carcinogenic gastric pathogen. *BMC Genomics* 11:164. doi:10.1186/1471-2164-11-164.
 15. Rivera-Amill V, Kim BJ, Seshu J, Konkel ME. 2001. Secretion of the virulence-associated *Campylobacter* invasion antigens from *Campylobacter jejuni* requires a stimulatory signal. *J. Infect. Dis.* 183:1607–1616.
 16. Shen Z, et al. 2009. Cytolethal distending toxin promotes *Helicobacter cinaedi*-associated typhlocolitis in interleukin-10-deficient mice. *Infect. Immun.* 77:2508–2516.
 17. Suerbaum S, et al. 2003. The complete genome sequence of the carcinogenic bacterium *Helicobacter hepaticus*. *Proc. Natl. Acad. Sci. U. S. A.* 100:7901–7906.
 18. Sugawara H, Ohyama A, Mori H, Kurokawa K. 2009. Microbial genome annotation pipeline (MiGAP) for diverse users, abstr S-001, p 1–2. Abstr. 20th Int. Conf. Genome Informatics, Kanagawa, Japan.
 19. Tomb JF, et al. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 388:539–547.
 20. Uçkay I, et al. 2006. Recurrent bacteremia with *Helicobacter cinaedi*: case report and review of the literature. *BMC Infect. Dis.* 6:86.