

# Genome Sequence of Extracellular-Protease-Producing *Alishewanella jeotgali* Isolated from Traditional Korean Fermented Seafood

Jaejoon Jung,<sup>a</sup> Jongsik Chun,<sup>b</sup> and Woojun Park<sup>a</sup>

Department of Environmental Science and Ecological Engineering, Korea University, Seoul, Republic of Korea,<sup>a</sup> and School of Biological Science, Seoul National University, Seoul, Republic of Korea<sup>b</sup>

*Alishewanella jeotgali* MS1<sup>T</sup> (= KCTC 22429<sup>T</sup> = JCM 15561<sup>T</sup>) was isolated from a traditional Korean fermented seafood, gajami sikhae (jeotgal), and has been reported as a novel species. *A. jeotgali* was proven to have extracellular proteolytic activity, which may play an important role in the fermentation environment of food containing fish flesh. Here, we present the genome sequence of *Alishewanella jeotgali* MS1<sup>T</sup> as the first sequenced strain in the genus *Alishewanella* and its taxonomic relatives.

Salty fermented seafood (jeotgal) is an essential part of the Korean diet. Because of the lack of starting inocula, the fermentation process relies on naturally occurring microorganisms that are already present in ingredients. Despite the important role of microorganisms in food fermentation, the diversity, abundance, and roles of microorganisms in such foods have not yet been elucidated. *Alishewanella jeotgali* MS1<sup>T</sup> (= KCTC 22429<sup>T</sup> = JCM 15561<sup>T</sup>) was isolated from a traditional Korean fermented seafood, gajami sikhae (jeotgal), and reported as a novel species (2). Stain MS1<sup>T</sup> can hydrolyze casein, form a biofilm, and produce pellicles (unpublished data).

The *A. jeotgali* MS1<sup>T</sup> genome was sequenced using a combination of an Illumina GA IIX genome analyzer (San Diego, CA) with a 100-bp paired-end library (25,425,184 reads) and the 454 GS FLX Titanium system (Roche Diagnostics, Branford, CT) with an 8-kb paired-end library (195,025 reads). The Illumina and 454 GS FLX reads were assembled using CLC Genomics Workbench 4.7.2 (CLCbio, Denmark) and GS assembler 2.5.1 (Roche Diagnostics, Branford, CT). The assembled sequences obtained from the 2 approaches resulted in 1 scaffold that consisted of 65 large contigs at 678.7-fold coverage. Genome annotation was achieved using Rapid Annotation in the Subsystem Technology (RAST) server (1) and the National Center for Biotechnology Information (NCBI) Prokaryotic Genome Automatic Annotation Pipeline (4, 5).

The *A. jeotgali* MS1<sup>T</sup> genome contains 3,844,563 bp with a G+C content of 50.66%, 3,669 coding sequences (CDS), 64 tRNA genes, and 6 rRNA genes. It does not have a plasmid. The RAST analysis showed that the *Pseudoalteromonas* species genome was closest to the *A. jeotgali* MS1<sup>T</sup> genome, followed by the *Alteromonas* and *Shewanella* genomes. A total of 2,202 CDS were classified using the Cluster of Orthologous Groups (COG) database (7). The most abundant groups of proteins were those that are involved in replication, recombination, and repair (COG category L [5.7%]) and in nucleotide transport and metabolism (COG category F [5.7%]). As the ability of the MS1<sup>T</sup> strain to hydrolyze gelatin and casein implies, the strain contains 2 extracellular proteases, which are homologues of the proteases of *Shewanella denitrificans* OS217 and *Idiomarina loihiensis* L2TR (reference 2 and unpublished data).

Currently, the genus *Alishewanella* contains 4 type species, which have been isolated from different habitats such as human fetus (8), tidal flat sediment (6), landfill soil (3), and fermented food (2). The fact that organisms in this genus have been isolated from various sources implies that *Alishewanella* species may have great adaptability

to diverse environments and unique genetic requirements for each habitat. *A. jeotgali* MS1<sup>T</sup> is the first strain whose genome has been sequenced from among all the known *Alishewanella* strains and taxonomic relatives such as *Rheinheimera* and *Alkalimonas*. To investigate the role of *A. jeotgali* MS1<sup>T</sup> in food fermentation and the ecology- and habitat-specific evolution of the genus *Alishewanella*, additional genome sequencing, comparative genomics, and analysis of food environments are being performed.

**Nucleotide sequence accession numbers.** This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under accession no. [AHTH00000000](http://dx.doi.org/10.1093/nar/38.11.3142). The version described in this paper is the first version, AHTH01000000.

## ACKNOWLEDGMENT

This work was supported by a grant from the Next-Generation BioGreen 21 Program (PJ00820803), Rural Development Administration, Republic of Korea.

## REFERENCES

1. Aziz RK, et al. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
2. Kim MS, et al. 2009. *Alishewanella jeotgali* sp. nov., isolated from traditional fermented food, and emended description of the genus *Alishewanella*. *Int. J. Syst. Evol. Microbiol.* 59:2313–2316.
3. Kim MS, Jo SK, Roh SW, Bae JW. 2010. *Alishewanella agri* sp. nov. isolated from landfill soil. *Int. J. Syst. Evol. Microbiol.* 60:2199–2203.
4. Klimke W, et al. 2009. The National Center for Biotechnology Information's protein clusters database. *Nucleic Acids Res.* 37:D216–D223.
5. Pruitt KD, Tatusova T, Klimke W, Maglott DR. 2009. NCBI reference sequences: current status, policy and new initiatives. *Nucleic Acids Res.* 37:D32–D36.
6. Roh SW, et al. 2009. *Alishewanella aestuarii* sp. nov. isolated from tidal flat sediment, and emended description of the genus *Alishewanella*. *Int. J. Syst. Evol. Microbiol.* 59:421–424.
7. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* 28:33–36.
8. Vogel BF, et al. 2000. Polyphasic taxonomic approach in the description of *Alishewanella fetalis* gen. nov., sp. nov., isolated from a human foetus. *Int. J. Syst. Evol. Microbiol.* 50:1133–1142.

Received 1 February 2012 Accepted 6 February 2012

Address correspondence to Woojun Park, [wpark@korea.ac.kr](mailto:wpark@korea.ac.kr).

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00153-12