

# Kinetic Studies of Iron Oxidation by Whole Cells of *Ferrobacillus ferrooxidans*

C. A. SCHNAITMAN,<sup>1</sup> M. S. KORCZYNSKI,<sup>2</sup> AND D. G. LUNDGREN

*Department of Bacteriology and Botany, Syracuse University, Syracuse, New York 13210*

Received for publication 3 May 1969

A colorimetric assay was developed for studying the kinetics of iron oxidation with whole cells of the chemoautotroph, *Ferrobacillus ferrooxidans*. The assay was more advantageous than the conventional method of Warburg manometry because of its simplicity, rapidity, and the small amount of cells required. The assay measured  $\text{Fe}^{3+}$  as a chloride complex which absorbs at 410 nm. Kinetic analysis showed the apparent  $K_m$  for iron oxidation to be  $5.4 \times 10^{-3}$  M in an unbuffered system and  $2.2 \times 10^{-3}$  M in the presence of  $\beta$ -alanine- $\text{SO}_4^{2-}$  buffer. Glycine and  $\beta$ -alanine buffers were used in the measurement of the pH optimum for iron oxidation; the optimum ranged from 2.5 to 3.8. The effect of pH was primarily on the  $V_{\max}$  while the  $K_m$  remained constant. Added  $\text{SO}_4^{2-}$  was found to stimulate iron oxidation by increasing the  $V_{\max}$  of iron oxidation by whole cells, but it did not affect the  $K_m$ . Results of assays of iron oxidation in systems containing various mole percentages of  $\text{SO}_4^{2-}$  and  $\text{Cl}^-$  indicated that  $\text{Cl}^-$  did not inhibit iron oxidation but that  $\text{SO}_4^{2-}$  was required. Sulfate could be partially replaced by  $\text{HPO}_4^{2-}$  and  $\text{HAsO}_4^{2-}$  but not by  $\text{BO}_3^-$ ,  $\text{MoO}_4^{2-}$ ,  $\text{NO}_3^-$ , or  $\text{Cl}^-$ ; formate and  $\text{MoO}_4^{2-}$  inhibited iron oxidation.

Iron-oxidizing bacteria (*Ferrobacillus-Thiobacillus* group) inhabit a narrow ecological zone where acid mine drainage, containing reduced iron and sulfur compounds, meets the aerobic atmosphere at the earth's surface. These organisms derive their sustenance by accelerating the oxidation of these reduced inorganic compounds and coupling the energy released to the reduction of carbon dioxide and the generation of reducing power.

The properties possessed by these autotrophic bacteria, permitting the oxidation of ferrous ion (or sulfur) in an acid environment (pH 2 to 4), may indicate a structural-functional relationship not found in heterotrophic organisms. Our interests lie in studying such relationships, and this paper is concerned with one phase of the problem, namely, a kinetic analysis of ferrous ion oxidation by intact cells. The latter is important in understanding any structural organization that might exist in the cell's iron oxidation enzyme system. Studies of growth patterns of bacteria as well as isolation and identification of cell substances can furnish only inferential information about the mechanics of iron oxidation. This paper reports on a kinetic analysis of ferrous ion oxidation and

shows that it follows Michaelis-Menten kinetics. Some of the parameters governing the rate of iron oxidation are discussed.

## MATERIALS AND METHODS

**Organism and culture procedure.** *F. ferrooxidans*, obtained from the culture of Leathen et al. (6), was grown on the 9K medium and in the manner previously described (9) in 16-liter glass carboys under forced aeration. Cells were harvested in a Sharples centrifuge after 48 hr of growth, washed three times with distilled water (pH 3.0), and stored in the cold (6 C) in acidified 9K salts or  $\beta$ -alanine- $\text{SO}_4^{2-}$  buffer (pH 3.6, 0.01 M). Stored cells were used within 4 days.

**Development of an assay system.** Initially, a direct colorimetric assay was used which measured the appearance of small amounts of  $\text{Fe}^{3+}$ , in the presence of variable anions, as a red complex formed with sodium thiocyanate (2). The reaction was stopped with 5 ml of 2 N HCl prior to analyzing for  $\text{Fe}^{3+}$ . The addition of HCl to tubes containing  $\text{Fe}^{3+}$  gave rise to a yellow color, presumably due to the formation of a ferric chloride complex, and it was possible to estimate the color spectrophotometrically at 410 nm. This  $\text{Fe}^{3+}$ -chloride complex formation served as a basis for the  $\text{Fe}^{3+}$  assay used in this report.

The assay procedure measuring the appearance of  $\text{Fe}^{3+}$  was as follows. Five acid-washed and thoroughly rinsed Bellco colorimeter tubes, each containing 3.8 ml of distilled water adjusted to pH 3.0 with  $\text{H}_2\text{SO}_4$ , were placed in a water bath at 35 C; 0.2 ml of a cell suspension (2 mg, dry weight of cells) was added to

<sup>1</sup> Present address: Department of Microbiology, University of Virginia Medical School, Charlottesville, Va.

<sup>2</sup> Present address: Department of Microbiology and Immunology, Duke University Medical Center, Durham, N.C. 27706.

each tube, and  $O_2$  was bubbled through the suspension via Pasteur pipettes. To each tube, 1 ml of  $FeSO_4 \cdot 7H_2O$  solution containing 2.5 mg of  $Fe^{2+}$  was added in rapid succession by use of repeating syringe. The oxidation was stopped at 0, 1, 3, 6, and 10 min by rapidly adding 5 ml of 2 N HCl. The HCl was added before the addition of  $Fe^{2+}$  ion for the zero-time tube. Tubes were removed and analyzed for  $Fe^{3+}$  at 410 nm in a Bausch & Lomb Spectronic 20 colorimeter. The intensity of the color was directly related to the amount of  $Fe^{3+}$  formed and agreed well with  $Fe^{3+}$  determinations by the thiocyanate assay method.

**Kinetic analysis of iron oxidation by whole cells.** Five test tubes, each containing 3.8 ml of distilled water adjusted to pH 3.0 with  $H_2SO_4$ , were placed in a water bath at 35 C. This temperature had been established as optimal for the system. To each tube, 0.2 ml of cell suspension was added and  $O_2$  was bubbled through the suspension via Pasteur pipettes. The rate of gas flowing through each tube was regulated by a needle valve to ensure a constant bubbling rate. At 15-sec intervals, 1 ml of an  $FeSO_4$  solution containing 2.5 mg (500  $\mu g/ml$ ) of  $Fe^{2+}$  was added to each tube. The oxidation of iron was stopped in individual tubes by the addition of 5 ml of 2 N HCl at intervals of 2, 3.25, 6.50, and 10.75 min. The development of the yellow ferric chloride complex was estimated as above. For these assays, a reaction tube which received HCl to stop the reaction at 1 min served as the blank; the assay was not considered linear for the first minute. The optical density (OD) at 410 nm was plotted versus reaction time, and the rate of change as  $OD_{410}$  per min was obtained from a line fitted to the OD values of the reaction tubes.

The effect of whole cells and  $Fe^{2+}$  substrate concentrations on the rate of iron oxidation was followed by varying the concentration of cells or of iron in the appropriate reaction system.

The pH of the reaction mixture of the assay system was varied by using different amino acid buffers. Glycine and  $\beta$ -alanine were dissolved in excess dilute  $H_2SO_4$  of known normality and back-titrated to the desired pH with KOH. The final concentrations of components in the assay system were:  $Fe^{2+}$ , 500  $\mu g/ml$ ; glycine or  $\beta$ -alanine, 0.05 M;  $SO_4^{2-}$ , 0.034 M. A variety of other buffers (Table 1) were also tested in order to determine whether they could replace the  $\beta$ -alanine- $SO_4^{2-}$  buffer.

The effect of  $SO_4^{2-}$  and  $Cl^-$  on iron oxidation was investigated. The assay system contained  $Fe^{2+}$  (200  $\mu g/ml$ , as  $FeSO_4 \cdot 7H_2O$ ); the  $SO_4^{2-}$  content was increased from 0.0036 to 0.05 M by the addition of  $K_2SO_4$ . The  $Cl^-$  levels were varied by preparing a series of assay systems in which all cations were the same and the anions consisted of various mole percentages of  $Cl^-$  and  $SO_4^{2-}$ . The test systems contained  $8.94 \times 10^{-3}$  M  $Fe^{2+}$ ,  $10^{-2}$  M  $\beta$ -alanine, and a total anion concentration of  $3.79 \times 10^{-2}$  M; the pH was adjusted to 3.2 with KOH.

All assay results were reproducible and all experiments were done in triplicate and with different batches of cells.

**Anionic replacement of  $SO_4^{2-}$  in the iron assay system.** The assay system to test for replacement of other anions contained:  $1.9 \times 10^{-2}$  M  $Cl^-$ ,  $4.5 \times 10^{-3}$  M  $Fe^{2+}$ , and  $5 \times 10^{-3}$  M  $\beta$ -alanine. The fact that  $Cl^-$  was not inhibitory at these levels and could not satisfy the anionic requirements for iron oxidation led to the use of a system containing  $Cl^-$  as the only anion to determine whether other anions could replace the  $SO_4^{2-}$  requirement for iron oxidation. The various test anions were added as the Na or K salt and were tested at  $2.5 \times 10^{-3}$  M, both in the presence and absence of  $2.5 \times 10^{-3}$  M  $SO_4^{2-}$ .

## RESULTS

The bubble tube-type of iron assay measuring the ferric chloride complex at 410 nm shows a linear increase in  $Fe^{3+}$  after a brief initial lag (Fig. 1); the lag could be due to the mixing requirement. The  $Fe^{3+}$  levels can be compared to those measured by the  $SCN^-$  assay, which also shows a linear response. However, the  $SCN^-$  assay has limitations which restrict its use for kinetic analysis; these limitations are not pertinent to this paper. The rate of iron oxidation ( $\Delta OD_{410}/min$ ) was found to be directly proportional to the concentration of cells used in the assay over the  $\Delta OD$  range of 0.02 to 0.085 (Fig. 2).

The effect of  $Fe^{2+}$  concentration (from 3.5 to 35.8 mM) on the rate of iron oxidation is shown in Fig. 3, and is expressed as a double reciprocal (Lineweaver-Burke) plot. The apparent  $K_m$  of this system was estimated to be  $5.4 \times 10^{-3}$  M.

The standard iron assay procedure and various




FIG. 1. Bubble tube assay for iron oxidation by intact cells of *F. ferrooxidans*. Iron oxidation was followed by measuring  $Fe^{3+}$  with the  $SCN^-$  complex and by measuring the  $Fe^{3+}$ -chloride complex at 410 nm.




FIG. 2. Rate of iron oxidation as a function of relative concentration of cells.




FIG. 3. Lineweaver-Burke plot of substrate concentration versus the rate of iron oxidation.

amino acid buffers were used to obtain more precise information concerning the effect of  $pH$  on iron oxidation by intact cells. Glycine- $SO_4^{2-}$  buffer was used over the  $pH$  range of 1.8 to 3.6, and  $\beta$ -alanine- $SO_4^{2-}$  buffer was used over the  $pH$  range of 3.2 to 4.8. Figure 4 illustrates the difference in the rate of iron oxidation in the two buffer systems. The  $pH$  optimum for  $Fe^{2+}$  oxidation was broad, extending from 2.4 to 3.6, and a rapid decrease in iron oxidation occurred above  $pH$  3.6 and below 2.0.

The effect of  $pH$  on iron oxidation was investigated further by observing its effect on the two kinetic parameters,  $K_m$  and  $V_{max}$  (Fig. 5). When




FIG. 4. Effect of  $pH$  on the rate of iron oxidation. Curve A, glycine- $SO_4^{2-}$  buffer; curve B,  $\beta$ -alanine- $SO_4^{2-}$  buffer.




FIG. 5. Effect of  $pH$  on the  $V_{max}$  and  $K_m$  of iron oxidation.

$\beta$ -alanine-sulfate buffer was used in the assay system over a  $pH$  range of 2.4 to 4.4 with the  $Fe^{2+}$  concentration varied from 3.5 to 17.9 mM for each  $pH$ , the  $K_m$  remained constant throughout most of the  $pH$  range. The change in  $V_{max}$  was more significant in terms of the  $pH$  optimum noted in Fig. 4, since it decreased below  $pH$  2.8 and above  $pH$  3.6. The  $K_m$  determined in the presence of  $\beta$ -alanine- $SO_4^{2-}$  buffer was  $2.2 \times 10^{-3}$  M.

Table 1 shows the effects of using various buffers in the assay system measuring  $Fe^{2+}$  oxidation by whole cells. Both formate and acetate buffer systems were inhibitory. Maleate, citrate, and malate buffers were also unsuitable, owing to inhibi-

TABLE 1. Buffers tested for use in the iron oxidation assay

Buffer system	pH tested	Results
Formate-formic acid <sup>a</sup>	3.2	Inhibition (70%) at $5 \times 10^{-5}$ M
Acetate-acetic acid <sup>a</sup>	3.6	Inhibition (55%) at $5 \times 10^{-2}$ M
Maleate-NaOH <sup>a</sup>	3.6	Slight auto-oxidation; no biological oxidation at $5 \times 10^{-2}$ M
Citrate-NaOH	3.6	Slight biological oxidation; large auto-oxidation at $5 \times 10^{-2}$ M
Malate-NaOH	3.6	Same as maleate
Glycine-SO <sub>4</sub> <sup>2-</sup>	2.1-3.7	No inhibition or stimulation; rate of oxidation the same as without buffer
$\beta$ -Alanine-SO <sub>4</sub> <sup>2-</sup>	3.2	Slight stimulation of oxidation over the control with no buffer

<sup>a</sup> Described by Long (7).

tion of the biological oxidation or to some auto-oxidation of iron. The slight stimulation of iron oxidation observed with  $\beta$ -alanine may be due to additional SO<sub>4</sub><sup>2-</sup> added to the system, but this did not occur in the case of glycine-SO<sub>4</sub><sup>2-</sup>.

The iron assay system was also used to investigate the effects of SO<sub>4</sub><sup>2-</sup> on iron oxidation. Addition of SO<sub>4</sub><sup>2-</sup> to the reaction mixture, ranging from 0.0036 to 0.05 M, approximately doubled the rate of iron oxidation (Fig. 6). Table 2 shows the effect of the addition of SO<sub>4</sub><sup>2-</sup>-containing salts. The four salts tested all caused a significant increase in iron oxidation. The low increase with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> is probably due to a slight inhibition caused by NH<sub>4</sub><sup>+</sup> ions.

Figure 7 shows the effect of SO<sub>4</sub><sup>2-</sup> on the  $K_m$  and  $V_{max}$  of the Fe<sup>2+</sup> oxidation reaction. Curve A is a reciprocal plot of an oxidation rate in an unbuffered system as previously described (Fig. 3), whereas curve B is a reciprocal plot of oxidation rates of the same cell suspension but the reaction mixture contained 0.035 M K<sub>2</sub>SO<sub>4</sub>. The addition of SO<sub>4</sub><sup>2-</sup> increased the velocity of the reaction but did not affect the  $K_m$ .

The effects of Cl<sup>-</sup> on the rate of iron oxidation are shown in Fig. 8, in which the rate of iron oxidation is plotted against the mole per cent Cl<sup>-</sup> in the assay system. A slight stimulation of iron oxidation occurred at low Cl<sup>-</sup> levels, but the rate decreased rapidly at high Cl<sup>-</sup> levels; no oxidation was observed when Cl<sup>-</sup> was the only anion present. Since these levels of Cl<sup>-</sup> do not inhibit iron oxidation, the data indicate a specific SO<sub>4</sub><sup>2-</sup> requirement.

The fact that Cl<sup>-</sup> was not inhibitory and that chloride could not satisfy the anionic requirements for iron oxidation led to the use of a system containing Cl<sup>-</sup> as the only anion, to determine whether other anions could replace SO<sub>4</sub><sup>2-</sup>. When various anions were added to the reaction system in the presence and absence of  $2.5 \times 10^{-3}$  M SO<sub>4</sub><sup>2-</sup>




FIG. 6. Effect of SO<sub>4</sub><sup>2-</sup> concentration on the rate of iron oxidation. The assay system was unbuffered and contained 200 μg of Fe<sup>2+</sup> per ml.

TABLE 2. Effects of various SO<sub>4</sub><sup>2-</sup>-containing salts on the rate of iron oxidation

SO <sub>4</sub> <sup>2-</sup> salt added	SO <sub>4</sub> <sup>2-</sup> in assay	ΔOD <sub>410</sub> /min
	M	
None	0.0036	0.033
K <sub>2</sub> SO <sub>4</sub>	0.041	0.066
MgSO <sub>4</sub>	0.041	0.060
Na <sub>2</sub> SO <sub>4</sub>	0.041	0.059
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	0.041	0.046

to test for both the inhibitory effect of anions and their ability to replace SO<sub>4</sub><sup>2-</sup>, it was noted that both HPO<sub>4</sub><sup>2-</sup> and HAsO<sub>4</sub><sup>2-</sup> could partially replace SO<sub>4</sub><sup>2-</sup> (Table 3). Both of these anions stimulated iron oxidation in the presence of SO<sub>4</sub><sup>2-</sup>, but the




FIG. 7. Double reciprocal plot showing the effect of added  $\text{SO}_4^{2-}$  on the kinetics of iron oxidation. Curve A represents the results obtained with no added  $\text{SO}_4^{2-}$ ; curve B represents the results of identical assays in which  $0.035 \text{ M K}_2\text{SO}_4$  was added to the assay system.




FIG. 8. Effect of varying mole fractions of  $\text{Cl}^-$  and  $\text{SO}_4^{2-}$  on iron oxidation. The total molarity of anion in the system was maintained constant ( $3.79 \times 10^{-2} \text{ M}$ ) but consisted of varying mole percentages of  $\text{SO}_4^{2-}$  and  $\text{Cl}^-$ .

stimulation was no greater than that observed when the equivalent amount of additional  $\text{SO}_4^{2-}$  was added. Borate appeared to be without effect, and it did not replace  $\text{SO}_4^{2-}$ . Nitrate did not replace  $\text{SO}_4^{2-}$  and was inhibitory in the presence of  $\text{SO}_4^{2-}$ ; molybdate did not replace  $\text{SO}_4^{2-}$  and was

TABLE 3. Effect of various anions on iron oxidation in a system containing only  $\text{Cl}^-$  and in the same system containing  $\text{Cl}^-$  plus  $\text{SO}_4^{2-}$

Anion added <sup>a</sup>	$\Delta\text{OD}_{410}/\text{min}$	
	No $\text{SO}_4^{2-}$	$2.5 \times 10^{-3} \text{ M SO}_4^{2-}$
$\text{HAsO}_4^{2-}$	.005	.033
$\text{HPO}_4^{2-}$	.015	.035
$\text{BO}_3^-$	.000	.031
$\text{NO}_3^-$	.000	.016
$\text{MoO}_4^{2-}$	.000	.000
$\text{SO}_4^{2-}$	.030	.035 <sup>b</sup>

<sup>a</sup> The assay system used to test anions other than  $\text{SO}_4^{2-}$  contained the following:  $5 \times 10^{-3} \text{ M}$   $\beta$ -alanine,  $1.9 \times 10^{-2} \text{ M Cl}^-$ , and  $4.5 \times 10^{-3} \text{ M Fe}^{2+}$ . The various test anions were added as the Na or K salt and were tested at  $2.5 \times 10^{-3} \text{ M}$  both in the presence and absence of  $2.5 \times 10^{-3} \text{ M SO}_4^{2-}$ .

<sup>b</sup> This assay contained a total of  $5 \times 10^{-3} \text{ M SO}_4^{2-}$ .

strongly inhibitory. These preliminary results demonstrate that the requirement for  $\text{SO}_4^{2-}$  in iron oxidation is quite specific.

## DISCUSSION

The assay procedure described satisfies the requirements for studying the reaction kinetics of the oxidation of ferrous to ferric iron by intact cells. Iron oxidation follows zero-order kinetics and the rate of the reaction is directly proportional to the concentration of cells added over a satisfactory range. The advantages of this assay are its simplicity and rapidity compared to Warburg respirometry. The assay system in its present form is not suited for studying iron oxidation by cell-free extracts, because of excessive forming; a good assay system for use with extracts remains to be developed.

An accurate estimation of the effects of pH on iron oxidation has been reported; such effects are important in establishing relationships between the organism and its natural acidic environment, as well as between substrate (ferrous ion) and "iron oxidase." The pH optimum for iron oxidation is broad; with intact cells, it is much lower than that reported for one of the enzymes involved in iron oxidation, namely,  $\text{Fe}^{2+}$ -cytochrome *c* reductase (1, 3). The reason for such a pH difference is unknown.

The major effect of pH is on the velocity ( $V_{\text{max}}$ ) and not on the  $K_m$  of the iron oxidation reaction. The effects of pH could be due to the ionization of groups around the active site of those enzymes involved in iron oxidation or due to a loss of iron-binding groups known to be present in the cell envelope (Bacteriol. Proc., p. 123, 1968).

Iron-oxidizing bacteria exhibit an interesting and somewhat unusual anion requirement in that  $\text{SO}_4^{2-}$  is required in substantial (not catalytic or nutrient) quantities for both growth and iron oxidation (5). This requirement may somehow be related to the fact that ferrobacilli can oxidize elemental sulfur to  $\text{SO}_4^{2-}$  (8, 10) and, in the cells' natural environment, high levels of reduced sulfur compounds are found. A role for  $\text{SO}_4^{2-}$  in the organism may be its involvement in the formation of high-energy sulfate compounds (5). Dugan and Lundgren (4) found that ferrous ions as well as sulfate ions are bound by the cell envelope of *F. ferrooxidans* and suggested that the binding of iron requires the participation of  $\text{SO}_4^{2-}$ . However, in light of the present kinetic studies, it doesn't appear likely that  $\text{SO}_4^{2-}$  affects the binding of  $\text{Fe}^{2+}$  at an enzymatic site, for the additions of various concentrations of  $\text{SO}_4^{2-}$  did not affect the  $K_m$  of the reaction. However, the studies do not rule out the possibility that sulfate might simply expose more enzymatic sites which could change the  $V_{\max}$  without affecting the  $K_m$ .

The partial replacement of  $\text{SO}_4^{2-}$  by  $\text{HPO}_4^{2-}$  and to a lesser extent by  $\text{HAsO}_4^{2-}$  indicates something about the nature of the anion requirement. At the pH of the assay, these compounds would probably exist as divalent anions, and the requirement may be for a suitable charge. This would explain the inability of  $\text{Cl}^-$  and  $\text{NO}_3^-$  to replace  $\text{SO}_4^{2-}$ . Also, since both  $\text{HPO}_4^{2-}$  and  $\text{HAsO}_4^{2-}$  form insoluble precipitates with ferric iron, this may in part explain their inability to completely replace the  $\text{SO}_4^{2-}$  requirement. Divalent anions could be required for the depolarization of positively charged sites on the cell surface which prevent the approach of the positively charged  $\text{Fe}^{++}$  substrate. Alternatively, they may also be needed for the maintenance of some essential configuration of the iron-cytochrome *c* reductase enzyme.

The assay system developed for studying the kinetics of iron oxidation has practical significance, for it permits the study of iron oxidation inhibitors. In this study, formate and molybdate were found to be good inhibitors of iron oxidation. These inhibitors may be of importance in controlling iron oxidation which results in a water pollution problem, namely, acid mine wastes.

#### ACKNOWLEDGMENT

The investigation was supported by the Federal Water Pollution Control Administration (WP-0548-03 and 14010 DAY).

#### LITERATURE CITED

1. Blaylock, B., and A. Nason. 1963. Electron transport systems of the chemoautotroph *Ferrobacillus ferrooxidans*. I. Cytochrome *c* containing iron oxidase. *J. Biol. Chem.* 238: 3453-3462.
2. Charlot, G., and P. Bezier. 1959. Quantitative inorganic analysis. John Wiley & Sons, Inc., New York.
3. Din, G. A., I. Suzuki, and H. Lees. 1967. Ferrous iron oxidation by *Ferrobacillus ferrooxidans*. Purification and properties of  $\text{Fe}^{++}$ -cytochrome *c* reductase. *Can. J. Microbiol.* 45:1523-1546.
4. Dugan, P. R., and D. G. Lundgren. 1965. Energy supply for the chemoautotroph *Ferrobacillus ferrooxidans*. *J. Bacteriol.* 89:825-834.
5. Lazaroff, N. 1963. Sulfate requirement for iron oxidation by *Thiobacillus ferrooxidans*. *J. Bacteriol.* 85:78-83.
6. Leathen, W. W., N. A. Kinsel, and S. A. Braley, Sr. 1956. *Ferrobacillus ferrooxidans*: a chemosynthetic autotrophic bacterium. *J. Bacteriol.* 72:700-704.
7. Long, C. 1961. *Biochemists handbook*, p. 30-41. D. VanNostrand Co., Princeton, N.J.
8. Margalith, P., M. Silver, and D. G. Lundgren. 1966. Sulfur oxidation by the iron bacterium *Ferrobacillus ferrooxidans*. *J. Bacteriol.* 92:1706-1709.
9. Silverman, M. P., and D. G. Lundgren. 1959. Studies on the chemosynthetic iron bacterium *Ferrobacillus ferrooxidans*. I. An improved medium and a harvesting procedure for securing high cell yields. *J. Bacteriol.* 77:642-647.
10. Unz, R., and D. G. Lundgren. 1961. A comparative nutritional study of three chemoautotrophic bacteria: *Ferrobacillus ferrooxidans*, *Thiobacillus ferrooxidans*, and *Thiobacillus thiooxidans*. *Soil Sci.* 92:302-313.